

Randomly Draw Subsets of Rows from Matrix in R

Fan Wang

2020-04-19

Contents

Drawly Random Rows	1
------------------------------	---

Drawly Random Rows

Go to the [RMD](#), [R](#), [PDF](#), or [HTML](#) version of this file. Go back to [fan's REconTools Package](#), [R Code Examples Repository \(bookdown site\)](#), or [Intro Stats with R Repository \(bookdown site\)](#).

Draw Random Subset of Sample

- r random discrete

We have a sample of N individuals in some dataframe. Draw without replacement a subset $M < N$ of rows.

```
# parameters, it_M < it_N
it_N <- 10
it_M <- 5

# Draw it_m from indexed list of it_N
set.seed(123)
ar_it_rand_idx <- sample(it_N, it_M, replace=FALSE)

# dataframe
df_full <- as_tibble(matrix(rnorm(4,mean=0,sd=1), nrow=it_N, ncol=4)) %>% rowid_to_column(var = "ID")

# random Subset
df_rand_sub_a <- df_full[ar_it_rand_idx,]

# Random subset also
df_rand_sub_b <- df_full[sample(dim(df_full)[1], it_M, replace=FALSE),]

# Print
# Display
kable(df_full) %>% kable_styling_fc()

kable(df_rand_sub_a) %>% kable_styling_fc()

kable(df_rand_sub_b) %>% kable_styling_fc()
```

Random Subset of Panel There are N individuals, each could be observed M times, but then select a subset of rows only, so each person is randomly observed only a subset of times. Specifically, there are 3 unique students with student ids, and the second variable shows the random dates in which the student showed up in class, out of the 10 classes available.

ID	V1	V2	V3	V4
1	0.1292877	0.4609162	0.1292877	0.4609162
2	1.7150650	-1.2650612	1.7150650	-1.2650612
3	0.4609162	0.1292877	0.4609162	0.1292877
4	-1.2650612	1.7150650	-1.2650612	1.7150650
5	0.1292877	0.4609162	0.1292877	0.4609162
6	1.7150650	-1.2650612	1.7150650	-1.2650612
7	0.4609162	0.1292877	0.4609162	0.1292877
8	-1.2650612	1.7150650	-1.2650612	1.7150650
9	0.1292877	0.4609162	0.1292877	0.4609162
10	1.7150650	-1.2650612	1.7150650	-1.2650612

ID	V1	V2	V3	V4
3	0.4609162	0.1292877	0.4609162	0.1292877
10	1.7150650	-1.2650612	1.7150650	-1.2650612
2	1.7150650	-1.2650612	1.7150650	-1.2650612
8	-1.2650612	1.7150650	-1.2650612	1.7150650
6	1.7150650	-1.2650612	1.7150650	-1.2650612

ID	V1	V2	V3	V4
5	0.1292877	0.4609162	0.1292877	0.4609162
3	0.4609162	0.1292877	0.4609162	0.1292877
9	0.1292877	0.4609162	0.1292877	0.4609162
1	0.1292877	0.4609162	0.1292877	0.4609162
4	-1.2650612	1.7150650	-1.2650612	1.7150650

```

# Define
it_N <- 3
it_M <- 10
svr_id <- 'student_id'

# dataframe
set.seed(123)
df_panel_rand <- as_tibble(matrix(it_M, nrow=it_N, ncol=1)) %>%
  rowid_to_column(var = svr_id) %>%
  uncount(V1) %>%
  group_by(!!sym(svr_id)) %>% mutate(date = row_number()) %>%
  ungroup() %>% mutate(in_class = case_when(rnorm(n(),mean=0,sd=1) < 0 ~ 1, TRUE ~ 0)) %>%
  filter(in_class == 1) %>% select(!!sym(svr_id), date) %>%
  rename(date_in_class = date)

# Print
kable(df_panel_rand) %>% kable_styling_fc()

```

student_id	date_in_class
1	1
1	2
1	8
1	9
1	10
2	5
2	8
2	10
3	1
3	2
3	3
3	4
3	5
3	6
3	9