



Performance Modeling of Computer Systems and Networks

Prof. Vittoria de Nitto Personè

Generating Continuous Random Variates

Università degli studi di Roma Tor Vergata
Department of Civil Engineering and Computer Science Engineering

Copyright © Vittoria de Nitto Personè, 2021

<https://creativecommons.org/licenses/by-nc-nd/4.0/>



(CC BY-NC-ND 4.0)

1

Prerequisite

We assume the knowledge of continuous random variables (sect.7.1).

In particular:

- $Uniform(a,b)$
- $Exponential(\mu)$
- $Normal(\mu,\sigma)$
- $Lognormal(n,b)$
- $Erlang(n,b)$
- $Student(n)$


Prof. Vittoria de Nitto Personè

2


2

Discrete Random Variates

$$F^*(u) = \min_x \{x : u < F(x)\}$$



Continuous Random Variates



Prof. Vittoria de Nitto Personè

3

3

Discrete Simulation
Generating Continuous Random Variates

Preliminary Definitions

The *inverse distribution function* (idf) of X is the function

$$F^{-1} : (0, 1) \rightarrow \mathcal{X}, \forall u \in (0, 1) \text{ as}$$

$$F^{-1}(u) = x$$

where $x \in \mathcal{X}$ is the unique possible value for $F(x) = u$

There is a one-to-one correspondence between possible values $x \in \mathcal{X}$ and cdf values $u = F(x) \in (0, 1)$


Prof. Vittoria de Nitto Personè

4

4

Continuous Random Variable idfs

- Unlike the a discrete random variable, the idf for a continuous random variable is a true inverse



- Can sometimes determine the idf in "closed form" by solving $F(x) = u$ for x

Prof. Vittoria de Nitto Personè

5

5

Examples

- if X is *Uniform*(a, b), $F(x) = (x-a)/(b-a)$ for $a < x < b$

$$x = F^{-1}(u) = a + (b-a)u \quad 0 < u < 1$$

- if X is *Exponential*(μ), $F(x) = 1 - \exp(-x/\mu)$ for $x > 0$

$$x = F^{-1}(u) = -\mu \ln(1-u) \quad 0 < u < 1$$

- if X is a continuous variable with possible value $0 < x < b$ and pdf $f(x) = 2x/b^2$, cdf $F(x) = (x/b)^2$

$$x = F^{-1}(u) = b\sqrt{u} \quad 0 < u < 1$$

Prof. Vittoria de Nitto Personè

6

6

Random Variate Generation By Inversion

- X is a continuous random variable with idf $F^{-1}(\cdot)$
- Continuous random variable U is *Uniform*(0,1)
- Z is the continuous random variable defined by $Z = F^{-1}(U)$

Theorem

Z and X are identically distributed

Algorithm 1

```
u = Random();
return F-1(u);
```

Prof. Vittoria de Nitto Personè

7

7

Inversion examples

- *Uniform*(a,b) Random Variate

```
u = Random();
return a + (b - a) * u;
```

- *Exponential*(μ) Random Variate

```
u = Random();
return -  $\mu \log(1-u)$ ;
```

Prof. Vittoria de Nitto Personè

8

8

Inversion algorithms

- Algorithms in the previous two examples are:
 - portable, exact, robust, efficient, clear, **synchronized and monotone**
- It is not always possible to solve for a continuous random variable idf explicitly by algebraic techniques
- Two other options may be available:
 - Use a function that accurately *approximates* $F^{-1}(\cdot)$
 - Determine the idf by solving $u = F(x)$ *numerically*
(see section 7.2.2)

Testing for correctness

- generate a sample of n random variates where n is large
- evaluate sample mean and standard deviation
- compare them with the theoretical values,
they should be *reasonably* close !!

This is not enough!! Different distributions can have
the same mean and standard deviation !!!

- generate a sample of n random variates and construct a k -bin continuous-data histogram with bin width δ
- f'' is the histogram density and $f(x)$ is the pdf

$$f'' \rightarrow f(x) \text{ as } n \rightarrow \infty \text{ and } \delta \rightarrow 0$$

- In practice, using a large but finite value of n and a small but non-zero value of δ , perfect agreement between f'' and f will not be achieved

Discrete case: natural sampling variability !
Continuous case: variability+binning !!

Truncation

- Let X be a continuous random variable with possible values \mathcal{X} and cdf $F(x) = \Pr(X \leq x)$
- Suppose we wish to restrict the possible values of X to $(a, b) \subset \mathcal{X}$

It is similar to, but simpler than truncation in the discrete-variable context

- X is $\leq a$ with probability $\Pr(X \leq a) = F(a)$
- X is $\geq b$ with probability $\Pr(X \geq b) = 1 - \Pr(X < b) = 1 - F(b)$
- X is between a and b with probability

$$\Pr(a < X < b) = \Pr(X < b) - \Pr(X \leq a) = F(b) - F(a)$$

2 cases for truncation

Case 1

if a and b are specified, the cdf of X can be used to determine the left-tail α , right-tail β truncation probabilities

$$\alpha = \Pr(X \leq a) = F(a) \quad \text{and} \quad \beta = \Pr(X > b) = 1 - F(b)$$

Case 2

if α and β are specified, the idf of X can be used to determine left and right truncation points

$$a = F^{-1}(\alpha) \quad \text{and} \quad b = F^{-1}(1 - \beta)$$

$$F(b) = 1 - \beta$$

Both transformations are exact !

Library `rvgs`

- Contains 7 continuous random variate generators
 - `double Chisquare(long n)`
 - `double Erlang(long n, double b)`
 - `double Exponential(double μ)`
 - `double Lognormal(double a, double b)`
 - `double Normal(double μ , double σ)`
 - `double Student(long n)`
 - `double Uniform(double a, double b)`