



Performance Modeling of Computer Systems and Networks

Prof. Vittoria de Nitto Personè

Interval Estimation

Università degli studi di Roma Tor Vergata
Department of Civil Engineering and Computer Science Engineering

Copyright © Vittoria de Nitto Personè, 2021
<https://creativecommons.org/licenses/by-nc-nd/4.0/>



1

model development

Algorithm 1.1: how to develop a model

1. Goals and objectives
2. *Conceptual* model (cm)
3. Convert cm into a *specification* model (sm)
1. Convert sm into a *computational* model (cptm)
2. Verify
3. Validate

Prof. Vittoria de Nitto Personè

2

2



Algorithm 1.2: using the resulting model

7. Design simulations experiments
 - What parameters should be varied?
 - perhaps many combinatoric possibilities
8. Make production runs
 - Record initial conditions, input parameters
 - Record statistical output
9. Analyze the output
 - Random components → statistical analysis (means, standard deviations, percentiles, histograms etc.)
10. Make decisions
 - The step9 results drive the decisions → actions
 - Simulation should be able to correctly predict the outcome of these actions (→ further refinements)
11. Document the results
 - summarize the gained insights in specific observations and conjectures useful for subsequent similar system models

Prof. Vittoria de Nitto Personè

3

3



- The accumulated average wait was printed every 20 job

Prof. Vittoria de Nitto Personè

4

4

Consider a sample x_1, x_2, \dots, x_n (continuous or discrete) with

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \quad s^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$$

Consider a piecewise constant sample path

$$x(t) = \begin{cases} x_1 & t_0 < t \leq t_1 \\ x_2 & t_1 < t \leq t_2 \\ \vdots & \vdots \\ x_n & t_{n-1} < t \leq t_n \end{cases}$$

$$\bar{x} = \frac{1}{\tau} \int_0^\tau x(t) dt = \frac{1}{t_n} \sum_{i=1}^n x_i \delta_i \quad s^2 = \frac{1}{\tau} \int_0^\tau (x(t) - \bar{x})^2 dt = \frac{1}{t_n} \sum_{i=1}^n (x_i - \bar{x})^2 \delta_i$$

Prof. Vittoria de Nitto Personè

5

5

Discrete Simulation
Interval Estimation

Central limit theorem

If X_1, X_2, \dots, X_n is an iid sequence of random variables (RVs) with

- common mean μ
- common standard deviation σ

and if \bar{X} is the (sample) mean of these RVs $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
then \bar{X} approaches a $Normal(\mu, \sigma / \sqrt{n})$
as $n \rightarrow \infty$

Prof. Vittoria de Nitto Personè

6


6

Sample Mean Distribution

- Choose one of the random variate generators in `rvgs` to generate a sequence of random variable samples with fixed sample size $n > 1$
- with the n -point samples indexed $j=1, 2, \dots$, the corresponding sample mean \bar{x} and sample standard deviation s can be calculated using Welford's algorithm

$$\underbrace{x_1, x_2, \dots, x_n}_{\bar{x}_1, s_1} \quad \underbrace{x_{n+1}, x_{n+2}, \dots, x_{2n}}_{\bar{x}_2, s_2} \quad \underbrace{x_{2n+1}, x_{2n+2}, \dots, x_{3n}}_{\bar{x}_3, s_3} \quad x_{3n+1}$$

- A continuous-data histogram can be created using program `cdh`



Prof. Vittoria de Nitto Personè

7

7

Properties of Sample Mean Histogram


If we denote with μ and σ the theoretical mean and standard deviation respectively of the random variates

- independent of n
 - the histogram mean is approximately μ
 - the histogram standard deviation is approximately σ / \sqrt{n}
- if n is sufficiently large,
 - the histogram density approximates the $Normal(\mu, \sigma / \sqrt{n})$ pdf

Prof. Vittoria de Nitto Personè

8

8

10000 n -point $Exponential(\mu)$ samples

9

9

Example

- The histogram density corresponding to the 36-point sample means is **closely** matched by the pdf of a $Normal(\mu, \sigma / \sqrt{n})$ RV
 for $Exponential(\mu)$ samples, $n=36$ is large enough for the sample mean to be approximately $Normal(\mu, \sigma / \sqrt{n})$
- The histogram density corresponding to the 9-point sample means matches **relatively well**, but with a skew to the left
 • $n=9$ is not large enough

Prof. Vittoria de Nitto Personè

10

10

Example (cont.)

- Essentially all of the sample means are within an interval of width of $4\sigma/\sqrt{n}$ centered about μ
- because $n \rightarrow \infty$ as $\sigma/\sqrt{n} \rightarrow 0$, if n is large, all the sample means will be close to μ
- In general:
 - the accuracy of the $Normal(\mu, \sigma/\sqrt{n})$ pdf approximation is dependent on the shape of a fixed population pdf
 - If the samples are drawn from a population with
 - a highly asymmetric pdf (like the $Exponential(\mu)$ pdf): n may need to be as large as 30 or more for good fit
 - a pdf symmetric about the mean (like the $Uniform(a,b)$ pdf): n as small as 10 or less may produce a good fit

Prof. Vittoria de Nitto Personè

11

11

Examples of Linear Data Transformations

- suppose x_1, x_2, \dots, x_n measured in seconds
 - to convert to minutes, let $x'_i = x_i/60$
($a=1/60, b=0$)

$$\bar{x}' = \frac{45}{60} = 0.75 \qquad s' = \frac{15}{60} = 0.25 \quad (\text{minutes})$$

- standardize data
($a=1/s, b=-\bar{x}/s$)

$$x'_i = \frac{1}{s}x_i - \frac{\bar{x}}{s} \qquad x'_i = \frac{x_i - \bar{x}}{s}$$

Then

$$\bar{x}' = 0 \qquad s' = 1$$

Used to avoid problems with very large (or small) valued samples

Prof. Vittoria de Nitto Personè

12


12

Standardized Sample Mean Distribution

We can standardize the sample means $\bar{x}_1, \bar{x}_2, \bar{x}_3, \dots$ by subtracting μ and dividing the result by σ/\sqrt{n} to form the standardized sample means z_1, z_2, z_3, \dots defined by

$$z_j = \frac{\bar{x}_j - \mu}{\sigma/\sqrt{n}} \quad j = 1, 2, 3, \dots$$

- Generate a continuous-data histogram for the standardized sample means by program cdh



Prof. Vittoria de Nitto Personè

13

13

Properties of Standardized Sample Mean Histogram

- independent of n
 - the histogram mean is approximately 0
 - the histogram standard deviation is approximately 1
- if n is sufficiently large,
 - the histogram density approximates the $Normal(0,1)$ pdf

Prof. Vittoria de Nitto Personè

14

14


t -Statistic Distribution

Definition

- each sample mean \bar{x}_j is a point estimate of μ
- each sample variance s_j^2 is a point estimate of σ^2
- each sample standard deviation s_j is a point estimate of σ

Want to replace *population* standard deviation σ with *sample* standard deviation s_j in standardization equation

$$z_j = \frac{\bar{x}_j - \mu}{\sigma / \sqrt{n}} \quad j = 1, 2, 3, \dots$$



$$\frac{s_j}{\sqrt{n-1}}$$

Prof. Vittoria de Nitto Personè


15

15

- Calculate the t -statistic

$$t_j = \frac{\bar{x}_j - \mu}{s_j / \sqrt{n-1}} \quad j = 1, 2, 3, \dots$$

- Generate a continuous-data histogram using `cdh`



Prof. Vittoria de Nitto Personè

16

16

Properties of t -statistic histogram


- if $n > 2$, the histogram mean is approximately 0
- if $n > 3$, the histogram standard deviation is approximately $\sqrt{(n-1)/(n-3)}$
- if n is sufficiently large, the histogram density approximates the pdf of a $Student(n-1)$ random variable

Prof. Vittoria de Nitto Personè

17

17

Example (cont.)



Prof. Vittoria de Nitto Personè

18

18

Example (cont.)

- The histogram mean and standard deviation are approximately 0.0 and $\sqrt{(n-1)/(n-3)} \approx 1.0$ respectively
- The histogram density corresponding to the 36-point sample means matches the pdf of a *Student*(35) RV relatively well
- The histogram density corresponding to the 9-point sample means matches the pdf of a *Student*(8) RV, but not as well

Prof. Vittoria de Nitto Personè

19

19

Interval Estimation

Theorem 2

If x_1, x_2, \dots, x_n is an independent random sample from a "source" of data with unknown mean μ , if \bar{x} and s are the mean and standard deviation of this sample, and if n is large, it is approximately true that

$$t = \frac{\bar{x} - \mu}{s/\sqrt{n-1}}$$

is a *Student*($n-1$) random variate

- provides the justification for estimating an interval that is likely to contain the mean μ
- as $n \rightarrow \infty$, the *Student*($n-1$) distribution becomes indistinguishable from *Normal*(0,1)

Prof. Vittoria de Nitto Personè

20


20

Suppose

- T is a $Student(n-1)$ random variable
- α is a "confidence parameter" with $0.0 < \alpha < 1.0$

Then there exists a corresponding positive real number t^*

$$\Pr(-t^* \leq T \leq t^*) = 1 - \alpha$$



Prof. Vittoria de Nitto Personè

21

21

Interval Estimation

- suppose μ is unknown. Since $t \approx Student(n-1)$

$$-t^* \leq \frac{\bar{x} - \mu}{s/\sqrt{n-1}} \leq t^*$$

will be approximately true with probability $1 - \alpha$

- right inequality: $\frac{\bar{x} - \mu}{s/\sqrt{n-1}} \leq t^* \Leftrightarrow \bar{x} - \mu \leq \frac{t^* s}{\sqrt{n-1}} \Leftrightarrow \bar{x} - \frac{t^* s}{\sqrt{n-1}} \leq \mu$
- left inequality: $-t^* \leq \frac{\bar{x} - \mu}{s/\sqrt{n-1}} \Leftrightarrow -\frac{t^* s}{\sqrt{n-1}} \leq \bar{x} - \mu \Leftrightarrow \mu \leq \bar{x} + \frac{t^* s}{\sqrt{n-1}}$

So, with probability $1 - \alpha$ (approximately),

$$\bar{x} - \frac{t^* s}{\sqrt{n-1}} \leq \mu \leq \bar{x} + \frac{t^* s}{\sqrt{n-1}}$$

Prof. Vittoria de Nitto Personè

22

22

Theorem 3

If

x_1, x_2, \dots, x_n is an independent random sample from a “source” of data with unknown mean μ

- if \bar{x} and s are the sample mean and sample standard deviation
- n is large

Then, given a confidence parameter α with $0.0 < \alpha < 1.0$, there exists an associated positive real number t^* such that

$$Pr\left(\bar{x} - \frac{t^* s}{\sqrt{n-1}} \leq \mu \leq \bar{x} + \frac{t^* s}{\sqrt{n-1}}\right) \cong 1 - \alpha$$

Prof. Vittoria de Nitto Personè


23

23

Example

- If $\alpha = 0.05$, we are 95% confident that μ lies somewhere between

$$\bar{x} - \frac{t^* s}{\sqrt{n-1}} \quad \text{and} \quad \bar{x} + \frac{t^* s}{\sqrt{n-1}}$$



- for a fixed sample size n and level of confidence $1 - \alpha$, use `rvms` to determine $t^* = \text{idfStudent}(n-1, 1 - \alpha/2)$
- ex. $n = 30, \alpha = 0.05 \rightarrow t^* = \text{idfStudent}(29, 0.975) \cong 2.045$

Prof. Vittoria de Nitto Personè

24

24

Definition

- The interval defined by the two endpoints $\bar{x} \pm \frac{t^* s}{\sqrt{n-1}}$ is a $(1-\alpha)\times 100\%$ confidence *interval estimate* for μ
- $(1-\alpha)$ is the *level of confidence* associated with this interval estimate and t^* is the *critical value* of t

Prof. Vittoria de Nitto Personè

25

25

Algorithm

To calculate an interval estimate for the unknown mean μ of the population from which a random sample x_1, x_2, \dots, x_n was drawn:

- pick a level of confidence $1-\alpha$ (typically $\alpha=0.05$)
- calculate the sample mean \bar{x} and standard deviation s (use Welford's algorithm)
- calculate the critical value $t^* = \text{idfStudent}(n-1, 1-\alpha/2)$
- calculate the interval endpoints $\bar{x} \pm \frac{t^* s}{\sqrt{n-1}}$

If n is sufficiently large, then you are $(1-\alpha)\times 100\%$ confident that the mean μ lies within the interval. The midpoint of the interval is \bar{x}


Prof. Vittoria de Nitto Personè

26

26

Tradeoff - Confidence Versus Sample Size

- For a fixed sample size
 - More confidence can be achieved only at the expense of a larger interval
 - A smaller interval can be achieved only at the expense of less confidence



Prof. Vittoria de Nitto Personè

27

27

Example

- The random sample of size $n = 10$:

1.051	6.438	2.646	0.805	1.505
0.546	2.281	2.822	0.414	1.307

is drawn from a population with unknown mean μ

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \quad s^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2 \quad s = \sqrt{s^2}$$

$$\bar{x} = 1.982$$

Prof. Vittoria de Nitto Personè

28

28

Example

- The random sample of size $n = 10$:


1.051	6.438	2.646	0.805	1.505
0.546	2.281	2.822	0.414	1.307

is drawn from a population with unknown mean μ

- $\bar{x} = 1.982$ and $s = 1.690$
- to calculate a 90% confidence interval estimate:
 - determine: $t^* = \text{idfStudent}(9, 0.95) \approx 1.833$
 - interval: $1.982 \pm (1.833)(1.690/\sqrt{9}) = 1.982 \pm 1.032$

$$\bar{x} \pm \frac{t^* s}{\sqrt{n-1}}$$

- we are approximately 90% confident that μ is between 0.950 and 3.014



Prof. Vittoria de Nitto Personè


29

29

Example (cont.)


- To calculate a 95% confidence interval estimate:
 - determine: $t^* = \text{idfStudent}(9, 0.975) \approx 2.262$
 - interval: $1.982 \pm (2.262)(1.690/\sqrt{9}) = 1.982 \pm 1.274$

- We are approximately 95% confident that μ is between 0.708 and 3.256



- To calculate a 99% confidence interval estimate:
 - determine: $t^* = \text{idfStudent}(9, 0.995) \approx 3.250$
 - interval: $1.982 \pm (3.250)(1.690/\sqrt{9}) = 1.982 \pm 1.832$

- We are approximately 99% confident that μ is between 0.150 and 3.814



- Note: $n=10$ is not large

Prof. Vittoria de Nitto Personè

30

30

1. starting from a sample x_1, x_2, \dots, x_n

- Program `estimate` automates the interval estimation process
- A typical application: estimate the value of an unknown population mean μ by using n replications to generate an independent random variate sample x_1, x_2, \dots, x_n
- Function `Generate()` represents a discrete-event or Monte Carlo simulation program that returns a random variate output x

Using the Generate Method

```
for (i = 1; i <= n; i++)
    xi = Generate();
return x1, x2, ..., xn;
```

- Given a level of confidence $1 - \alpha$, program `estimate` can be used with x_1, x_2, \dots, x_n to compute an interval estimate for μ

Prof. Vittoria de Nitto Personè

31

31

`estimate.c`

```
#include <math.h>
#include <stdio.h>
#include "rvms.h"
#define LOC 0.95

/* level of confidence, */
/* use 0.95 for

95% confidence */
int main(void)
{ long n = 0; /* counts data points */
  double sum = 0.0;
  double mean = 0.0;
  double data;
  double stdev;
  double u, t, w;
  double diff;
  while (!feof(stdin)) { /* use Welford's one-pass method */
    scanf("%lf\n", &data); /* to calculate the sample mean */
    n++; /* and standard deviation */
    diff = data - mean;
    sum += diff * diff * (n - 1.0) / n;
    mean += diff / n;
  }
  stdev = sqrt(sum / n)
```

Prof. Vittoria de Nitto Personè

32

32

$$t^* = \text{idfStudent}(n-1, 1 - \alpha/2)$$

```

if (n > 1) {
    u = 1.0 - 0.5 * (1.0 - LOC);      /* interval parameter */
    t = idfStudent(n - 1, u);         /* critical value of t */
    w = t * stdev / sqrt(n - 1);      /* interval half width */
    printf("\nbased upon %ld data points", n);
    printf(" and with %d%% confidence\n", (int) (100.0 * LOC + 0.5));
    printf("the expected value is in the interval");
    printf("%10.2f +/- %6.2f\n", mean, w); }
else
    printf("ERROR - insufficient data\n");
return (0);}


```

Prof. Vittoria de Nitto Personè

33

33

Tradeoff - Confidence Versus Sample Size



The only way to make the interval smaller without lessening the level of confidence is to increase the sample size

$$\bar{x} \pm \frac{t^* s}{\sqrt{n-1}}$$

- Good news: with simulation, we can collect more data
- Bad news: interval size decreases with \sqrt{n} , not n

Prof. Vittoria de Nitto Personè


34

34

How Much More Data Is Enough?

- How large should n be to achieve an interval estimate $\bar{x} \pm w$ where w is user-specified?
- Answer: Use Welford's Algorithm with the algorithm p. 28 to iteratively collect data until a specified interval width is achieved

Note: if n is large then t^* is essentially independent of n




35

35

Asymptotic Value of t^*

- The asymptotic (large n) value of t^* is

$$t_{\infty}^* = \lim_{n \rightarrow \infty} \text{idfStudent}(n-1, 1-\alpha/2) = \text{idfNormal}(0.0, 1.0, 1-\alpha/2)$$
- Unless α is very close to 0.0, if $n > 40$, the asymptotic value t_{∞}^* can be used
- If $n > 40$ and wish to construct a 95% confidence interval estimate, $t_{\infty}^* = 1.960$ can be used in the algorithm on p.28



36

36

Example

- Given a reasonable guess for s and a user-specified half-width parameter w , if t_{∞}^* is used in place of t^*

n can be determined by solving $w = \frac{t^* s}{\sqrt{n-1}}$ for n :

$$n = \left\lceil \left(\frac{t_{\infty}^* s}{w} \right)^2 \right\rceil + 1$$

provided $n > 40$

- For example, if $s=3.0$ and want to estimate μ with 95% confidence to within ± 0.5 , a value of $n = 139$ should be used

Prof. Vittoria de Nitto Personè

37

37

Example

$$n = \left\lceil \left(\frac{t_{\infty}^* s}{w} \right)^2 \right\rceil + 1$$

- If a reasonable guess for s is not available, w can be specified as a proportion of s thereby eliminating s from the previous equation
- For example, if w is 10% of s and 95% confidence is desired, $n = 385$ should be used to estimate μ to within $\pm w$

$$(w/s = 0.1)$$

See in the book algorithm 8.1.2 to obtain confidence interval starting from the sample x_1, x_2, \dots, x_n or from the half-width parameter w respectively

Prof. Vittoria de Nitto Personè

38

38

The meaning of confidence

Incorrect:

“For this 95% confidence interval, the probability that μ is within this interval is 0.95”

- Why incorrect?
 - μ is not a random variable; it is constant (but unknown)
 - the interval endpoints are random

Correct:

“If I create many 95% confidence intervals, approximately 95% of them should contain μ ”


Prof. Vittoria de Nitto Personè

39

39

Example

- 100 samples of size $n=9$ drawn from $Normal(6,3)$ population
- For each sample, construct a 95% confidence interval
- 95 intervals contain $\mu=6$
- Three intervals “too low”, two intervals “too high”



Prof. Vittoria de Nitto Personè

40

40

Exercise

- Exercises 8.1.1, 8.1.5
- Consider case study 1 or case study 2, at your choice. Derive the sample mean histogram from one run (as in the picture in slide 8) and for two different sizes for the samples. Compare the obtained results with reference to the Exponential sample mean histograms seen in this lecture (slide p.10).

Prof. Vittoria de Nitto Personè

41