



Performance Modeling of Computer Systems and Networks

Prof. Vittoria de Nitto Personè

Finite-Horizon and Infinite-Horizon Statistics

Università degli studi di Roma Tor Vergata
Department of Civil Engineering and Computer Science Engineering

Copyright © Vittoria de Nitto Personè, 2021

<https://creativecommons.org/licenses/by-nc-nd/4.0/>



(CC BY-NC-ND 4.0)

1

Simulation studies

Algorithm 1.2: using the resulting model

7. Design simulations experiments

- What parameters should be varied?
- perhaps many combinatoric possibilities

8. Make production runs

- Record initial conditions, input parameters
- Record statistical output

9. Analyze the output

- Random components → statistical analysis (means, standard deviations, percentiles, histograms etc.)

10. Make decisions

- The step9 results drive the decisions → actions
- Simulation should be able to correctly predict the outcome of these actions (→ further refinements)

11. Document the results

- summarize the gained insights in specific observations and conjectures useful for subsequent similar system models

Prof. Vittoria de Nitto Personè

2

2

Central limit theorem

If X_1, X_2, \dots, X_n is an iid sequence of random variables (RVs) with

- common mean μ
- common standard deviation σ

and if \bar{X} is the (sample) mean of these RVs

then \bar{X} approaches a $Normal(\mu, \sigma/\sqrt{n})$ as $n \rightarrow \infty$

Theorem 2

If x_1, x_2, \dots, x_n is an independent random sample from a "source" of data with unknown mean μ , if \bar{x} and s are the mean and standard deviation of this sample, and if n is large, it is approximately true that

$$t = \frac{\bar{x} - \mu}{s/\sqrt{n-1}}$$
is a $Student(n-1)$ random variate

Theorem 3

If x_1, x_2, \dots, x_n is an independent random sample from a "source" of data with unknown mean μ

- if \bar{x} and s are the sample mean and sample standard deviation
- n is large

Then, given a confidence parameter α with $0.0 < \alpha < 1.0$, there exists an associated positive real number t^* such that

$$Pr\left(\bar{x} - \frac{t^* s}{\sqrt{n-1}} \leq \mu \leq \bar{x} + \frac{t^* s}{\sqrt{n-1}}\right) \cong 1 - \alpha$$

Prof. Vittoria de Nitto Personè

3

3

Algorithm

To calculate an interval estimate for the unknown mean μ of the population from which a random sample x_1, x_2, \dots, x_n was drawn:


- pick a level of confidence $1 - \alpha$ (typically $\alpha = 0.05$)
- calculate the sample mean \bar{x} and standard deviation s (use Welford's algorithm)
- calculate the critical value $t^* = \text{idfStudent}(n-1, 1 - \alpha/2)$
- calculate the interval endpoints $\bar{x} \pm \frac{t^* s}{\sqrt{n-1}}$

If n is sufficiently large, then you are $(1 - \alpha) \times 100\%$ confident that the mean μ lies within the interval. The midpoint of the interval is \bar{x}

Prof. Vittoria de Nitto Personè

4

4



5

```

....
while ((t.arrival < STOP) || (number > 0)) {
    t.next= Min(t.arrival, t.completion); /* next event time */
    ....

printf(" ... jobs", index);
printf(" average interarrival time ..", t.last / index);
printf(" average wait ...", area.node / index);
printf(" average delay ...", area.queue / index);
printf(" average service time ...", area.service / index);
printf(" average # in the node ... ", area.node / t.current);
printf(" average # in the queue .. ", area.queue / t.current);
printf(" utilization ....", area.service / t.current);
printf(" jobs remaining in the node= ....", number);

```

Prof. Vittoria de Nitto Personè

6

STOP=200
seed (9 digits or less) >> 1234

for 103 jobs
average interarrival time = 1.93
average wait = 7.34
average delay = 5.60
average service time ... = 1.74
average # in the node ... = 3.69
average # in the queue .. = 2.81
utilization = 0.87
jobs remaining in the node= 0

for 89 jobs
average interarrival time = 2.22
average wait = 4.19
average delay = 2.61
average service time ... = 1.58
average # in the node ... = 1.88
average # in the queue .. = 1.17
utilization = 0.71
jobs remaining in the node= 0


for 99 jobs
average interarrival time = 2.01
average wait = 7.54
average delay = 5.79
average service time ... = 1.75
average # in the node ... = 3.76
average # in the queue .. = 2.89
utilization = 0.87
jobs remaining in the node= 4

for 88 jobs
average interarrival time = 2.25
average wait = 4.23
average delay = 2.64
average service time ... = 1.59
average # in the node ... = 1.88
average # in the queue .. = 1.17
utilization = 0.71
jobs remaining in the node= 1

Prof. Vittoria de Nitto Personè

7

7



seeds

	1234	5678	4321	
average wait	7.34	4.19	2.65	} STOP=200 about 100 jobs
theoretical value	7.54	4.23	2.68	

8 s


If the program is executed multiple times varying only the rngs initial seed from replication to replication,

- the average wait in the node will vary significantly
- for most replications, the average wait will not be close to the steady-state average wait
- the initial conditions affect the results


Prof. Vittoria de Nitto Personè

8

8



9



10

Def. *Steady-state statistics*

Steady-state system statistics are those statistics, **if they exist**, that are produced by simulating the operation of a *stationary* discrete-event system for an effectively infinite length of time

A *finite-horizon* discrete-event simulation is one for which the simulated operational time is finite

An *infinite-horizon* discrete-event simulation is one for which the simulated operational time is effectively infinite

- Transient system statistics are those statistics that are produced by a finite-horizon discrete-event simulation
- Steady-state statistics are produced by an infinite-horizon simulation
- The initial conditions affect finite-horizon statistics
- The initial conditions do not affect infinite-horizon statistics: after enough time, the system loses memory of its initial state

Another Important Distinction

- ✧ In an infinite-horizon simulation, the system “environment” is assumed to remain static
If the system is a single-server service node, both the arrival rate and the service rate are assumed to remain constant in time
- ✧ In a finite-horizon simulation, no need to assume a static environment

Relative Importance of Two Statistics

- The “traditional” view: steady-state statistics are most important
 - ✧ Steady-state statistics are better understood because they are much more easy to analyze mathematically
 - ✧ It is frequently difficult to accurately model initial conditions and non-stationary system parameters
- The “pragmatic” view: transient statistics are most important because steady-state is just a convenient fiction
- Depending on the application, both transient and steady-state statistics may be important

Relative Importance of Two Statistics

- Important to decide which statistics best characterize the system's performance

one of the most important skills:

the ability to decide, on a system-by-system basis,
which kind of statistics best characterizes the
 system's performance.

Steady-state or Transient Statistics

Consider a bank that opens at 9 AM and closes at 5 PM.

A finite-horizon simulation over the **8-hour** period produces **transient** statistics valuable in determining the optimal staffing of tellers throughout the day.

Consider a fast food restaurant with a drive-up window that experiences a lunch rush period between 11:45 AM and 1:15 PM with an arrival rate that remains constant over the rush period.

This **90-minute** period could be simulated for a much longer time period, producing **steady-state** statistics which might be valuable for estimating the average wait time at the drive-up window.

Initial and Terminal Conditions

- **Finite-horizon** discrete-event simulations are also known as *terminating simulations*
 - ✧ In program ssq4, the *system state* is idle at the beginning and at the end of the simulation
 - ✧ The *terminal condition* is specified by the “close the door” time
 - ✧ The *system state* of sis4 is the current and on-order inventory levels; these states are the same at the beginning and at the end of the simulation
 - ✧ The *terminal condition* is specified by the number of time intervals
- **Infinite-horizon** discrete-event simulations (*non-terminating simulations*) must be terminated; typically done using whatever stopping conditions are most convenient
 - ✧ The steady-state statistics are based on such a huge amount of data that a few “non-steady-state” data points accumulated at the beginning and the end of the simulation should have no significant impact (bias) on output statistics

Prof. Vittoria de Nitto Personè

17

17

Formal Representation

- The state variable $X(\cdot)$ is known formally as a *stochastic process*
 - The typical objective of a **finite-horizon** simulation of this system would be to estimate the time-averaged *transient* statistic

random variable $\rightarrow \bar{X}(\tau) = \frac{1}{\tau} \int_0^\tau X(t) dt$

where $\tau > 0$ is the terminal time

- The typical objective of an **infinite-horizon** simulation of this system would be to estimate the time-averaged *steady-state* statistics

is not a random variable $\rightarrow \bar{x} = \lim_{\tau \rightarrow \infty} \bar{X}(\tau) = \lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int_0^\tau X(t) dt$

Prof. Vittoria de Nitto Personè

18

18

Replication

- If a discrete-event simulation is repeated, varying only the rngs initial states from run to run, each run of the simulation program is a *replication* and the totality of replications is said to be *ensemble*
- Replications are used to generate *estimates* of the same transient statistic
- The initial seed for each replication should be chosen to be no replication-to-replication overlap
- The standard way is to use the final state of each rngs stream from *one* replication as the initial state for the *next* replication accomplished by calling PlantSeeds once outside the main replication loop

Independent Replications and Interval Estimation

Suppose the *finite-horizon* simulation is replicated n times, each time generating a state time history $x_i(t)$

$$\bar{x}_i(\tau) = \frac{1}{\tau} \int_0^\tau x_i(t) dt$$

where $i = 1, 2, \dots, n$ is the replication index

Each data point $\bar{x}_i(\tau)$ is an independent observation of the random variable $\bar{X}(\tau)$

If n is large enough, the pdf of $\bar{X}(\tau)$ can be estimated from a histogram of the $\bar{x}_i(\tau)$

Independent Replications and Interval Estimation

In practise, it is usually only the expected value $E[\bar{X}(\tau)]$ that is desired. A *point* estimate of this transient statistic is available as an *ensemble average*, even if n is not large

$$\frac{1}{n} \sum_{i=1}^n \bar{x}_i(\tau)$$


An interval estimate for $E[\bar{X}(\tau)]$ can be calculated
Use the interval estimation technique from Section 8.1

Example 8.3.6

A modified version of ssq2 was used to produce **20 replications**

- 100 jobs processed through M/M/1 service node
 - Node is initially idle
 - Arrival rate is = 1.0
 - Service rate is = 1.25
- The resulting 20 observations of the **average wait** in the node:

from program estimate 95%-confidence-interval: we are 95% confident that if we were to do millions of replications the ensemble average would be somewhere between 2.29 and 4.37



Example 8.3.7

- The modified version of program ssq2 was used to produce **60 more replications**
- Consistent with \sqrt{n} rule, expect two-fold decrease in the width of the interval estimate
- Based on 80 replications, the resulting 95% confidence interval estimate was 3.25 ± 0.39 (2.86, 3.64)



In this case 100 jobs are not enough to produce steady-state statistics

the bias of the initially idle state is still evident in the transient statistic


Prof. Vittoria de Nitto Personè

23

23

Example 8.3.8

- As a continuation of Example 8.3.6, the number of **jobs per replication** was increased from 100 to 1000
- 20 replications were used to produce 20 observations of the average wait in the node (3.45, 4.19)



Relative to Example 8.3.6, much more symmetric sample mean in Example 8.3.8 (2.29, 4.37)



Prof. Vittoria de Nitto Personè

24

24

Example 8.3.8

- The 1000-jobs per replication results are more consistent with the underlying theory of interval estimation
 - Requires a sample mean distribution that is approximately $Normal(\mu, \sigma / \sqrt{n})$
 - Sample mean distribution is centered on (unknown) population
- 1000 jobs may achieve steady-state; 100 jobs cannot

Prof. Vittoria de Nitto Personè

25