

Problem 2.2.2

2.2-2. The trapezoidal rule (modified Euler method) for numerical integration approximates the integral of a function $x(t)$ by summing trapezoid areas as shown in Fig. P2.2-2. Let $y(t)$ be the integral of $x(t)$.




Figure P2.2-2 Trapezoidal rule for numerical integration.

Figure P2.2-2 Full Alternative Text

a. Write the difference equation relating $y[(k+1)T]$, $y(kT)$, $x[(k+1)T]$, and $x(kT)$ for this rule.

b. Show that the transfer function for this integrator is given by

$$\frac{Y(s)}{X(s)} = \frac{(1/2)(s+1)}{s-1}$$

(a)

$$y[(k+1)T] = y(kT) + \int_{kT}^{(k+1)T} x(t) dt \quad \approx \frac{T}{2} [x(kT) + x((k+1)T)]$$

D.F. \rightarrow

$$= y(kT) + \frac{T}{2} [x(kT) + x((k+1)T)] \quad \#$$

or

$$y[k+1] = y[k] + \frac{T}{2} (x[k] + x[k+1]) \quad \#$$

(b) $(z \text{ advance})$

$$\mathcal{Z}\{y((k+1)T)\} = z \cdot Y(z)$$

$$\mathcal{Z}\{y(kT)\} = Y(z)$$

$$\mathcal{Z}\{x((k+1)T)\} = z X(z)$$

$$\mathcal{Z}\{x(kT)\} = X(z)$$

$$\therefore z Y(z) = Y(z) + \frac{T}{z} (X(z) + z X(z))$$

$$(z-1) Y(z) = \frac{T}{z} (1+z) X(z)$$

$$\frac{Y(z)}{X(z)} = \frac{T}{z} \frac{(z+1)}{(z-1)} \#$$

Problem 2.2.3

2.2-3.

- a. The transfer function for the right-side rectangular-rule integrator was found in Problem 2.2-1 to be $Y(z)/X(z) = Tz/(z - 1)$. We would suspect that the reciprocal of this transfer function should yield (an approximation to a differentiator.) That is, if $w(kT)$ is a numerical derivative of $x(t)$ at $t = kT$,

$$\frac{W(z)}{X(z)} = \frac{z - 1}{Tz}$$

Write the difference equation describing this differentiator.

- b. Draw a figure similar to those in Fig. P2.2-1 illustrating the approximate differentiation.

- c. Repeat part (a) for the left-side rule, where $W(z)/X(z) = T/(z - 1)$.

- d. Repeat part (b) for the differentiator of part (c).

I use Matlab

$$(a) \frac{W(z)}{X(z)} = \frac{(z-1)}{Tz} = \frac{(1-z^{-1})}{T}$$

$$\Rightarrow W(z) \cdot T = X(z) - z^{-1}X(z)$$

\int inverse z

$$\Rightarrow W(kT) \cdot T = x(kT) - x((k-1)T)$$

$$\Rightarrow W(kT) = [x(kT) - x((k-1)T)] / T \quad \#$$

$$(b) \frac{W(z)}{X(z)} = \frac{T}{(z-1)} = \frac{Tz^{-1}}{(1-z^{-1})}$$

$$\Rightarrow T \cdot z^{-1}X(z) = W(z) - z^{-1}W(z)$$

\int inverse z

$$\Rightarrow T \cdot x[(k-1)T] = w[kT] - w[(k-1)T]$$

But, it's the integrator. I can't do part (d). (∵ problem requests ^adifferentiator)

Awww, I think the question should be like using $\frac{Y(z)}{X(z)} = \frac{T}{(z-1)}$

to describe the integrator, then $\frac{W(z)}{X(z)} = \frac{(z-1)}{T}$ is the differentiator.

let do it according to the question above.

$$\frac{W(z)}{X(z)} = \frac{(z-1)}{T} \Rightarrow T \cdot W(z) = X(z)(z-1) = zX(z) - X(z)$$



$\int z^{-1}$

$$Tw(kT) = x((k+1)T) - x(kT)$$

$$\Rightarrow w(kT) = [x((k+1)T) - x(kT)] / T \quad \#$$

(b)(d)

$$x[k] = x[k] - x[k-1] \quad \#$$



$$x[k] = x[k+1] - x[k] \quad \#$$

Problem 2.6.3

2.6-3. Given the difference equation

$$x(k) - x(k-1) + x(k-2) = e(k)$$

where $e(k) = 1$ for $k \geq 0$.

- Solve for $x(k)$ as a function of k , using the z -transform. Give the values of $x(0)$, $x(1)$, and $x(2)$.
- Verify the values $x(0)$, $x(1)$, and $x(2)$, using the power-series method.
- Verify the values $x(0)$, $x(1)$, and $x(2)$ by solving the difference equation directly.
- Will the final-value property give the correct value for $x(\infty)$?

(a)

$$x(k) - x(k-1) + x(k-2) = e(k) = 1 \text{ (unit step)}$$

$$X(z) - z^{-1}X(z) + z^{-2}X(z) = E(z) = \frac{z}{z-1}$$

$$X(z)(1 - z^{-1} + z^{-2}) = \frac{z}{z-1}$$

$$X(z) = \frac{z^3}{(z-1)(z^2-z+1)} \quad \left(\text{pole} = 1, \frac{1 \pm \sqrt{3}j}{2} \right)$$

$$\frac{X(z)}{z} = \frac{z^2}{(z-1)(z^2-z+1)} = \frac{A}{(z-1)} + \frac{Bz+C}{(z^2-z+1)}$$

$$= \frac{A(z^2-z+1) + Bz(z-1) + C(z-1)}{(z-1)(z^2-z+1)} \quad \begin{cases} A+B=1 \\ -A-B+C=0 \\ A-C=0 \end{cases} \Rightarrow \begin{cases} C=1 \\ A=C \\ \Rightarrow B=0 \end{cases}$$

$$\Rightarrow X(z) = \frac{z}{(z-1)} + \frac{z}{z^2-z+1} \xrightarrow{z^{-1}} x(k) = 1 + 2\sqrt{3}^{-1} \sin\left(\frac{\pi k}{3}\right) \quad \begin{cases} x(0) = 1 \\ x(1) = 1 + \frac{2}{\sqrt{3}} \cdot \frac{\sqrt{3}}{2} = 2 \\ x(2) = 1 + \frac{2}{\sqrt{3}} \cdot \frac{\sqrt{3}}{2} = 2 \end{cases}$$

(b) power series method:

$$X(z) = \frac{z^3}{(z-1)(z^2-z+1)} = \frac{z^3}{z^3 - 2z^2 + z - 1}$$

$$\begin{array}{r} 1 + \frac{2}{z} + \frac{2}{z^2} + \frac{1}{z^3} + \dots \\ z^3 - 2z^2 + z - 1 \overline{) z^3} \\ \underline{z^3 - 2z^2 + z - 1} \\ 2z^2 - 2z + 1 \\ \underline{2z^2 - 4z + 4 - \frac{2}{z}} \\ 2z - 3 + \frac{2}{z} \\ \underline{2z - 4 + \frac{4}{z} - \frac{2}{z^2}} \\ 1 - \frac{2}{z} + \frac{2}{z^2} \\ \underline{1 - \frac{2}{z} + \frac{2}{z^2} - \frac{1}{z^3}} \\ \frac{1}{z^3} \end{array}$$

∴ first term of the sequence are: $\{1, 2, 2, 1, \dots\}$

$$\Rightarrow x(0)=1, x(1)=2, x(2)=2, \dots \#$$

(c) solve directly

$$x(k) = x(k-1) - x(k-2) + 1$$

let initial conditions: $x(-1) = x(-2) = 0$

$$\text{then } \begin{cases} x(0) = 1 \\ x(1) = x(0) - x(-1) + 1 = 2 \\ x(2) = x(1) - x(0) + 1 = 2 \end{cases} \#$$

(d) Final Value Theorem is:

$$\lim_{k \rightarrow \infty} x(k) = \lim_{z \rightarrow 1} (z-1)E(z) = \lim_{z \rightarrow 1} (z-1) \cdot \frac{z^3}{(z-1)(z^2-z+1)} \\ = \frac{z^3}{(z^2-z+1)} = 1. \quad \text{has poles} = e^{\frac{j\pi}{3}}, e^{-\frac{j\pi}{3}}$$

but $(|e^{\frac{j\pi}{3}}|^2 + |e^{-\frac{j\pi}{3}}|^2)^{\frac{1}{2}} = \text{magnitude} = 1$ (it's on unit circle, not inside)

also, $\lim_{k \rightarrow \infty} x[k]$ does NOT exist since the sequence

$x[k] = (1, 2, 2, 0, 0, \dots) \Rightarrow$ it's **NOT CONVERGENCE** #

to sum up: **NO** the FVT is NOT Applicable.

Problem 2.7.1

2.7-1. (a) Find $e(0)$, $e(1)$, and $e(10)$ for

$$E(z) = \frac{0.1}{z(z-0.9)}$$

using the inversion formula.

$$e_k = \frac{1}{2\pi j} \oint E(z) z^{k-1} dz = \sum \text{Res} \{ E(z) z^{k-1} \}$$

(b) Check the value of $e(0)$ using the initial-value property. $\lim_{k \rightarrow 0} x[k] = \lim_{z \rightarrow \infty} X(z)$

(c) Check the values calculated in part (a) using partial fractions.

$$E(z) = z \cdot \frac{P(z)}{Q(z)}$$

(d) Find $e(k)$ for $k = 0, 1, 2, 3$, and 4 if $\mathcal{Z}[e(k)]$ is given by

$$E(z) = \frac{1.98z}{(z^2 - 0.9z + 0.9)(z - 0.8)(z^2 - 1.2z + 0.27)}$$

$(z - 0.9)(z - 0.3)$

(e) Find a function $e(t)$ which, when sampled at a rate of 10 Hz ($T = 0.1s$), results in the transform $E(z) = 2z/(z - 0.8)$.

(f) Repeat part (e) for $E(z) = 2z/(z + 0.8)$.

$$\Rightarrow A = -\frac{1}{9}, B = \frac{1}{9}$$

(g) From parts (e) and (f), what is the effect on the inverse z-transform of changing the sign on a real pole?

(a) $E(z) = (0.1) \cdot \frac{1}{z(z-0.9)} \Rightarrow E(z) z^{k-1} = 0.1 \cdot \frac{z^{k-1}}{z(z-0.9)} = 0.1 \cdot \frac{z^k}{z^2(z-0.9)}$

it has pole $z = 0.9$ and repeated poles at $z = 0$ (order 2)

$$e[k] = \frac{1}{2\pi j} \oint E(z) z^{k-1} dz = \sum \text{Res} \{ E(z) z^{k-1} \}$$

$k=0$ ($e[0]$) * At $z = 0.9$: $\text{Res} \{ E(z) z^{k-1} \} = \lim_{z \rightarrow 0.9} [(z - 0.9) E(z) z^{k-1}]$
 (singularity of order 1) $= \lim_{z \rightarrow 0.9} (z - 0.9) \frac{0.1 z^0}{z^2(z-0.9)} = 0.1 \cdot (0.9)^{-2}$ (a) $\therefore e[0] = \sum \text{Res} \{ E(z) z^{k-1} \}$
 $= (a) + (b) = 0$ #

* At $z = 0$: $\text{Res} \{ E(z) z^{k-1} \} = \lim_{z \rightarrow 0} \frac{1}{(2-1)!} \frac{d^{2-1}}{dz^{2-1}} [(z-0)^2 \cdot 0.1 \frac{z^0}{z^2(z-0.9)}]$
 ($n=2$ (order)) $= \lim_{z \rightarrow 0} \frac{1}{1} \cdot \frac{d}{dz} \left[\frac{z^0}{(z-0.9)} \cdot (0.1) \right] = \lim_{z \rightarrow 0} \left(\frac{-0.1}{(z-0.9)^2} \right) = -0.1 \cdot (0.9)^{-2}$ (b)

$$(k=1) e[1]: E(z) \cdot z^{k-1} = \frac{0.1}{z(z-0.9)} \cdot 1$$

$$z = 0.9: \lim_{z \rightarrow 0.9} (z-0.9) \cdot \frac{0.1}{z(z-0.9)} = \frac{0.1}{0.9} = \frac{1}{9} \quad (a)$$

$$z = 0: \lim_{z \rightarrow 0} \frac{1}{(z-1)!} \cdot \frac{d^{z-1}}{dz^{z-1}} \left[(1-z-0) \cdot \frac{0.1}{z(z-0.9)} \right] = \frac{-1}{9} \quad (b)$$

$$\therefore e[1] = \sum \text{Res } E(z) = (a) + (b) = 0 \quad \#$$

$$(k=10) e[10]: E(z) \cdot z^{k-1} = \frac{0.1 \cdot z^9}{z(z-0.9)} = \frac{0.1 \cdot z^8}{(z-0.9)}$$

$$z = 0.9: \lim_{z \rightarrow 0.9} (z-0.9) \cdot \frac{0.1 \cdot z^8}{(z-0.9)} = 0.1 \cdot 0.9^8$$

$$z = 0: \lim_{z \rightarrow 0} \frac{1}{(z-1)!} \cdot \frac{d^{z-1}}{dz^{z-1}} \left[(1-z-0) \cdot \frac{0.1 \cdot z^8}{(z-0.9)} \right] = 0$$

$$\therefore e[10] = \sum \text{Res} \{ E(z) z^9 \} = 0.1 (0.9)^8 \quad \#$$

(b)

$$\text{IVT: } e(0) = \lim_{z \rightarrow 0} E(z) = \lim_{z \rightarrow 0} \left(\frac{0.1}{z(z-0.9)} \right) = 0 \Rightarrow \text{verified } \checkmark \quad \#$$

(c)

$$E(z) = \frac{0.1}{z(z-0.9)} = \frac{A}{z} + \frac{B}{z-0.9}$$

$$= A(z-0.9) + Bz$$

$$\begin{cases} A+B=0 \\ -0.9A=0.1 \end{cases} \Rightarrow A = \frac{-0.1}{0.9} \Rightarrow A = -\frac{1}{9}, B = \frac{1}{9}$$

$$\text{then: } E(z) = \frac{-1}{9} \frac{1}{z} + \frac{1}{9} \frac{1}{(z-0.9)}$$

$$z^{-1} \downarrow$$

$$e[k] = \frac{1}{9} (0.9^{k-1} u[k-1] - \delta[k-1])$$

verify =

$$k=0: e[0] = \frac{1}{9} (0.9^{-1} u[-1] - \delta[-1]) = 0 + 0 = 0 \quad \checkmark$$

$$k=1: e[1] = \frac{1}{9} (0.9^0 u[0] - \delta[0]) = \frac{1}{9} (1-1) = 0 \quad \checkmark$$

$$k=10: e[10] = \frac{1}{9} (0.9^9 u[9] - \delta[9]) = \frac{1}{9} (0+0.9^9) = (0.1) \cdot 0.9^8 \quad \checkmark \quad \#$$

(d) since the denominator is order 5, and try to divide the numerator of order 1. As a result, the coefficient of first three terms should be zero by power series method $\Rightarrow e(0) = e(1) = e(2) = e(3) = 0$

then, $e(4)$ is equal to 1.98 because we need to minus

$$\text{the numerator } \Rightarrow \begin{cases} e(0) = e(1) = e(2) = e(3) = 0 \\ e(4) = 1.98 \end{cases}$$

(the first nonzero term start from k

$$\text{where } k = \text{degree(Denominator)} - \text{degree(Nominator)} = 4 \quad \#$$

(e) find $e(t)$, with $T = 10 \text{ Hz} = 0.1 \text{ sec}$ and $E(z) = \frac{2z}{(z-0.8)}$

$$\Rightarrow E(z) = 2 \cdot \frac{z}{(z-0.8)} \xrightarrow{z^{-1}} e[k] = 2(0.8)^k, k=0,1,\dots$$

$$0.8 = e^{10s} \Rightarrow \log_e 0.8 = 10s \Rightarrow s = 10^{-1} \log_e(0.8)$$

$$\therefore e(t) = e[kT] = 2(0.8)^k = 2 \cdot e^{(10s)t}, \text{ where } s = 10^{-1} \log_e(0.8) \quad \#$$

(f) again, find $e(t)$, with $T = 10 \text{ Hz} = 0.1 \text{ sec}$ but $E(z) = \frac{2z}{(z+0.8)}$

$$\Rightarrow E(z) = 2 \cdot \frac{z}{(z+0.8)} \xrightarrow{z^{-1}} e[k] = 2(0.8)^k (-1)^k, k=0,1,\dots$$





$$0.8 = e^{10s} \Rightarrow \log_e 0.8 = 10s \Rightarrow s = 10^{-1} \log_e(0.8)$$

$$(-1)^k = \cos(\pi k) = \cos\left(\frac{\pi t}{0.1}\right)$$

$$\therefore e(t) = e[kT] = 2(0.8)^k (-1)^k = 2 \cdot e^{(10s)t} \cdot \cos(10\pi t), \text{ where } s = 10^{-1} \log_e(0.8) \quad \#$$

(g) is in the next page!

[9] I use Matlab to plot (e) and (f) for understanding



Change a real pole from $+0.8$ to -0.8 causes the inverse z-transfer is multiplied by $(-1)^k$, which means it causes the corresponding time signal to oscillate at the highest freq. to be captured by the sampler. #