

# Machine Learning

## Lecture 01-2: Basics of Information Theory

Nevin L. Zhang

lzhang@cse.ust.hk

Department of Computer Science and Engineering  
The Hong Kong University of Science and Technology

# Outline


## 1 Jensen's Inequality

## 2 Entropy

## 3 Divergence


## 4 Mutual Information

# Concave functions



- A function  $f$  is **concave** on interval  $I$


# Concave functions



- A function  $f$  is **concave** on interval  $I$  if for any  $x, y \in I$ ,

$$\lambda f(x) + (1 - \lambda)f(y) \leq f(\lambda x + (1 - \lambda)y) \text{ for any } \lambda \in [0, 1]$$

# Concave functions




- A function  $f$  is **concave** on interval  $I$  if for any  $x, y \in I$ ,

$$\lambda f(x) + (1 - \lambda)f(y) \leq f(\lambda x + (1 - \lambda)y) \text{ for any } \lambda \in [0, 1]$$

Weighted average of function is upper bounded by function of weighted average.

# Concave functions




- A function  $f$  is **concave** on interval  $I$  if for any  $x, y \in I$ ,

$$\lambda f(x) + (1 - \lambda)f(y) \leq f(\lambda x + (1 - \lambda)y) \text{ for any } \lambda \in [0, 1]$$

Weighted average of function is upper bounded by function of weighted average.

It is strictly concave

# Concave functions




- A function  $f$  is **concave** on interval  $I$  if for any  $x, y \in I$ ,

$$\lambda f(x) + (1 - \lambda)f(y) \leq f(\lambda x + (1 - \lambda)y) \text{ for any } \lambda \in [0, 1]$$

Weighted average of function is upper bounded by function of weighted average.

It is **strictly concave** if the equality holds only when  $x=y$ .

# Concave functions



- A function  $f$  is **concave** on interval  $I$  if for any  $x, y \in I$ ,

$$\lambda f(x) + (1 - \lambda)f(y) \leq f(\lambda x + (1 - \lambda)y) \text{ for any } \lambda \in [0, 1]$$

Weighted average of function is upper bounded by function of weighted average.

It is **strictly concave** if the equality holds only when  $x=y$ .



# Jensen's Inequality

## Theorem (1.1)

*Suppose function  $f$  is concave on interval  $I$ . Then*

- *For any  $p_i \in [0, 1]$ ,  $\sum_{i=1}^n p_i = 1$  and  $x_i \in I$ .*

# Jensen's Inequality

## Theorem (1.1)

*Suppose function  $f$  is concave on interval  $I$ . Then*

- *For any  $p_i \in [0, 1]$ ,  $\sum_{i=1}^n p_i = 1$  and  $x_i \in I$ .*

$$\sum_{i=1}^n p_i f(x_i) \leq f\left(\sum_{i=1}^n p_i x_i\right)$$

# Jensen's Inequality

## Theorem (1.1)

*Suppose function  $f$  is concave on interval  $I$ . Then*

- *For any  $p_i \in [0, 1]$ ,  $\sum_{i=1}^n p_i = 1$  and  $x_i \in I$ .*

$$\sum_{i=1}^n p_i f(x_i) \leq f\left(\sum_{i=1}^n p_i x_i\right)$$

*Weighted average of function is upper bounded by function of weighted average.*

# Jensen's Inequality

## Theorem (1.1)

*Suppose function  $f$  is concave on interval  $I$ . Then*

- *For any  $p_i \in [0, 1]$ ,  $\sum_{i=1}^n p_i = 1$  and  $x_i \in I$ .*

$$\sum_{i=1}^n p_i f(x_i) \leq f\left(\sum_{i=1}^n p_i x_i\right)$$

*Weighted average of function is upper bounded by function of weighted average.*

- *If  $f$  is strictly CONCAVE, the equality holds iff  $p_i \times p_j \neq 0$  implies  $x_i = x_j$ .*

# Jensen's Inequality

## Theorem (1.1)

*Suppose function  $f$  is concave on interval  $I$ . Then*

- *For any  $p_i \in [0, 1]$ ,  $\sum_{i=1}^n p_i = 1$  and  $x_i \in I$ .*

$$\sum_{i=1}^n p_i f(x_i) \leq f\left(\sum_{i=1}^n p_i x_i\right)$$


*Weighted average of function is upper bounded by function of weighted average.*

- *If  $f$  is strictly CONCAVE, the equality holds iff  $p_i \times p_j \neq 0$  implies  $x_i = x_j$ .*

Exercise: Prove this (using induction).


# Logarithmic function

- The logarithmic function is concave in the interval  $(0, \infty)$ :



# Logarithmic function

- The logarithmic function is concave in the interval  $(0, \infty)$ :




- Hence

$$\sum_{i=1}^n p_i \log(x_i) \leq \log\left(\sum_{i=1}^n p_i x_i\right) \quad 0 \leq x_i$$

# Logarithmic function

- The logarithmic function is concave in the interval  $(0, \infty)$ :



- Hence


$$\sum_{i=1}^n p_i \log(x_i) \leq \log\left(\sum_{i=1}^n p_i x_i\right) \quad 0 \leq x_i$$

- In words, exchanging  $\sum_i p_i$  with  $\log$  increases quantity.



# Logarithmic function

- The logarithmic function is concave in the interval  $(0, \infty)$ :



- Hence

$$\sum_{i=1}^n p_i \log(x_i) \leq \log\left(\sum_{i=1}^n p_i x_i\right) \quad 0 \leq x_i$$

- In words, **exchanging  $\sum_i p_i$  with  $\log$  increases quantity**. Or, swapping expectation and logarithm increases quantity:

$$E[\log x] \leq \log E[x].$$

# Outline

- 1 Jensen's Inequality
- 2 Entropy
- 3 Divergence
- 4 Mutual Information

# Entropy

- The **entropy** of a random variable  $X$ :

$$H(X) = \sum_x P(X) \log \frac{1}{P(X)}$$

# Entropy

- The **entropy** of a random variable  $X$ :

$$H(X) = \sum_x P(X) \log \frac{1}{P(X)} = -E_P[\log P(X)]$$

with convention that  $0 \log(1/0) = 0$ .

# Entropy

- The **entropy** of a random variable  $X$ :

$$H(X) = \sum_x P(X) \log \frac{1}{P(X)} = -E_P[\log P(X)]$$

with convention that  $0 \log(1/0) = 0$ .

- Base of logarithm is 2, unit is bit.

# Entropy

- The **entropy** of a random variable  $X$ :

$$H(X) = \sum_x P(X) \log \frac{1}{P(X)} = -E_P[\log P(X)]$$

with convention that  $0 \log(1/0) = 0$ .

- Base of logarithm is 2, unit is bit.
- Sometimes, also called the entropy of the distribution,  $H(P)$ .

# Entropy

- The **entropy** of a random variable  $X$ :

$$H(X) = \sum_x P(X) \log \frac{1}{P(X)} = -E_P[\log P(X)]$$

with convention that  $0 \log(1/0) = 0$ .

- Base of logarithm is 2, unit is bit.
- Sometimes, also called the entropy of the distribution,  $H(P)$ .
- $H(X)$  **measures the amount of uncertainty about  $X$ .**

# Entropy

- The **entropy** of a random variable  $X$ :

$$H(X) = \sum_X P(X) \log \frac{1}{P(X)} = -E_P[\log P(X)]$$

with convention that  $0 \log(1/0) = 0$ .

- Base of logarithm is 2, unit is bit.
- Sometimes, also called the entropy of the distribution,  $H(P)$ .
- $H(X)$  **measures the amount of uncertainty about  $X$ .**

For real-valued variable, replace  $\sum_X \dots$  with  $\int \dots dx$ .



# Entropy

Example:

- $X$  — result of coin tossing
- $Y$  — result of dice throw
- $Z$  — result of randomly pick a card from a deck of 54

# Entropy

Example:

- $X$  — result of coin tossing
- $Y$  — result of dice throw
- $Z$  — result of randomly pick a card from a deck of 54
- Which one has the highest uncertainty?

# Entropy

Example:

- $X$  — result of coin tossing
- $Y$  — result of dice throw
- $Z$  — result of randomly pick a card from a deck of 54
- Which one has the highest uncertainty?
- Entropy:

$$H(X) = \frac{1}{2} \log 2 + \frac{1}{2} \log 2 = 1(\log 2)$$

$$H(Y) = \frac{1}{6} \log 6 + \dots + \frac{1}{6} \log 6 = \log 6$$

$$H(Z) = \frac{1}{54} \log 54 + \dots + \frac{1}{54} \log 54 = \log 54$$

Indeed we have:

$$H(X) < H(Y) < H(Z).$$

# Entropy


- $X$  binary. The chart on the right shows  $H(X)$  as a function of  $p=P(X=1)$ .

# Entropy

- $X$  binary. The chart on the right shows  $H(X)$  as a function of  $p=P(X=1)$ .
- The higher  $H(X)$  is, the more uncertainty about the value of  $X$

# Entropy

- $X$  binary. The chart on the right shows  $H(X)$  as a function of  $p=P(X=1)$ .
- The higher  $H(X)$  is, the more uncertainty about the value of  $X$



# Entropy

## Proposition (1.2)

- $H(X) \geq 0$

# Entropy

## Proposition (1.2)

- $H(X) \geq 0$
- $H(X) = 0$  iff  $P(X=x) = 1$  for some  $x \in \Omega_X$ .



# Entropy

## Proposition (1.2)

- $H(X) \geq 0$
- $H(X) = 0$  iff  $P(X=x) = 1$  for some  $x \in \Omega_X$ . i.e. iff no uncertainty.

# Entropy

## Proposition (1.2)

- $H(X) \geq 0$
- $H(X) = 0$  iff  $P(X=x) = 1$  for some  $x \in \Omega_X$ . i.e. iff no uncertainty.
- $H(X) \leq \log(|X|)$  with equality iff  $P(X=x)=1/|X|$ .

# Entropy

## Proposition (1.2)

- $H(X) \geq 0$
- $H(X) = 0$  iff  $P(X=x) = 1$  for some  $x \in \Omega_X$ . i.e. iff no uncertainty.
- $H(X) \leq \log(|X|)$  with equality iff  $P(X=x)=1/|X|$ .

# Entropy

## Proposition (1.2)

- $H(X) \geq 0$
- $H(X) = 0$  iff  $P(X=x) = 1$  for some  $x \in \Omega_X$ . i.e. iff no uncertainty.
- $H(X) \leq \log(|X|)$  with equality iff  $P(X=x)=1/|X|$ .  
*Uncertainty is the highest in the case of uniform distribution.*

**Proof:** Because  $\log$  is concave, by Jensen's inequality:

# Entropy

## Proposition (1.2)

- $H(X) \geq 0$
- $H(X) = 0$  iff  $P(X=x) = 1$  for some  $x \in \Omega_X$ . i.e. iff no uncertainty.
- $H(X) \leq \log(|X|)$  with equality iff  $P(X=x)=1/|X|$ .  
*Uncertainty is the highest in the case of uniform distribution.*

**Proof:** Because  $\log$  is concave, by Jensen's inequality:

$$H(X) = \sum_x P(X) \log \frac{1}{P(X)}$$

# Entropy

## Proposition (1.2)

- $H(X) \geq 0$
- $H(X) = 0$  iff  $P(X=x) = 1$  for some  $x \in \Omega_X$ . i.e. iff no uncertainty.
- $H(X) \leq \log(|X|)$  with equality iff  $P(X=x)=1/|X|$ .  
*Uncertainty is the highest in the case of uniform distribution.*

**Proof:** Because  $\log$  is concave, by Jensen's inequality:

$$\begin{aligned} H(X) &= \sum_x P(X) \log \frac{1}{P(X)} \\ &\leq \log \sum_x P(X) \frac{1}{P(X)} \end{aligned}$$

# Entropy

## Proposition (1.2)

- $H(X) \geq 0$
- $H(X) = 0$  iff  $P(X=x) = 1$  for some  $x \in \Omega_X$ . i.e. iff no uncertainty.
- $H(X) \leq \log(|X|)$  with equality iff  $P(X=x)=1/|X|$ .  
*Uncertainty is the highest in the case of uniform distribution.*

**Proof:** Because  $\log$  is concave, by Jensen's inequality:

$$\begin{aligned} H(X) &= \sum_x P(X) \log \frac{1}{P(X)} \\ &\leq \log \sum_x P(X) \frac{1}{P(X)} = \log |X| \end{aligned}$$

# Conditional entropy

- The **conditional entropy** of  $Y$  given event  $X=x$ :



# Conditional entropy

- The **conditional entropy** of  $Y$  given event  $X=x$ :
  - Entropy of the conditional distribution  $P(Y|X = x)$ , i.e.

# Conditional entropy

- The **conditional entropy** of  $Y$  given event  $X=x$ :
  - Entropy of the conditional distribution  $P(Y|X=x)$ , i.e.

$$H(Y|X=x) = \sum_Y P(Y|X=x) \log \frac{1}{P(Y|X=x)}$$

# Conditional entropy

- The **conditional entropy** of  $Y$  given event  $X=x$ :
  - Entropy of the conditional distribution  $P(Y|X=x)$ , i.e.

$$H(Y|X=x) = \sum_Y P(Y|X=x) \log \frac{1}{P(Y|X=x)}$$

The uncertainty that remains about  $Y$  when  $X$  is known to be  $y$ .

# Conditional entropy

- The **conditional entropy** of  $Y$  given event  $X=x$ :
  - Entropy of the conditional distribution  $P(Y|X=x)$ , i.e.

$$H(Y|X=x) = \sum_Y P(Y|X=x) \log \frac{1}{P(Y|X=x)}$$

The uncertainty that remains about  $Y$  when  $X$  is known to be  $y$ .

- It is possible that  $H(Y|X=x) > H(Y)$

# Conditional entropy

- The **conditional entropy** of  $Y$  given event  $X=x$ :
  - Entropy of the conditional distribution  $P(Y|X=x)$ , i.e.

$$H(Y|X=x) = \sum_Y P(Y|X=x) \log \frac{1}{P(Y|X=x)}$$

The uncertainty that remains about  $Y$  when  $X$  is known to be  $y$ .

- It is possible that  $H(Y|X=x) > H(Y)$ 
  - Intuitively  $X=x$  might contradicts our prior knowledge about  $Y$  and increase our uncertainty about  $Y$

# Conditional entropy

- The **conditional entropy** of  $Y$  given event  $X=x$ :
  - Entropy of the conditional distribution  $P(Y|X=x)$ , i.e.

$$H(Y|X=x) = \sum_Y P(Y|X=x) \log \frac{1}{P(Y|X=x)}$$

The uncertainty that remains about  $Y$  when  $X$  is known to be  $y$ .

- It is possible that  $H(Y|X=x) > H(Y)$ 
  - Intuitively  $X=x$  might contradicts our prior knowledge about  $Y$  and increase our uncertainty about  $Y$
  - Exercise: Give example.

# Conditional Entropy

- The **conditional entropy** of  $Y$  given variable  $X$ :

# Conditional Entropy

- The **conditional entropy** of  $Y$  given variable  $X$ :

$$H(Y|X) = \sum_x P(X = x)H(Y|X=x)$$



# Conditional Entropy

- The **conditional entropy** of  $Y$  given variable  $X$ :

$$\begin{aligned} H(Y|X) &= \sum_x P(X=x) H(Y|X=x) \\ &= \sum_x P(X) \sum_Y P(Y|X) \log \frac{1}{P(Y|X)} \end{aligned}$$

# Conditional Entropy

- The **conditional entropy** of  $Y$  given variable  $X$ :

$$\begin{aligned} H(Y|X) &= \sum_x P(X=x) H(Y|X=x) \\ &= \sum_x P(X) \sum_Y P(Y|X) \log \frac{1}{P(Y|X)} \\ &= \sum_{X,Y} P(X,Y) \log \frac{1}{P(Y|X)} \end{aligned}$$

# Conditional Entropy

- The **conditional entropy** of  $Y$  given variable  $X$ :

$$\begin{aligned} H(Y|X) &= \sum_x P(X=x) H(Y|X=x) \\ &= \sum_x P(X) \sum_Y P(Y|X) \log \frac{1}{P(Y|X)} \\ &= \sum_{X,Y} P(X,Y) \log \frac{1}{P(Y|X)} \\ &= -E[\log P(Y|X)] \end{aligned}$$

# Conditional Entropy

- The **conditional entropy** of  $Y$  given variable  $X$ :

$$\begin{aligned}
 H(Y|X) &= \sum_x P(X=x) H(Y|X=x) \\
 &= \sum_x P(X) \sum_Y P(Y|X) \log \frac{1}{P(Y|X)} \\
 &= \sum_{X,Y} P(X,Y) \log \frac{1}{P(Y|X)} \\
 &= -E[\log P(Y|X)]
 \end{aligned}$$

The average uncertainty that remains about  $X$  when  $Y$  is known.

# Conditional Entropy

- The **conditional entropy** of  $Y$  given variable  $X$ :

$$\begin{aligned}
 H(Y|X) &= \sum_x P(X=x)H(Y|X=x) \\
 &= \sum_x P(X) \sum_Y P(Y|X) \log \frac{1}{P(Y|X)} \\
 &= \sum_{X,Y} P(X,Y) \log \frac{1}{P(Y|X)} \\
 &= -E[\log P(Y|X)]
 \end{aligned}$$

The average uncertainty that remains about  $X$  when  $Y$  is known.

# Outline

- 1 Jensen's Inequality
- 2 Entropy
- 3 Divergence**
- 4 Mutual Information

# Kullback-Leibler divergence

- **Relative entropy** or **Kullback-Leibler divergence**

# Kullback-Leibler divergence

- **Relative entropy** or **Kullback-Leibler divergence**
  - Measures how much a distribution  $Q(X)$  differs from a "true" probability distribution  $P(X)$ .



# Kullback-Leibler divergence

## ■ **Relative entropy** or **Kullback-Leibler divergence**

- Measures how much a distribution  $Q(X)$  differs from a "true" probability distribution  $P(X)$ .
- **K-L divergence** of  $Q$  from  $P$  is defined as follows:

$$KL(P||Q) = \sum_x P(X) \log \frac{P(X)}{Q(X)}$$

# Kullback-Leibler divergence

## ■ Relative entropy or Kullback-Leibler divergence

- Measures how much a distribution  $Q(X)$  differs from a "true" probability distribution  $P(X)$ .
- **K-L divergence** of  $Q$  from  $P$  is defined as follows:

$$KL(P||Q) = \sum_X P(X) \log \frac{P(X)}{Q(X)}$$

$$0 \log \frac{0}{0} = 0 \text{ and } p \log \frac{p}{0} = \infty \text{ if } p \neq 0$$

# Kullback-Leibler divergence

## ■ Relative entropy or Kullback-Leibler divergence

- Measures how much a distribution  $Q(X)$  differs from a "true" probability distribution  $P(X)$ .
- **K-L divergence** of  $Q$  from  $P$  is defined as follows:

$$KL(P||Q) = \sum_X P(X) \log \frac{P(X)}{Q(X)}$$

$$0 \log \frac{0}{0} = 0 \text{ and } p \log \frac{p}{0} = \infty \text{ if } p \neq 0$$

# Kullback-Leibler divergence

## ■ Relative entropy or Kullback-Leibler divergence

- Measures how much a distribution  $Q(X)$  differs from a "true" probability distribution  $P(X)$ .
- **K-L divergence** of  $Q$  from  $P$  is defined as follows:

$$KL(P||Q) = \sum_X P(X) \log \frac{P(X)}{Q(X)}$$

$$0 \log \frac{0}{0} = 0 \text{ and } p \log \frac{p}{0} = \infty \text{ if } p \neq 0$$

# Kullback-Leibler divergence

Theorem (1.2)

(**Gibbs' inequality**)

$$KL(P, Q) \geq 0$$

*with equality holds iff  $P$  is identical to  $Q$*

# Kullback-Leibler divergence

## Theorem (1.2)

(**Gibbs' inequality**)

$$KL(P, Q) \geq 0$$

*with equality holds iff  $P$  is identical to  $Q$*

**Proof:**

$$\sum_X P(X) \log \frac{P(X)}{Q(X)}$$

# Kullback-Leibler divergence

## Theorem (1.2)

(**Gibbs' inequality**)

$$KL(P, Q) \geq 0$$

*with equality holds iff  $P$  is identical to  $Q$*

**Proof:**

$$\sum_X P(X) \log \frac{P(X)}{Q(X)} = - \sum_X P(X) \log \frac{Q(X)}{P(X)}$$

# Kullback-Leibler divergence

## Theorem (1.2)

(**Gibbs' inequality**)

$$KL(P, Q) \geq 0$$

with equality holds iff  $P$  is identical to  $Q$

**Proof:**

$$\begin{aligned} \sum_X P(X) \log \frac{P(X)}{Q(X)} &= - \sum_X P(X) \log \frac{Q(X)}{P(X)} \\ &\geq - \log \sum_X P(X) \frac{Q(X)}{P(X)} \end{aligned}$$



# Kullback-Leibler divergence

## Theorem (1.2)

(**Gibbs' inequality**)

$$KL(P, Q) \geq 0$$

with equality holds iff  $P$  is identical to  $Q$

**Proof:**

$$\begin{aligned} \sum_X P(X) \log \frac{P(X)}{Q(X)} &= - \sum_X P(X) \log \frac{Q(X)}{P(X)} \\ &\geq - \log \sum_X P(X) \frac{Q(X)}{P(X)} \quad \text{Jensen's inequality} \end{aligned}$$

# Kullback-Leibler divergence

## Theorem (1.2)

(**Gibbs' inequality**)

$$KL(P, Q) \geq 0$$

with equality holds iff  $P$  is identical to  $Q$

**Proof:**

$$\begin{aligned} \sum_X P(X) \log \frac{P(X)}{Q(X)} &= - \sum_X P(X) \log \frac{Q(X)}{P(X)} \\ &\geq - \log \sum_X P(X) \frac{Q(X)}{P(X)} && \text{Jensen's inequality} \\ &= - \log \sum_X Q(X) = 0. \end{aligned}$$

# Kullback-Leibler divergence

## Theorem (1.2)

(**Gibbs' inequality**)

$$KL(P, Q) \geq 0$$

with equality holds iff  $P$  is identical to  $Q$

**Proof:**

$$\begin{aligned} \sum_X P(X) \log \frac{P(X)}{Q(X)} &= - \sum_X P(X) \log \frac{Q(X)}{P(X)} \\ &\geq - \log \sum_X P(X) \frac{Q(X)}{P(X)} && \text{Jensen's inequality} \\ &= - \log \sum_X Q(X) = 0. \end{aligned}$$

KL divergence between  $P$  and  $Q$  is larger than 0 unless  $P$  and  $Q$  are identical.

# Cross Entropy

- Entropy:  $H(P) = \sum_x P(X) \log \frac{1}{P(X)} = -E[\log P(x)]$

# Cross Entropy

- Entropy:  $H(P) = \sum_X P(X) \log \frac{1}{P(X)} = -E[\log P(x)]$

- **Cross entropy:**

$$H(P, Q) = \sum_X P(X) \log \frac{1}{Q(X)} = -E_P[\log Q(X)]$$

- Relationship with KL:

$$KL(P||Q) = \sum_X P(X) \log \frac{P(X)}{Q(X)}$$

# Cross Entropy

■ Entropy:  $H(P) = \sum_X P(X) \log \frac{1}{P(X)} = -E[\log P(x)]$

■ **Cross entropy:**

$$H(P, Q) = \sum_X P(X) \log \frac{1}{Q(X)} = -E_P[\log Q(X)]$$

■ Relationship with KL:

$$KL(P||Q) = \sum_X P(X) \log \frac{P(X)}{Q(X)} = E_P[\log P(X)] - E_P[\log Q(X)]$$

# Cross Entropy

■ Entropy:  $H(P) = \sum_X P(X) \log \frac{1}{P(X)} = -E[\log P(x)]$

■ **Cross entropy:**

$$H(P, Q) = \sum_X P(X) \log \frac{1}{Q(X)} = -E_P[\log Q(X)]$$

■ Relationship with KL:

$$\begin{aligned} KL(P||Q) &= \sum_X P(X) \log \frac{P(X)}{Q(X)} = E_P[\log P(X)] - E_P[\log Q(X)] \\ &= H(P, Q) - H(P) \end{aligned}$$

Or,

$$H(P, Q) = KL(P||Q) + H(P)$$

# A corollary

## Corollary (1.1)

### (Gibbs Inequality)

$$H(P, Q) \geq H(P), \text{ or}$$



# A corollary

## Corollary (1.1)

### (Gibbs Inequality)

$$\begin{aligned} H(P, Q) &\geq H(P), \text{ or} \\ \sum_x P(X) \log Q(X) &\leq \sum_x P(X) \log P(X) \end{aligned}$$

# A corollary

## Corollary (1.1)

### (Gibbs Inequality)

$$\begin{aligned} H(P, Q) &\geq H(P), \text{ or} \\ \sum_X P(X) \log Q(X) &\leq \sum_X P(X) \log P(X) \end{aligned}$$

In general, let  $f(X)$  be a non-negative function. Then

$$\sum_X f(X) \log Q(X) \leq \sum_X f(X) \log P^*(X)$$

where  $P^*(X) = f(X) / \sum_X f(X)$ .

# Unsupervised Learning

- Unknown true distribution  $P(\mathbf{x})$ .

$$P(\mathbf{x}) \xrightarrow{\text{sampling}} \mathcal{D} = \{\mathbf{x}_i\}_{i=1}^N \xrightarrow{\text{learning}} Q(\mathbf{x})$$

# Unsupervised Learning

- Unknown true distribution  $P(\mathbf{x})$ .

$$P(\mathbf{x}) \xrightarrow{\text{sampling}} \mathcal{D} = \{\mathbf{x}_i\}_{i=1}^N \xrightarrow{\text{learning}} Q(\mathbf{x})$$

- Objective:
  - **Minimizing KL:**  $KL(P||Q)$

# Unsupervised Learning

- Unknown true distribution  $P(\mathbf{x})$ .

$$P(\mathbf{x}) \xrightarrow{\text{sampling}} \mathcal{D} = \{\mathbf{x}_i\}_{i=1}^N \xrightarrow{\text{learning}} Q(\mathbf{x})$$

- Objective:

- **Minimizing KL:**  $KL(P||Q)$
- Same as **minimizing cross entropy:**  $H(P, Q)$

# Unsupervised Learning

- Unknown true distribution  $P(\mathbf{x})$ .

$$P(\mathbf{x}) \xrightarrow{\text{sampling}} \mathcal{D} = \{\mathbf{x}_i\}_{i=1}^N \xrightarrow{\text{learning}} Q(\mathbf{x})$$

- Objective:

- **Minimizing KL:**  $KL(P||Q)$
- Same as **minimizing cross entropy:**  $H(P, Q)$
- Approximating the cross entropy using data:

$$H(P, Q) = - \int P(\mathbf{x}) \log Q(\mathbf{x}) d\mathbf{x}$$

# Unsupervised Learning

- Unknown true distribution  $P(\mathbf{x})$ .

$$P(\mathbf{x}) \xrightarrow{\text{sampling}} \mathcal{D} = \{\mathbf{x}_i\}_{i=1}^N \xrightarrow{\text{learning}} Q(\mathbf{x})$$

- Objective:

- **Minimizing KL:**  $KL(P||Q)$
- Same as **minimizing cross entropy:**  $H(P, Q)$
- Approximating the cross entropy using data:

$$\begin{aligned} H(P, Q) &= - \int P(\mathbf{x}) \log Q(\mathbf{x}) d\mathbf{x} \\ &\approx -\frac{1}{N} \sum_{i=1}^N \log Q(\mathbf{x}_i) \end{aligned}$$

# Unsupervised Learning

- Unknown true distribution  $P(\mathbf{x})$ .

$$P(\mathbf{x}) \xrightarrow{\text{sampling}} \mathcal{D} = \{\mathbf{x}_i\}_{i=1}^N \xrightarrow{\text{learning}} Q(\mathbf{x})$$

- Objective:

- **Minimizing KL:**  $KL(P||Q)$
- Same as **minimizing cross entropy:**  $H(P, Q)$
- Approximating the cross entropy using data:

$$\begin{aligned} H(P, Q) &= - \int P(\mathbf{x}) \log Q(\mathbf{x}) d\mathbf{x} \\ &\approx -\frac{1}{N} \sum_{i=1}^N \log Q(\mathbf{x}_i) \\ &= -\frac{1}{N} \log Q(\mathcal{D}) \end{aligned}$$

- Same as **maximizing likelihood:**  $\log Q(\mathcal{D})$ .



# Supervised Learning

- Unknown true distribution  $P(\mathbf{x}, y)$ , where  $y$  is **label** of input  $\mathbf{x}$ .

$$P(\mathbf{x}, y) \xrightarrow{\text{sampling}} \mathcal{D} = \{\mathbf{x}_i, y_i\}_{i=1}^N \xrightarrow{\text{learning}} Q(y|\mathbf{x})$$

# Supervised Learning

- Unknown true distribution  $P(\mathbf{x}, y)$ , where  $y$  is **label** of input  $\mathbf{x}$ .

$$P(\mathbf{x}, y) \xrightarrow{\text{sampling}} \mathcal{D} = \{\mathbf{x}_i, y_i\}_{i=1}^N \xrightarrow{\text{learning}} Q(y|\mathbf{x})$$

- Objective:
  - **Minimizing cross (conditional) entropy:**

$$H(P, Q) = - \int P(\mathbf{x}, y) \log Q(y|\mathbf{x}) d\mathbf{x} dy$$

# Supervised Learning

- Unknown true distribution  $P(\mathbf{x}, y)$ , where  $y$  is **label** of input  $\mathbf{x}$ .

$$P(\mathbf{x}, y) \xrightarrow{\text{sampling}} \mathcal{D} = \{\mathbf{x}_i, y_i\}_{i=1}^N \xrightarrow{\text{learning}} Q(y|\mathbf{x})$$

- Objective:
  - **Minimizing cross (conditional) entropy:**

$$\begin{aligned} H(P, Q) &= - \int P(\mathbf{x}, y) \log Q(y|\mathbf{x}) d\mathbf{x} dy \\ &\approx - \frac{1}{N} \sum_{i=1}^N \log Q(y_i|\mathbf{x}_i) \end{aligned}$$

# Supervised Learning

- Unknown true distribution  $P(\mathbf{x}, y)$ , where  $y$  is **label** of input  $\mathbf{x}$ .

$$P(\mathbf{x}, y) \xrightarrow{\text{sampling}} \mathcal{D} = \{\mathbf{x}_i, y_i\}_{i=1}^N \xrightarrow{\text{learning}} Q(y|\mathbf{x})$$

- Objective:

- **Minimizing cross (conditional) entropy:**

$$\begin{aligned} H(P, Q) &= - \int P(\mathbf{x}, y) \log Q(y|\mathbf{x}) d\mathbf{x} dy \\ &\approx - \frac{1}{N} \sum_{i=1}^N \log Q(y_i|\mathbf{x}_i) \end{aligned}$$

- Same as **maximizing loglikelihood**:  $\sum_{i=1}^N \log Q(y_i|\mathbf{x}_i)$ ,
- Or **minimizing the negative loglikelihood (NLL)**:  
 $-\sum_{i=1}^N \log Q(y_i|\mathbf{x}_i)$

# Jensen-Shannon divergence

- KL is not symmetric:  $KL(P||Q)$  usually is not equal to reverse KL  $KL(Q||P)$ .

# Jensen-Shannon divergence

- KL is not symmetric:  $KL(P||Q)$  usually is not equal to reverse KL  $KL(Q||P)$ .
- **Jensen-Shannon divergence** is one symmetrized version of KL:

$$JS(P||Q) = \frac{1}{2}KL(P||M) + \frac{1}{2}KL(Q||M)$$

where  $M = \frac{P+Q}{2}$

# Jensen-Shannon divergence

- KL is not symmetric:  $KL(P||Q)$  usually is not equal to reverse KL  $KL(Q||P)$ .
- **Jensen-Shannon divergence** is one symmetrized version of KL:

$$JS(P||Q) = \frac{1}{2}KL(P||M) + \frac{1}{2}KL(Q||M)$$

where  $M = \frac{P+Q}{2}$

- Properties:
  - $0 \leq JS(P||Q) \leq \log 2$
  - $JS(P||Q) = 0$  if  $P = Q$
  - $JS(P||Q) = \log 2$  if  $P$  and  $Q$  has disjoint support.

# Outline

- 1 Jensen's Inequality
- 2 Entropy
- 3 Divergence
- 4 Mutual Information**



# Mutual information

- The **mutual information** of  $X$  and  $Y$ :

# Mutual information

- The **mutual information** of  $X$  and  $Y$ :

$$I(X; Y) = H(X) - H(X|Y)$$

# Mutual information

- The **mutual information** of  $X$  and  $Y$ :

$$I(X; Y) = H(X) - H(X|Y)$$

- Average reduction in uncertainty about  $X$  from learning the value of  $Y$ ,

# Mutual information

- The **mutual information** of  $X$  and  $Y$ :

$$I(X; Y) = H(X) - H(X|Y)$$

- Average reduction in uncertainty about  $X$  from learning the value of  $Y$ , or
- Average amount of information  $Y$  conveys about  $X$ .

# Mutual information and KL Divergence

- Note that:

$$I(X; Y) = \sum_X P(X) \log \frac{1}{P(X)} - \sum_{X,Y} P(X, Y) \log \frac{1}{P(X|Y)}$$

# Mutual information and KL Divergence

■ Note that:

$$\begin{aligned} I(X; Y) &= \sum_X P(X) \log \frac{1}{P(X)} - \sum_{X,Y} P(X, Y) \log \frac{1}{P(X|Y)} \\ &= \sum_{X,Y} P(X, Y) \log \frac{1}{P(X)} - \sum_{X,Y} P(X, Y) \log \frac{1}{P(X|Y)} \end{aligned}$$

# Mutual information and KL Divergence

■ Note that:

$$\begin{aligned} I(X; Y) &= \sum_X P(X) \log \frac{1}{P(X)} - \sum_{X,Y} P(X, Y) \log \frac{1}{P(X|Y)} \\ &= \sum_{X,Y} P(X, Y) \log \frac{1}{P(X)} - \sum_{X,Y} P(X, Y) \log \frac{1}{P(X|Y)} \\ &= \sum_{X,Y} P(X, Y) \log \frac{P(X|Y)}{P(X)} \end{aligned}$$

# Mutual information and KL Divergence

■ Note that:

$$\begin{aligned}
 I(X; Y) &= \sum_X P(X) \log \frac{1}{P(X)} - \sum_{X,Y} P(X, Y) \log \frac{1}{P(X|Y)} \\
 &= \sum_{X,Y} P(X, Y) \log \frac{1}{P(X)} - \sum_{X,Y} P(X, Y) \log \frac{1}{P(X|Y)} \\
 &= \sum_{X,Y} P(X, Y) \log \frac{P(X|Y)}{P(X)} \\
 &= \sum_{X,Y} P(X, Y) \log \frac{P(X, Y)}{P(X)P(Y)} \quad \text{equivalent definition}
 \end{aligned}$$



# Mutual information and KL Divergence

■ Note that:

$$\begin{aligned}
 I(X; Y) &= \sum_X P(X) \log \frac{1}{P(X)} - \sum_{X,Y} P(X, Y) \log \frac{1}{P(X|Y)} \\
 &= \sum_{X,Y} P(X, Y) \log \frac{1}{P(X)} - \sum_{X,Y} P(X, Y) \log \frac{1}{P(X|Y)} \\
 &= \sum_{X,Y} P(X, Y) \log \frac{P(X|Y)}{P(X)} \\
 &= \sum_{X,Y} P(X, Y) \log \frac{P(X, Y)}{P(X)P(Y)} \quad \text{equivalent definition} \\
 &= KL(P(X, Y) || P(X)P(Y))
 \end{aligned}$$

■ Due to equivalent definition:

# Mutual information and KL Divergence

- Note that:

$$\begin{aligned}
 I(X; Y) &= \sum_X P(X) \log \frac{1}{P(X)} - \sum_{X,Y} P(X, Y) \log \frac{1}{P(X|Y)} \\
 &= \sum_{X,Y} P(X, Y) \log \frac{1}{P(X)} - \sum_{X,Y} P(X, Y) \log \frac{1}{P(X|Y)} \\
 &= \sum_{X,Y} P(X, Y) \log \frac{P(X|Y)}{P(X)} \\
 &= \sum_{X,Y} P(X, Y) \log \frac{P(X, Y)}{P(X)P(Y)} \quad \text{equivalent definition} \\
 &= KL(P(X, Y) || P(X)P(Y))
 \end{aligned}$$

- Due to equivalent definition:

$$I(X; Y) = H(X) - H(X|Y) = I(Y; X) = H(Y) - H(Y|X)$$

# Property of Mutual information

## Theorem (1.3)

$$I(X; Y) \geq 0$$

*with equality holds iff  $X \perp Y$ .*

# Property of Mutual information

## Theorem (1.3)

$$I(X; Y) \geq 0$$

*with equality holds iff  $X \perp Y$ .*

Interpretation:  $X$  and  $Y$  are independent iff  $X$  contains no information about  $Y$  and vice versa.

# Property of Mutual information

## Theorem (1.3)

$$I(X; Y) \geq 0$$

*with equality holds iff  $X \perp Y$ .*

Interpretation:  $X$  and  $Y$  are independent iff  $X$  contains no information about  $Y$  and vice versa.

**Proof:** Follows from previous slide and Theorem 1.2.

# Conditional Entropy Revisited

## Theorem (1.4)

$H(X|Y) \leq H(X)$  with equality holds iff  $X \perp Y$

# Conditional Entropy Revisited

## Theorem (1.4)

$H(X|Y) \leq H(X)$  with equality holds iff  $X \perp Y$

Observation reduces uncertainty in average except for the case of independence.

# Conditional Entropy Revisited

## Theorem (1.4)

$H(X|Y) \leq H(X)$  with equality holds iff  $X \perp Y$

Observation reduces uncertainty in average except for the case of independence.

**Proof:** Follows from Theorem 1.3.



# Mutual information and Entropy

- From definition of mutual information

$$I(X; Y) = H(X) - H(X|Y)$$

# Mutual information and Entropy

- From definition of mutual information

$$I(X; Y) = H(X) - H(X|Y)$$

and the chain rule,

$$H(X, Y) = H(Y) + H(X|Y)$$

# Mutual information and Entropy

- From definition of mutual information

$$I(X; Y) = H(X) - H(X|Y)$$

and the chain rule,

$$H(X, Y) = H(Y) + H(X|Y)$$

we get

# Mutual information and Entropy

- From definition of mutual information

$$I(X; Y) = H(X) - H(X|Y)$$

and the chain rule,

$$H(X, Y) = H(Y) + H(X|Y)$$

we get

$$H(X) + H(Y) = H(X, Y) + I(X; Y)$$

# Mutual information and Entropy

- From definition of mutual information

$$I(X; Y) = H(X) - H(X|Y)$$

and the chain rule,

$$H(X, Y) = H(Y) + H(X|Y)$$

we get

$$H(X) + H(Y) = H(X, Y) + I(X; Y)$$

$$I(X; Y) = H(X) + H(Y) - H(X, Y)$$

# Mutual information and Entropy

- From definition of mutual information

$$I(X; Y) = H(X) - H(X|Y)$$

and the chain rule,

$$H(X, Y) = H(Y) + H(X|Y)$$

we get

$$H(X) + H(Y) = H(X, Y) + I(X; Y)$$


$$I(X; Y) = H(X) + H(Y) - H(X, Y)$$

- Consequently

- $H(X, Y) \leq H(X) + H(Y)$  with equality holds iff  $X \perp Y$ .

# Mutual information and entropy

Venn Diagram: Relationships among joint entropy, conditional entropy, and mutual information



$$H(X) + H(Y) = H(X, Y) + I(X; Y)$$

$$I(X; Y) = H(X) - H(X|Y)$$

$$I(Y; X) = H(Y) - H(Y|X)$$

# Conditional Mutual information

- The **conditional mutual information** of  $X$  and  $Y$  given  $Z$ :



# Conditional Mutual information

- The **conditional mutual information** of  $X$  and  $Y$  given  $Z$ :

$$I(X; Y|Z) = H(X|Z) - H(X|Y, Z)$$

# Conditional Mutual information

- The **conditional mutual information** of  $X$  and  $Y$  given  $Z$ :

$$I(X; Y|Z) = H(X|Z) - H(X|Y, Z)$$

- Average amount of information  $Y$  conveys about  $X$  given  $Z$ .

# Conditional mutual information and KL Divergence

Note:

$$I(X; Y|Z) = \sum_{X,Z} P(X, Z) \log \frac{1}{P(X|Z)} - \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Y, Z)}$$

# Conditional mutual information and KL Divergence

Note:

$$\begin{aligned} I(X; Y|Z) &= \sum_{X,Z} P(X, Z) \log \frac{1}{P(X|Z)} - \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Y, Z)} \\ &= \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Z)} - \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Y, Z)} \end{aligned}$$

# Conditional mutual information and KL Divergence

Note:

$$\begin{aligned}
 I(X; Y|Z) &= \sum_{X,Z} P(X, Z) \log \frac{1}{P(X|Z)} - \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Y, Z)} \\
 &= \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Z)} - \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Y, Z)} \\
 &= \sum_{X,Y,Z} P(X, Y, Z) \log \frac{P(X|Y, Z)}{P(X|Z)} \quad \text{equivalent definition}
 \end{aligned}$$

# Conditional mutual information and KL Divergence

Note:

$$\begin{aligned}
 I(X; Y|Z) &= \sum_{X,Z} P(X, Z) \log \frac{1}{P(X|Z)} - \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Y, Z)} \\
 &= \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Z)} - \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Y, Z)} \\
 &= \sum_{X,Y,Z} P(X, Y, Z) \log \frac{P(X|Y, Z)}{P(X|Z)} \quad \text{equivalent definition} \\
 &= \sum_Z P(Z) \sum_{X,Y} P(X, Y|Z) \log \frac{P(X, Y|Z)}{P(X|Z)P(Y|Z)}
 \end{aligned}$$

# Conditional mutual information and KL Divergence

Note:

$$\begin{aligned}
 I(X; Y|Z) &= \sum_{X,Z} P(X, Z) \log \frac{1}{P(X|Z)} - \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Y, Z)} \\
 &= \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Z)} - \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Y, Z)} \\
 &= \sum_{X,Y,Z} P(X, Y, Z) \log \frac{P(X|Y, Z)}{P(X|Z)} \quad \text{equivalent definition} \\
 &= \sum_Z P(Z) \sum_{X,Y} P(X, Y|Z) \log \frac{P(X, Y|Z)}{P(X|Z)P(Y|Z)} \\
 &= \sum_Z P(Z) KL(P(X, Y|Z), P(X|Z)P(Y|Z))
 \end{aligned}$$

# Conditional mutual information and KL Divergence

Note:

$$\begin{aligned}
 I(X; Y|Z) &= \sum_{X,Z} P(X, Z) \log \frac{1}{P(X|Z)} - \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Y, Z)} \\
 &= \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Z)} - \sum_{X,Y,Z} P(X, Y, Z) \log \frac{1}{P(X|Y, Z)} \\
 &= \sum_{X,Y,Z} P(X, Y, Z) \log \frac{P(X|Y, Z)}{P(X|Z)} \quad \text{equivalent definition} \\
 &= \sum_Z P(Z) \sum_{X,Y} P(X, Y|Z) \log \frac{P(X, Y|Z)}{P(X|Z)P(Y|Z)} \\
 &= \sum_Z P(Z) KL(P(X, Y|Z), P(X|Z)P(Y|Z)) \geq 0.
 \end{aligned}$$



# Property of conditional mutual information

## Theorem (1.5)

$$I(X; Y|Z) \geq 0$$

$$H(X|Z) \geq H(X|Y, Z)$$

*with equality hold iff  $X \perp Y|Z$ .*

Interpretation:

- More observations reduce uncertainty on average except for the case of conditional independence.

# Property of conditional mutual information

## Theorem (1.5)

$$I(X; Y|Z) \geq 0$$

$$H(X|Z) \geq H(X|Y, Z)$$

*with equality hold iff  $X \perp Y|Z$ .*

Interpretation:

- More observations reduce uncertainty on average except for the case of conditional independence.
- $X$  and  $Y$  are independently given  $Z$

# Property of conditional mutual information

## Theorem (1.5)

$$I(X; Y|Z) \geq 0$$

$$H(X|Z) \geq H(X|Y, Z)$$

with equality hold iff  $X \perp Y|Z$ .

Interpretation:

- More observations reduce uncertainty on average except for the case of conditional independence.
- $X$  and  $Y$  are independently given  $Z$  iff  $X$  contain no information about  $Y$  given  $Z$  and vice versa:

$$X \perp Y|Z \equiv I(X; Y|Z) = 0.$$

# Property of conditional mutual information

## Theorem (1.5)

$$I(X; Y|Z) \geq 0$$

$$H(X|Z) \geq H(X|Y, Z)$$

with equality hold iff  $X \perp Y|Z$ .

Interpretation:

- More observations reduce uncertainty on average except for the case of conditional independence.
- $X$  and  $Y$  are independently given  $Z$  iff  $X$  contain no information about  $Y$  given  $Z$  and vice versa:

$$X \perp Y|Z \equiv I(X; Y|Z) = 0.$$

Another characterization of conditional independence.