

Assignment-based Subjective Questions

1. From your analysis of the categorical variables from the dataset, what could you infer about their effect on the dependent variable?

2.

Analysis of the categorical variables from shows that bike rental rates are likely to be higher in

- Demand is higher summer and the fall season especially when the weather is Clear and pleasant.
- Rental is higher during months of June, July, August, September, and October
- Sat and Wed has the highest demand.
- Bikes have increased from 2018 to 2019.
- Additionally, we could confirm that bike rental is higher on holidays especially for Casual users.

3. Why is it important to use `drop_first=True` during dummy variable creation?


Usage of `drop_first=True` when creating dummy variables is a common practice in machine learning to improve model stability, interpretability, and computational efficiency while avoiding issues related to multicollinearity. However, it's essential to be aware of the specific requirements of your problem and the algorithms you're using, as there may be cases where including all dummy variables is appropriate.

4. Looking at the pair-plot among the numerical variables, which one has the highest correlation with the target variable?

- The temp variable has the highest correlation with the target variable.
- Causal and Registered variables have a high correlation with target variable but since the summation of them make the target variable it is being ignored


5. How did you validate the assumptions of Linear Regression after building the model on the training set?

- Linear relationship between independent and dependent variables




Most the data points are around the Actual Vs Predicted Line

- Error distribution of residuals



No specific pattern is being observed which means that Error terms are independent of each other.

- Distribution of Error Terms



Model is distributed with mean zero which means it is observing the Linear Regression rules

6. Based on the final model, which are the top 3 features contributing significantly towards explaining the demand of the shared bikes?

The top 3 variables are:

- **Temp:** Temperature and Clear environment seems to prompt users to take on more biking activity
- **yr:** Business seems to be growing organically year on year.
- **season:** Summer and Fall has higher demand.

General Subjective Questions

1. Explain the linear regression algorithm in detail.

Linear regression is a fundamental supervised machine learning algorithm used for modeling the relationship between a dependent variable (also known as the target or outcome) and one or more independent variables (predictors or features) by fitting a linear equation to the observed data. The goal of linear regression is to find the best-fitting linear relationship that can be used for prediction and understanding the dependencies between variables. Linear regression explained in detail:

1. **Simple Linear Regression:**

Simple linear regression is used when there is only one independent variable that affects the dependent variable. The linear equation for simple linear regression is:

$$Y = \theta_0 + \theta_1 X + \epsilon$$

2. **Multiple Linear Regression:**

Multiple linear regression extends the concept to situations where there are multiple independent variables. The linear equation for multiple linear regression is:

$$Y = \theta_0 + \theta_1 X_1 + \theta_2 X_2 + \dots + \theta_p X_p + \epsilon$$

3. **Fitting the Model:**

The goal in linear regression is to estimate the values of $\theta_0, \theta_1, \theta_2, \dots, \theta_p$ that minimize the sum of squared errors between the predicted values (\hat{Y}) and the actual values of the dependent variable in the training data. This is typically done using the method of least squares, which minimizes the sum of the squared residuals (the differences between the predicted and actual values).

4. **Assumptions:**

- The relationship between the independent variables and the dependent variable is linear.
- The error terms (ϵ) are independent of each other.
- The variance of the error terms is constant across all levels of the independent variables.
- The error terms follow a normal distribution.
- Independent variables are not highly correlated with each other.

5. **Interpretation:**

Linear regression allows for the interpretation of the coefficients $\theta_0, \theta_1, \theta_2, \dots, \theta_p$. For example, θ_1 represents the change in the dependent variable Y for a one-unit change in X_1 , holding all other variables constant.

6. **Evaluation:**

To assess the performance of a linear regression model, various metrics can be used, such as Mean Squared Error (MSE), R-squared (R^2), and others, to measure how well the model fits the data and makes predictions.

7. **Predictions:**

Once the model is trained, it can be used to make predictions on new or unseen data by plugging in the values of the independent variables into the linear equation.

8. **Extensions:**

Linear regression has several extensions and variations, including ridge regression, lasso regression, and polynomial regression, which address issues like multicollinearity and allow for more flexible modeling.

Linear regression is a powerful and interpretable tool commonly used for tasks such as predicting house prices, analyzing the impact of variables on an outcome, and understanding relationships in data. However, it has its limitations, and its effectiveness depends on the assumptions being met and the nature of the data being modeled.

2. Explain the Anscombe's quartet in detail.

Anscombe's quartet is a famous example in statistics and data visualization created by Francis Anscombe in 1973 to demonstrate that summary statistics alone may not provide a complete understanding of the data. It consists of four datasets, each containing 11 data points, with two numerical variables (X and Y). What makes Anscombe's quartet remarkable is that these four datasets have nearly identical simple descriptive statistics (such as means, variances, and correlation coefficients) for X and Y, yet they have vastly different graphical representations and relationships between the variables.

It emphasizes the importance of data visualization and graphical exploration thus revealing hidden patterns, outliers, and relationships that might not be apparent from summary statistics alone. This quartet is often used in statistics and data science education to emphasize the importance of visualizing data before drawing conclusions or making decisions based on statistical summaries.

3. What is Pearson's R?

The Pearson correlation coefficient (PCC) is a correlation coefficient that measures linear correlation between two sets of data. It is the ratio between the covariance of two variables and the product of their standard deviations; thus, it is essentially a normalized measurement of the covariance, such that the result always has a value between -1 and 1.

An absolute value of exactly 1 implies that a linear equation describes the relationship between X and Y perfectly, with all data points lying on a line. The correlation sign is determined by the regression slope: a value of +1 implies that all data points lie on a line for which Y increases as X increases, and vice versa for -1. A value of 0 implies that there is no linear dependency between the variables.

the formula for ρ can also be written as

$$\rho_{X,Y} = \frac{E[XY] - E[X]E[Y]}{\sqrt{E[X^2] - (E[X])^2} \sqrt{E[Y^2] - (E[Y])^2}}.$$

4. What is scaling? Why is scaling performed? What is the difference between normalized scaling and standardized scaling?

Scaling is a technique performed in pre-processing during building a machine learning model to standardize the independent feature variables in the dataset in a fixed range. The dataset could have several features which are highly ranging between high magnitudes and units. If there is no scaling performed on this data, it leads to incorrect modelling as there will be some mismatch in the units of all the features involved in the model. The difference between normalization and standardization is that while normalization brings all the data points in a range between 0 and 1, standardization replaces the values with their Z scores.

5. You might have observed that sometimes the value of VIF is infinite. Why does this happen?

Reasons why the VIF might become infinite:

- Perfect multicollinearity: Perfect multicollinearity occurs when two or more independent variables in a regression model are perfectly correlated with each other. In this case, one variable can be exactly predicted by a linear combination of the others, leading to an R^2 value of 1 and an infinite VIF.
- Nearly perfect multicollinearity: Even if multicollinearity is not perfect but very high, the R^2 value can still approach 1, resulting in a very large VIF.
- Too small a sample size: When you have a small sample size relative to the number of independent variables in the model, it can lead to unstable estimates and high VIF values, including infinity.

6. What is a Q-Q plot? Explain the use and importance of a Q-Q plot in linear regression.

The quantile-quantile plot is a graphical method for determining whether two samples of data came from the same population or not. A q-q plot is a plot of the quantiles of the first data set against the quantiles of the second data set.

The importance of a Q-Q plot in linear regression lies in its ability to assess the normality assumption of residuals visually and quantitatively. If the Q-Q plot shows a relatively straight line, it provides evidence that the residuals are normally distributed, which is one of the key assumptions of linear regression. On the other hand, if the Q-Q plot shows significant deviations from a straight line, it suggests that the normality assumption may not hold, and you may need to consider transformations or other methods to address outliers in your data before drawing conclusions from your regression analysis.