


# Crop Physiology



AGRIMoon.Com

All About Agriculture...

## CROP PHYSIOLOGY

Author

**TNAU**



**AGRIMOOON.COM**

All About Agriculture...


# Index

Lecture	Lecture Name	Page No
<b>1</b>	Introduction - Importance of crop physiology in agriculture.	5-9
<b>2</b>	Role and significance of water - diffusion, imbibition, osmosis and its significance, plasmolysis.	10-20
<b>3</b>	Field capacity, Available soil water and permanent wilting point	21-27
<b>4</b>	Absorption of water – mode of water absorption – active and Passive absorption and factors affecting absorption	28-40
<b>5</b>	Translocation of solutes - phloem and xylem transport.	41-48
<b>6</b>	Transpiration - types - Steward's theory of mechanism - significance, factors affecting transpiration and guttation - antitranspirants.	49-62
<b>7</b>	Mineral nutrition - introduction - criteria of essentiality of elements - macro, secondary and micronutrients - soil less culture - sand and hydroponics.	63-67
<b>8</b>	Mechanism of uptake - physiological role of nutrients.	68-78
<b>9</b>	Foliar diagnosis - nutritional and physiological disorders - foliar nutrition-fertigation	79-92
<b>10</b>	Photosynthesis - requirements of photosynthesis - light, CO <sub>2</sub> , pigments and H <sub>2</sub> O.	93-110
<b>11</b>	Photosynthetic pathways - c <sub>3</sub> , c <sub>4</sub> and cam	111-126
<b>12</b>	Respiration - Glycolysis, TCA and Pentose Phosphate Pathway.	127-131
<b>13</b>	Krebs' cycle / citric acid cycle /tca cycle	132-144
<b>14</b>	Protein and fat synthesis	145-150
<b>15</b>	Photoperiodism - short day, long day and day neutral plants - phytochrome. Role of phytochrome in flowering and regulation of flowering.	151-156
<b>16</b>	Transmission of stimulus - theories of flowering.	157-164

<b>17</b>	Source sink relationship - yield components - harvest index and its importance	165-168
<b>18</b>	Plant growth	169-173
<b>19</b>	Growth analysis	174-179
<b>20</b>	Plant growth regulators	180-194
<b>21</b>	Practical application of plant growth regulators in crop productivity	195-197
<b>22</b>	Environmental stresses	198-211
<b>23</b>	Seed germination	212-215
<b>24</b>	Abscission and senescence	216-219
<b>25</b>	Global warming - physiological effects on crop Productivity	220-229

## 01. INTRODUCTION

The spectacular diversity of plant size and form is familiar to everyone. In nature all plants carry out similar physiological processes. As primary producers, plants convert solar energy to chemical energy. Being non motile, plants must grow toward light, and they must have efficient vascular systems for movement of water, mineral nutrients, and photosynthetic products throughout the plant body. Green land plants must also have mechanisms for avoiding desiccation.



**Figure 1. Principal Parts of a Vascular Plant**

The meaning of Plant Physiology refers to “the science of properties and functions in normal conditions”. The aim of the Plant Physiology has been described as early as the early 20th Century by the Russian Plant Physiologist, V.I. Palladin as : “Which is to gain a complete and thorough knowledge of all the Phenomena occurring in plants, to analyse complex life processes. So as to interpret them in terms of simpler one and reduce them finally to the principles of physics and chemistry”. Nevertheless, Noggle and fritz (1983) described the Plant Physiology as “the science concerned with processes and functions, the response of plants to changes in environment and the growth and development that results from responses


Crop physiology is concerned with the processes and functions of the crops at cellular, sub-cellular and whole plant levels in response to environmental variables and growth. In short, **physiology is the study of functional aspects of crop plants.**

### Cell

Plants are multicellular organisms composed of millions of cells with specialized functions. At maturity, such specialized cells may differ greatly from one another in their structures. However, all plant cells have the same basic eukaryotic organization: They contain a nucleus, a cytoplasm, and sub cellular organelles, and they are enclosed in a membrane that defines their boundaries.

In plants, cell migrations are prevented because each walled cell and its neighbor are cemented together by a **middle lamella**. As a consequence, plant development unlike animal development, depends solely on patterns of cell division and cell enlargement.



Plant cells have two types of walls: primary and secondary. **Primary cell walls** are typically thin and are characteristic of young, growing cells. **Secondary cell walls** are thicker and stronger than primary walls and are deposited when most cell enlargement has ended. Secondary cell walls owe their strength and toughness to **lignin**, a brittle, glue-like material. The evolution of lignified secondary cell walls provided plants with the structural reinforcement necessary to grow vertically above the soil and to colonize the land.



## Plant anatomy


There are two categories of seed plants, gymnosperms and angiosperms. **Gymnosperms** are the less advanced type. **Angiosperms**, the more advanced type of seed plant which dominate the landscape. About 250,000 species are known, but many more remain to be characterized. The major innovation of the angiosperms is the flower; hence they are referred to as **flowering plants**.

Three major tissue systems are found in **flowering plants**; in all plant organs contain dermal tissue, ground tissue, and vascular tissue. The vegetative body is composed of three organs: **leaf**, **stem**, and **root**. The primary function of a leaf is photosynthesis, that of the stem is support, and that of the root is anchorage and absorption of water and minerals. Leaves are attached to the stem at **nodes**, and the region of the stem between two nodes is termed the **internode**. The stem together with its leaves is commonly referred to as the **shoot**.



**Plant parts** (Source: Plant Physiology by Taiz and Zeiger)

Plant growth is concentrated in localized regions of cell division called **meristems**. Nearly all nuclear divisions (mitosis) and cell divisions (cytokinesis) occur in these meristematic regions. In a young plant, the most active meristems are called **apical meristems**; they are located at the tips of the stem and the root. At the nodes, **axillary buds** contain the apical meristems for branch shoots. Lateral roots arise from the **pericycle**, an internal meristematic tissue. Proximal and overlapping the meristematic regions are zones of cell elongation in which cells increase dramatically in length and width.



Cells usually differentiate into specialized types after they elongate. The phase of plant development that gives rise to new organs and to the basic plant form is called **primary growth**. Primary growth results from the activity of apical meristems, in which cell division is followed by progressive cell enlargement, typically elongation. After elongation in a given region is complete, **secondary growth** may occur. Secondary growth involves two lateral meristems: the **vascular cambium** (plural *cambia*) and the **cork cambium**. The vascular cambium gives rise to secondary xylem (wood) and secondary phloem. The cork cambium produces the periderm, consisting mainly of cork cells.

The architecture, mechanics, and function of plants depend crucially on the structure of the cell wall. The wall is secreted and assembled as a complex structure that varies in form and composition as the cell differentiates. Without a cell wall, plants would be very different organisms from what we know. Indeed, the plant cell wall is essential for many processes in plant growth, development, maintenance, and reproduction:

- Plant cell walls determine the mechanical strength of plant structures, allowing those structures to grow to great heights.
- Cell walls glue cells together, preventing them from sliding past one another. This constraint on cellular movement contrasts markedly to the situation in animal cells, and it dictates the way in which plants develop

- A tough outer coating enclosing the cell, the cell wall acts as a cellular "exoskeleton" that controls cell shape and allows high turgor pressures to develop.
- Plant morphogenesis depends largely on the control of cell wall properties because the expansive growth of plant cells is limited principally by the ability of the cell wall to expand.
- The cell wall is required for normal water relations of plants because the wall determines the relationship between the cell turgor pressure and cell volume
- The bulk flow of water in the xylem requires a mechanically tough wall that resists collapse by the negative pressure in the xylem.
- The wall acts as a diffusion barrier that limits the size of macromolecules that can reach the plasma membrane from outside, and it is a major structural barrier to pathogen invasion.

Much of the carbon that is assimilated in photosynthesis is channeled into polysaccharides in the wall. During specific phases of development, these polymers may be hydrolyzed into their constituent sugars, which may be scavenged by the cell and used to make new polymers


## 02. Role and significance of water

Water is said to be the liquid of life. Because, life is originated in organs, environmental and in the course of evolution it became fully dependent upon water in a number of ways. Water is one of the most plentiful chemicals available in the earth and the chemical formula is  $H_2O$ . It is a tiny V-shaped molecule contains three atoms do not stay together as the hydrogen atoms are constantly exchanging between water molecules


The water molecule consists of an oxygen atom covalently bonded to two hydrogen atoms. The two O—H bonds form an angle of  $105^\circ$  (Figure ). Because the oxygen atom is more **electronegative** than hydrogen, it tends to attract the electrons of the covalent bond. This attraction results in a partial negative charge at the oxygen end of the molecule and a partial positive charge at each hydrogen.

Water has special properties that enable it to act as a solvent and to be readily transported through the body of the plant. These properties derive primarily from the polar structure of the water molecule.

- The Polarity of water molecules gives rise to hydrogen bonds
- The Polarity of water makes an excellent solvent
- The Thermal properties of water result from hydrogen bonding
- The Cohesive and adhesive properties of water are due to hydrogen bonding



- **Solute:** type of molecule dissolved in another type of substance; that substance is called a...
- **Solvent:** substance that dissolves the solute



### **Importance of water to plants**

- Water typically constitutes 80 to 95% of the mass of growing plant tissues.
- Water is the main constituent of protoplasm comprising up to about 90-95 per cent of its total weight. In the absence of water, protoplasm becomes inactive and is even killed.
- Different organic constituents of plants such as carbohydrates proteins, nucleic acid and enzymes etc. Lose their physical and chemical properties in the absence of water.
- Water participates directly in many metabolic processes. Inter conversion of carbohydrates and organic acids depend upon hydrolysis and condensation reaction.
- Water increases the rate of respiration. Seeds respire fast in the presence of water.
- Water is the source of hydrogen atom for the reduction of  $\text{CO}_2$  in the reaction of photosynthesis.


## CONCENTRATION EXAMPLES

- High solute concentration: lots of sugar dissolved in a relatively small amount of water
- Low solute concentration: little sugar dissolved in a relatively high amount of water



- Water acts as a solvent and acts as a carrier for many substance. It forms the medium in which several reactions take place.
- Water present in the vacuoles helps in maintaining the turgidity of the cells which is a must for proper activities of life and to maintain this form and structure.
- Water helps in translocation of solutes
- In tropical plants, water plays a very important role of thermal regulation against high temperature.
- The elongation phase of cell growth depends on absorption of water.

### *“Dissociation” of water*



## Properties of water

1. Solvent for electrolyte & non electrolyte
2. High specific heat
3. High latent heat of vaporization (540 cal g-1)
4. Cohesive and Adhesive Properties
5. High surface tension
6. High Tensile Strength
7. Stabilizes temperature
8. Transparent to visible radiation
9. Low viscosity

### CONCENTRATION

- Concentration refers to how much of some substance is present, compared to another substance.
- For instance, a high solute concentration has a relatively high amount of solute and low amount of solvent.

### HOW IT HELPS IN PLANTS?


WATER PLAYS A CRUCIAL ROLE in the life of the plant. For every gram of organic matter made by the plant, approximately 500 g of water is absorbed by the roots, transported through the plant body and lost to the atmosphere. Even slight imbalances in this flow of water can cause water deficits and severe malfunctioning of many cellular processes. Thus, every plant must delicately balance its uptake and loss of water.

### REMEMBER!


- Solutes can be many different kinds of molecules (sugars, gases, nutrients, proteins, and lipids)
- Solvents can vary as well (solids, liquids, or gases), but are usually H<sub>2</sub>O

## Diffusion, osmosis and imbibitions

### Diffusion: molecules moving randomly



### Diffusion



Solute transport is from the left to the right; movement of the solutes is due to the concentration gradient ( $dC/dx$ ).

The movement of materials in and out of the cells in plants takes place in a solution or gaseous form. Although the exact process of this is not very clear, three physical processes are usually involved in it. They are diffusion, osmosis and imbibition.

The movement of particles or molecules from a region of higher concentrations to a region of lower concentration is called as diffusion. The rate of diffusion of gases is faster than liquids or solutes. The diffusion particles have a certain pressure called as the diffusion pressure which is directly proportional to the number as concentration of the diffusing particles. These forms the diffusion takes place always from a region of higher diffusion pressure to a region of lower diffusion pressure (i.e) along a diffusion pressure gradient. The rate of diffusion increases if,

- i) Diffusion pressure gradient is steeper
- ii) Temperature is increased

- iii) Density of the differing particles is lesser
- iv) Medium through which diffusion occurs is less concentrated.

Diffusion of more than one substance at the same time and place may be at different rates and in different direction, but is independent of each other. A very common example of this is the gaseous exchange in plants.

Beside osmotic diffusion the above mentioned simple diffusion also plays a very important role in the life of the plants.


- It is an essential step in the exchange of gases during respiration and photosynthesis
- During passive salt uptake, the ions are absorbed by diffusion
- It is important in stomatal transpiration as the last step in the pollen, where diffusion of water vapour from the interrelation space into the outer atmosphere occurs through open stomata.

### Osmosis

The diffusion of solvent molecules into the solution through a semi permeable membrane is called as osmosis (some times called as *Osmotic diffusion*). In case there are two solutions of different concentration separated by the semi permeable membrane, the diffusion of solvent will take place from the less concentrated suitable into the more concentrated solution till both the solutions attain equal concentration.


## What happens in osmosis

- ❖ Osmosis is the diffusion of water across selectively permeable membranes.




### Osmotic pressure

As a result of the separation of solution from its solvent (or) the two solutions by the semi permeable membrane, a pressure is developed in solution to the pressure by dissolved solutes in it. This is called as osmotic pressure (O.P). OP is measured in terms of atmospheres and is directly proportional to the concentration of dissolved solutes in the solution. More concentration solution has higher O.P. O.P of a solution is always higher than its pure solvent.



During osmosis, the movement of solvent molecules taken place from the solution whose osmotic pressure is lower (i.e less concentration as hypotonic) into the solution whose osmotic pressure is higher (i.e, more concentrated as hypertonic). Osmotic diffusion of solvent molecules will not take place if the two solutions separated by the semipermeable membrane are of equal concentration having equal *Osmotic pressures* (i.e., they are isotonic). In plant cells, plasma membrane and tonoplast act as selectively permeable or differentially permeable membrane.



### End-osmosis

If a living plant cell is placed in water or hypotonic solution whose O.P is lower than cell sap, water enters the cell sap by osmosis and the process is called end osmosis. As a result of entry of water with the cell sap, a pressure is developed which press the protoplasm against the cell wall and become turgid. This pressure is called a turgor pressure.

Consequence of the turgor pressure is the wall pressure which is exerted by the elastic cell wall against the expanding protoplasm. At a given time, turgor pressure (T.P) equals the wall pressure (W.P).

$$T.P = W.P$$

## Exosmosis

If on the other hand, the plant cell is placed in hypertonic solution (whose O.P is higher than cell sap) the water move out the cell sap into the outer solution and the cell becomes flaccid. This process is known as exosmosis. Cell (or) tissue will remain as such in isotonic solution.


## Significance of osmosis in plants

1. Large quantities of water are absorbed by roots from the soil by osmosis
2. Cell to cell movement of water and other substances dissolve is involves osmosis
3. Opening and closing of stomata depend upon the turgor pressure of guard cells
4. Due to osmosis, the turgidity of the cells and hence the shape or form of them organs is maintained.
5. The resistance of plants to drought and frost increases with increase in osmotic pressure to later cells
6. Turgidity of the cells of the young seedling allows them to come out of the soil.

**hypotonic - solution whose osmotic pressure is lower ( less concentration)**

**hypertonic - solution whose osmotic pressure is higher ( more concentration)**

**isotonic - diffusion of solvent molecules will not take place**



### Imbibition

Certain substances if placed in a particular liquid absorb it and swell up. For example, when some pieces of grass or dry wood or dry seeds are placed in water they absorb the water quickly and swell up considerably so that their volume is increased. These substances are called as imbibants and the phenomenon as imbibition, certain force of attraction is existing between imbibants and the involved substance. In plants, the hydrophilic colloids *viz.*, protein and carbohydrates such as starch, cellulose and pectic substance have strong alteration towards water.

Imbibition plays a very important role in the life of plants. The first step in the absorption of water by the roots of higher plants is the imbibition of water by the cell walls of the root hairs. Dry seeds require water by imbibition for germination.

As a result of imbibition, a pressure is developed which is called as imbibition pressure or matric potential ( $\psi_m$ ). It is analogous to the osmotic potential of a solution. With reference to pure water, the values of  $\psi_m$  are always negative. The water potential of an imbibant is equal to its matric potential plus any turgor or other pressure (pressure potential) which may be imposed upon the imbibant.

$$\psi_w = \psi_m + \psi_p$$


If the imbibant is unconfined to turgor or such pressure, the equation can be significant to

$$\psi_w = \psi_m$$

## Plasmolysis

When a plant cell or tissue is placed in a hypertonic solution water comes out from the cell sap into the outer solution of exosmosis and the protoplasm begins to shrink or contract. The protoplasm separates from the cell wall and assumes a spherical form and this phenomenon is called plasmolysis. Incipient plasmolysis is the stage where protoplasm begins to contract from the cell wall. If a plasmolysed cell in tissue is placed in water, the process of endosmosis takes place. Water enters into the cell sap, the cell becomes turgid and the protoplasm again assumes its normal shape and position. This phenomenon is called deplasmolysis.

### Diagrammatic view of normal plant cell and plasmolysed plant cell




### Advantages of plasmolysis

1. It indicates the semi-permeable nature of the plasma membrane.
2. It is used to determine the osmotic pressure of the cell sap.
3. Plasmolysis is used in salting of meat and fishes. Addition of concentrated sugar solution to jam and jellies checks the growth of fungi and bacteria which become plasmolysed in concentrated solution.

### 03. Field capacity, Available soil water and permanent wilting point


#### Field capacity or water holding capacity of the soil

After heavy rain fall or irrigation of the soil some water is drained off along the slopes while the rest percolates down in the soil. Out of this water, some amount of water gradually reaches the water table under the force of gravity (gravitational water) while the rest is retained by the soil. This amount of water retained by the soil is called as field capacity or water holding capacity of the soil.



Field capacity is affected by soil profiles soil structure and temperature. The effective depth of a soil, as determined by physical and chemical barriers, together with the clay content of the soil within that depth, determine the water holding capacity of the profile, and how much of the water is available to plants. Effective soil depth varies between plant species. Wheat is used as the benchmark plant in this assessment. Available water holding

capacity rankings are estimated from soil texture, structure and stone content within the potential root zone of a wheat plant.



Water-holding capacity is controlled primarily by soil texture and organic matter. Soils with smaller particles (silt and clay) have a larger surface area than those with larger sand particles, and a large surface area allows a soil to hold more water. In other words, a soil with a high percentage of silt and clay particles, which describes fine soil, has a higher water-holding capacity. The table illustrates water-holding-capacity differences as influenced by texture. Organic matter percentage also influences water-holding capacity. As the percentage increases, the water-holding capacity increases because of the affinity organic matter has for water.

It is the water content of the soil after downward drainage of gravitational water. It is the capillary capacity of a soil. It is the upper limit of soil water storage for the plant growth. At field capacity, the soil water potential is  $-0.1$  to  $-0.3$  bars.

Texture	Field Capacity	Wilting point	Available water
Coarse sand	0.6	0.2	0.4
Fine sand	1.0	0.4	0.6
Loamy sand	1.4	0.6	0.8
Sandy loam	2.0	0.8	1.2
Light sandy clay loam	2.3	1.0	1.3
Loam	2.7	1.2	1.5
Sandy clay loam	2.8	1.3	1.5
Clay loam	3.2	1.4	1.8
Clay	4.0	2.5	1.5
Self-mulching clay	4.5	2.5	2.0

## Water potential

Every component of a system possesses free energy capable of doing work under constant temperature conditions. For non-electrolytes, free energy / mole is known as chemical potential. With respect to water, the chemical potential of water is called as water potential. The chemical potential is denoted by a Greek letter Psi ( $\psi$ ).

For pure water, the water potential is Zero. The presence of solute particles will reduce the free energy of water or decrease the water potential. Therefore it is expressed in negative value.

It is therefore, water potential of solution is always less than zero so in negative value.

For solutions, water potential is determined by three internal factors i.e.

$$\psi_w = \psi_w + \psi_s + \psi_p$$

$\psi_s$  = is the solute potential or osmotic potential

$\psi_p$  = pressure potential or turgor potential

$\psi_w$  = is the matric potential. Matric potential can be measured for the water molecules adhering on the soil particles and cell wall.

In plant system, the matric potential is disregarded.

Therefore,

$$\psi_w = \psi_s + \psi_p$$

### Osmotic pressure

Osmotic pressure is equivalent to osmotic potential but opposite in sign. Osmotic pressure in a solution results due to the presence of solutes and the solutes lower the water potential. Therefore osmotic pressure is a quantitative index of the lowering of water potential in a solution and using thermodynamic terminology is called as osmotic potential.

Osmotic pressure and osmotic potential are numerically equal but opposite in sign.

Osmotic pressure has positive sign

Osmotic potential has negative sign ( $\psi_s$ )

For eg.

$$I_A OP = 20 \text{ atm.}$$

$$\psi_w = -20 \text{ atm}$$

### Turgor pressure

In plant cell, the turgor pressure results due to the presence of water molecules is turgor pressure. The potential created by such pressures is called pressure potential ( $\psi_p$ )

In a normal plant cell, the water potential

$$\psi_w = \psi_s + \psi_p \quad - \text{partially turgid cell}$$

(High)

$$\psi_w = \text{Zero} \quad - \text{Fully turgid cell}$$

(Highest)

$$\psi_w = \psi_s \quad - \text{Flaccid cell or plasmolysed cell}$$

(Lowest)

## Water relation

Water forms the major constituent of living (cells) things and the cells originated in a highly aqueous medium and all the vital processes of the life are carried out in it. Besides, water predominately acts as a source of hydrogen to plants and is released by the photolysis of water during photosynthesis.

In living tissue, water is the medium for many biochemical reactions and extraction process. Inorganic nutrients, photosynthesis, bases and hormones are all transported in aqueous solution. Evaporation of water can control the temperature of leaf on canopy soil nutrients are available to plant roots only when dissolved in water. In short, water is essential for life and plays a unique role in virtually all biological process.

### Example:

There are 2 cells A and B in contact with each other, cell A has a pressure potential (turgor pressure) of 4 bars and certain sap with an osmotic potential of -12 bars.

Cell B has pressure potential of 2 bars and certain sap with osmotic potential of -5 bars.

Then,

$$\begin{aligned}\psi_w \text{ of cell A} &= \psi_s + \psi_p \\ &= -12 + (+4) \\ &= -8 \text{ bars}\end{aligned}$$

$$\begin{aligned}\psi_w \text{ of cell B} &= -5 + (+2) \\ &= -3 \text{ bars}\end{aligned}$$

Hence, water will move from cell B to cell A (i.e., towards lower or more negative water potential) with a force of  $(-8 - (-3)) = -5$  bars.

### Diffusion Pressure Déficit (DPD) (Suction pressure)

Diffusion pressure of a solution is always lower than its pure solvent. The difference between the diffusion pressure of the solution and its solvent at a particular temperature and

atmosphere conditions is called as diffusion pressure deficit (D.P.D). If the solution is more concentrated D.P.D increases but it decreases with the dilution of the solution,

D.P.D of the cell sap or the cells is a measure of the ability of the cells to absorb water and hence is often called as the suction pressure (S.P). It is related with osmotic pressure (O.P) and turgor pressure (T.P) of cell sap and also the wall pressure (W.P) as follows.

$$\text{D.P.D. (S.P)} = \text{O.P} - \text{W.P}$$

But

$$(\text{W.P}) = \text{T.P}$$

$$\text{D.P.D} = \text{O.P} - \text{T.P}$$

Due to the entry of the water the osmotic pressure of the cell sap decreases while its turgor pressure is increased so much so that in a fully turgid cell T.P equals the O.P

$$\text{O.P} = \text{T.P} = \text{D.P.D} = \text{O}$$

$$\text{In fully plasmolysed cells: } \text{T.P} = \text{O}$$

$$\text{So } \text{D.P.D} = \text{O.P}$$

D.P.D. in case of plant cells is not directly proportional to their osmotic pressure or the concentration of the cell sap but depend both on O.P and T.P. Higher osmotic pressure of the cell sap is usually accompanied by lower turgor pressure so that its D.P.D is greater and water enters into it. But sometimes it is possible that two cells are in contact with each other one having higher O.P and also higher turgor pressure than the other cell and still its does not draw water. It is because of its lower D.P.D., no matter is O.P is higher.

Cell a

$$\text{O.P} = 25 \text{ atm.} \longrightarrow \text{O.P} = 30 \text{ atm}$$

$$\text{T.P} = 15 \text{ atm.}$$

$$\text{D.P.D} = 10 \text{ atm.}$$

Cell b

$$\text{T.P} = 10 \text{ atm.} \quad \text{A}$$

$$\text{D.P.D} = 30 \text{ atm.}$$

Cell a		Cell b
O.P = 35 atm.	←	O.P = 40 atm
T.P = 10 atm.		T.P = 20 atm.
D.P.D = 25 atm.		D.P.D = 20 atm.

Entry of water into the cell depends on D.P.D and not on O.P only

## ABSORPTION OF WATER – MODE OF WATER ABSORPTION – ACTIVE AND PASSIVE ABSORPTION AND FACTORS AFFECTING ABSORPTION.



### PRELIMINARY OF WATER POTENTIAL

Most organisms are comprised of at least 70% or more water. Some plants, like a head of lettuce, are made up of nearly 95% water. When organisms go dormant, they lose most of their water. For example, seeds and buds are typically less than 10% water, as are desiccated rotifers, nematodes and yeast cells. Earth is the water planet (that's why astronomers get so excited about finding water in space). Water is the limiting resource for crop productivity in most agricultural systems

### LEARN MORE ABOUT WATER POTENTIAL

- In general, water always moves down its water potential gradient from areas of higher water potential to areas of lower water potential.
- Water potential is typically measured as the amount of pressure needed to stop the movement of water.
- The unit used to express this pressure is the megapascal (MPa).

The three factors that most commonly determine water potential are

<p><b>1) Solute concentration</b></p> <p>As solute concentration increases, water potential decreases. This is why water diffuses from regions of lower total solute concentration...</p>  <p>...to regions of higher total solute concentration.</p>	<p><b>2) Pressure</b></p> <p>The flow of water is also affected by pressure.</p>  <p>positive pressure increases water potential...</p>  <p>...while negative pressure decreases it.</p>	<p><b>3) Gravity</b></p> <p>Water flows in response to gravity because water at higher elevations has more potential energy, and thus a higher water potential ...</p>  <p>...than water at lower elevations.</p>
--	--	--

### WHAT IS WATER POTENTIAL?

**Water potential** is the potential energy of water relative to pure free water (e.g. deionized water) in reference conditions. It quantifies the tendency of water to move from one area to another due to osmosis, gravity, mechanical pressure, or matrix effects including surface tension. Water potential is measured in units of pressure and

is commonly represented by the Greek letter  $\Psi$  (Psi). This concept has proved especially useful in understanding water movement within plants, animals, and soil.

### Components of water potential

Much different potential affect the total water potential and sum of these potentials determines the overall water potential and the direction of water flow:

$$\Psi = \Psi_0 + \Psi_s + \Psi_p + \Psi_g + \Psi_v + \Psi_m$$

Where:

- $\Psi_0$  is the reference correction,
- $\Psi_s$  is the solute potential,
- $\Psi_p$  is the pressure potential,
- $\Psi_g$  is the gravimetric component,
- $\Psi_v$  is the potential due to humidity, and
- $\Psi_m$  is the potential due to matrix effects (e.g., fluid cohesion and surface tension.)

## COMPONENT OF WATER POTENTIAL

### 1. Pressure potential

Pressure potential is based on mechanical pressure, and is an important component of the total water potential within plant cells. Pressure potential is increased as water enters a cell. As water passes through the cell wall and cell membrane, it increases the total amount of water present inside the cell, which exerts an outward pressure that is retained by the structural rigidity of the cell wall.

The pressure potential in a living plant cell is usually positive. In plasmolysed cells, pressure potential is almost zero. Negative pressure potentials occur when water is pulled through an open system such as a plant xylem vessel. Withstanding negative pressure potentials (frequently called tension) is an important adaptation of xylem vessels.

### 2.Solute potential

Pure water is usually defined as having a solute potential ( $\Psi_s$ ) of zero, and in this case, solute potential can never be positive. The relationship of solute concentration (in molarity) to solute potential is given by the van 't Hoff equation:

$$\Psi_s = -miRT$$

Where

$m$  - The concentration in molarity of the solute,

$i$  - The van 't Hoff factor, the ratio of amount of particles in solution to amount of formula units dissolved,

$R$  - The ideal gas constant, and  $T$  is the absolute temperature.

### **3. Matrix potential**

When water is in contact with solid particles (e.g., clay or sand particles within soil) adhesive intermolecular forces between the water and the solid can be large and important. The forces between the water molecules and the solid particles in combination with attraction among water molecules promote surface tension and the formation of menisci within the solid matrix. Force is then required to break these menisci. The magnitude of matrix potential depends on the distances between solid particles--the width of the menisci and the chemical composition of the solid matrix. In many cases, matrix potential can be quite large and comparable to the other components of water potential discussed above.

It is worth noting that matrix potentials are very important for plant water relations. Strong (very negative) matrix potentials bind water to soil particles within very dry soils. Plants then create even more negative matrix potentials within tiny pores in the cell walls of their leaves to extract water from the soil and allow physiological activity to continue through dry periods.

### **4. Gravity ( $\Psi_g$ ):**

Contributions due to gravity which is usually ignored unless referring to the tops of tall trees.

## **ABSORPTION OF WATER**

We know from a very early age that plants obtain water through their roots, though it is not perhaps until our school biology lessons that we learn of the important role that water plays in the process of photosynthesis. Most of the water absorption is carried out by the younger part of the roots. Just behind the growing tip of a young root is the piliferous region, made up of hundreds of projections of the epidermal tissue, the root hairs.


## **STRUCTURE INVOLVED IN WATER ABSORPTION**

In higher plants water is absorbed through root hairs which are in contact with soil water and form a root hair zone a little behind the root tips. Root hairs are tubular hair like prolongations of the cells of the epidermal layer (when epidermis bears root hairs it is also known as piliferous layer of the roots. The walls of root hairs are permeable and consist of pectic substances and cellulose which are strongly hydrophilic in nature root hairs contain vacuoles filled with cell sap. When roots elongate, the older root hairs die and new root hairs are developed so that they are in contact with fresh supplies of water in the soil. Lateral Movement of water is achieved through root. This can described as follows:

## ROOTS

Often roots are overlooked, probably because they are less visible than the rest of the plant. However, it's important to understand plant root systems (Fig 1) because they have a pronounced effect on a plant's size and vigor, method of propagation, adaptation to soil types, and response to cultural practices and irrigation.

Fig 1. Diagrammatically the internal structure of a typical root



Roots typically originate from the lower portion of a plant or cutting. They have a root cap, but lack nodes and never bear leaves or flowers directly. Their principal functions are to absorb nutrients and moisture, anchor the plant in the soil, support the stem, and store food. In some plants, they can be used for propagation.

## STRUCTURE OF ROOTS

Internally, there are three major parts of a root (Fig 2):

- The **meristem** is at the tip and manufactures new cells; it is an area of cell division and growth.
- Behind the meristem is the **zone of elongation**. In this area, cells increase in size through food and water absorption. As they grow, they push the root through the soil.
- The **zone of maturation** is directly beneath the stem. Here, cells become specific tissues such as epidermis, cortex, or vascular tissue.

A root's **epidermis** is its outermost layer of cells (Fig 2). These cells are responsible for absorbing water and minerals dissolved in water. **Cortex** cells are involved in moving water from the epidermis to the **vascular tissue** (xylem and phloem) and in storing food. Vascular tissue is located in the center of the root and conducts food and water.

Fig 2. Cross section of roots





Fig 3. Structure of root hair



Externally, there are two areas of importance: the root cap and the root hairs (Figure 3). The **root cap** is the root's outermost tip. It consists of cells that are sloughed off as the root grows through the soil. Its function is to protect the root meristem.

**Root hairs** are delicate, elongated epidermal cells that occur in a small zone just behind the root's growing tip. They generally appear as fine down to the naked eye. Their function is to increase the root's surface area and absorptive capacity. Root hairs usually live 1 or 2 days. When a plant is transplanted, they are easily torn off or may dry out in the sun.

## **WATER MOVEMENT MECHANISM IN PLANTS**


In plants, following two pathways are involved in the water movement. They are

- (1) Apoplastic pathway
- (2) Symplastic pathway
- (3) Transmembrane pathway

### **1. Apoplastic pathway (Fig 4)**

The apoplastic movement of water in plants occurs exclusively through the **cell wall** without crossing any membranes. The cortex receive majority of water through apoplastic way as loosely bound cortical cells do not offer any resistance. But the movement of water in root beyond cortex apoplastic pathway is blocked by casparyan strip present in the endodermis.

**Fig 4**



## 2. Symplastic pathway (Fig 5)

The movement of water from one cell to other cell through the **plasmodesmata** is called the symplastic pathway of water movement. This pathway comprises the network of cytoplasm of all cells inter-connected by plasmodesmata.





Fig 5

## 3. Transmembrane pathway (Fig 6)


In plant roots, water movement from soil till the endodermis occurs through apoplastic pathway i.e. only through cell wall. The casparyan strips in the endodermis are made-up of wax -like substance suberin which blocks water and solute movement through the cell wall of the endodermis. As a result water is forced to move through cell membranes and may cross the tonoplast of vacuole. This movement of water through cell membranes is called transmembrane pathway.

Fig 6.




Following schematic diagram showing the apoplastic and symplastic pathway of water movement through root (Fig 7)


Fig 5



**Apoplastic (Red) and symplastic (Blue) and transmembrane (green) pathways of movement of substances in a plant cell**



With the help of the following schematic arrow flow chart, you can understand the path of water from soil to root xylem.



## MECHANISM OF WATER ABSORPTION

### 1. Active absorption of water

In this process the root cells play active role in the absorption of water and metabolic energy released through respiration is consumed active absorption may be of two kinds.

### Steps involved in the active osmotic absorption of water

First step in osmotic the osmotic absorption of water is the imbibition of soil water by the hydrophilic cell walls of root hairs. Osmotic pressure of the cell sap of root hairs is usually higher than the OP of the soil water. Therefore, the DPD and suction presume in the root hairs become higher and water from the cell walls enters into them through plasma membrane by osmotic diffusion. As a result, OP, suction pressure and DPD of root hairs how become lower, while their turgor pressure is increased.

Now the cortical cells adjacent to root hairs have high OP, SP & DPD in comparison to the root hairs.

Therefore, water is drawn into the adjacent cortical cells from root hairs by osmotic diffusion. In the same way, by cell to cell osmotic diffusion gradually reaches the inner most cortical cells and the endodermis.


Osmotic diffusion of water into endodermis takes place through special thin walled passage cells because the other endodermis cells have casparyan strips on thin walls which are impervious to water.


Water from endodermis cells is drawn into the cells of pericycle by osmotic diffusion which now become turgid and their suction pressure is decreased.

In the last step, water is drawn into xylem from turgid pericycle cells (In roots the vascular bundles are radical and protoxylem elements are in contact with pericycle). It is because in the absence of turgor pressure of the xylem vessels, the SP of xylem vessels become higher than SP of the cells of the pericycle when water enters into xylem from pericycle a pressure is developed in the xylem of roots which can raise the water to a certain height in the xylem. This pressure is called as root pressure.

### (A) Osmotic absorption

Water is absorbed from the soil into the xylem of the roots according to osmotic gradient.






### (B) Non-osmotic absorption

Water is absorbed against the osmotic gradient. Sometimes, it has been observed that absorption of water takes place even when OP of soil water is high than OP of cell sap. This type of absorption which is non-osmotic and against the osmotic gradient requires the expenditure of metabolic energy probably through respiration.

### 2. Passive absorption of water

It is mainly due to transpiration, the root cells do not play active role and remain passive.

#### STEPS:



## **2. Passive absorption of water**

Passive absorption of water takes place when rate of transpiration is usually high. Rapid evaporation of water from the leaves during transpiration creates a tension in water in the xylem of the leaves. This tension is transmitted to water in xylem of roots through the xylem of stream and water rises upward to reach the transpiring surfaces. As the results soil water enters into the cortical cells through root hairs to reach the xylem of roots to maintain the supply of water. The force of this entry of water is created in leaves due to rapid transpiration and hence, the root cells remain passive during this process.

## **External factors affecting absorption of water**

### **1. Available soil water**

Sufficient amount of water should be present in the soil in such form which can easily be absorbed by the plants. Usually the plants absorb capillary water i.e water present in films in between soil particles other forms of water in the soil eg. Hygroscopic water, combined water, gravitational water etc. is not easily available to plants.

Increased amount of water in the soil beyond a certain limit results in poor aeration of the soil which retards metabolic activities of root cells like respiration and hence, the rate of water absorption is also retarded.

### **2. Concentration of soil solution**

Increased concentration of soil solution (due to presence of more salts in the soil) results in higher OP. If OP of soil solution will become higher than the OP of cell sap in root cells, the water absorption particularly the osmotic absorption of water will be greatly suppressed. Therefore, absorption of water is poor in alkaline soils and marshes.

### **3. Soil air**

Absorption of water is retarded in poorly aerated soils because in such soils deficiency of O<sub>2</sub> and consequently the accumulation of CO<sub>2</sub> will retard the metabolic activities of roots like respiration. This also inhibits

rapid growth and elongation of the roots so that they are deprived of fresh supply of water in the soil. Water logged soils are poorly aerated and hence, are physiologically dry. They are not good for absorption of water.

#### **4. Soil temperature**

Increase in soil temperature up to about 30°C favours water absorption. At higher temperature water absorption is decreased. At low temperature also water absorption decreased so much so that at about 0°C, it is almost decreased. This is probably because at low temperature.

1. The viscosity of water and protoplasm is increased
2. Permeability of cell membrane is decreased
3. Metabolic activity of root cells are decreased
4. Root growth and elongation of roots are checked.

#### **Quiz**

1. Roots have a root cap, but lack nodes and never bear leaves or flowers directly
2. The meristem is at the tip and manufactures new cells; it is an area of cell division and growth.
3. Behind the meristem is the zone of elongation
4. The zone of maturation is directly beneath the stem. Here, cells become specific tissues such as epidermis, cortex, or vascular tissue.
5. Root hairs are delicate, elongated epidermal cells that occur in a small zone just behind the root's growing tip.
6. The movement of water from one cell to other cell through the plasmodesmata is called the symplastic pathway of water movement.
7. The caspian strips in the endodermis are made-up of wax -like substance suberin which blocks water and solute movement through the cell wall of the endodermis.



## 05. TRANSLOCATION OF SOLUTES

### **Translocation of organic solutes**

The movement of organic food materials or the solutes in soluble form one place to another in higher plants is called as translocation of organic solutes

### **Directions of translocation**

Translocation of organic solutes may take place in the following directions.

#### **1. Downward translocation**

Mostly, the organic material is manufactured by leaves and translocated downward to stem and roots for consumption and storage.

#### **2. Upward translocation**

It takes place mainly during the germination of seeds, tubers etc. When stored food after being converted into soluble form is supplied to the upper growing part of the young seedling till it has developed green leaves.

Upward translocation of solutes also takes place through stem to young leaves, buds and flowers which are situated at the tip of the branch.

#### **3. Lateral translocation**

Radical translocation of organic solutes also takes place in plants from the cells of the pith to cortex.

### **Path of the translocation of organic solutes**

#### **1. Path of downward translocation**

Downward translocation of the organic solutes takes place through phloem. This can be proved by the ringing experiment.

#### **2. Path of upward translocation**

Although translocation of organic solutes take place through phloem, but under certain conditions it may take place through xylem.

#### **3. Path of lateral translocation**

Lateral translocation from pith to cortex takes place through medullary rays.

## Mechanism of translocation

Various theories have been put forward to explain the mechanism of phloem conduction. Among them Munch's (1930) hypothesis is most convincing.

### Munch's mass flow or pressure flow hypothesis

According to this hypothesis put forward by Munch (1930) and others, the translocation of organic solutes takes place through phloem along a gradient of turgor pressure from the region of higher concentration of soluble solutes (supply end) to the region of lower concentration (consumption end). The principle involved in this hypothesis can be explained by a simple physical system as shown in Fig.

Two membranes X and Y permeable only to water and dipping in water are connected by a tube T to form a closed system membrane X contains more concentrated sugar solution than in membrane Y.

Due to higher osmotic pressure of the concentrated sugar solution in the membrane X, water enters into it so that its turgor pressure is increased. The increase in turgor pressure results in mass flow of sugar solution to membrane Y through the T till the concentration of sugar solution in both the membranes is equal.

In the above system it could be possible to maintain continuous supply of sugars in membrane X and its utilization on conversion into insoluble form in membrane Y, the flow of sugar solution from X to Y will continue indefinitely.

According to this theory, a similar analogous system for the translocation of organic solutes exists in plants. As a result of photosynthesis, the mesophyll cells in the leaves contain high concentration of organic food material in them in soluble form and correspond to membrane X or supply end.


The cells of stem and roots where the food material is utilized or converted into insoluble form correspond to membrane Y or consumption end. While the sieve tubes in phloem which are placed and to end correspond to the tube T.

Mesophyll cells draw water from the xylem of the leaf due to higher osmotic pressure and suction presume of their sap so that their turgor pressure is increased. The turgor presume in the cells of stem and the roots is comparatively low and hence, the soluble organic solutes begin to flow en mass from mesophyll through phloem down to the cells of stem and the roots under the gradient of turgor presume. In the stem and the roots, the organic solutes are either consumed or converted into insoluble form and the excess water is released into xylem through cambium.

## **XYLEM TRANSPORT**

### **ASCENT OF SAP**

The water after being absorbed by the roots is distributed to all parts of the plants. In order to reach the topmost part of the plant, the water has to move upward through the stem. The upward movement of water is called as Ascent of sap.



Ascent of sap can be studied under the following two headings.

1. Path of ascent of sap
2. Mechanism of ascent of sap.

### 1. Path of ascent of sap

Ascent of sap takes place through xylem. It can be shown by the experiment.

A leafy twig of Balsam plant (it has semi transpiration stem) is cut under water (to avoid entry of air bubble through the cut end of the stem) and placed in a beaker containing water with some Eosine (a dye) dissolved in it.

After sometimes coloured lines will be seen moving upward in the stem. If sections of stem are cut at this time, only the xylem elements will appear to be filled with coloured water.

## 2. Ringing experiment

A leafy twig from a tree is cut under water and placed in a beaker filled with water. A ring of bark is removed from the stem. After sometime it is observed that the leaves above the ringing part of the stem remain fresh and green. It is because water is being continuously supplied to the upper part of the twig through xylem.

## B. Mechanism of ascent of sap

In small trees and herbaceous plants, the ascent of sap can be explained easily, but in tall trees like Eucalyptus and conifers reaching a height of 300-400 feet), where water has to rise up to the height of several hundred feet, the ascent of sap, it feet, becomes a problem. To explain the mechanism of Ascent of sap, a number of theories have been put forward.

- a. vital theory
- b. root pressure theory
- c. physical force theory
- d. transpiration pull and cohesion of water theory

## A. Vital theories

According to vital theories, the ascent of sap is under the control of vital activities in the stem.

1. According to Godlewski (1884) – Ascent of sap takes place due to the pumping activity xylem tissues which are living.
2. According to Bose (1923) – upward translocation of water takes place due to pulsatory activity of the living cells of the inner most cortical layer just outside the endodermis.

## B. Root pressure theory

Although, root pressure which is developed in the xylem of the roots can raise water to a certain height but does not seem to be an effective force in ascent of sap due to the following reasons. Magnitude of root pressure is very low (about 2 atmos).

Even in the absence of root pressure, ascent of sap continues. For example, when leafy twig is cut under water and placed in a beaker full of water it remains fresh and green for sufficient long time.

### C. Physical force theories

Various physical forces may be involved in ascent of sap.

#### 1. Atmospheric pressure

This does not seem to be convincing because

it cannot act on water present in xylem in roots

Incase it is working, and then also it will not be able to raise water beyond 34.

#### 2. Imbibition

Sachs (1878) supported the view that ascent of sap could take place by imbibition through the walls of xylem. But imbibitional force is insignificant in the A. of sap because it takes place through the lumen of xylem elements and not through walls.

#### 3. Capillary force

In plants the xylem vessels are placed one above the other forming a sort of continuous channel which can be compared with long capillary tubes and it was thought that as water rises in capillary tube due to capillary force in the same manner ascent of sap takes place in the xylem.

### D. Transpiration pull and cohesion of water theory


This theory was originally proposed by Dixon and Jolly (1894) later supported and elaborated by Dixon (1924). This theory is very convincing and has now been widely supported by many workers.

Although H- bond is very weak (Containing about 5 K –cal – energy) but they are present in enormous numbers as incase of water, a very strong mutual force of attraction or cohesive force develops between water molecules and hence they remain in the form of a continuous water column in the xylem. The magnitude of this force is very high (up to 350 atm), therefore the continuous water column in the xylem cannot be broken easily due to the

force of gravity or other abstractions offered by the internal tissues in the upward movement of water.

The adhesive properties of water i.e. attractions between the water molecules and the containers walls (here the walls of xylem) further ensure the continuity of water column in xylem.

When transpiration takes place in the leaves at the upper parts of the plant, water evaporates from the intercellular spaces of the leaves to the outer atmosphere through stomata. More water is released into the intercellular spaces from mesophyll cells. In turn, the mesophyll cells draw water from the xylem of the leaf. Due to all this, a tension is created in the xylem elements of the leaves. This tension is transmitted downward to water in xylem elements of the root through the xylem of petiole and stem and the water is pulled upward in the form of continuous unbroken water column to reach the transpiring surfaces up to the top of the plant





## 06. TRANSPERSION

Although large quantities of water are absorbed by plant from the soil but only a small amount of it is utilized. The excess of water is lost from the aerial parts of plants in the form of water vapours. This is called as transpiration.

Transpiration is of three types

### **1. Stomatal transpiration**

Most of the transpiration takes place through stomata. Stomata are usually confined in more numbers on the lower sides of the leaves. In monocots. Eg. Grasses they are equally distributed on both sides. While in aquatic plants with floating leaves they are present on the upper surface.

### **2. Cuticular transpiration**

Cuticle is impervious to water, even though, some water may be lost through it. It may contribute a maximum of about 10% of the total transpiration.

### **3. Lenticular transpiration**

Some water may be lost by woody stems through lenticells which is called as lenticular transpiration.

#### **Mechanism of stomatal transpiration**

The mechanism of stomatal transpiration which takes place during the day time can be studied in three steps.

- i. Osmotic diffusion of water in the leaf from xylem to intercellular space above the stomata through the mesophyll cells.
- ii. Opening and closing of stomata (stomatal movement)
- iii. Simple diffusion of water vapours from intercellular spaces to other atmosphere through stomata.
  - ◆ Inside the leaf the mesophyll cells are in contact
  - ◆ With xylem, and on the other hand with intercellular space above the stomata

- When mesophyll cells draw water from the xylem they become turgid and their diffusion pressure deficit (DPD) and osmotic pressure (OP) decreases with the result that they release water in the form of vapour in intercellular spaces close to stomata by osmotic diffusion. Now in turn, the O.P and D.P.D of mesophyll cells become higher and hence, they draw water from xylem by osmotic diffusion.


### Opening and closing of stomata (Stomatal movement)

The stomata are easily recognized from the surrounding epidermal cells by their peculiar shape. The epidermal cells that immediately surround the stomata may be similar to other epidermal cells or may be different and specialized. In the latter case, they are called as subsidiary cells.

The guard cells differ from other epidermal cells also in containing chloroplasts and peculiar thickening on their adjacent surface (in closed stomata) or on surfaces.

Consequent to an increase in the osmotic pressure (OP) and diffusion pressure deficit (DPD) of the guard cells (which is due to accumulation of osmotically active substances), osmotic diffusion of water from surrounding epidermal cells and mesophyll

## Control of Stomatal Opening and Closing



cells into guard cells follows. This increase the turgor pressure (TP) of the guard cells and they become turgid. The guard cells swell, increase in length and their adjacent thickened surfaces starch forming a pore and thus the stomata open.

On the other hand, when OP and DPD of guard cells decrease (due to depletion of osmotically active substances) relative to surrounding epidermal and mesophyll cells, water is released back into the latter by osmotic diffusion and the guard cells become flaccid. The thickened surfaces of the guard cells come close to each other, thereby closing the stomatal pore and stomata.

Osmotic diffusion of water into guard cells occur when their osmotic pressure increases and water potential decreases (i.e become more negative) related to those of surrounding epidermal and mesophyll cells. The guard cells become flaccid when their osmotic pressure decreases relative to the surrounding cells (Movement of water takes place from a region of higher water potential to a region of lower water potential).

These may be several different agents or mechanisms which control stomatal movements.

Hydrolysis of starch into sugars in guard cells

Synthesis of sugars or organic acids in them

The active pumping of  $K^+$  ions in the guard.

### **1. Hydrolysis of starch into sugars in guard cells**


#### **Starch – sugar Inter conversion theory**

This classical theory is based on the effect of pH on starch phosphorylase enzyme which reversibly catalyses the conversion of starch + inorganic phosphate into glucose -1 phosphate.

During the day, pH is high in guard cells. This favours hydrolysis of starch (which is insoluble into glucose -1- phosphate (which is soluble) so that osmotic pressure is increased in guard cells.

Consequently water enters, into the guard cells by osmotic diffusion from the surrounding epidermal and mesophyll cells. Guard cells become turgid and the stomata open.

During dark, reverse process occurs. Glucose 1-phosphate is converted back into starch in the guard cells thereby decreasing osmotic pressure. The guard cell release water, become flaccid and stomata become closed.




According to Steward (1964), the conversion of starch and inorganic phosphate into glucose-1-phosphate does not cause any appreciable change in the osmotic pressure because the inorganic phosphate and glucose-1-phosphate are equally active osmotically.

In this scheme he has suggested that,

Glucose-1-phosphate should be further converted into glucose and inorganic phosphate for the opening of stomata.

Metabolic energy in the form of ATP would be required for the closing of stomata which probably comes through respiration.



## 2. Synthesis of sugars or organic acids in Guard cells

During day light photosynthesis occurs in guard cells as they contain chloroplast. The soluble sugars formed in this process may contribute in increasing the osmotic potential of guard cells and hence resulting in stomatal opening. However, very small amounts of soluble sugars (osmotically active) have been extracted from the guard cells which are insufficient to affect water potential.

As a result of photosynthesis CO<sub>2</sub> concentration in guard cells decreases which leads to increased pH up of organic acids, chiefly malic acid during this period in guard cells. The formation of malic acid would produce proton that could operate in an ATP-driven proton K<sup>+</sup> exchange pump moving protons into the adjacent epidermal cells and K ions into guard cells and thus may contribute in increasing the osmotic pressure of the guard cells and leading to stomatal opening.

Reverse process would occur in darkness.

## 3. ATP –Driven proton (H<sup>+</sup>) – K exchange pump mechanism in Guard cells

According to this mechanism, there is accumulation of K<sup>+</sup> ions in the guard cells during day light period. The protons (H<sup>+</sup>) are ‘pumped out’ from the guard cells into the adjacent epidermal cells and in exchange K<sup>+</sup> ions are mediated through ATP and thus are an active process. ATP is generated in non-cyclic photophosphorylation in photosynthesis in the guard cells. The ATP required in ion exchange process may also come through respiration.

The accumulation of K ion is sufficient enough to significantly decrease the water potential of guard cells during day light. Consequently, water enters into them from the adjacent epidermal and mesophyll cells thereby increasing their turgor pressure and opening the stomatal pore.

Reverse situation prevails during dark when stomata are closed. There is no accumulation of ‘K’ in g cells in dark.

(iii) The last step in the mechanism of transpiration is the simple diffusion of water vapours from the intercellular spaces to the atmosphere through open stomata. This is because the intercellular spaces are more saturated with moisture in comparison to the outer atmosphere in the vicinity of stomata.

### **Significance of Transpiration**

Plants waste much of their energy in absorbing large quantities of water and most of which is ultimately lost through transpiration.

Some people think that – Transpiration is advantageous to plants.

Others regard it as an unavoidable process which is rather harmful.

### **Advances of transpiration**

#### **1. Role of movement of water**

Plays an important role in upward movement of water i.e. Ascent of sap in plants.

#### **2. Role in absorption and translocation of mineral salts**

Absorption of water and mineral salts are entirely independent process. Therefore transpiration has nothing to do with the absorption of mineral salts.

However, once mineral salts have been absorbed by the plants, their further translocation and distribution may be facilitated by transpiration through translocation of water in the xylem elements.


#### **Transpiration from the leaf regulates by**

#### **3. Role of regulation of temperature**

Some light energy absorbed by the leaves is utilized in photosynthesis; rest is converted into heat energy


1. difference in water vapor concentration between the leaf air spaces and the external air
2. diffusional resistance ( $r$ ) of this pathway
3. leaf stomatal resistance ( $rs$ )
4. leaf boundary layer resistance
5. control of stomatal apertures by the guard cells

which raises their temperature. Transpiration plays an important role in controlling the temperature of the plants. Rapid evaporation of water from the aerial parts of the plant through transpiration brings down their temperature and thus prevents them from excessive heating.




### Transpiration as a necessary evil

1. When the rate of transpiration is high and soil is deficient in water, an internal water deficit is created in the plants which may affect metabolic processes
2. Many xerophytes have to develop structural modification and adaptation to check transpiration.



3. Deciduous trees have to shed their leaves during autumn to check loss of water.

But, in spite of the various disadvantages, the plants cannot avoid transpiration due to their peculiar internal structure particularly those of leaves. Their internal structure although basically meant for gaseous exchange for respiration, P.S. etc. is such that it cannot check the evaporation of water. Therefore, many workers like Curtis (1926) have called transpiration as necessary evil.



## Factors affecting transpiration rate

### A. External factors

#### 1. Atmospheric humidity

In humid atmosphere, (when relative humidity) is high), the rate of transpiration decreases. It is because atmosphere is more saturated with moisture and retards the diffusion of water vapour from the intercellular spaces of the leaves to the outer atmosphere through stomata.

In dry atmosphere, the RH is low and the air is not saturated with moisture and hence, the rate of transpiration increases.

#### 2. Temperature

An increase in temperature brings about an increase in the rate of transpiration by

1. lowering the relative humidity
2. Opening of stomata widely

#### 3. Wind

- i. When wind is stagnant (not blowing), the rate of transpiration remains normal
- ii. When the wind is blowing gently, the rate of transpiration increases because it removes moisture from the vicinity of the transpiration parts of the plant thus facilitating the diffusion of water vapour from the intercellular spaces of the leaves to the outer atmosphere through stomata.
- iii. When the wind is blowing violently, the rate of transpiration decreased because it creates hindrance in the outward diffusion of water vapours from the transpiring part and it may also close the stomata.

#### **4. Light**

Light increases the rate of transpiration because,

In light stomata open; It increases the temperature

In dark, due to closure of stomata, the stomatal transpiration is almost stopped.

#### **5. Available soil water**

Rate of transpiration will decrease if there is not enough water in the soil in such form which can be easily absorbed by the roots.


#### **6. CO<sub>2</sub>**

An increase in CO<sub>2</sub> concentration in the atmosphere (Over the usual concentration) more so inside the leaf, leads towards stomatal closure and hence it retards transpiration.

### **B. Internal factors**


#### **1. Internal water conditions**

It is very essential for transpiration. Deficiency of water in the plants will result in decrease of transpiration rate. Increase rate of transpiration containing for longer periods often create internal water deficit in plants because absorption of water does not keep pace with it.



## 2. Structural features

The number, size, position and the movement of stomata affect rate of transpiration. In dark stomata are closed and stomatal transpiration is checked. Sunken stomata help in reducing the rate of stomatal transpiration. In xerophytes the leaves are reduced in size or may even fall to check transpiration. Thick cuticle on presence of wax coating on exposed parts reduces cuticles transpiration.





### Antitranspirants

A number of substances are known which when applied to the plants retard their transpiration. Such substances are called as antitranspirants. Some examples of antitranspirants are colourless plastics, silicone, oils, low viscosity waxes, phenyl mercuric acetate, abscisic acid,  $\text{CO}_2$ , etc. Colourless plastic, silicone oils and low viscosity waxes belong to one group as these are sprayed on the leaves, form after film which is permeable to  $\text{O}_2$  and  $\text{CO}_2$  but not to water.

Fungicide phenyl mercuric acetate, when applied in low concentration ( $10^{-4} \text{ m}$ ), it exercised a very little toxic effect on leaves and resulted in partial closure of stomatal pores for a period of two weeks. Similarly ABA a plant hormone also induces stomatal closure.  $\text{CO}_2$  is an effective antitranspirant. A little rise in  $\text{CO}_2$  concentration from the natural 0.03% to 0.05% induces partial closure of stomata. Its higher concentration cannot be used which results in complete closure of stomata affecting adversely the photosynthesis and respiration.

### GUTTATION



In some plants such as garden nasturtium, tomato, colocasia etc, water drops ooze out from the uninjured margins of the leaves where a main vein ends. This is called as guttation and takes place usually early in the morning when the rate of absorption and root pressure are high while the transpiration is very low.

The phenomenon of guttation is associated with the presence of special types of stomata at the margins of the leaves which are called as **water stomata or hydathodes**. Each hydathode consists of a water pore which remains permanently open.

Below this there is a small cavity followed by a loose tissue called as epithem. This epithem is in close association with the ends of the vascular elements of veins. Under high root pressure the water is given to the epithem by the xylem of the veins. From epithem

water is released into the cavity. When this cavity is completely filled with watery solution, the later begins to ooze out in the form of watery drops through the water pore.

#### Difference between transpiration and Guttation

Transpiration	Guttation
1. Water is lost from aerial parts of plants in the form of invisible water vapours	Watery solution oozes out from uninjured margins of aerial leaves only
2. Transpiration occurs mostly through stomata. It may also takes place through cuticle and lenticels	It occurs only through hydathodes (water stomata)
3. It takes place throughout the day, its rate being maximum at noon.	It takes place only early in the morning when root pressure and the rate of water absorption are higher

## 07. MINERAL NUTRITION

The term, *mineral nutrient* is generally used to refer to an inorganic ion obtained from the soil and required for plant growth. The chemical form in which elements are applied to plants is called as *nutrient*. Nutrition may be defined as the supply and absorption of chemical compounds needed for plant growth and metabolism.

The nutrients indispensable for the growth and development of higher plants are obtained from three sources viz., atmosphere, water and soil. The atmosphere provides carbon and oxygen as carbon dioxide. Carbon is reduced during photosynthesis and oxygen is utilized during aerobic respiration. Soil provides the mineral ions.

### **Essential elements**

The term essential mineral element was proposed by Arnon and Stout (1939). These are the composition of both macro and microelements, in the absence of any one of these elements the plant cannot maintain its normal growth and develops deficiency symptoms, affects metabolism and die prematurely. Of the many elements that have been detected in plant tissues, only 16 are essential for all higher plants. They are C, H, O, N, P, K, Ca, Mg, S, Zn, Cu, Fe, Mn, B, Cl and Mo. In the absence of each of the essential elements, plants develop deficiency symptoms characteristic of the deficient element and die prematurely.

### **Macronutrients**

The nutrient elements which are required for the growth of plants relatively in larger quantities are called as *major nutrients* or *macronutrients*. The major elements required for growth of plants are C, H, O, N, P, K, Ca, Mg and S. Among these nutrients, C, H and O are taken up by the plants from the atmosphere and water. The N, P, K, Ca, Mg and S are taken up by the plants from the soil and they are applied in the form of chemical fertilizers either through the soil or foliage.

### **Micronutrients**

The nutrient elements which are required comparatively in small quantities are called as *minor* or *micro nutrients* or trace elements. The micronutrients required for the plant growth are Zn, Cu, Fe, Mn, Mo, B and Cl.

### Tracer elements or labeled elements

The nutrient elements that are required for plants are sometimes labeled and used to study their movement or tracing out the involvement of such nutrients in metabolism in different organs of plants, are called as *tracer elements*. They may either be stable or radioactive types and they are also called as *isotopic elements*.

E.g. Stable isotopes:  $^{15}\text{N}$ ,  $^{12}\text{C}$ ,  $^{31}\text{P}$

Radio active :  $^{14}\text{C}$ ,  $^{32}\text{P}$ ,  $^{65}\text{Zn}$ ,  $^{56}\text{Fe}$ ,  $^{60}\text{Co}$ , etc.

### Hidden hunger

When the plants are not able to meet their requirement either one or more of these essential elements, the plants will undergo starvation for such elements. At the initial stage of deficiency of such elements plants will not show any characteristic symptoms which could be exhibited morphologically and due to want of those elements some activities of plants would rather be affected and the internal deficiency is called as *Hidden hunger*.

### General role of essential elements

In general, an element is essential to the life of a higher green plant for one or more of the following three reasons.

1. It may perform a nutritive role by being a component of one or more of the major classes of plant constituents.
2. It may be a catalytic role either as an action for of an enzyme or as an integral component of an enzyme.
3. It may function as a free ion and thereby exert a balancing role in maintaining electro-neutrality within plant cells (e.g. Potassium).

### Criteria for essentiality of elements

The demonstration of the essentially several elements (macro and micronutrients), especially, micronutrients is rather very difficult. In view of the technical difficulties associated with demonstrating the essentiality of elements required in very small amounts, Arnon and Stout (1939) suggested the adoption of the following three criteria of essentiality for judging the exact status of a mineral in the nutrient of a plant.

1. The element must be essential for normal growth or reproduction and the plant processes cannot proceed without it.
2. The element cannot be replaced by another element.
3. The requirement must be direct i.e., not the result of some indirect effect such as relieving toxicity caused by some other substance.

Another recent suggestion to the criteria of essentiality is that some elements might better be called *functional or metabolic elements* rather than essential elements. This is intended to indicate that an element that is metabolically active, functional or metabolic may or may not be essential. For example in chlorine-bromine, chlorine is designated as a functional element rather than an essential element as chlorine can be substituted with bromine.

Based on the mobility in phloem, elements are also classified into three types.

- |                      |                     |
|----------------------|---------------------|
| 1. Mobile elements   | : N, K, P, S and Mg |
| 2. Immobile elements | : Ca, Fe and B      |
| 3. Intermediate      | : Zn, Mn, Cu, Mo    |

### **Functions of elements**

*Protoplasmic elements* : N, P, S


*Balancing elements* : Ca, Mg, K – counteract to toxic effects of other minerals  
by causing ionic balance.

*Frame work elements* : C, H<sub>2</sub>O – as they are the constituents of carbohydrates that form cell walls.

*Catalytic elements* : Mn, Cu, Mg, etc.

### **SOIL LESS GROWTH OR HYDROPONICS**

The practice of growing plants in nutrient enriched water without soil is called as soil less growth or hydroponics. However, the term hydroponics is now being applied to plants rooted in sand, gravel or other similar matter which is soaked with a recycling flow of nutrient – enriched water.



According to a recent limited nations report on hydroponics: In area of tropics, where the water deficiency is the limiting factor in crop production, the soil less methods hold out much promise because of the more economical use of water.

The report also indicated that in some areas, lack of fertile soil or very thin soil layers may also move soil less methods worth serious consideration.

Besides these the other advantages of growing cucumbers, egg plants, peppers, lettuces, spinach and other vegetables hydroponically under controlled environment are

1. The regulation of nutrients

2. Control of pests and diseases
3. Reduction of labour cost
4. Sometimes quicker yield

But there are two main drawbacks of hydroponics farming.

1. Firstly the cost of setting up the system is very high
2. Secondly it requires skills and knowledge of its operation

## 08. MECHANISM OF UPTAKE - PHYSIOLOGICAL ROLE OF NUTRIENTS

### Mechanism

Previously it was thought that absorption of mineral salts takes place along with water absorption. But it is now understood that mineral salt absorption and water absorption are two different processes.

**Soil serves as a main source of mineral salts of **ionic forms** in clay crystals have a central nucleus called **micelle****


The micelles are negatively charged and maintain a balance, they attract and hold positively charged ions on the surface

**cationic forms**


K, Mg, Ca, Fe, Mn, Cu, Zn and Co

**anionic forms and N,P,B,S and Cl.**

These ions are either in the form of loosely absorbed ions or firmly absorbed ions on the colloidal particles



Mineral salts are absorbed from the soil solution in the form of ions. They are chiefly absorbed through the meristematic regions of the roots near the tips.




Plasma membrane of the root cells is not permeable to all the ions. It is selectively permeable. All the ions of the same salt are not absorbed at equal rate but leads unequal absorption of ions. First step in the absorption of mineral salts is the process of Ion exchange which does not require metabolic energy.

The further processes of the absorption of mineral salts may be of two types.

1. Passive and 2. Active

**Root interception** – roots obtain nutrients by physically contacting nutrients in soil solution or on soil surfaces;


- roots contact ~1% of soil volume;
- mycorrhizal infection of root increase root-soil contact




## 1. Passive absorption

When the concentration of mineral salts is higher in the outer solution than in the cell sap of the root cells, the mineral salts are absorbed according to the concentration gradient by simple process of diffusion. This is called as passive absorption because it does not require expenditure of metabolic energy.

**Ion exchange**




The ions adsorbed on the surface of the plasma membrane of the root cells may be exchanged with the ions of same sign from external solution for eg. The cation  $K^+$  of the external soil solution may exchanged with  $H^+$  ions adsorbed on the surface of the plasma membrane. Similarly anion may be exchanged with  $OH^-$  ions. There are two theories regarding the mechanism of ion exchange.



### 1. Contact exchange theory


According to this theory the ions adsorbed or the surface of root cells and clay particles are not held tightly but oscillate within small volume of space. If the roots and clay particles are in close contact with each other, the oscillation volume of ions adsorbed on root surface may overlap by the oscillation volume of ions adsorbed on clay particles, and the ions adsorbed on clay particle may be exchanged with the ions adsorbed on root surface directly without first being dissolved in soil solution.



### 2. Carbonic acid exchange theory

According to this theory, the CO<sub>2</sub> released during respiration of root cells combines with water to form carbonic acid (H<sub>2</sub>CO<sub>3</sub>). Carbonic acid dissociates into H<sup>+</sup> and an anion HCO<sub>3</sub> in soil solution. These H<sup>+</sup> ions may be exchanged for cations adsorbed on the clay particles. The cations thus released into the soil solution from the clay particles, may be

adsorbed on root cells in exchange for  $H^+$  ions or as in ion pairs with bicarbonate. Thus, the soil solution plays an important role in carbonic acid exchange theory.



## 2. Active absorption of mineral salts

It has been observed that the cell sap in plants accumulates large quantities of mineral salts ions against the concentration gradient. The accumulation of mineral salts against to concentration gradient is an active process which involves the expenditure of metabolic energy through respiration. The active absorption of mineral salts involves the operation of a carrier compound present in the plasma membrane of the cells.

### The carrier concept

According to this theory, the plasma membrane is impermeable to free ions. But some compounds present in it acts as carrier and combines with ions to form carrier- ion-complex which can move across the membrane. On the inner side of the membrane this complex leaves releasing ions into the cell while the carrier goes back to the outer surface to pick up fresh ions. They are two hypotheses based on the carrier concept to explain the mechanism of active salt absorption. Although they are not universally accepted.

#### 1. Lundegardhs cytochrome pump theory

Lundegardh and Burstrom (1933) believed that there was a definite correlation between respiration and anion absorption. Thus when a plant is transferred from water to a


salt solution the rate of respiration increases. This increase in rate of respiration over the normal respiration has been called as anion respiration or salt respiration.

Lundegardh (1954) proposed cytochrome pump theory which is based on the following assumptions.

1. The mechanism of anion and cation absorption is different
2. Anions are absorbed through cytochrome chain by an active process.  
(Cytochromes are ion – porphyrin proteins that act as enzymes and helps in electron transfer during respiration).
3. Cations are absorbed passively.


According to this theory

- 1) Dehydrogenase reactions on inner side of the membrane give rise to protons ( $H^+$ ) and electrons ( $e^-$ ).
- 2) The electrons travel over the cytochrome chain towards outside the membrane, so that the Fe of the cytochrome becomes reduced ( $Fe^{++}$ ) on the outer surface and oxidized ( $Fe^{+++}$ ) on the inner surface.
- 3) On the outer surface, the reduced cytochrome is oxidized by oxygen releasing the electron ( $e^-$ ) and taking an anion ( $A^-$ ).
- 4) The electron thus released unites with  $H^+$  and oxygen to form water
- 5) The anion ( $A^-$ ) travels over the cytochrome chain towards inside.
- 6) On the inner surface the oxidized cytochrome becomes reduced by taking an electron produced through the dehydrogenase reactions and the anion ( $A$ ) is released.
- 7) As the result of anion absorption, a cation ( $M$ ) moves passively from outside to inside to balance the anion.



## 2. Bennert – Clark's protein Lecithin Theory

In 1856, Bennet – Clark suggested that because the cell membranes chiefly consist of phospholipids and proteins and certain enzymes seem to be located on them, the carrier could be a protein associated with the phosphatide called as lecithin. He also assumed the presence of different phosphatides to correspond with the number of known competitive groups of cations and anions.



According to this theory

1. Phosphate group in the phosphatide is regarded as the active centre binding the cations and the basic choline group as the anion binding centre.
2. The ions are liberated on the inner surface of the membrane by decomposition of lecithin by the enzyme lecithinase.


3. The regeneration of the carrier lecithin form phosphatidic acid and choline takes place in the presence of the enzyme choline acetylase and choline esterase and ATP. The latter acts as a source of energy.

### Donnans' Equilibrium


The accumulation of ions inside the cells without involving expenditure of the metabolic energy can be explained to some extent by Donnan's equilibrium theory.

According to this theory there are certain pre existing ions inside the cell which cannot diffuse outside through membrane. Such ions are called as in diffusible or fixed ions. However, the membrane is permeable to both anions and cations of the outer solutions.

Suppose there are certain fixed anions in the cell which is in contact with outer solution containing anions and cations. Normally equal number of anions and cations would have diffused into the cell through an electrical potential to balance each other, but to balance the fixed anions more cations will diffuse into the cell. This equilibrium is known as Donnan's equilibrium. In this particular case, there would be an accumulation of cations inside the cell.



If however, there are fixed cations inside the cell, the Donnan's equilibrium will result in the accumulation of anions inside the cell.



## Specific roles of essential mineral elements

### A. Macronutrients

#### 1. Nitrogen

- Nitrogen is important constituent of proteins, nucleic acids, porphyrins (chlorophylls & cytochromes) alkaloids, some vitamins, coenzymes etc
- Thus N plays very important role in metabolism, growth, reproduction and heredity.

#### 2. Phosphorus

- It is important constituent of nucleic acids, phospholipids, coenzymes NADP, NADPH<sub>2</sub> and ATP
- Phospholipids along with proteins may be important constituents of cell membranes
- P plays important role in protein synthesis through nucleic acids and ATP
- Through coenzymes NAD, NADP and ATP, it plays important role in energy transfer reactions of cell metabolism eg. Photosynthesis, respiration and fat metabolism etc.

#### Potassium

- Although potassium is not a constituent of important organic compound in the cell, it is essential for the process of respiration and photosynthesis
- It acts as an activator of many enzymes involved in carbohydrate metabolism and protein synthesis
- It regulates stomatal movement

- Regulates water balance

### **Calcium**

- It is important constituent of cell wall
- It is essential in the formation of cell membranes
- It helps to stabilize the structure of chromosome
- It may be an activation of many enzymes

### **Magnesium**

- It is very important constituent of chlorophylls
- It acts as activation of many enzymes in nucleic acid synthesis and carbohydrate metabolism
- It plays important role in binding ribosomal particles during protein synthesis.

### **Sulphur**

- It is important constituent of some amino acids (cystine, cysteine and methionine) with which other amino acids form the protein
- S helps to stabilize the protein structure
- It is also important constituent of vitamin i.e biotin, thiamine and coenzyme A
- Sulphhydryl groups are necessary for the activity of many enzymes.

### **Iron**

- Important constituent of iron porphyrin – proteins like cytochromes, peroxidase, catalases, etc.
- It is essential for chlorophyll synthesis
- It is very important constituent of ferredoxin which plays important role in photochemical reaction in photosynthesis and in biological nitrogen fixation.

### **Micro nutrients**

#### **Zinc**

- It is involved in the biosynthesis of growth hormone auxin (indole 3 acetic acid)

- It acts activator of many enzymes like carbonic anhydrase and alcohol dehydrogenase, etc.

### **Manganese**

- It is an activator of many respiratory enzymes
- It is also an activator of the enzyme nitrite reductase
- It is necessary for the evolution of oxygen (photolysis) during photosynthesis

### **Copper**

- It is an important constituent of plastocyanin (copper containing protein)
- It is also a constituent of several oxidizing enzymes.

### **Boron**

- Boron facilitates the translocation of sugars by forming sugar borate complex.
- It involves in cell differentiation and development since boron is essential for DNA synthesis
- Also involves in fertilization, hormone metabolism etc.

### **Molybdenum**

- It is constituent of the enzyme nitrate reductase and thus plays an important role in nitrogen metabolism
- It is essential for flower formation and fruit set.

## 09. Foliar diagnosis - Nutritional and Physiological disorders

### A. Foliar diagnosis - Symptoms

#### Nitrogen

- Plant growth is stunted because protein content cell division and cell enlargement are decreased
- N deficiency causes chlorosis of the leave i.e yellowing older leaves are affected first.
- In many plants eg. Tomato, the stem, petiole and the leaf veins become purple coloured due to the formation of anthocyanin pigments.



#### Phosphorus

- P deficiency may cause premature leaf fall
- Dead necrotic areas are developed on leave or fruits

- Leaves may turn to dark green to blue green colour. Sometimes turn to purplish colour due to the synthesis and accumulation of anthocyanin pigments.



## Potassium

- Mottled chlorosis of leaves occurs
- Neurotic areas develop at the tip and margins of the leaf
- Plants growth remains stunted with shortening of internodes.




## Calcium

- Calcium deficiency causes disintegration of growing meristematic regions of root, stem and leaves
- Chlorosis occurs along the margins of the younger leaves
- Malformation of young leaves takes place




## Magnesium

- Mg deficiency causes mottled chlorosis with veins green and leaf tissues yellow or white appearing first on older leaves
- Dead neurotic patches appear on the leaves
- In cotton Mg deficiency leads to reddening of leaves and disorder is called as reddening in cotton.



## Sulphur


- Deficiency causes chlorosis of the leaves
- Tips and margins of the leaf roll in ward
- Stem becomes hard due to the development of sclerenchyma.



## Micronutrients

### Iron

Iron deficiency causes chlorosis of young leaves which is usually interveinal.



### Zinc

- Zinc deficiency causes chlorosis of the young leaves which starts from tips and the margins
- The size of the young leaves is very much reduced. This disorder is called as 'little leaf disease'
- Stalks will be very short.



### Manganese

- The young leaves are affected by mottled chlorosis
- Veins remain green
- Small necrotic spots developed on the leaves with yellow strips

### Copper

- Copper deficiency causes necrosis of the tip of the young leaves
- It also causes die-back of citrus and fruit trees
- Also causes reclamation disease or white tip disease of cereals and leguminous plants.



## Boron

- Boron deficiency causes death of shoot tip
- Flower formation is suppressed
- Root growth is stunted
- The other diseases caused by B deficiency is
- Heart rot of beet
- Stem crack of celery
- Brown heart of cabbage
- Water core of turnip
- Internal cork formation in apple
- Hen and chicken in grapes



### Molybdenum

- Molybdenum deficiency causes interveinal chlorosis of older leaves
- Flower formation is inhibited
- Causes whiptail disease in cauliflower plants.



## Foliar Nutrition

Foliar nutrition is fertilizing certain crop plants through aerial spraying.

### Mechanism

Penetration of the spray solution or nutrient solution occurs through cuticle the layer of polymerized wax which occurs on outer surface of the epidermal cells of leaves. After penetration in the cuticle, further penetration take place through fine, thread like semimicroscopic structure called ectodesmata. This extends through the outer epidermal cell wall, from the inner surface of the cuticle to the plasma membrane. When the substance reaches plasma membrane of an epidermal cell, it will be observed by mechanism similar to those which operate in root cells.

1. Foliar nutrition may serve as a mean of applying supplemental macronutrients during critical growth periods when it is impracticable to apply fertilizers to soil. Eg. Unusual period of dry weather.
2. Foliar nutrition may afford a remedy for the time lag between soil applied and plant absorbed. Time is too long because of fast growing rates.

## NUTRITIONAL DISORDERS

When a nutrient element insufficiency (deficiency and/or toxicity) occurs, visual symptoms may or may not appear, although normal plant development will be slowed. When visual symptoms do occur, such symptoms can frequently be used to identify the source of the insufficiency.

## Deficiency Symptoms

- Stunted or reduced growth of the entire plant with the plant itself either remaining green or lacking an over-all green color with either the older or younger leaves being light green to yellow in color.
- Chlorosis of leaves, either interveinal or of the whole leaf itself, with symptoms either on the younger and/or older leaves, or both (chlorosis due to the loss or lack of chlorophyll production).
- Necrosis or death of a portion (margins or interveinal areas) of a leaf, or the whole leaf, usually occurring on the older leaves.
- Slow or stunted growth of terminals (rosetting), the lack of terminal growth, or death of the terminal portions of the plant.
- A reddish purpling of leaves, frequently more intense on the under side of older leaves due to the accumulation of anthocyanin (Mottling)

*Chlorosis* is caused by the deficiency of mineral elements such as Mn, K, Zn, Fe, Mg, S and N. *Mottling* is caused due to the deficiencies of N, Mg, P, S and *Necrosis* due to the deficiency of Mg, K, Zn, Ca and Mo.

## Toxicity Symptoms

Visual symptoms of toxicity may not always be the direct effect of the element in excess on the plant, but the effect of the excess element on one or more other elements. For example, an excessive level of potassium (K) in the plant can result in either magnesium (Mg) and/or calcium (Ca) deficiency, excess phosphorus (P) can result in a zinc (Zn) deficiency and excess Zn in an iron (Fe) deficiency.

These effects would compare to elements, such as boron (B), chlorine (Cl), copper (Cu), and manganese (Mn), which create visual symptoms that are the direct effect of an excess of that element present in the plant.

Some elements, such as aluminum (Al) and copper (Cu) can affect plant growth and development due to their toxic effect on root development and function.

## **Hidden Hunger**

In some instances, a nutrient element insufficiency may be such that no symptoms of stress will visually appear with the plant seeming to be developing normally. This condition has been named hidden hunger, a condition that can be uncovered by means of either a plant analysis and/or tissue test.

A hidden hunger occurrence frequently affects the final yield and the quality of the product produced. For grain crops, the grain yield and quality may be less than expected; for fruit crops, abnormalities, such as blossomed rot and internal abnormalities may occur, and the post harvest characteristics of fruits and flowers will result in poor shipping quality and reduced longevity. Another example is potassium (K) insufficiency in corn, a – deficiency that is not evident until at maturity when plants easily

## **PHYSIOLOGICAL DISORDERS**

Physiological disorder is the abnormal growth pattern or abnormal external or internal conditions of fruits due to adverse environmental conditions such as deviation from normal state of temperature, light, moisture, nutrient, harmful gases and inadequate supply of growth regulators.

### **Disorders associated with low temperature**

#### **1. Leaf chlorosis and frost banding**

Chlorosis was caused by a disruption of chloroplasts caused by winter cold. Green chlorophyll pigments are often converted into yellow pigment. Leaf may appear with distinct bleached bands across the blade of young plants called frost banding e.g.: sugarcane, wheat and barley.

#### **2. Leaf necrosis and malformations**

Spring frost causes various types and degree of injury including cupping, crinkling finishing and curling of leaves of apple trees and

stone fruits. The distortion is caused by death of the developed tissues before the expansion of leaves.

### **3. Stem disorders**

Frost cracks develop when tree trunk or limbs lost their heat too rapidly. The outer layer of bark and wood cool most rapidly and subjected to appreciable tension causing marked shrinkage and cracking following a sudden temperature drop. Affected timber is of poor quality.

## **Disorders associated with high temperature**

### **1. Leaf scorch**

High temperature causes leaf scorch directly or indirectly by stimulating excessive evaporation and transpiration. Tip burn of potato is a widespread example for this disorder.

### **2. Sunscald**

In leaf vegetable crops like lettuce and cabbage, when leaves on the top of the head are exposed to intense heat, water soaked lesions or blistered appearance occur. These irregular shaped areas become bleached and parched later.

### **3. Water core**

In fruit crop like Tomato, exposure to high temperature causes death of the outer cells of fruit skin. Subsequently corky tissue occurs beneath the skin, with watery appearance of the flesh near the core of the fruits faster. Often light stress is coupled with heat stress e.g. sun scald of bean, sun burning of soybean and cowpea. In flower crop like chrysanthemum, increase in light intensity affects flower bud formation. Reproduction phase does not commence and modified into leaf like bracts.

## Disorders caused by light stress

Adverse light intensity causes impaired growth and reduced vigour. Subsequently leaves gradually lose green colour, turning pale green to yellow, stems may dieback little every year. Insufficient light limits photosynthesis, causing food reserves to be depleted.

## Identification of Physiological Disorders and Corrective Measures

Crop	Malady	Corrective measure
Rice	Severe chlorosis of leaves	1% super phosphate and 0.5% ferrous Sulphate
Rice	Irregular flowering and chaffiness multiple deficiency of nutrients	1% super phosphate and magnesium sulphate.
Rice	Tip drying and marginal scoring and Browning	1% super phosphate and 0.5% zinc sulphate.
Maize	Chlorosis	A spray solution containing 0.5% ferrous sulphate and 0.5% urea.
Maize	'White bud' yellowing in the bud leaves only	0.5% zinc sulphate spray with 1% urea.
Maize	Tip drying and marginal scoring pinkish colouration of lower leaves	1% super phosphate and 0.5% zinc sulphate.
Maize	Marginal scorching and yellowing.	0.5% ferrous sulphate and 1% urea
	Irregular drying of tips and margins	25 kg of zinc sulphate / ha
Sorghum	Chlorosis of younger leaves	Spray of 0.5% ferrous sulphate with


		0.5%urea and 0.5% ammonium sulphate
<b>Cowpea</b>	Water soaked necrotic spots on leaf surface. Root growth very much restricted in 10-12 days old seedling	Spray containing sulphate and zinc sulphate 0.1% and 0.1% urea
<b>Groundnut</b>	Chlorosis of terminal leaves	0.5% ferrous sulphate and urea 1%

## 10. PHOTOSYNTHESIS

*Photosynthesis is a vital physiological process where in the chloroplast of green plants synthesizes sugars by using water and carbon dioxide in the presence of light.*

Photosynthesis literally means *synthesis with the help of light* i.e. plant synthesize organic matter (carbohydrates) in the presence of light.

Photosynthesis is sometimes called as carbon assimilation (assimilation: absorption into the system). This is represented by the following traditional equation.



During the process of photosynthesis, the light energy is converted into chemical energy and is stored in the organic matter, which is usually the carbohydrate. One molecule of glucose for instance, contains about 686 K Calories energy. CO<sub>2</sub> and water constitute the raw material for this process and oxygen and water are formed as the by products during photosynthesis. *Stephen Hales* (1727) first explained the relationship between sunlight and leaves and *Sachs* (1887) established that starch was the visible product of photosynthesis.




FIGURE 7.2 Electromagnetic spectrum. Wavelength ( $\lambda$ ) and frequency ( $\nu$ ) are inversely related. Our eyes are sensitive to only a narrow range of wavelengths of radiation, the visible region, which extends from about 400 nm (violet) to about 700 nm (red). Short-wavelength (high-frequency) light has a high energy content; long-wavelength (low-frequency) light has a low energy content.

### Photosynthetic apparatus

The chloroplast in green plants constitutes the photosynthetic apparatus. In higher plants, the chloroplast is discoid in shape, 4-6  $\mu$  in length and 1-2 $\mu$  thick. The chloroplast is bounded by two unit membranes of approximately 50 $\text{\AA}$  thickness and consists of lipids and


proteins. The thickness of the two membranes including the space enclosed by them is approximately 300°A (1 Angstrom: 0.1 cm).

Internally, the chloroplast is filled with a hydrophilic matrix called as *stroma* embedded with *grana*. Each grana consists of 5-25 disk shaped grana lamellae (thylakoid) placed one above the other like the stack of coins. Each grana lamella of thylakoid encloses a space called *loculus* and the thylakoid membrane consists of alternating layer of lipids and proteins. Some of the grana lamella of thylakoid of grana are connected with thylakoid of other grana by somewhat thinner *stroma lamella or fret membrane*. Chlorophyll and other photosynthetic pigments are confined to grana. The chlorophylls are the site of photochemical reactions.

### Photosynthetic pigments

Photosynthetic pigments are of three types; Chlorophylls, Carotenoids and Phycobilins.

- Chlorophylls and Carotenoids are insoluble in water and can be extracted only with organic solvents such as acetone, petroleum ether and alcohol.
- Phycobilins are soluble in water
- Carotenoids include carotenes and xanthophylls. The xanthophylls are also called as *carotenols*.



### **Chlorophylls (green pigments)**

Chlorophylls are magnesium porphyrin compounds. The porphyrin ring consists of four pyrrol rings joined together by CH bridges. Long chain C atoms called as phytol chain is attached to porphyrin ring at pyrrol ring IV.

The chemical structure of chlorophyll *a* and chlorophyll *b* are well established. The molecular formula for chlorophyll *a*:  $C_{55}H_{72}O_5N_4$  Mg and chlorophyll *b*:  $C_{55}H_{70}O_6N_4$  Mg. Both of them consist of Mg porphyrin head which is hydrophilic and a phytol tail which is

lipophilic. The two chlorophylls differ because in chlorophyll *b* there is a –CHO group instead of CH<sub>3</sub> group at the 3<sup>rd</sup> C atom in pyrrol ring II.

Chlorophyll is formed from protochlorophyll in light. The protochlorophyll lacks 2H atoms one each at 7<sup>th</sup> and 8<sup>th</sup> C atoms in pyrrol ring IV.

### **Carotenoids (yellow or orange pigments)**

1. Carotenes: Carotenes are hydrocarbons with a molecular formula C<sub>40</sub>H<sub>56</sub>
2. Xanthophylls (carotenols)

They are similar to carotenes but differ in having two oxygen atoms in the form of hydroxyl or carboxyl group. The molecular formula is C<sub>40</sub>H<sub>56</sub>O<sub>2</sub>. The role of Carotenoids is absorption of light energy and transfer the light energy to chlorophyll *a* molecules. They also play a very important role in preventing photodynamic damage within the photosynthetic apparatus. Photodynamic damage is caused by O<sub>2</sub> molecules which is very reactive and is capable of oxidizing whole range of organic compounds such as chlorophylls and thereby making them unfit for their normal physiological function.

### **Phycobilins (red and blue pigments)**

These also contain four pyrrol rings but lack Mg and the phytol chain.

### **Location of photosynthetic pigments in chloroplast**

The photosynthetic pigments are located in grana portions of the chloroplast. They are present in the thylakoid membrane or membrane of grana lamella. The membrane of thylakoid is made up of proteins and lipids or the membrane consists of both lipid layer and protein layer. The hydrophilic *heads* of the chlorophyll molecules remain embedded in the protein layer while lipophilic phytol tail in the lipid layer. The other pigments are thought to be present along with chlorophyll molecules.


### **Distribution of photosynthetic pigments in plant kingdom**

Pigments	Distribution in plant kingdom
<b>Chlorophylls</b>	

Chlorophyll <i>a</i>	All photosynthesizing plants except bacteria
Chlorophyll <i>b</i>	Higher plants and green algae
Chlorophyll <i>c</i>	Diatoms and brown algae
Chlorophyll <i>d</i>	Red algae
Bacteria chlorophylls <i>a, b, c, d &amp; e</i>	Purple and green bacteria
<b>Carotenoids</b>	
Carotenes ( $\alpha$ and $\beta$ )	Higher plants and algae
Xanthophylls	Higher plants and algae
Lutein	Green leaves and Green and Red algae
Violaxanthin	Green leaves
Fucoxanthin	Brown algae
<b>Phycobilins</b>	
Phycocyanins	Blue green algae and red algae
Phycoerythrins	Blue green algae and Red algae
Allophycocyanin	Blue – green and Red algae

### Light

The chief source of light energy for photosynthesis is sun. The solar radiation or solar energy passes through the space and reaches the earth in the form of *electromagnetic radiation* with waves of varying lengths. The various portions of electromagnetic spectrum are gamma rays, ultraviolet rays, visible rays and infrared rays. The wavelength of these rays ranges from 280 nm to 1000 nm.




- |              |                                      |
|--------------|--------------------------------------|
| Below 280 nm | - X rays, Gamma rays and Cosmic rays |
| 280-390 nm   | - Ultra violet radiation             |
| 400-510 nm   | - Blue light                         |
| 510-610 nm   | - Green light                        |
| 610-700 nm   | - Red light                          |
| 700-1000 nm  | - Far red light (IR)                 |
- Visible light (PAR)  
(VIBGYOR)

Photosynthetic pigments absorb light energy only in the visible part of the spectrum. The earth receives only about 40% (or about  $5 \times 10^{20}$  K cal) of the total solar energy. The rest is either absorbed by the atmosphere or scattered into the space. Only about 1% of the total solar energy received by the earth is absorbed by the pigments and utilized in photosynthesis.

### Absorption spectra of chlorophyll

The absorption of different wavelengths of light by a particular pigment is called *absorption spectrum*. Chlorophylls absorb maximum light in the violet blue and red part of the spectrum. The absorption peaks of chlorophyll *a* are 410 and 660; for chlorophyll *b* 452 and 642. Carotenoids absorb light energy in blue and blue green part of the spectrum.



### **Transfer of light energy absorbed by accessory pigments to chlorophyll *a***

All pigments except chlorophyll *a* are called as *accessory pigments or antenna pigments*. The light energy absorbed by accessory pigments is transferred to chlorophyll *a* molecule. The transfer of light energy from accessory pigments to chlorophyll *a* is called as *resonance or Forster transfer* and takes part in primary photochemical reaction in photosynthesis. Chlorophyll *a* molecules also absorb light energy directly. As a result of absorbing the light energy, the chlorophyll molecule gets *excited*.

### **Excited states of atoms or molecules (*fluorescence and phosphorescence*)**

The normal state of the chlorophyll molecule or atom is called as *ground state or singlet state*. When an electron of a molecule or an atom absorbs a quantum of light, it is raised to a higher energy level which is called as *excited second singlet state*. This state is unstable and has a life time of  $10^{-12}$  seconds.

The electron comes to the next higher energy level by the loss of some of its extra energy in the form of heat. This higher energy level is called as *excited first singlet state* and is also unstable with a half life of  $10^{-9}$  seconds. From the first singlet state, the excited electron may return to the ground state in two ways viz., either losing its remaining extra energy in the form of heat or in the form of radiant energy. The second process is called *fluorescence*. The chlorophyll molecules exit the extra energy in the form of fluorescent light when they are exposed to incident light. Fluorescent light is of longer wavelength than the incident light.

The excited molecule or the atom may also lose its excitation energy by internal conversion and comes to another excited state called as *triplet state* which is meta stable with a half life of  $10^{-3}$  seconds. From the triplet state, the excited molecule or the atom may return to the ground state in three ways.


- (i) By losing its remaining extra energy in the form of heat
- (ii) By losing extra energy in the form of radiant energy (*phosphorescence*) and the chlorophyll molecules emit phosphorescent light even after the incident radiant light is cut off. The phosphorescent light is of longer wavelength than incident light and also fluorescent light.

- (iii) Electrons carrying the extra energy may be expelled from the molecule and is consumed in some further photochemical reaction and the fresh normal electron returns to the molecule.

### Quantum requirement and quantum yield

Light rays consist of tiny particles called *photons* and the energy carried by a photon is called *quantum*. The number of photons (quantum) required to release one molecule of oxygen in photosynthesis is called *quantum requirement*. On the other hand, the number of oxygen molecules released per photon of light in photosynthesis is called as *quantum yield*. The quantum yield is always in fraction of one.

Warburg found minimum quantum requirement for photosynthesis as four. It is because the reduction of one molecule of CO<sub>2</sub> by two molecules of H<sub>2</sub>O requires the transfer of 4H atoms. The transfer of each H atoms from H<sub>2</sub>O to CO<sub>2</sub> requires one photon or quantum of light.



(CH<sub>2</sub>O) in the above equation represent 1/6 of the carbohydrate molecule such as glucose. One molecule of glucose contains 686 K. cal of energy. Therefore, 1/6 glucose molecule contains 686/6 i.e., approximately 112 K.cal energy. It is also known that the rate of photosynthesis is maximum at red light and each photon of red light contains about 40 K cal. of energy. This would suggest that the efficiency with which the plants can convert light energy into chemical energy is 112 / 40 x 4: 70%, which indeed is very high.

According to Emerson and his coworkers, photosynthesis is a very complicated process and is not so efficient to convert all the light energy into chemical energy. There is a considerable loss of light energy absorbed during photosynthesis and therefore the minimum quantum requirement for photosynthesis as suggested by Emerson and coworkers are 8-10. Considering that the quantum requirement for photosynthesis is 8-10, the quantum yield would accordingly be 1/8 to 1/10 (0.125 to 0.10)

## Mechanism of Photosynthesis

### Photo systems (Two pigment systems)

The discovery of red drop and the Emerson's enhancement effect led the scientists to suggest that photosynthesis is driven by two photochemical processes. These processes are associated with two groups of photosynthetic pigments called as *pigment system I* and *pigment system II*. Wavelength of light shorter than 680 nm affect both the pigments systems while wavelength longer than 680 nm affect only pigment system I.

In green plants, pigment system I contains chlorophyll *a*, *b* and carotene. In this pigment system, a very small amount of chlorophyll *a* absorbing light at 700 nm, known as P700 however constitutes the reaction centre of *photosystem I*.

The pigment system II contains chlorophyll *b* and some forms of chlorophyll *a* (such as chlorophyll *a* 662, chlorophyll *a* 677 and chlorophyll *a* 679) and xanthophylls. A very small amount of special form of chlorophyll called P680 constitute the reaction centre of pigment system II. Carotenoids are present in both the pigment systems

The two pigment systems I and II are interconnected by a protein complex called cytochrome b<sub>6</sub>-f complex. The other intermediate components of electron transport chain viz., plastoquinone (PQ) and plastocyanin (PC) act as mobile electron carriers between the complex and either of the two pigment systems. The light energy absorbed by other pigment is ultimately trapped by P700 and P680 forms of chlorophyll *a* which alone take part in further photochemical reaction.

Pigment system I (PSI) complex consists of 200 chlorophylls, 50 Carotenoids and a molecule of chlorophyll *a* absorbing light at 700 nm(P700) and this constitute the reaction centre of photosystem I. Pigment system II (PSII) complex consists of 200 chlorophylls, 50 Carotenoids and a mole of chlorophyll *a* absorbing light at 680 nm, called P 680 at the centre. This constitutes the reaction centre of pigment system II.

### Photosynthetic units – The Quantasomes

Emerson and Arnold (1932) showed that about 2500 chlorophyll molecules are require fixing one molecule of CO<sub>2</sub> in photosynthesis. This number of chlorophyll molecules was called the *chlorophyll unit* but the name was subsequently changed to *photosynthetic*

*unit.* However, since the reduction or fixation of one  $\text{CO}_2$  molecule requires about 10 quanta of light, it is assured that 10 flashes of light are required to yield one  $\text{O}_2$  molecule or reduction of one molecule of  $\text{CO}_2$ . Thus each individual unit would contain 1/10<sup>th</sup> of 2500 i.e., 250 molecules.

### Action spectrum

The pigments present in plants or any living organism have the ability to absorb radiant energy to carry out photo physiological reactions. It is difficult to decide which specific pigment is actually associated with the particular photochemical reactions. Hence, a common procedure to identify the pigment involved in a particular photoreaction is to determine the action spectrum i.e. measuring the rate of the particular photoreaction.

Once the action spectrum for a photo physiological reaction is determined, the next step is to compare this action spectrum with absorption spectrum of a pigment.

Two pigments, A and B were isolated from the same plant and their absorption spectra were determined. Pigment A has a peak in absorption at 395 nm and the pigment B at 660 nm. The close correspondence between the absorption spectrum and the action spectrum of pigment B strongly supports that Pigment B is responsible for absorbing radiant energy to drive this photoreaction.

### Mechanism of photosynthesis

The biosynthesis of glucose by the chloroplast of green plants using water and  $\text{CO}_2$  in the presence of light is called photosynthesis. Photosynthesis is a complex process of synthesis of organic food materials. It is a complicated oxidation- reduction process where water is oxidized and  $\text{CO}_2$  is reduced to carbohydrates. The mechanism of photosynthesis consists of two parts.

1. Light reaction / Primary photochemical reaction / Hill's reaction/ Arnon's cycle
2. Dark reaction / Black man's reaction / Path of carbon in photosynthesis.

#### **1. Light reaction or Primary photochemical reaction or Hill's reaction**

In light reaction, ATP and  $\text{NADPH}_2$  are produced and in the dark reaction,  $\text{CO}_2$  is reduced with the help of ATP and  $\text{NADPH}_2$  to produce glucose. The light reaction is called primary photochemical reaction as it is induced by light. Light reaction is also called as Hill's

reaction as Hill proved that chloroplast produce  $O_2$  from water in the presence of light. It is also called as Arnon's cycle because Arnon showed that the  $H^+$  ions released by the breakdown of water are used to reduce the coenzyme NADP to NADPH. Light reaction includes photophosphorylation as ATP is synthesized in the presence of light. The reaction takes place only in the presence of light in *grana* portion of the chloroplast and it is faster than dark reaction. The chlorophyll absorbs the light energy and hence the chlorophyll is called as *photosystem* or *pigment system*. Chlorophylls are of different types and they absorb different wavelengths of light. Accordingly, chlorophylls exist in two photo systems, Photosystem I (PSI) and Photosystem II (PS II). Both photo systems are affected by light with wavelengths shorter than 680nm, while PS I is affected by light with wavelengths longer than 680nm.

### The components of photo systems

Photosystem I	Photosystem II
Chlorophyll <i>a</i> 670	Chlorophyll <i>a</i> 660
Chlorophyll <i>a</i> 680	Chlorophyll <i>a</i> 670
Chlorophyll <i>a</i> 695	Chlorophyll <i>a</i> 680 or P680
Chlorophyll <i>a</i> 700 or P700	Chlorophyll <i>b</i>
Chlorophyll <i>b</i>	Phycobilins
Carotenoids	Xanthophylls
P700 form of Chlorophyll <i>a</i> is the active reaction centre	P680 form of Chlorophyll <i>a</i> is the active reaction centre

The light reaction can be studied under the following headings.

#### i. Absorption of light energy by chloroplast pigments

Different chloroplast pigments absorb light in different regions of the visible part of the spectrum.


#### ii. Transfer of light energy from accessory pigments to chlorophyll *a*

All the photosynthetic pigments except chlorophyll *a* are called as accessory or antenna pigments. The light energy absorbed by the accessory pigments is transferred by resonance to chlorophyll *a* which alone can take part in photochemical reaction. Chlorophyll

*a* molecule can also absorb the light energy directly. In pigment system I, the photoreaction centre is P700 and in pigment system II, it is P680.


### iii. Activation of chlorophyll molecule by photon of light

When P700 or P680 forms of chlorophyll *a* receives a photon (quantum) of light, becomes an excited molecule having more energy than the ground state energy. After passing through the unstable second singlet state and first singlet stage the chlorophyll molecules comes to the meta stable triplet state. This excited state of chlorophyll molecule takes part further in primary photochemical reaction i.e. the electron is expelled from the chlorophyll *a* molecule.



### iv. Photolysis of water and O<sub>2</sub> evolution (oxidation of water)


These processes are associated with pigment system II and are catalyzed by Mn<sup>++</sup> and Cl<sup>-</sup> ions. When pigment system II is active i.e it receives the light, the water molecules split into OH<sup>-</sup> and H<sup>+</sup> ions (*Photolysis of water*). The OH<sup>-</sup> ions unite to form some water molecules again and release O<sub>2</sub> and electrons.



### v. Electron transport and production of assimilatory powers (NADPH<sub>2</sub> and ATP)

It has already been observed that when chlorophyll molecule receives the photon of light, an electron is expelled from the chlorophyll *a* molecule along with extra energy. This electron after traveling through a number of electron carriers is utilized for the production of NADPH<sub>2</sub> from NADP and also utilized for the formation of ATP molecules from ADP and inorganic phosphate (Pi). The transfer of electrons through a series of coenzymes is called *electron transport* and the process of formation of ATP from ADP and Pi using the energy of electron transport is called as *photosynthetic phosphorylation or photophosphorylation*. The types of Phosphorylation include *cyclic and non- cyclic*.

## Cyclic electron transport and cyclic photophosphorylation




The electrons released from photosystem I goes through a series of coenzymes and returns back to the same photosystem I. This electron transport is called *cyclic electron transport*. The synthesis of ATP occurring in cyclic electron transport is called *cyclic photophosphorylation*. The cyclic electron transport involves only pigment system I. This situation is created when the activity of pigment system II is blocked. Under this condition,

1. Only pigment system I remain active
2. Photolysis of water does not take place
3. Blockage of noncyclic ATP formation and this causes a drop in CO<sub>2</sub> assimilation in dark reaction
4. There is a consequent shortage of oxidized NADP

Thus, when P700 molecule is excited in pigment system I by absorbing a photon (quantum) of light, the ejected electron is captured by ferredoxin via FRS. From ferredoxin, the electrons are not used up for reducing NADP to NADPH + H<sup>+</sup> but ultimately it falls back

to the P700 molecule via number of other intermediate electron carriers. The electron carriers are probably cytochrome  $b_6$ , cytochrome  $f$  and plastocyanin.

During this electron transport, phosphorylation of ADP molecule to form ATP molecule take place at two places i.e., between ferredoxin and cytochrome  $b_6$  and between cytochrome  $b_6$  and cytochrome  $f$ . Thus, two ATP molecules are produced in this cycle. Since the electron ejected from P700 molecule is cycled back, the process has been called as *cyclic electron transport* and the accompanying phosphorylation as the *cyclic photophosphorylation*.



### Significance of cyclic photophosphorylation

1. During cyclic electron transport and phosphorylation, photolysis of water, O<sub>2</sub> evolution and reduction of NADP do not take place.
2. The electron returns or cycles back to original position in the P700 form of chlorophyll *a*. Here, chlorophyll molecule serves both as donor and acceptor of the electron.
3. It generates energy rich ATP molecules at two sites and as such cannot drive dark reactions of photosynthesis

On the other hand, non- cyclic photophosphorylation does not produce sufficient ATP in relation to NADPH to operate the dark phase of photosynthesis. Therefore, the deficiency of ATP molecule in non-cyclic photophosphorylation is made up by the operations of cyclic photophosphorylation.

Secondly, the cyclic photophosphorylation may be an important process in providing ATP for photosynthesis and other processes such as synthesis of starch, proteins, lipids, nucleic acids and pigments within the chloroplast.

### Non cyclic photophosphorylation

The electron released from photosystem II goes through a series of enzymes and Co-enzymes to photosystem I. This is called non cyclic electron transport and the Synthesis of ATP in non cyclic electron transport is called non- cyclic photo phosphorylation. The main function of non cyclic electron transport is to produce the assimilatory powers such as NADPH<sub>2</sub> and ATP and the process occurs in photosystem I and II.

This process of electron transport is initiated by the absorption of a photon (quantum) of light by P680 form of chlorophyll *a* molecule in the pigment system II, which gets excited and an electron is ejected from it so that an electron deficiency or a hole is left behind in the P680 molecule.

The ejected electron is trapped by an unknown compound known as Q. From Q, the electron passes downhill along a series of compounds or intermediated electron carriers such as cytochrome *b*<sub>6</sub>, plastoquinone, cytochrome *f* and a copper containing plastocyanin and ultimately received by pigment system I. At one place during electron transport i.e.

between plastoquinone and cytochrome *f*, one molecule of ATP is formed from ADP and inorganic phosphate.

Now, when a photon of light is absorbed by P700 form of chlorophyll molecule in the pigment system I, this gets excited and an electron is ejected from it. This ejected electron is trapped by FRS (Ferredoxin Reducing Substance) and it is then transferred to a non-heme iron protein called ferredoxin. From ferredoxin, electron is transferred to NADP so that NADP is reduced to  $\text{NADPH} + \text{H}^+$

The hole in pigment system I has been filled by electron coming from pigment system II. But, the hole or an electron deficiency in pigment system II is filled up by the electron coming from photolysis of water where, water acts as electron donor.

In this scheme of electron transport, the electron ejected from pigment system II did not return to its place of origin, instead it is taken up by pigment system I. Similarly, the electron ejected from pigment system I did not cycle back and was consumed in reducing NADP. Therefore, this electron transport has been called as *non-cyclic electron transport* and accompanying phosphorylation as *non-cyclic photophosphorylation*.

The non cyclic electron transport (photophosphorylation) takes the shape of Z and hence it is called by the name Z-scheme. Non cyclic photophosphorylation and  $\text{O}_2$  evolution are inhibited by CMU (3-(4'-Chlorophyl) – 1-1dimethyl urea and 3-(3-4-dichlorophenyl)-1, 1-dimethyl urea (DCMU).

### **Significance of non cyclic electron transport**

1. It involves PS I and PSII
2. The electron expelled from P680 of PSII is transferred to PS I and hence it is a non cyclic electron transport.
3. In non cyclic electron transport, photolysis of water (Hill's reaction and evolution of  $\text{O}_2$ ) takes place.
4. Phosphorylation (synthesis of ATP molecules) takes place at only one place.
5. The electron released during photolysis of water is transferred to PS II.
6. The hydrogen ions ( $\text{H}^+$ ) released from water are accepted by NADP and it becomes  $\text{NADPH}_2$

7. At the end of non cyclic electron transport, energy rich ATP, assimilatory power NADPH<sub>2</sub> and oxygen from photolysis of water are observed.
8. The ATP and NADPH<sub>2</sub> are essential for the dark reaction wherein, reduction of CO<sub>2</sub> to carbohydrate takes place.

**Comparison of cyclic and non cyclic electron transport and photophosphorylation in chloroplasts**

<b>Cyclic electron transport and photo phosphorylation</b>		<b>Non cyclic electron transport and photo phosphorylation</b>
1	Associated with pigment system I	Associated with pigment system I and II
2	The electron expelled from chlorophyll molecule is cycled back	The electron expelled from chlorophyll molecule is not cycled back. But, its loss is compensated by electron coming from photolysis of water
3	Photolysis of water and evolution of O <sub>2</sub> do not take place	Photolysis of water and evolution of O <sub>2</sub> take place
4	Phosphorylation takes place at two places	Phosphorylation takes place at only one place
5	NADP <sup>+</sup> is not reduced	NADP <sup>+</sup> is reduced to NADPH <sup>+</sup> + H <sup>+</sup>

**Significance of light reaction**

1. Light reaction takes place in chlorophyll in the presence of light.
2. During light reaction, the assimilatory powers ATP and NADPH<sub>2</sub> are synthesized.
3. The assimilatory powers are used in dark reaction for the conversion of CO<sub>2</sub> into sugars.
4. Photolysis of water occurs in light reaction. The H<sup>+</sup> ions released from water are used for the synthesis of NADPH<sub>2</sub>
5. Plants release O<sub>2</sub> during light reaction

**Red drop and Emerson's enhancement effect**

Robert Emerson noticed a sharp decrease in quantum yield at wavelength greater than 680 nm, while determining the quantum yield of photosynthesis in *chlorella* using

monochromatic light of different wavelengths. Since this decrease in quantum yield took place in the red part of the spectrum, the phenomenon was called as *red drop*.

Later, they found that the inefficient far-red light beyond 680 nm could be made fully efficient if supplemented with light of shorter wavelength (blue light). The quantum yield from the two combined beams of light was found to be greater than the sum effects of both beams used separately. This enhancement of photosynthesis is called as *Emerson's Enhancement*.

## 11. PHOTOSYNTHETIC PATHWAYS - C<sub>3</sub>, C<sub>4</sub> AND CAM

### Dark reaction or Blackman's reaction or Path of carbon in photosynthesis


This is the second step in the mechanism of photosynthesis. The chemical processes of photosynthesis occurring independent of light is called *dark reaction*. It takes place in the stroma of chloroplast. The dark reaction is purely enzymatic and it is slower than the light reaction. The dark reactions occur also in the presence of light. In dark reaction, the sugars are synthesized from CO<sub>2</sub>. The energy poor CO<sub>2</sub> is fixed to energy rich carbohydrates using the energy rich compound, ATP and the assimilatory power, NADPH<sub>2</sub> of light reaction. The process is called carbon fixation or carbon assimilation. Since Blackman demonstrated the existence of dark reaction, the reaction is also called as *Blackman's reaction*. In dark reaction two types of cyclic reactions occur

1. Calvin cycle or C<sub>3</sub> cycle
2. Hatch and Slack pathway or C<sub>4</sub> cycle

### Calvin cycle or C<sub>3</sub> cycle


It is a cyclic reaction occurring in the dark phase of photosynthesis. In this reaction, CO<sub>2</sub> is converted into sugars and hence it is a process of carbon fixation. The Calvin cycle was first observed by Melvin Calvin in chlorella, unicellular green algae. Calvin was awarded Nobel Prize for this work in 1961. Since the first stable compound in Calvin cycle is a 3 carbon compound (3 phosphoglyceric acid), the cycle is also called as C<sub>3</sub> cycle. The reactions of Calvin's cycle occur in three phases.

1. Carboxylative phase
2. Reductive phase
3. Regenerative phase




### 1. Carboxylative phase

Three molecules of  $\text{CO}_2$  are accepted by 3 molecules of 5C compound viz., ribulose diphosphate to form three molecules of an unstable intermediate 6C compound. This reaction is catalyzed by the enzyme, carboxy dismutase




The three molecules of the unstable 6 carbon compound are converted by the addition of 3 molecules of water into six molecules of 3 phosphoglyceric acid. This reaction is also catalyzed by the enzyme carboxy mutase.




3 phosphoglyceric acid (PGA) is the first stable product of dark reaction of photosynthesis and since it is a 3 carbon compound, this cycle is known as C3 cycle.

## 2. Reductive phase

Six molecules of 3PGA are phosphorylated by 6 molecules of ATP (produced in the light reaction) to yield 6 molecules of 1-3 diphospho glyceric acid and 6 molecules of ADP. This reaction is catalyzed by the enzyme, Kinase





Six molecules of 1, 3 diphosphoglyceric acid are reduced with the use of 6 molecules of NADPH<sub>2</sub> (produced in light reaction) to form 6 molecules of 3 phospho glyceraldehyde. This reaction is catalysed by the enzyme, triose phosphate dehydrogenase.




## 3. Regenerative phase

In the regenerative phase, the ribose diphosphate is regenerated. The regenerative phase is called as *pentose phosphate pathway* or *hexose monophosphate shunt*. It involves the following steps.

- Some of the molecules of 3 phospho glyceraldehyde into dihydroxy acetone phosphate. Both 3 phospho glyceraldehyde and dihydroxy acetone phosphate then unite in the presence of the enzyme, aldolase to form fructose, 1-6 diphosphate.




- Fructose 6 phosphate is converted into fructose 6 phosphate in the presence of phosphorylase




- Some of the molecules of 3 phospho glyceraldehyde instead of forming hexose sugars are diverted to regenerate ribulose 1-5 diphosphate




- 3 phospho glyceraldehyde reacts with fructose 6 phosphate in the presence of enzyme transketolase to form erythrose 4 phosphate (4C sugar) and xylulose 5 phosphate(5C sugar)




5. Erythrose 4 phosphate combines with dihydroxy acetone phosphate in the presence of the enzyme aldolase to form sedoheptulose 1,7 diphosphate(7C sugar)





6. Sedoheptulose 1, 7 diphosphate loses one phosphate group in the presence of the enzyme phosphatase to form sedoheptulose 7 phosphate.




7. Sedoheptulose phosphate reacts with 3 phospho glyceraldehyde in the presence of transketolase to form xylulose 5 phosphate and ribose 5 phosphate ( both % c sugars)




8. Ribose 5 phosphate is converted into ribulose 1, 5 diphosphate in the presence of enzyme, phosphopentose kinase and ATP. Two molecules of xylulose phosphate are also converted into one molecule of ribulose monophosphate. The ribulose monophosphate is phosphorylated by ATP to form ribulose diphosphate and ADP, thus completing Calvin cycle.



In the dark reaction,  $\text{CO}_2$  is fixed to carbohydrates and the  $\text{CO}_2$  acceptor ribulose diphosphate is regenerated. In Calvin cycle, 12 NADPH<sub>2</sub> and 18 ATPs are required to fix 6  $\text{CO}_2$  molecules into one hexose sugar molecule (fructose 6 phosphate).





Schematic diagram of light reaction and Calvin cycle

### C4 cycle or Hatch and Slack pathway

It is the alternate pathway of C3 cycle to fix CO<sub>2</sub>. In this cycle, the first formed stable compound is a 4 carbon compound viz., oxaloacetic acid. Hence it is called C4 cycle. The path way is also called as Hatch and Slack as they worked out the pathway in 1966 and it is also called as C4 dicarboxylic acid pathway. This pathway is commonly seen in many grasses, sugar cane, maize, sorghum and amaranthus.


The C4 plants show a different type of leaf anatomy. The chloroplasts are dimorphic in nature. In the leaves of these plants, the vascular bundles are surrounded by bundle sheath of larger parenchymatous cells. These bundle sheath cells have chloroplasts. These chloroplasts of bundle sheath are larger, lack grana and contain starch grains. The chloroplasts in mesophyll cells are smaller and always contain grana. This peculiar anatomy of leaves of C4 plants is called Kranz anatomy. The bundle sheath cells are bigger and look like a ring or wreath. Kranz in German means wreath and hence it is called Kranz anatomy. The C4 cycle involves two carboxylation reactions, one taking place in chloroplasts of mesophyll cells and another in chloroplasts of bundle sheath cells. There are four steps in Hatch and Slack cycle:

1. Carboxylation
2. Breakdown

3. Splitting
4. Phosphorylation


### 1. Carboxylation

It takes place in the chloroplasts of mesophyll cells. Phosphoenolpyruvate, a 3 carbon compound picks up CO<sub>2</sub> and changes into 4 carbon oxaloacetate in the presence of water. This reaction is catalysed by the enzyme, phosphoenol pyruvate carboxylase.




### 2. Breakdown

Oxaloacetate breaks down readily into 4 carbon malate and aspartate in the presence of the enzyme, transaminase and malate dehydrogenase.



### 3. Splitting


In the sheath cells, malate and aspartate split enzymatically to yield free CO<sub>2</sub> and 3 carbon pyruvate. The CO<sub>2</sub> is used in Calvin's cycle in the sheath cell.




The second Carboxylation occurs in the chloroplast of bundle sheath cells. The CO<sub>2</sub> is accepted by 5 carbon compound ribulose diphosphate in the presence of the enzyme, carboxy dismutase and ultimately yields 3 phosphoglyceric acid. Some of the 3 phosphoglyceric acid is utilized in the formation of sugars and the rest regenerate ribulose diphosphate.

### 4. Phosphorylation

The pyruvate molecule is transferred to chloroplasts of mesophyll cells where, it is phosphorylated to regenerate phosphoenol pyruvate in the presence of ATP. This reaction is catalysed by pyruvate phosphokinase and the phosphoenol pyruvate is regenerated.



In the Kranz anatomy of the leaves, the C3 and C4 cycles of carboxylation are linked and this is due to the Kranz anatomy of the leaves. The C4 plants are more efficient in photosynthesis than the C3 plants. The enzyme, phosphoenol pyruvate carboxylase of the C4 cycle is found to have more affinity for CO<sub>2</sub> than the ribulose diphosphate carboxylase of the C3 cycle in fixing the molecular CO<sub>2</sub> in organic compound during Carboxylation.



### Crassulacean Acid Metabolism (CAM) cycle or the dark fixation of CO<sub>2</sub> in succulents

CAM is a cyclic reaction occurring in the dark phase of photosynthesis in the plants of Crassulaceae. It is a CO<sub>2</sub> fixation process wherein, the first product is malic acid. It is the third alternate pathway of Calvin cycle, occurring in mesophyll cells. The plants exhibiting CAM cycle are called CAM plants. Most of the CAM plants are succulents e.g., Bryophyllum, Kalanchoe, Crassula, Sedium, Kleinia etc. It is also seen in certain plants of Cactus e.g. Opuntia, Orchid and Pine apple families.

CAM plants are usually succulents and they grow under extremely xeric conditions. In these plants, the leaves are succulent or fleshy. The mesophyll cells have larger number of chloroplasts and the vascular bundles are not surrounded by well defined bundle sheath cells. In these plants, the stomata remain open during night and closed during day time. The CAM plants are adapted to photosynthesis and survival under adverse xeric conditions. CAM plants are not as efficient as C<sub>4</sub> plants in photosynthesis. But they are better suited to conditions of extreme desiccation.

CAM involves two steps:


1. Acidification
2. Deacidification

#### Acidification

In darkness, the stored carbohydrates are converted into phosphoenol pyruvic acid by the process of Glycolysis. The stomata in CAM plants are open in dark and they allow free diffusion of  $\text{CO}_2$  from the atmosphere into the leaf. Now, the phosphoenolpyruvic acid carboxylated by the enzyme phosphoenol pyruvic acid carboxylase and is converted in to oxaloacetic acid.




The oxaloacetic acid is then reduced to malic acid in the presence of the enzyme malic dehydrogenase. The reaction requires  $\text{NADPH}_2$  produced in Glycolysis.



The malic acid produced in dark is stored in the vacuole. The malic acid increases the acidity of the tissues.

### Deacidification


During day time, when the stomata are closed, the malic acid is decarboxylated to produce pyruvic acid and evolve carbon dioxide in the presence of the malic enzyme. When the malic acid is removed, the acidity decreases the cells. This is called deacidification. One molecule of  $\text{NADP}^+$  is reduced in this reaction.



The pyruvic acid may be oxidized to  $\text{CO}_2$  by the pathway of Kreb's cycle or it may be reconverted to phosphoenol pyruvic acid and synthesize sugar by C3 cycle. The  $\text{CO}_2$  released by deacidification of malic acid is accepted by ribulose diphosphate and is fixed to carbohydrate by C3 cycle.

CAM is a most significant pathway in succulent plants. The stomata are closed during day time to avoid transpiration loss of water. As the stomata are closed,  $\text{CO}_2$  cannot enter into the leaves from the atmosphere. However, they can carry out photosynthesis during the day time with the help of  $\text{CO}_2$  released from organic acids. During night time, organic acids are synthesized in plenty with the help of  $\text{CO}_2$  released in respiration and the  $\text{CO}_2$  entering from

the atmosphere through the open stomata. Thus, the CO<sub>2</sub> in dark acts as survival value to these plants.



### Comparison of the plants of C<sub>3</sub> and C<sub>4</sub> cycle

	C <sub>3</sub> Plant	C <sub>4</sub> Plant
1.	Only C <sub>3</sub> cycle is found	Both C <sub>4</sub> and C <sub>3</sub> cycles are found.
2.	The efficiency of CO <sub>2</sub> absorption at low concentration is far less and hence, they are less efficient.	The efficiency of CO <sub>2</sub> absorption at low concentration is quite high and hence, they are more efficient plants.
3.	The CO <sub>2</sub> acceptor is Ribulose-1, 5-diphosphate.	The CO <sub>2</sub> acceptor is phospho enol pyruvate.
4.	The first stable product is phospho glyceric acid (PGA).	Oxaloacetate (OAA) is the first stable product.
5.	Plants show one type of chloroplast (monomorphic type).	Plants show dimorphic type of chloroplast. The chloroplast of parenchymatous bundle sheath is different from that of mesophyll cells (dimorphic type). The chloroplasts in bundle sheath cell are centripetally

		arranged and lack grana. Leaves show <i>Kranz type</i> of anatomy.
6.	In each chloroplast, two pigment systems (Photosystem I and II) are present.	In the chloroplasts of bundle sheath cells, the photosystem II is absent. Therefore, these are dependent on mesophyll chloroplasts for the supply of NADPH + H <sup>+</sup> .
7.	The Calvin cycle enzymes are present in mesophyll chloroplast. Thus, the Calvin cycle occurs.	Calvin cycle enzymes are absent in mesophyll chloroplasts. The cycle occurs only in the chloroplasts of bundle sheath cells.
8.	The CO <sub>2</sub> compensation point is 50-150 ppm CO <sub>2</sub> .	The CO <sub>2</sub> compensation point is 0-10 ppm CO <sub>2</sub> .
9.	Photorespiration is present and easily detectable.	Photorespiration is present only to a slight degree or absent.
10.	The CO <sub>2</sub> concentration inside leaf remains high (about 200 ppm).	The CO <sub>2</sub> concentration inside the leaf remains low (about 100 ppm).
11.	The <sup>13</sup> C/ <sup>12</sup> C ratio in C-containing compounds remains relatively low (both <sup>13</sup> CO <sub>2</sub> and <sup>12</sup> CO <sub>2</sub> are present in air).	The ratio is relatively high, i.e. C <sub>4</sub> plants are more enriched with <sup>13</sup> C than C <sub>3</sub> plants.
12.	Net rate of photosynthesis in full sunlight (10,000 – 12,000 ft. c.) is 15-25 mg. of CO <sub>2</sub> per dm <sup>2</sup> of leaf area per hour.	It is 40-80 mg. of CO <sub>2</sub> per dm <sup>2</sup> of leaf area per hour. That is, photosynthetic rate is quite high. The plants are efficient.
13.	The light saturation intensity reaches in the range of 1000-4000 ft. c.	It is difficult to reach saturation even in full sunlight.
14.	Bundle sheath cells are unspecialized.	The bundle sheath cells are highly developed with unusual construction of organelles.
15.	The optimum temperature for the process is 10-25°C.	In these plants, it is 30-45°C and hence, they are warm climate plants. At this temperature, the rate of photosynthesis is double than that is in C <sub>3</sub> plants.

16.	18 ATPs are required to synthesize one glucose molecule.	30 ATPs are required to synthesize one glucose molecule.
-----	--	--

## Factors affecting photosynthesis

### I. External factors

#### 1. Light

It is the most important factor of photosynthesis. Any kind of artificial light such as electric light can induce photosynthesis. Out of the total solar energy, only 1-2 % is used for photosynthesis and the rest is used for other metabolic activities. The effect of light on photosynthesis can be studied under three categories.

##### a. *Light intensity*

Wolkoff (1966) found that the rate of photosynthesis is directly proportional to light intensity. But the extremely high light intensities do not favor for higher photosynthetic rates. The high light intensity which fails to accelerate photosynthesis is called light saturation intensity. Of the light falling on a leaf, about 80 per cent is absorbed, 10 per cent is reflected and 10 % is transmitted. The rate of photosynthesis is greater in intense light than in diffused light. The plants are grouped into two types on the basis of light requirement.

- i. Heliophytes (Sun plants)
- ii. Sciophytes (Shade plants)

At a specific light intensity, the amount of CO<sub>2</sub> used in photosynthesis and the amount of CO<sub>2</sub> released in respiration are volumetrically equal. This specific light intensity is known as *light compensation point*.

At very high light intensity, beyond a certain point, the photosynthetic cells exhibit *photo oxidation*. This phenomenon is called *solarisation* and a result of this, inactivation of chlorophyll molecules, bleaching of chlorophyll molecules and even inactivation of some enzymes take place resulting in the destruction of whole photosynthetic apparatus. In general, low light intensity favours stomatal closure and in turn reduced rate of photosynthesis.

##### b. *Light quality (wavelength)*

Photosynthesis occurs only in the visible part of the light spectrum i.e., between 400 and 700 nm. The maximum rate of photosynthesis occurs at red light followed by blue light.

The green light has minimum effect and photosynthesis cannot take place either in the infrared or in the ultraviolet light.

### *c. Light duration*

In general tropical plants get 10-12 hours of light per day and this longer period of light favours photosynthesis.

## **2. Carbon dioxide**

CO<sub>2</sub> is one of the raw materials required for photosynthesis. If the CO<sub>2</sub> concentration is increased at optimum temperature and light intensity, the rate of photosynthesis increases. But, it is also reported that very high concentration of CO<sub>2</sub> is toxic to plants inhibiting photosynthesis.

## **3. Temperature**

The rate of photosynthesis increases by increase in temperature up to 40 °C and after this, there is reduction in photosynthesis. High temperature results in the denaturation of enzymes and thus, the dark reaction is affected. The temperature requirement for optimum photosynthesis varies with the plant species. For example, photosynthesis stops in many plants at 0 °C but in some conifers, it can occur even at -35 °C. Similarly photosynthesis stops beyond 40-50 °C in certain plants; but certain bacteria and blue green algae can perform photosynthesis even at 70 °C.

## **4. Water**

Water has indirect effect on the rate of photosynthesis although it is one of the raw materials for the process. The amount of water utilized in photosynthesis is quite small and even less than 1 per cent of the water absorbed by a plant. Water rarely acts as a limiting factor for photosynthesis. During water scarcity, the cells become flaccid and the rate of photosynthesis might go down.

## **5. Oxygen**

Oxygen is a byproduct of photosynthesis and an increase in the O<sub>2</sub> concentration in many plants results in a decrease in the rate of photosynthesis. The phenomenon of inhibition of photosynthesis by O<sub>2</sub> was first discovered by Warburg (1920) in green alga Chlorella and this effect is known as Warburg's effect. This is commonly observed in C3 plants.

In plants, there is a close relationship between Warburg's effect and photorespiration. The substrate of photorespiration is glycolate and it is synthesized from some intermediates of Calvin's cycle. In plants that show Warburg's effect, increased O<sub>2</sub> concentration result in diversion of these intermediates of Calvin cycle into the synthesis of glycolate, thereby showing higher rate of photorespiration and lower photosynthetic productivity.

## **6. Mineral elements**

The elements like Mg, Fe, Cu, Cl, Mn, P etc are involved in the key reactions of photosynthesis and hence, the deficiency of any of these nutrients caused reduction in photosynthesis.

## **7. Chlorophyll content**

It is very much essential to trap the light energy. In 1929, Emerson found direct relationship between the chlorophyll content and rate of photosynthesis. In general, the chlorophyll sufficient plants are green in colour showing efficient photosynthesis. The chlorotic leaves due to irregular synthesis of chlorophyll or breakdown of chlorophyll pigment exhibit inefficient photosynthesis.

## **8. Leaf**

The leaf characters such as leaf size, chlorophyll content, number of stomata. Leaf orientation and leaf age are some of the factors that are responsible for photosynthesis. The maximum photosynthetic activity is usually seen in the physiologically functional and full size leaves (usually third/fourth leaf from the tip of the shoot system).

## **9. Carbohydrates**

If the accumulated carbohydrates are not translocated, the photosynthetic rate is reduced and respiration is increased. Sugar is converted into starch and gets accumulated in the chloroplasts. This reduces the effective surface in the chloroplast and the rate of photosynthesis is decreased.

## **10. Phytohormones**

Treharne (1970) reported first that photosynthesis may be regulated by plant hormone system. He found that gibberellic acid and cytokinin increase the carboxylating activity and photosynthetic rates. Meidner (1967) also reported that kinetin @ 3µm causes 12 per cent increase in photosynthesis within one hour of the treatment.

## PHOTORESPIRATION

The excessive respiration that takes place in green cells in the presence of light is called as photorespiration. Decker (1955) discovered the process and it is also called as C2 cycle as the 2 carbon compound glycolic acid acts as the substrate in photorespiration. In general, respiration takes place under both light and dark conditions. However in some plants, the respiration is more in light than in dark. It is 3-5 times higher than the rate of respiration in dark. Photorespiration is carried out only in the presence of light. But the normal respiration is not light dependent and it is called dark respiration.


In photorespiration, temperature and oxygen concentration play an important role. Photorespiration is very high when the temperature is between 25 and 30 °C. The rate of photorespiration increases with the increase in the concentration of oxygen. Three cell organelles namely chloroplast, peroxisome and mitochondria are involved in the photorespiration. This kind of respiration is seen in plants like cotton, pulses, capsicum, peas, tomato, petunia soybean, wheat, oats, paddy, chlorella etc and it is absent in grasses.

### **Mechanism**

1. In the presence of excess oxygen and low CO<sub>2</sub>, ribulose 1,5 diphosphate produced in the chloroplast during photosynthesis is split into 2 phospho glycolic acid and 3 phospho glyceric acid by the enzyme, ribulose 1,5 diphosphate oxygenase.
2. The 3 phospho glyceric acid enters the Calvin cycle.
3. In the next step, phosphate group is removed from 2 phosphoglycolic acid to produce glycolic acid by the enzyme, phosphatase.
4. Glycolic acid then comes out of chloroplast and enters the peroxisome. Here, it combines with oxygen to form glyoxylic acid and hydrogen peroxide. This reaction is catalyzed by the enzyme, glycolic acid oxidase. Hydrogen peroxide is toxic and it is broken down into water and oxygen by the enzyme, Catalase. Photorespiration is an oxidation process. In this process, glycolic acid is converted into carbohydrate and CO<sub>2</sub> is released as the by product. As glycolic acid is oxidized in photorespiration, it is also called as glycolate metabolism.
5. The glyoxylic acid is converted into glycine by the addition of one amino group with the help of the enzyme, amino transferase.

6. Now, the glycine is transported from the peroxisome into the mitochondria. In the mitochondria, two molecules of glycine condense to form serine and liberate carbon dioxide and ammonia.
7. Amino group is removed from serine to form hydroxyl pyruvic acid in the presence of the enzyme, transaminase.
8. Hydroxy pyruvic acid undergoes reduction with the help of NADH to form glyceric acid in the presence of enzyme alpha hydroxyl acid reductase.
9. Finally, regeneration of 3 phosphoglyceric acid occurs by the phosphorylation of glyceric acid with ATP. This reaction is catalyzed by the enzyme, Kinase.
10. The 3 phosphoglyceric acid is an intermediate product of Calvin cycle. If it enters the chloroplast, it is converted into carbohydrate by photosynthesis and it is suppressed nowadays with the increased CO<sub>2</sub> content in the atmosphere.

## PHOTORESPIRATION



### Significance of photorespiration

1. Photorespiration helps in classifying the plants


Generally, photorespiration is found in C3 plants and absent in C4 plants.

2. Carbon dioxide is evolved during the process and it prevents the total depletion of CO<sub>2</sub> in the vicinity of chloroplasts.

2. The process causes oxidation of glycolic acid which arises as an unwanted byproduct of photosynthesis. The glycolic acid after oxidation is converted into carbohydrate but the remainder is converted into CO<sub>2</sub>.
3. Photorespiration uses energy in the form of ATP and reduced nucleotides, but normal respiration yields ATP and reduced nucleotides.
4. It is believed that photorespiration was common in earlier days when CO<sub>2</sub> content was too low to allow higher rates.

## 12. RESPIRATION

The cellular oxidation or break down of carbohydrates into  $\text{CO}_2$  and  $\text{H}_2\text{O}$ , and release of energy is called as *respiration*. It is a reverse process of photosynthesis. In respiration, the oxidation of various organic food substances like carbohydrates, fats, proteins etc, may take place. Among these, glucose is the commonest.




This oxidation process is not so simple and does not take place in one step. Breakdown of glucose involves many steps releasing energy in the form of ATP molecules and also forming a number of carbon compounds (intermediates). Respiration is a vital process that occurs in all living cells of the plant and the most actively respiring regions are floral buds, vegetative buds, germinating seedlings, stem and root apices.

### **Types of respiration**

Degradation of organic food for the purpose of releasing energy can occur with or without the participation of oxygen. Hence, respiration can be classified into two types; aerobic and anaerobic respiration.

### **Aerobic respiration**

Aerobic respiration takes place in the presence of oxygen and the respiratory substrate gets completely oxidized to carbon dioxide and water as end products.



(Glucose)

This type of respiration is of common occurrence and it is often used as a synonym of respiration.

### **Anaerobic respiration**

It takes place in the absence of oxygen and the respiratory substrate is incompletely oxidized. Some other compounds are also formed in addition to carbon dioxide. This type of respiration is of rare occurrence but, common among microorganisms like yeasts.



Glucose      Ethanol

### **Respiratory substrate**

A respiratory substrate is an organic substance which can be degraded to produce energy which is required for various activities of the cell. The respiratory substrates include carbohydrates, fats, organic acids, protein etc.

### ***Carbohydrates***

The carbohydrates constitute the most important respiratory substrate and the common amongst them are starch, sucrose, glucose and fructose. The complex carbohydrates are first hydrolyzed to simple sugars and then they are utilized.

Starch → Disaccharides → Hexoses

### ***Fats***

The fats are important storage food in seeds. Nearly 80 per cent of the angiosperms have fats as the main storage food in their seeds. At the time of seed germination, large amount of fats are converted into carbohydrates while the remaining fats are utilized in respiration. Fats are first broken down to glycerol and fatty acids. The fatty acids are broken down to acetyl coenzyme by  $\beta$ -oxidation. The acetyl coenzyme enters Kreb's cycle for further degradation and releases energy. Glycerol can directly enter the respiratory channel via glyceraldehyde.

### ***Organic acids***

Organic acids normally do not accumulate in plants to any appreciable extent except in the members of the family, Crassulaceae. Organic acids are oxidized under aerobic conditions to carbon dioxide and water.

### ***Proteins***

Under normal conditions, proteins are used up as respiratory substrate only in seeds rich in storage proteins. In vegetative tissues, proteins are consumed only under starvation. The proteins are hydrolyzed to form amino acids. Later, the amino acids undergo deamination forming organic acids and the organic acids can enter Kreb's cycle directly.

### **Mechanism of Respiration**

1. Glycolysis
2. Aerobic breakdown of pyruvic acid (Kreb's cycle)
3. Electron Transport System/ Terminal oxidation / oxidative phosphorylation
5. Pentose phosphate pathway

## A. GLYCOLYSIS / EMBDEN – MEYER HOF – PARANAS (EMP) PATHWAY

Glycolysis can take place even in the absence of O<sub>2</sub>. One molecule of the 6 carbon compound, glucose is broken down through a series of enzyme reactions into two 3-carbon compounds, the pyruvic acid. Glycolysis takes place in the cytoplasm and it does not require oxygen. Hence it is an anaerobic process.


- Glucose molecules react with ATP molecules in the presence of the enzyme hexokinase to form glucose -6- phosphate.




- Glucose-6-phosphate is isomerised into fructose-6-phosphate in the presence of phospho hexose isomerase.



- Fructose-6-phosphate reacts with one molecule of ATP in the presence of phospho hexo kinase forming fructose 1, 6-disphosphate.




- Fructose 1, 6 diphosphate is converted into two trioses, 3-phospho glyceraldehyde and dihydroxy acetone phosphate in the presence of aldolase.



- 3-phosphoglyceraldehyde reacts with H<sub>3</sub>PO<sub>4</sub> and forms 1,3-diphosphoglyceraldehyde where, the reaction is non –enzymatic.


- 1, 3-Diphosphoglyceraldehyde is oxidized to form 1,3- diphosphoglycerate in the presence of triose-phosphate dehydrogenase and coenzyme NAD<sup>+</sup>. The NAD<sup>+</sup> acts as hydrogen acceptor and reduced to NADH<sup>+</sup> + H<sup>+</sup> in the reaction.



- 1, 3-Diphosphoglycerate reacts with ADP in the presence of phosphoglyceric transphorylase (kinase) to form 3 phosphoglyceric acid and ATP.



7. 3, Phosphoglycerate → 2, Phosphoglycerate acid is isomerized into 2 phosphoglyceric acid in the presence of the enzyme, phospho glycero mutase



8. 2 phosphoglyceric acid is converted into 2-phosphoenolpyruvic acid in the presence of enolase.



9. 2 phospho enol pyruvic acid reacts with ADP to form one molecule each of pyruvic acid and ATP in the presence of pyruvate kinase.




Glycolysis or EMP pathway is common in both aerobic and anaerobic respiration.

### The overall glycolytic process can be summarized as follows



- Thus there is a gain of  $4 - 2 = 2$  ATP molecules per hexose sugar molecule oxidized during this process.
- Besides this, 2 molecules of reduced coenzyme  $\text{NADH}_2$  are also produced per molecule of hexose sugar in glycolysis.
- During aerobic respiration, these two  $\text{NADH}_2$  are oxidized via the electron transport chain to yield 3 ATP molecules each. Thus 6 ATP molecules are formed.

# Glycolysis




## Summary of Glycolysis




### 13. KREBS' CYCLE / CITRIC ACID CYCLE /TCA CYCLE

The pyruvic acid produced in glycolysis enters into Krebs' cycle for further oxidation. Krebs' cycle is also known as citric acid cycle or Tri carboxylic acid (TCA) cycle. This aerobic process takes place in mitochondria where necessary enzymes are present in matrix.


11. Pyruvic acid reacts with CoA and NAD and is oxidatively decarboxylated. One molecule of CO<sub>2</sub> is released and NAD is reduced. Pyruvic acid is converted into acetyl CoA.




12. Acetyl-CoA condenses with oxaloacetic acid in the presence of condensing enzyme and water molecule to form citric acid. CoA becomes free.




13. Citric acid is dehydrated in the presence of aconitase to form cis – aconitic acid



14. Cis-aconitic acid reacts with one molecule of water to form Isocitric acid




15. Iso-citric acid is oxidized to oxalo succinic acid in the presence of Isocitric dehydrogenase. NADP is reduced to NADPH<sub>2</sub> in the reaction.




Isocitric acid + NADP


16. Oxalo succinic acid is decarboxylated in the presence of oxalo succinic decarboxylase to form  $\alpha$  - ketoglutaric acid and a second molecule of  $\text{CO}_2$  is released.




17.  $\alpha$  - ketoglutaric acid reacts with CoA and NAD in the presence of  $\alpha$  - ketoglutaric acid dehydrogenase complex and is oxidatively decarboxylated to form succinyl CoA and a third mole of  $\text{CO}_2$  is released. NAD is reduced in the reaction.




18. Succinyl CoA reacts with water molecule to form succinic acid. CoA becomes free and one molecule of GDP (Guanosine diphosphate) is phosphorylated in presence of inorganic phosphate to form one molecule of GTP.




19. Succinic acid is oxidized to fumaric acid in the presence of succinic dehydrogenase and co enzyme FAD is reduced in this reaction.




20. One mole of  $\text{H}_2\text{O}$  is added to Fumaric acid in the presence of fumarase to form malic acid.



21. In the last step, malic acid is oxidized to oxaloacetic acid in the presence of malic dehydrogenase and one molecule of coenzyme i.e. NAD is reduced.



## KREBS CYCLE or TCA CYCLE



### **Pentose phosphate pathway (PPP) / Hexose mono phosphate (HMP) shunt/ Phosphogluconate pathway / Warburg and Dicken's pathway**

The pentose phosphate pathway occurs in the cytoplasm outside the mitochondria and it is an alternative pathway to glycolysis and Kreb's cycle. The presence of some compounds like iodoacetate, fluorides, arsenates etc. inhibit some steps in glycolysis and that leads to the alternate pathway. This pathway was discovered by Warburg and Dicken (1938). This pathway does not produce ATP but it produces another form of energy called reducing power in the form of NADPH. It is not oxidized in the electron transport system but, it serves as hydrogen and electron donor in the biosynthesis of fatty acids and steroids. The pentose phosphate pathway consists of two distinct phases. In the first phase, hexose is converted into pentose and in the second phase, pentose is reconverted into hexose.

In the process, oxidation of glucose 6 phosphate leads to the formation of 6 phosphogluconic acid (pentose phosphate). Since glucose is directly oxidized without entering glycolysis, it is called as *direct oxidation*.



It provides ribose sugars for the synthesis of nucleic acids and is also required for shikimic acid pathway. Although ATP is not produced, NADPH is produced and serves as hydrogen and electron donor in the biosynthesis of fatty acids and steroids. The pathway is also called as *phosphogluconate pathway* as the first product in this pathway is phosphogluconate.

## **OXIDATIVE PHOSPHORYLATION**

### **C. TERMINAL OXIDATION OF THE REDUCED COENZYMES / ELECTRON TRANSPORT SYSTEM AND OXIDATIVE PHOSPHORYLATION**

The last step in aerobic respiration is the oxidation of reduced coenzymes produced in glycolysis and Krebs' cycle by molecular oxygen through FAD, UQ (ubiquinone), cytochrome b, cytochrome c, cytochrome a and cytochrome  $a_3$  (cytochrome oxidase).

Two hydrogen atoms or electrons from the reduced coenzyme ( $\text{NADH}_2$  or  $\text{NADPH}_2$ ) travel through FAD and the cytochromes and ultimately combines with  $1/2\text{O}_2$  molecule to produce one molecule of  $\text{H}_2\text{O}$ . This is called as *terminal oxidation*.

The terminal oxidation of each reduced coenzyme requires  $1/2\text{O}_2$  molecule and 2H atoms (i.e.  $2\text{e}^- + 2\text{H}^+$ ) to produce one  $\text{H}_2\text{O}$  molecule. Except for flavoproteins (like FAD)

and ubiquinone (UQ) which are hydrogen carriers, the other components of electron transport chain (cytochromes) are only electron carriers i.e. they cannot give or take protons ( $H^+$ )

During the electron transport, FAD and the iron atom of different cytochromes get successively reduced ( $Fe^{++}$ ) and oxidized ( $Fe^{+++}$ ) and enough energy is released in some places which is utilized in the photophosphorylation of ADP molecules in the presence of inorganic phosphate to generate energy rich ATP molecules. Since, this oxidation accompanies phosphorylation; it is called as *oxidative phosphorylation*.

One molecule of ATP with 7.6 Kcal.energy is synthesized at each place when electrons are transferred from

1. Reduced  $NADH_2$  or  $NADPH_2$  to FAD
2. Reduced cytochrome b to cytochrome c
3. Reduced cytochrome a to cytochrome  $a_3$

Thus, oxidation of one molecule of reduced  $NADH_2$  or  $NADPH_2$  will result in the formation of 3 ATP molecules while the oxidation of  $FADH_2$  lead to the synthesis of 2 ATP molecules.

According to the most recent findings, although in eukaryotes terminal oxidation of mitochondrial NADH / NADPH results in the production of 3 ATP molecules but that of extra mitochondrial NADH / NADPH yields only 2 ATP molecules. Therefore, the two reduced coenzyme molecules (NADH) produced per hexose sugar molecule during Glycolysis will yield only  $2 \times 2 : 4$  ATP molecules instead of 6 ATP molecules. Complete oxidation of a glucose molecule (hexose sugar) in aerobic respiration results in the net gain of 36 ATP molecules in most eukaryotes.

One glucose molecule contains about 686 Kcal. Energy and 38 ATP molecules will have 273.6 Kcal energy. Therefore about 40% ( $273.6/686$ ) energy of the glucose molecule is utilized during aerobic breakdown and the rest is lost as heat. Since huge amount of energy is generated in mitochondria in the form of ATP molecules, they are called as *Power Houses of the cell*.

ATP molecules contain energy in terminal pyrophosphate bonds. When these energy rich bonds break, energy is released and utilized in driving various other metabolic processes of the cell.

#### **Differences between oxidative phosphorylation and Photophosphorylation**

	Oxidative phosphorylation	Photophosphorylation
--	---------------------------	----------------------

1	It occurs during respiration	Occurs during photosynthesis
2	Occurs inside the mitochondria (inner membrane of cristae)	Occurs inside the chloroplast (in the thylakoid membrane)
3	Molecular O <sub>2</sub> is required for terminal oxidation	Molecular O <sub>2</sub> is not required
4	Pigment systems are not involved	Pigment systems, PSI and PSII are involved
5	It occurs in electron transport system	Occurs during cyclic and non cyclic electron transport
6	ATP molecules are released to cytoplasm and used in various metabolic reactions of the cell	ATP molecules produced are utilized for CO <sub>2</sub> assimilation in the dark reaction of photosynthesis

### Efficiency of respiration

The total energy content of one molecule of glucose is 686 Kcal. Out of this energy, available free energy is 673.6 Kcal and the energy content of ATP molecule is calculated as 7.3 Kcal. The efficiency of respiration may be expressed as follows.

$$\frac{\text{Kcal of energy conserved in ATP}}{\text{Total free energy available}} \times 100$$

$$\text{Efficiency of respiration: } \frac{38 \times 7.3}{673.6} \times 100: 41\%$$

$$\text{Efficiency of aerobic respiration : } \frac{38 \times 7.3}{673.6} \times 100: 41\%$$

$$\text{Efficiency of anaerobic respiration: } \frac{2 \times 7.3}{47} \times 100: 31\%$$

$$\text{Efficiency of fermentation : } \frac{2 \times 7.3}{40} \times 100: 36.5\%$$

### Respiratory quotient


The ratio of the volume of CO<sub>2</sub> released to the volume of O<sub>2</sub> taken during respiration is called as respiratory quotient and is denoted as RQ

$$RQ = \frac{\text{Volume of CO}_2}{\text{Volume of O}_2}$$

### Value of RQ

The value of RQ depends upon the nature of the respiratory substrate and the amount of O<sub>2</sub> present in respiratory substrate.

- When **carbohydrates** such as hexose sugars are oxidized in respiration, the value of RQ is 1 or unity because volume of CO<sub>2</sub> evolved equals to the volume of O<sub>2</sub> absorbed.



Glucose

$$RQ = \frac{\text{volume of CO}_2}{\text{volume of O}_2} = \frac{6}{6} = 1 \text{ or unity}$$

- When **fats** are the respiratory substrate, the value of RQ becomes less than one because fats are poorer in O<sub>2</sub> in comparison to carbon and they require more O<sub>2</sub> for their oxidation,



Tripalmitin

$$RQ = \frac{\text{volume of CO}_2}{\text{volume of O}_2} = \frac{102}{145} = 0.7$$

(Fats are oxidized in respiration usually during the germination of fatty seeds).

- When **organic acids** are oxidized in respiration, the value of RQ becomes more than one. It is because organic acids are rich in O<sub>2</sub> and require less O<sub>2</sub> for their oxidation.



Malic acid

$$RQ = \frac{\text{volume of CO}_2}{\text{volume of O}_2} = \frac{4}{3} = 1.3$$

### Energy budgeting

Stages	Gain of	Consumption	Net gain of
--------	---------	-------------	-------------

	ATP	of ATP	ATP
Glycolysis			
1) Glucose → Glucose 6 PO <sub>4</sub>		1	
2) Fructose 6 PO <sub>4</sub> → Fructose 1,6 di PO <sub>4</sub>		1	
3) 1,3 diphosphoglyceraldehyde → 1,3 diphospho glyceric acid	6		
4) 1,3 diphospho glyceric acid → 3 phosphoglyceric acid	2		
5) 2 phosphoenol pyruvic acid → Pyruvic acid	2		
Total	10	-2	8
Kreb's cycle			
6) Pyruvic acid → Acetyl CoA	3		
7) Isocitric acid → Oxalosuccinic acid	3		
8) ketoglutaric acid → Succinyl CoA	3		
9) Succinyl Co A → Succinic Acid	1		
10) Succinic acid → Fumaric Acid	2		
11) Malic acid → Oxaloacetic acid	3		
Total ATP mol. produced per Pyruvic acid	15		15
Total ATP mol. produced for 2 Pyruvic acids	15 x 2:30		30
Grand Total	40	-2	8+ 30 = 38

## FACTORS AFFECTING RESPIRATION

### A. External factors

#### 1. Temperature

Temperature has profound influence on the rate of respiration. Optimum temperature for respiration is about 30°C, minimum 0°C and maximum about 45°C. At low temperature, the respiratory enzymes becomes inactive, consequently the rate of respiration falls. It is due to this fact that the quality of fruits and vegetables stored at low temperature does not deteriorate. At very high temperature, respiration slows down and may even be stopped due to denaturation of the respiratory enzymes.

#### 2. Oxygen

In complete absence of O<sub>2</sub>, anaerobic respiration takes place while aerobic respiration stops. In higher plants, the anaerobiosis produces large amount of alcohol which is toxic to plants. If some amount of O<sub>2</sub> is available, anaerobic respiration slows down and aerobic respiration starts. The concentration of O<sub>2</sub> at which aerobic respiration is optimum and anaerobic respiration is stopped, is called as *extinction point*.

It is observed that under anaerobic conditions, much more sugar is taken up per quantity of yeast present than it is consumed in the presence of oxygen. The inhibition on the rate of carbohydrate breakdown by oxygen is called as *Pasteur's effect*.

### **3. Carbon dioxide**

Higher concentration of CO<sub>2</sub> in the atmosphere especially in the poorly aerated soil has retarding effect on the rate of respiration.

### **4. Inorganic salts**

If a plant or tissue is transferred from water to salt solution, the rate of respiration increases (called as *salt respiration*).

### **5. Water**

Proper hydration of cells is essential for respiration. Rate of respiration decreases with decreased amount of water, so much so, that in dry seeds, the respiration is at its minimum. It is because in the absence of a medium, the respiratory enzymes become inactive.

### **6. Light**

The effect of light is indirect on the rate of respiration through the synthesis of organic food matter in photosynthesis.

### **7. Wound or injury**

Injury or wounds result in increased respiration as the plants in such a state require more energy which comes from respiration. The wounded cells become more meristematic to form new cells for healing the wound.

## **Internal factors**

### **1. Protoplasmic factors**

The amount of protoplasm in the cell and its state of activity influence the rate of respiration.

- The rate of respiration is higher in young meristematic cells which divide actively and requires more energy. Such cells have greater amount of protoplasm and no vacuoles.

- In old mature tissues, the rate of respiration is lower because of lesser amount of active protoplasm

## 2. Concentration of respiratory substrate

Increased concentration of respirable food material brings about an increase in the rate of respiration.

Under starvation conditions, such as in etiolated leaves, the rate of respiration slows down considerably. If such etiolated leaves are supplied with sucrose solution for few days even in dark conditions, the rate of respiration increases.

### Differences between Photorespiration and Dark respiration

	Photorespiration	Dark /Mitochondrial respiration
1	It occurs in the presence of light	It occurs in the presence of both light and dark.
2	The substrate is glycolate	The respiratory substrate may be carbohydrate, fat or protein.
3	It occurs in chloroplast, peroxisome and mitochondria	The process occurs in the cytoplasm and mitochondria
4	It occurs in temperate plants like, wheat and cotton ( mainly in C3 plants)	It occurs in C4 plants ( maize and sugar cane)
5	It occurs in the green tissues of plants	It occurs in all the living plants( both green and non green )
6	The optimum temperature is 25- 35°C	It is not temperature sensitive
7	This process increases with increased CO <sub>2</sub> concentration.	This process saturates at 2-3 percent O <sub>2</sub> in the atmosphere and beyond this concentration there is no increase.
8	Hydrogen peroxide is formed during the reaction	Hydrogen peroxide is not formed.
9	ATP molecules are not produced,	Several ATP molecules are produced.
10	Reduced coenzymes such as NADPH <sub>2</sub> , NADH <sub>2</sub> and FADH <sub>2</sub> are not produced.	Reduced coenzymes such as NADPH <sub>2</sub> , NADH <sub>2</sub> and FADH <sub>2</sub> are produced.
11	One molecule of ammonia is released	No ammonia is produced

	per molecule of CO <sub>2</sub> released.	
12	Phosphorylation does not occur	Oxidative phosphorylation occurs.

### Differences between respiration and photosynthesis

	Respiration	Photosynthesis
1	It is catabolic process resulting in the destruction of stored food	It is an anabolic process resulting in the manufacture of food.
2	Light is not essential for the process	Light is very much essential
3	Oxygen is absorbed in the process	Oxygen is liberated
4	Carbon dioxide and water are produced	Carbon dioxide is fixed to form carbon containing compound
5	Potential energy is converted into Kinetic energy	Light energy is converted into chemical energy (potential energy)
6	Glucose and oxygen are the raw materials	Carbon dioxide and water are the raw materials
7	Energy is released during respiration and hence it is an exothermic process.	Energy is stored during photosynthesis and hence it is an endothermic process
8	Reduction in the dry weight	Gain in the dry weight
9	Chlorophyllous tissues are not necessary	Chlorophyllous tissues are essential for the process

### Differences between aerobic respiration and fermentation

	Aerobic respiration	Fermentation
1	It occurs in all living cells of the plants throughout the day and night	Occurs outside the plant cells and in certain microorganisms
2	It takes place in the presence of oxygen	Absence of oxygen
3	The end products are CO <sub>2</sub> and H <sub>2</sub> O	End products are CO <sub>2</sub> and alcohol or other organic acids
4	It is not toxic to plants	It is toxic to plants
5	Complete oxidation of food material is	Incomplete oxidation is observed

	observed	
6	Large amount of energy (673 kCal) is released per glucose molecule	Very small amount of energy (21 kCal) is released per glucose molecule
7	The complete oxidation yields 38 ATP molecules	The incomplete oxidation in fermentation yields only two ATP molecules
8	The enzyme, zymase is not required but many other enzymes and coenzymes are required	Zymase is required in the case of carbohydrates

## 14. PROTEIN AND FAT SYNTHESIS

### **Biosynthesis of protein**

Protein is a complex organic nitrogenous substance found in all living tissues of plants and animals. They are polymer of amino acids in linear order. Synthesis of protein may take place from amino acids produced by direct amination of organic acids or by degradation of protein. Former is known as *primary protein synthesis* while the latter is called *secondary protein synthesis*. Protein synthesis occurs in pre – DNA synthesis phase ( $G_1$  phase) of cell cycle.

Biosynthesis of protein takes place in prokaryotes as well as in eukaryotes. Kinds of protein to be synthesized depend upon the gene (DNA segment). Gene is continuous uninterrupted sequence of nucleotides which codes for a single polypeptide chain. Now it is believed that the sequence of some eukaryotic genes is found to be interrupted by nucleotides that are not represented with the amino acid sequence of protein. They are non-coding (silent). Genes control all metabolic processes by synthesizing proteins (enzymes).

Structure of an eukaryotic gene showing Exon (coding part) and Intron (non-coding part).

Mechanism of protein synthesis: Protein synthesis takes place in following two stages:

- I. Transcription
- II. Translation

**Transcription:** Transcription occurs throughout inter phase and continues up to early prophase of cell division. It is primary stage of protein synthesis. When DNA produces DNA the process is called replication but when DNA produces RNA the process is called transcription. In the former case DNA is duplicated while in the latter case protein is synthesized. During transcription RNA is synthesized on DNA template. Here, information or order contained in DNA is passed on the mRNA for synthesis of particular protein. The information is in coded form and consists of three nitrogenous bases (triplet codons). The part of DNA responsible for synthesis of mRNA is which leads to one polypeptide chain is

called cistron (functional gene). New strands of mRNA are synthesized on DNA template making use of RNA nucleotides present in surroundings in 5' 3' director just like DNA chains.

Single DNA strand serves as template for RNA polymerase and synthesizes RNA. DNA strand which serves as template for transcription is called the sense strand. The complementary strand is antisense strand. In SV – 40 viruses both strands of DNA are transcribed and it is called symmetrical transcription, but when only one strand is transcribed it is called asymmetrical transcription. Former type of transcription also occurs in Polymer virus DNA and in the mitochondria genome.

#### Mechanism of transcription

Transcription involves following events:

- I. Uncoiling of DNA molecule
- II. Synthesis and action of enzyme RNA polymerase
- III. Synthesis of hn RNA / mRAN.

**Uncoiling of DNA molecule:** As per “nucleosome model” of chromosome. Chromosomes are ‘matarmala’ like beaded structure. Beads are separated by string and are making up of DNA and histone protein. Histone proteins are of 5 kinds: H2A, H2B, H3, H4 and H5 or H1. First four constitute the bead core and H1 links two beads. The DNA molecules are wrapped on histone protein cores and linker protein core in beaded and linker regions respectively.

**Structure and function of enzyme RNA polymerase:** RNA polymerase is a holoenzyme. Core particle consists of sub units  $\alpha'\beta$ ,  $\beta'$  and  $\psi$ . Cofactor consists of sigma factor. For functional RNA polymerase formation the two (core enzyme and sigma factor) unite. Sigma factor recognized correct start signal at DNA template and core enzyme continues transcription. Sigma factor dissociates after initiation of transcription to adjure with other core enzyme of RNA polymerase. In prokaryotes RNA polymerase is only one type while in eukaryotes.

**Production of mRNA / hnRNA:** In prokaryotes where nucleus is not well organized mRNA is the direct product of transcription, while in eukaryotes the direct product of transcription is hnRNA (heteronucleic RNA) and mRNA is derived from hnRNA by cutting and splicing. HnRNA has coding and non-coding sequences. Coding sequences are interrupted by non-coding sequences. Non coding sequences are removed by splicing (cutting) by endonuclease enzyme and coding sequences are ligased together to form mRNA. The spliced non-coding sequences are degraded within nucleus. It never goes out of nucleus. Thus, only fraction of hnRNA is translocated to cytoplasm from nucleus via nuclear pore.

In eukaryotes migration of mRNA from nucleus to cytoplasm via nuclear pore occurs through poly a tail. According to another view, ribosome's pull the mRNA from nucleus to cytoplasm. Now mRNA gets established in cytoplasm.

**Translation:** Translation is a process in which order (message) given by DNA to mRNA for synthesis of particular protein is implemented (conveyed). Genetic information concealed in mRNA directs the synthesis of particular protein. These orders are in coded form. This coded information (expressed through codons) is recognized by tRNA having anticodons. Anticodons are opposite to codons (codons and anticodons are complementary to each other).

**Mechanism of translation:** It involves following events:

- I. Activation and selection of amino acids
- II. Transfer of amino acids to tRNA molecules
- III. Formation of protein synthesizing apparatus and chain initiation
- IV. Chain elongation
- V. Chain termination

## FAT SYNTHESIS

Fat synthesis can be studied under the following heads: -

### 1. Fat synthesis of Glycerol

There may be different methods of the formation of glycerol in plants, but one of the very common methods is from dihydroxy acetone phosphate which is an intermediate of glycolysis. Dihydroxyacetone phosphate is first reduced to  $\alpha$ -glycerophosphate by the enzyme glycerol – 3 – phosphate dehydrogenase. Co-enzyme NADH<sub>2</sub> is oxidized in this reaction.  $\alpha$ -glycerophosphate is then hydrolysed by *glycerol phosphatase* to liberate phosphoric acid and forming glycerol.

## 2. Synthesis of Fatty acids

Long chain saturated fatty acids\* are synthesized in plants from active two carbon units, the acetyl – CoA ( $\text{CH}_3\text{CO}.\text{CoA}$ ). Although the reactions of  $\beta$  – oxidation of fatty acids are reversible, the fatty acids are not formed simply by the reverse reactions of  $\beta$  – oxidation. Synthesis of fatty acids from  $\text{CH}_3\text{CO}.\text{CoA}$  takes place step by step. In each step the fatty acid chain is increased by two carbon atoms. Each step involves two reactions –

- (i) In the first reaction which takes place in the presence of acetyl – CoA carboxylase, acetyl – CoA combines with  $\text{CO}_2$  to form malonyl – CoA (malonic acid is 3 – C compound). ATP provides energy while  $\text{Mn}^{++}$  and biotin are required as co-factors.
- (ii) Malonyl CoA reacts with another molecule of  $\text{CH}_3\text{CO}.\text{CoA}$  in the presence of fatty acid synthetase and Coenzyme NADPH<sub>2</sub> to form Coenzyme – A derivative butric acid (butyric acid contains 4 – atoms). One mol. Of  $\text{CO}_2$ ,  $\text{H}_2\text{O}$  and CoA are released while NADPH<sub>2</sub> oxidised in the reaction.

Butyryl CoA, in the next step will combine with malonyl CoA to form CoA derivative of fatty acid containing 6-C atoms. This process is repeated till Coenzymes-A derivative of long chain fatty acid (which may contain up to 16-18C atoms) is produced.

(As a matter of fact the enzyme fatty acid synthetase is not simple but a complex of many enzymes (multienzyme complex) and an acyl carrier protein called as ACP\*\*. And actually the reaction (ii) described above only summarises a number of reactions involved in the synthesis of fatty acid from acetyl – CoA which can be grouped under 3 categories

**Initiation reaction:** In this reaction acetyl CoA transfers its acetyl group to one of the – SH groups of multienzyme complex i.e. fatty acid synthetase.

Unsaturated fatty acids are synthesized by denaturation of saturated fatty acid. ACP is similar to CoA in having phosphopantetheine as the functional unit in their structures. In CoA, it is esterified to Adenosine 3, 5 – bisphosphate but in ACP, it is esterified to serine of a protein chain consisting of 81 amino acids.

### Chain elongation reactions

Six different reactions involved here are (i) malonyl transfer, (ii) condensation, (iii) reduction, (iv) dehydration, (v) reduction and (vi) acyl transfer. Chain elongation starts with the transfer of malonyl group from malonyl – CoA to second – SH group of the multienzyme complex. Then, there is a condensation of the latter so that a 4 – C unit is produced. This unit by next three reactions i.e. reduction, dehydration and reduction is converted into saturated 4 – C unit (i.e. butyryl – CoA). In acyl transfer reaction the fatty acid residue is transferred back to the – SH group to which the acetyl group was transferred in initiation reaction. The cycle is repeated again and again with malonyl transfer, condensation etc. till the fatty acid residue consists of up to 16 - 18 C atoms. Each such turn elongates fatty acid chain by 2 – C atoms. Details of chain elongation reactions are given below

### Termination reaction

When the fatty acid residue has attained a desired length the chain elongation stops at reaction (v) and the cycle is not repeated. The acyl group instead of being transferred to the – SH of the enzyme is transferred to – SH group of Co-enzyme A (CoASH) molecule. Thus, CoA derivate of the fatty acid is produced which can then be utilized in fat synthesis. The enzyme becomes free.

It is believed that during this process of fatty acid synthesis, the acyl group of fatty acid bound to the – SH group of ACP. The latter then passes it from one enzyme of the complex to the other.

### (3) Condensation of fatty acids and Glycerol

The fats or triglycerides are synthesized not from glycerol and free fatty acids but from  $\alpha$ -glycerophosphate and CoA derivatives of fatty acid, i.e. fatty acyl CoA residues. First, there is acylation of  $\alpha$ -glycerophosphate by two fatty acyl-CoA molecule to form phosphatidic acid. Now dephosphorylation occurs in the presence of phosphatase and a deglyceride is formed. The acylation of the free-OH groups of diglyceride completes the biosynthesis of triglyceride or fat.


## 15. PHOTOPERIODISM

Photoperiodism is the phenomenon of physiological changes that occur in plants in response to relative length of day and night (i.e. photoperiod). The response of the plants to the photoperiod, expressed in the form of flowering is also called as photoperiodism. The phenomenon of photoperiodism was first discovered by Garner and Allard (1920). Depending upon the duration of photoperiod, the plants are classified into three categories.

1. Short day plants (SDP)
2. Long day plants (LDP)
3. Day neutral plants (DNP)

### 1. *Short day plants*


#### SHORT DAY PLANTS



These plants require a relatively short day light period (usually 8-10 hours) and a continuous dark period of about 14-16 hours for subsequent flowering. These plants are also known as long-night plants

E.g. Rice, coffee, soybean, tobacco and chrysanthemum

- In short day plants, the dark period is critical and must be continuous. If this dark period is interrupted with a brief exposure of red light (660-665 nm wavelength), the short day plant will not flower.
- Maximum inhibition of flowering with red light occurs at about the middle of critical dark period.
- However, the inhibitory effect of red light can be overcome by a subsequent exposure with far-red light (730-735 mm wavelength)
- Interruption of the light period with red light does not have inhibitory effect on flowering in short day plants.
- Prolongation of the continuous dark period initiates early flowering.




## 2. Long day plants

These plants require longer day light period (usually 14-16 hours) in a 24 hours cycle for subsequent flowering. These plants are also called as short night plants.

E.g. Wheat, radish, cabbage, sugar beet and spinach.

- In long day plants, light period is critical
- A brief exposure of red light in the dark period or the prolongation of light period stimulates flowering in long day plants.

## LONG DAY PLANTS



### ***3. Day neutral plants***

These plants flower in all photoperiod ranging from 5 hours to 24 hours continuous exposure.

E.g. Tomato, cotton, sunflower, cucumber, peas and certain varieties of tobacco.

During recent years, intermediate categories of plants such as *long short day plants* and *short long day plants* have also been recognized.

#### *i. Long short day plants*

These are short day plants but must be exposed to long days during early periods of growth for subsequent flowering. E.g. Bryophyllum.

#### *ii. Short –long day plants*

These are long day plants but must be exposed to short day during early periods of growth for subsequent flowering. E.g. certain varieties of wheat and rye.

### Differences between short day and long day plants

	<b>Short day plant</b>	<b>Long day plant</b>
1	Plants flower when photoperiod is less than the critical day length	Plants flower when photoperiod is more than the critical day length
2	Interruption during light period with darkness does not inhibit flowering	Interruption during light period with darkness inhibit flowering
3	Flowering is inhibited if the long dark period is interrupted by a flash of light	Flowering occurs if the long dark period is interrupted by a flash of light
4	Long continuous and uninterrupted dark period is critical for flowering	Dark period is not critical for flowering
5	Flowering does not occur under alternating cycles of short day and short light period.	Flowering occurs under alternating cycles of short day followed by still shorter dark periods

### Phytochrome

It is observed that that a brief exposure with red light during critical dark period inhibits flowering in a short day plant and this inhibitory effect can be reversed by a subsequent exposure with far-red light. Similarly, prolongation of the critical light period or the interruption of the dark period stimulates flowering in long-day plants.

This inhibition of flowering in short day plant and stimulation of flowering in long day plants involves the operation of a proteinaceous pigment called *phytochrome*. It is present in the plasma membrane of cells and it has two components, chromophore and protein. Phytochrome is present in roots, coleoptiles, stems, hypocotyls, cotyledons, petioles, leaf blades, vegetative buds, flower tissues, seeds and developing fruits of higher plants.

The pigment, phytochrome exists in two different forms i.e., red light absorbing form which is designated as *Pr* and far red light absorbing form which is designated as *Pfr*. These two forms of the pigment are photo chemically inter convertible. When *Pr* form of the pigment absorbs red light (660-665 nm), it is converted into *Pfr* form. When *Pfr* form of the pigment absorbs far red light (730-735 nm), it is converted into *Pr* form. The *Pfr* form of pigment gradually changes into *Pr* form in dark.

It is considered that during day time, the *Pfr* form of the pigment is accumulated in the plants which are inhibitory to flowering in short day plants but is stimulatory in long day plants. During critical dark period in short day plants, this form gradually changes into *Pr* form resulting in flowering. A brief exposure with red light will convert this form again into *Pfr* form thus inhibiting flowering.

Reversal of the inhibitory effect of red light during critical dark period in SDP by subsequent far-red light exposure is because, the *Pfr* form after absorbing far-red light (730-354 nm) will again be converted back into *Pr* form.

Prolongation of critical light period or the interruption of the dark period by red-light in long day plants will result in further accumulation of the *Pfr* form of the pigment, thus stimulating flowering in long-day plants.

#### **Differences between Pr and Pfr forms of phytochrome**

	<b>Pr form</b>	<b>Pfr form</b>
1	It is blue green in colour	It is light green in colour
2	It is an inactive form of phytochrome and it does not show phytochrome mediated responses	It is an active form of phytochrome and hence shows phytochrome mediated responses
3	It has maximum absorption in red region (about 680nm)	It has maximum absorption in far-red region (about 730nm)
4	It can be converted into Pfr form in red region (660-665nm)	It can be converted into Pr form in far red region (730-735nm)
5	It is found diffused throughout the cytosol	It is found in discrete areas of cytosol
6	The Pr form contains many double bonds in pyrrole rings	The Pfr form contains rearranged double bonds in all pyrrole rings

#### **Significance of photoperiodism**

Photoperiodism is an example for *physiological preconditioning*. The stimulus is given at one time and the response is observed after months. Exposure to longer photoperiods hastens flowering (E.g). In wheat, the earing is hastened. During long light exposure, *Pr* form

is converted into *Pfr* form and flowering is initiated. If dark period is greater, *Pfr* is converted into *Pfr* form that inhibits flowering.

The important phytochrome mediated photo responses in plants include photoperiodism, seed germination, sex expression, bud dormancy, rhizome formation, leaf abscission, epinasty, flower induction, protein synthesis, pigment synthesis, auxin catabolism, respiration and stomatal differentiation.

## 16. TRANSMISSION OF STIMULUS - THEORIES OF FLOWERING.

### Photoperiodic Induction

The influence of the length of day and night on the initiation of flowering is called *photoperiodic induction* or *photo induction*.

Plants may require one or more inductive cycle for flowering. An appropriate photoperiod in 24 hours cycle constitutes one inductive cycle. If a plant which has received sufficient inductive cycle is subsequently placed under unfavourable photoperiod, it will still flower.

Flowering will also occur if a plant receives inductive cycles after intervals of unfavourable photoperiods (i.e. discontinuous inductive cycle). This persistence of photoperiodic after effect is called as photoperiodic induction.

- An increase in the number of inductive cycles results in early flowering of the plant. For instance, xanthium (a short day plant) requires only one inductive cycle and normally flowers after about 64 days. It can be made to flower even after 13 days if it has received 4-8 inductive cycle. In such case number of flowers is also increased.
- Continuous inductive cycles promote early flowering than discontinuous inductive cycle.

Some of the examples of plants which requires more than one inductive cycle for subsequent flowering are,

Biloxi soybean (SDP) - 2 inductive cycles

Salvia (SDP) - 17 Inductive cycles

Plantago (LDP) - 25 Inductive cycles

### Critical day length

*Maryland mammoth* tobacco and xanthium are short day plants, but the *Maryland mammoth* tobacco is induced to flower when the photoperiod is shorter than 12 hours (12L /12D) whereas, xanthium is induced to flower when the photoperiod is shorter than 15.5 hours (15.5L /8.5D). The photoperiod required to induce flowering is referred to as the *critical day length*. Hence, the critical day length for *Maryland mammoth* tobacco and

xanthium are 12 and 15.5 hours respectively. A short day plant is one that flowers on photoperiods shorter than the critical day length.

Long day plants, on the other hand, are induced to flower on photoperiods longer than critical day length. For example, the critical day length for *Hyoscyamus niger* is 11 hours (11L /13D) and it is induced to flower on photoperiods longer than 11 hours.

Suppose, xanthium and *Hyoscyamus niger* are exposed to a photoperiod of 14 hours of light and 10 hours of darkness (14L/10D), flowering will be induced in both plants. Xanthium, a short-day plant, will flower because 14L /10D photoperiod is shorter than critical day length of 15.5 hours. *Hyoscyamus*, a long-day plant, will flower because 14L/10D is longer than the critical day length of 11 hours.

### **Perception of photoperiodic stimulus and presence of a floral hormone**

- Photoperiodic stimulus is perceived by the leaves and a floral hormone is produced in the leaves which are then translocated to the apical tip, subsequently causing initiation of floral primordia.
- Photoperiodic stimulus perceived by the leaves can be shown by a simple experiment on cocklebur (xanthium), a short day plant. Cocklebur plant will flower if it has previously been kept under short day conditions. If the plant is defoliated and kept under short day condition, it will not flower. Flowering will also occur if all the leaves of the plant except one leaf have been removed.
- If the cocklebur plant whether intact or defoliated is kept under long day condition it will not flower. But if even one of its leaves is exposed to short day condition and the rest are under long day condition, flowering will occur.
- The photoperiodic stimulus is transmitted from one branch of the plant to another branch. For example, if in a two branched cocklebur plant one branch is exposed to short day and the other to long day photoperiod, flowering occurs on both the branches.
- Flowering also occurs if one branch is kept under long day conditions and other branch from which all the leaves except one have been removed is exposed to short day condition. However, if one branch is exposed to long photoperiod and the other

has been defoliated, under short day conditions, flowering will not occur in any of the branches.

### **Flowering stimulus: Florigen**

The flowering stimulus is produced in leaves and translocated to apical and lateral meristems where flower formation is initiated. Chailakhyan (1937) called the flowering stimulus or flowering hormone as *Florigen*.

Flowering stimulus is similar in long day plants and short day plants. This can be proved by a grafting experiment and can be translocated from one plant to another.

Maryland mammoth tobacco, a short day plant and *Hyoscyamus niger*, a long day plant, are grafted so that the leafy shoots of both the species are available for experiment. If the grafted plants are exposed to either long day or short day conditions, both partners flower. If grafting union is not formed, the flowering stimulus is not translocated from one partner to another partner.

### **Theories of Flowering**

1. Bunning's hypothesis
2. Chailakhyan's hypothesis

#### ***Bunning's hypothesis:***

Bunning (1958) assumes the presence of endogenous rhythms (Oscillator which consist of two half cycles. The first half cycle occurs in day and is called *photophilous phase*. During this, anabolic process predominates including flowering in plants. The other half cycle is dark, sensitive and is called *skotophilous phase*. In this, catabolic process (dehydration of starch) predominates.

SD plants have a critical day length of 9 hours. This period falls within the photophilous phase. Light during scotophil phase will inhibit photo process initiated during photophase. The L.D. plants have a critical day length of 15 hours and some light falls in the skoto philous phase. Under these conditions in L.D. plants will flower. In S.D. plants

oscillator is present close to skoto philous phase, while in L.D. plants it is close to *photo philous phase*.

#### ***Chilakhyan's hypothesis:***

This hypothesis assumes that flowering hormone – florigen is a complex of two types of substances – gibberellin and anthesins. Gibberellin is essential for growth of the plant stems and anthesins are required for flower formation.

According to him, flowering in all annual seed plants requires two phases: (i) *Floral stem formation phase* (ii) *Flower formation phase*. First phase involves increased carbohydrate metabolism and respiration with increased content of GA in leaves. Second phase requires intensive nitrogen metabolism, higher content of anthesins in leaves and nucleic acid metabolites in stem buds.

Long day conditions favour the first phase while short day conditions favour second phase. In long day plants gibberellins are critical, while anthesins are critical in short day plants. However, anthesin is hypothetical; it has not been isolated as yet.

## **VERNALISATION**

The cold treatment given to plant buds, seeds or seedlings for promoting early flowering is known as *Vernalisation*. In short, the chilling treatment for induction of early flowering is called Vernalisation.

Besides an appropriate photoperiod, certain plants require a low temperature treatment during their early stages of the life for subsequent flowering in the later stages. This low temperature treatment requirement was termed vernalization by Lysenko (1928). Due to vernalization, the vegetative period of the plant is cut short resulting in an early flowering. In nature, vernalisation takes place in the seed stage in annuals like winter rye (*Secale cereale*). The biennials and many perennials respond to cold treatment at a very late stage. E.g. Henbane, apples etc.

## **Perception of cold stimulus and presence of floral hormone**

The cold stimulus is perceived by the apical meristems. The perception of the cold stimulus results in the formation of a floral hormone which is transmitted to other parts of the plant. In certain cases, the cold stimulus may even be transmitted to another plant across a graft union.

For instance, if a vernalized henbane plant is grafted to an unvernalized henbane plant, the later also flowers. This is due to the induction of the plant to produce a hormone named as *Vernalin* by Melchers (1939).

## **Conditions necessary for vernalization**

### *1. Age of the plant*

The age of the plant is an important factor in determining the responsiveness of the plant to the cold stimulus and it differs in different species. In cereals like winter wheat, the vernalization is effective only if the germinating seeds have received cold temperature treatment for sufficient time.

While in the case of biennial variety of henbane (*Hyoscyamus niger*), the plant will respond to the cold treatment, only if they are at rosette stage and completed at least 10 days of growth.

### *2. Appropriate low temperature and duration of the exposure*

Most suitable temperature for vernalizing the plants ranges between 1-6°C. The effectiveness of low temperature treatment decreases from 0 to 4°C. Low temperature at about -6°C is completely ineffective. Similarly at high temperatures from 7°C onwards, the response of the plants is decreased. Temperature of about 12-14°C is almost effective in vernalizing the plant. Besides an appropriate low temperature, a suitable duration of the cold treatment is essential for vernalization. Depending upon the degree of temperature and in different species this period may vary, but usually the duration of the chilling treatment is about one and half months or more.

### 3. Oxygen

The vernalization is an aerobic process and requires metabolic energy. In the absence of O<sub>2</sub>, cold treatment becomes completely ineffective.

### 4. Water

Sufficient amount of water is also essential for vernalization. Vernalization of the dry seed is not possible.

## Mechanism of Vernalization

There are two main theories to explain the mechanism of vernalisation.

### 1. Phasic developmental theory

This theory was proposed by Lysenko (1934) as follows.


- (i) The growth (increase in size) and development (i.e. progressive change in the characteristic of the new organs) are two distinct phenomena.
- (ii) According to this theory, the process of the development of an annual seed plant consists of a series of phases which must occur in some predetermined sequence.
- (iii) Commencement of any of these phases will take place only when the preceding phase has been completed.
- (iv) The phases require different external conditions for the completion such as light and temperature.
- (v) Vernalization accelerates the thermo phase i.e. that phase of development which is dependent upon temperature.

Thus, in winter wheat, low temperature is required for the completion of first thermo phase. After this, the next phase that is dependent upon light (photo phase) starts. Vernalization of winter wheat accelerates the first thermo phase so that there is an early swing from vegetative to reproductive phase or flowering.

### 2. Hormonal theories

It has already been described that vernalization probably involves the formation of a floral hormone called as *vernalin*. Based on this fact, many hypothetical schemes have been

proposed by different workers from time to time. The first hormonal theory proposed by Long and Melchers (1947) is schematically shown below.



According to this scheme, the precursor A is converted into a thermo labile compound B during cold treatment. Under normal conditions B changes into C which ultimately causes flowering. But at higher temperature B is converted into D and flowering does not take place (devernalization).

## Devernalization

The positive effect of the low temperature treatment on the vernalization of the plant can be counteracted by subsequent high temperature. This is called devernalization. The devernalized plant can again be vernalized by subsequent low temperature treatment.

## Vernalization and Gibberellins

The gibberellins are known to replace the low temperature requirement in certain biennial plants such as henbane, where the plant normally remains vegetative and retains its rosette habit during the first growing season and after passing through the winter period flowers in the next season. The gibberellins cause such plants to flower even during the first year.

## Significance of vernalization

1. Vernalization shortens the vegetative period of the plant
2. It increases cold resistance of the plants
3. Vernalization increases the resistance of plants to fungal diseases.
4. It is a physiological process that substitutes or compensates the effect of thermo phase.
5. In biennials, vernalisation induces early flowering and early fruit setting.
6. A non vernalised shoot apex can be induced to flower by grafting the plant with a vernalised plant.

## 17. SOURCE SINK RELATIONSHIP

### **Source**

1. It is the regions of photoassimilates production
2. Export photoassimilates
3. Chlorophyllous tissues
4. Leaves, stipules, fruit wall, young stem, pedicel, awns, peduncle, calyx, bract etc

### **Sink**

1. Regions of photoassimilates consumption
2. Import photoassimilates
3. Growing regions
4. Storage organs – Fruit and Seed

### **Source strength**

1. Source Size x Source activity
2. Differences in CO<sub>2</sub> fixation (Rubisco & PEP Case)
3. Leaf characters – size, thickness, mesophyll size, compaction, vascular bundle
4. Carrying capacity of sieve element (temp., H<sub>2</sub>O, nutrients, hormone)

### **Sink strength**

1. Sink size x Sink activity
2. Potential capacity of the sink to accumulate assimilates
3. Competition among different sink

### **Source sink interaction**

1. Source sink equilibrium
2. Small surplus source for stress
3. High source size during sink differentiation
4. Improve strength by activity
5. Synchrony of sink organ development

6. Increased HI is reached – increase DMA
7. Reduce photorespiration in C3 plants

Evans (1983)

- Reduced growth of non harvestable organ
- Prolonged faster storage
- Enhanced competition of storage organ
- Enhanced competition of regulatory process
- Reduced stem weight and height
- Reduced root weight with adequate nutrient and H<sub>2</sub>O
- Improved agronomic support (avoid biotic & abiotic stress)
- Hormonal regulation
- Developmental plasticity (small surplus source for stress)

### **Efficient system**

1. Quick export of photoassimilates to avoid end product inhibition
2. Efficient root system
3. More photosynthetic rate
4. Optimum LAI (4 to 6)
5. High photosynthetic rate & high DMA

### **Blackman's law of limiting factor**

1. A process is controlled by several factors
2. The phase of the process is limited by slowest factor
3. Compensation mechanism working under canopy level

### **Dry matter accumulation (DMA)**

G x E interaction; nutrients; CO<sub>2</sub> fixation rate (path way); photorespiration; vascular network; LAI & LAD; source-sink limiting condition; root-shoot balance  
HI

$$\text{Ye} = \text{Yb} \times h$$

$$\text{/ HI} = \{\text{Yield}_{(\text{Eco})}/\text{Yield}_{(\text{Biol})}\} \times 100$$

### **Improve Harvest index (HI)**

- Increase biomass production (DMA)
- Synchronized development of reproductive organ
- Reproductively determinate
- High source strength at the time of sink differentiation
- Reduced growth of non harvestable organ
- Reduced leaf growth at reproductive stage with high LAD
- Optimum LAI and early peak LAI
- More prolonged and faster storage, enhanced competitiveness among of the storage organ
- High photosynthetic rate
- Improved HI by increased size and number of sink organ
- Decline in duration of Vegetative growth and increased duration of Reproductive growth.

### **Limitations**

- Source: wheat, rice, pulses, oilseeds
- Sink: bajra, ragi
- Transport: sorghum, maize (green leaf at harvest; senescence of phloem Parenchyma)

#### **Sink limitation:**

- Late anthesis (Long Vegetative phase)
- Indeterminate (Vegetative & Reproductive growth)
- Vegetative growth at Reproductive phase
- Less sink number and size
- Hormonal imbalance
- Any Stress
- Multi-sink demand (nodules supply 25 – 75 % of N demand)

#### **Source limitation:**

- Low canopy photosynthesis
- Low optimum LAI
- Slow peak LAI (lag vegetative growth)
- Low LAD at filling

Early leaf senescence

Stress – nutrients, water

### **Plant Growth Regulators (PGRs)**

ABA inhibit sucrose uptake in source (Loading)

Auxin promotes source uptake

Starch accumulation in chloroplast inhibit photosynthesis

ABA in leaves causes closure of stomata (Inhibit CO<sub>2</sub> fixation)

Cytokinin delays senescence of source and sink

Cytokinin in sink increases photoassimilates import

Ethylene induces senescence process.

## 18. PLANT GROWTH

Growth is defined as a vital process that brings about a permanent and irreversible change in any plant or its part in respect to its size, form, weight and volume. Growth is restricted only to living cells and is accomplished by metabolic processes involving synthesis of macromolecules, such as nucleic acids, proteins, lipids and polysaccharides at the expense of metabolic energy.

Growth at cellular level is also accompanied by the organization of macromolecules into assemblages of membranes, plastids, mitochondria, ribosome and other cell organelles. Cells do not definitely increase in size but divide, giving rise to daughter cells. An important process during cell division is synthesis and replication of nuclear DNA in the chromosomes, which is then passed into the daughter cells. Therefore, the term growth is used to denote an increase in size by cell division and cell enlargement, together with the synthesis of new cellulose materials and the organization of cellular organelles.

### **Growth regions**

Typical growth regions in plants are the apices of shoot and root. Such growing regions are known as apical meristems, primary meristems or regions of primary growth. These apical meristems are responsible for the increase in length, differentiation of various appendages and formation of plant tissues.

### **Phases of growth**

Growth is not a simple process. It occurs in meristematic regions where the meristematic cell has to pass through the following 3 phases.

1. Cell formation phase
2. Cell elongation phase
3. Cell differentiation (cell maturation)

The cell formation phase is represented by meristematic zone and cell enlargement phase by cell elongation zone.

The dividing meristematic cells are thin walled and have dense protoplasm with a large nucleus and with or without very small vacuoles. The intercellular spaces are also absent. The newly formed cells after the first phase of cell division have to pass through the second phase of cell enlargement. During the second phase of cell elongation on account of large quantities of solutes inside the growing cell, water enters the cell due to osmotic effect


resulting in the increased turgidity and expansion and dilation of the thin and elastic cell wall. This phase also results in appearance of large vacuoles.

In the last phase or cell maturation, the secondary walls are laid down and cell matures and gets differentiated into permanent tissue.

### Growth curve

Growth curve is a graph obtained by plotting the growth rate of a plant against time factor. The growth rate of a cell, a plant organ, a whole plant or the whole life cycle of plant is measured in terms of length, size, area, volume or weight. It has been found that different growth phases result in 'S' shaped curve or sigmoid curve. In initial stages during the phase of cell formation, the growth rate increases slowly while it increases rapidly during the phase of cell elongation or cell enlargement and again slows down during the phase of cell maturation.

## GROWTH RATE PHASES - GROWTH CURVE



The period during which the course of growth takes place is known as grand period of growth. Thus, in a standard growth curve, three well marked regions can be observed, the initial growth stage (lag phase), the grand period of growth (exponential or log phase) and the steady stage (maturity stage or senescence or stationary phase). The overall growth may be affected by external or internal factors but the S- shaped curve of grand period of growth is

never influenced. This growth curve suits well to the entire life of an annual plant when measured in terms of dry weight against time.

Early growth of the plant is limited by the amount of food reserves in the seed. When the emerged seedlings develop an adequate root system and enough leaf surfaces to support vigorous photosynthesis and anabolism, a period of rapid increase in size is possible.

High metabolic rates are not maintained indefinitely and eventually processes are set in motion that leads to cessation of growth. The factors responsible for the decrease in growth are competition for essential metabolites, growth substances, water, light or the accumulation of inhibitors, toxic substances or waste materials.

Blackman (1919) suggested that the growth of the plants can be represented by equation.

$$W_1 = W_0 e^{rt}$$

Where,  $W_1$  is the final size (Wt, ht etc) after time  $t$ .  $W_0$  is the initial size at the beginning of the time period.  $r$  is the rate at which plant substance is laid down during time  $t$  and  $e$  is the base of natural logarithm. Blackman pointed out that equation describes the way in which money placed at compound interest increases with time; the term compound interest law is used to describe such phenomenon. In banks, compound interest is usually applied quarterly or annually so that the increase in amount occurs as a jump. With plant system, compound interest is applied continuously and size increase follows a smooth curve.

From the equation, the final size of an organism ( $W_1$ ) depends on the initial size ( $W_0$ ). Larger seed give a larger plant.

In addition, equation shows that plant size also depends on the magnitude of  $r$ , the relative growth rate. Blackman suggested that  $r$  might be used as a measure of the ability of the plant to produce new plant material and called  $r$  as the *efficiency index*. The plants with high efficiency index could be expected to outperform plants with low efficiency index.

### **Measurement of growth**

The measurement of growth is possible in terms of either increase in weight or increase in volume or area. The common and simplest method for the measurement of

growth can be a direct method by which the growth is measured by a scale at regular intervals from beginning to end. The other methods that can be used are horizontal microscope, auxanometers.

### **Factors influencing growth**

Growth is affected by all factors that affect the activity of protoplasm. Both physiological and environmental factors such as water, minerals, photosynthesis, respiration, climate and edaphic factors significantly influence the growth. In general, factors can be grouped into external and internal factors.

#### **External factors**

##### **1. Light**

It has direct effect on photosynthesis and transpiration. Light in terms of intensity, quality and periodicity influence the growth very much.

*Light intensity:* A weak light promotes shortening of internodes and affects expansion of leaf. Very weak light reduces the rate of over all growth and also photosynthesis due to poor development of chlorophyll and higher rate of water loss from the plant.

*Light quality:* The different wavelengths of light have different responses to growth. In blue violet radiation, the internodal growth is pronounced while green colour light promotes the expansion of leaves as compared to complete spectrum of visible light. The red light favours the growth while infra red and UV is detrimental to growth.

*Light duration:* There is remarkable effect of the duration of light on the growth. The induction and suppression of flowering depend on duration

##### **2. Temperature**

The plants have different temperature requirements based on the region where they are grown. In general, best growth takes place between 28 and 33 C. and it varies from temperate to tropical conditions. The optimum temperature requirement is essential for seed germination, growth, metabolic activities, flowering and yield.

##### **3. Oxygen**

The growth of the plant is directly proportional to the amount of oxygen which is essential for respiration during which the food materials are oxidized to release energy.

#### **4. Carbon dioxide**

It is one of the major factors that influence the photosynthesis. The rate of photosynthesis increases as the availability of CO<sub>2</sub> increases while other factors are not limiting.

#### **5. Water**

Water is an essential factor for growth. It is essential for uptake of nutrients, translocation of nutrients and food materials, regulating transpiration and for various physiological processes like photosynthesis, respiration and enzymatic activities.

#### **6. Nutrients and food materials**

The rate of growth is directly proportional to the availability of nutrients and food materials. The shortage of food supply affects the growth as it provides the growth material to the growing region and also it provides the potential energy to the growing region.

#### **Internal factors**

1. Growth hormones and their availability
2. Resistance to climatic, edaphic and biological stresses
3. Photosynthetic rate and respiration
4. Assimilate partitioning and nitrogen content
5. Chlorophyll and other pigments
6. Source-sink relationship and enzyme activities

## 19. GROWTH ANALYSIS

Growth analysis can be used to account for growth in terms that have functional or structural significance. The type of growth analysis requires measurement of plant biomass and assimilatory area (leaf area) and methods of computing certain parameters that describe growth. The growth parameters that are commonly used in agricultural research and the name of the scientists who proposed the parameters are given below.

LAI	-	Williams (1946)
LAR	-	Radford (1967)
LAD	-	Power <i>et al.</i> (1967)
SLA	-	Kvet <i>et al.</i> (1971)
SLW	-	Pearce <i>et al.</i> (1968)
NAR	-	Williams (1946)
CGR	-	Watson (1956)
RGR	-	Williams (1946)
HI	-	Nichiporovich (1951)

### i. Leaf Area

This is the area of photosynthetic surface produced by the individual plant over a period of interval of time and expressed in  $\text{cm}^2 \text{ plant}^{-1}$ .

### ii. Leaf Area Index (LAI)

Williams (1946) proposed the term, Leaf Area Index (LAI). It is the ratio of the leaf of the crop to the ground area over a period of interval of time. The value of LAI should be optimum at the maximum ground cover area at which crop canopy receives maximum solar radiation and hence, the TDMA will be high.

Total leaf area of a plant

$$\text{LAI} = \frac{\text{Total leaf area of a plant}}{\text{Ground area occupied by the plant}}$$

### iii. Leaf Area Ratio (LAR)

The term, Leaf Area Ratio (LAR) was suggested by Radford (1967), expresses the ratio between the area of leaf lamina to the total plant biomass or the LAR reflects the

leafiness of a plant or amount of leaf area formed per unit of biomass and expressed in  $\text{cm}^{-2}$   $\text{g}^{-1}$  of plant dry weight.

$$\text{LAR} = \frac{\text{Leaf area per plant}}{\text{Plant dry weight}}$$

#### iv. Leaf Weight Ratio (LWR)

It was coined by (Kvet *et al.*, 1971) Leaf weight ratio is expressed as the dry weight of leaves to whole plant dry weight and is expressed in  $\text{g g}^{-1}$ .

$$\text{LWR} = \frac{\text{Leaf dry weight}}{\text{Plant dry weight}}$$

#### v. Leaf Area Duration (LAD)

To correlate dry matter yield with LAI, Power *et al.* (1967) integrated the LAI with time and called as Leaf Area Duration. LAD takes into account, both the duration and extent of photosynthetic tissue of the crop canopy. The LAD is expressed in days.

$$\text{LAD} = \frac{L_1 + L_2}{2} \times (t_2 - t_1)$$

$L_1$  = LAI at the first stage  
 $L_2$  = LAI at the second stage,  $(t_2 - t_1)$  = Time interval in days

#### vi. Specific Leaf Area (SLA)

Specific leaf area is a measure of the leaf area of the plant to leaf dry weight and expressed in  $\text{cm}^2\text{g}^{-1}$  as proposed by Kvet *et al.* (1971).

$$\text{SLA} = \frac{\text{Leaf area}}{\text{Leaf weight}}$$

Hence, if the SLA is high, the photosynthesizing surface will be high. However no relationship with yield could be expected.

#### vii. Specific Leaf Weight (SLW)

It is a measure of leaf weight per unit leaf area. Hence, it is a ratio expressed as g cm<sup>-2</sup> and the term was suggested by Pearce *et al.* (1968). More SLW/unit leaf area indicates more biomass and a positive relationship with yield can be expected.

$$\text{SLW} = \frac{\text{Leaf weight}}{\text{Leaf area}}$$

#### viii. Absolute Growth Rate (AGR)

AGR is the function of amount of growing material present and is influenced by the environment. It gives Absolute values of biomass between two intervals. It is mainly used for a single plant or single plant organ e.g. Leaf growth, plant weight etc.

$$\text{AGR} = \frac{h_2 - h_1}{t_2 - t_1} \text{ cm day}^{-1}$$

Where, h<sub>1</sub> and h<sub>2</sub> are the plant height at t<sub>1</sub> and t<sub>2</sub> times respectively.

#### ix. Net Assimilation Rate (NAR)

The term, NAR was used by Williams (1946). NAR is defined as dry matter increment per unit leaf area or per unit leaf dry weight per unit of time. The NAR is a measure of the average photosynthetic efficiency of leaves in a crop community.

$$\text{NAR} = \frac{(W_2 - W_1)}{(t_2 - t_1)} \times \frac{(\log_e L_2 - \log_e L_1)}{(L_2 - L_1)}$$

Where, W<sub>1</sub>and W<sub>2</sub> is dry weight of whole plant at time t<sub>1</sub> and t<sub>2</sub> respectively

L<sub>1</sub> and L<sub>2</sub> are leaf weights or leaf area at t<sub>1</sub> and t<sub>2</sub> respectively

t<sub>1</sub> – t<sub>2</sub> are time interval in days

NAR is expressed as the grams of dry weight increase per unit dry weight or area per unit time ( $\text{g g}^{-1}\text{day}^{-1}$ )

#### **x. Relative Growth Rate (RGR)**

The term was coined by Williams (1946). Relative Growth Rate (RGR) expresses the total plant dry weight increase in a time interval in relation to the initial weight or Dry matter increment per unit biomass per unit time or grams of dry weight increase per gram of dry weight and expressed as unit dry weight / unit dry weight / unit time ( $\text{g g}^{-1}\text{day}^{-1}$ )

$$\text{RGR} = \frac{\log_e W_2 - \log_e W_1}{t_2 - t_1}$$

Where,  $W_1$  and  $W_2$  are whole plant dry weight at  $t_1$  and  $t_2$  respectively  
 $t_1$  and  $t_2$  are time interval in days

#### **xi. Crop Growth Rate (CGR)**

The method was suggested by Watson (1956). The CGR explains the dry matter accumulated per unit land area per unit time ( $\text{g m}^{-2} \text{ day}^{-1}$ )

$$\text{CGR} = \frac{(W_2 - W_1)}{\rho (t_2 - t_1)}$$

Where,  $W_1$  and  $W_2$  are whole plant dry weight at time  $t_1 - t_2$  respectively

$\rho$  is the ground area on which  $W_1$  and  $W_2$  are recorded.

CGR of a species are usually closely related to interception of solar radiation

#### **xii. Total dry matter production (TDMP) and its distribution**

The TDMP is the biomass accumulated by the whole plant over a period of interval of time and its distribution (allocation) to different parts of the plant such as roots, stems, leaves and the economic parts which controls the sink potential.

#### **xiii. Translocation percentage (TP)**

The term translocation percentage indicates the quantum of photosynthates translocated from source (straw) to the grain (panicle/grains) from flowering to harvest.

$$TP = \frac{\text{Straw weight at flowering} - \text{straw weight at harvest}}{\text{Panicle weight at flowering} - \text{panicle weight at harvest}}$$

#### xiv. Light extinction coefficient

It is the ratio of light intercepted by crop between the top and bottom of crop canopy to the LAI.

$$K = \frac{\log_e I / I_o}{LAI}$$

Where,  $I_o$  and  $I$  are the light intensity at top and bottom of a population with LAI

#### xv. Light Transmission Ratio (LTR)

It is expressed as the ratio of quantum of light intercepted by crop canopy at top to the bottom. Light intensity is expressed in K lux or  $W m^{-2}$

$$LTR = I / I_o$$

Where,  $I$  : light intercepted at the bottom of the crop canopy

$I_o$ : light intercepted at the top of the crop canopy

#### xvi. Dry Matter Efficiency (DME)

It is defined as the percent of dry matter accumulated in the grain from the total dry matter produced over the crop growth period.

$$DME = \frac{\text{Grain yield}}{\text{TDMP}} \times \frac{100}{\text{Duration of crop}}$$

#### xvii. Unit area efficiency (UAE)

It is expressed as the quantum of grain yield produced over a unit land area for a specified crop growth period.

$$\text{UAE} = \frac{\text{Grain yield}}{\text{Land area}} \times \frac{1}{\text{Duration of crop}}$$

### xviii) Harvest Index

The harvest index is expressed as the percent ratio between the economic yield and total biological yield and was suggested by Nichiporovich (1951).

$$\text{HI} = \frac{\text{Economic yield}}{\text{Total biological yield}} \times 100$$

## 20. PLANT GROWTH REGULATORS

Plant growth regulators or phytohormones are organic substances produced naturally in higher plants, controlling growth or other physiological functions at a site remote from its place of production and active in minute amounts. Thimmann (1948) proposed the term *Phyto hormone* as these hormones are synthesized in plants. *Plant growth regulators* include auxins, gibberellins, cytokinins, ethylene, growth retardants and growth inhibitors. Auxins are the hormones first discovered in plants and later gibberellins and cytokinins were also discovered.

### **Hormone**

An endogenous compound, which is synthesized at one site and transported to another site where it exerts a physiological effect in very low concentration. But ethylene (gaseous nature), exert a physiological effect only at a near a site where it is synthesized.

Classified definition of a hormone does not apply to ethylene.

### **Plant growth regulators**

- Defined as organic compounds other than nutrients, that affects the physiological processes of growth and development in plants when applied in low concentrations.
- Defined as either natural or synthetic compounds that are applied directly to a target plant to alter its life processes or its structure to improve quality, increase yields, or facilitate harvesting.

### **Plant Hormone**

When correctly used, is restricted to naturally occurring plant substances, there fall into five classes. Auxin, Gibberellins, Cytokinin, ABA and ethylene. Plant growth regulator includes synthetic compounds as well as naturally occurring hormones.

### **Plant Growth Hormone**

The primary site of action of plant growth hormones at the molecular level remains unresolved.

#### **Reasons**

- Each hormone produces a great variety of physiological responses.
- Several of these responses to different hormones frequently are similar.
- The response of a plant or a plant part to plant growth regulators may vary with the variety of the plant.
- Even a single variety may respond differently depending on its age, environmental conditions and physiological state of development (especially its natural hormone content) and state of nutrition. There are always exceptions for a general rule suggesting the action of a specific growth regulator on plants.
- There are several proposed modes of action in each class of plant hormone, with substantial arguments for and against each mode.


#### **Hormone groups**

Auxin	-	Substances generally resembles IAA and has the ability to stimulate the elongation of coleoptiles.
Gibberellins	-	are diterpenoids, which have the ability to elongate the stem of green seedlings especially certain dwarf and rosette types.
Cytokinin	-	Usually substituted Adenines, which resembles zeatin (Naturally occurring cytokinin in <i>Zea mays</i> ) and have the ability to stimulate cytokinensis in cultures of tobacco cells.
Ethylene	-	Gaseous regulator that stimulate is diametric growth in the apices of dicot seedlings.
Inhibitors	-	are regulators of growth, which originally depress the

#### **Auxins**

Auxins are a group of phytohormones produced in the shoot and root apices and they migrate from the apex to the zone of elongation. Auxins promote the growth along the longitudinal axis of the plant and hence the name (auxeing : to grow). The term, auxin was introduced by Kogl and Haagen- Smit (1931). Went (1928) isolated auxin from the *Avena* coleoptile tips by a method called *Avena coleoptile or curvature test* and concluded that no growth can occur without auxin. Auxins are widely distributed through out the plant however, abundant in the growing tips such as coleoptile tip, buds, root tips and leaves. Indole Acetic Acid (IAA) is the only naturally occurring auxin in plants. The synthetic auxins include,

### Avena Curvature Test



IBA : Indole Butyric Acid

NAA : Naphthalene Acetic acid

MENA: Methyl ester of Naphthalene acetic acid

MCPA: 2 Methyl 4 chloro phenoxy acetic acid

TIBA : 2, 3, 5 Tri iodo benzoic acid

2, 4-D : 2, 4 dichloro phenoxy acetic acid

2, 4, 5-T: 2, 4, 5 – Trichloro phenoxy acetic acid

Natural auxins may occur in the form of either *free auxins*- which freely move or diffuse out of the plant tissues readily or *bound auxins*- which are released from plant tissues only after hydrolysis, autolysis or enzymolysis.

### **Physiological effects of auxin**


#### **1. Cell division and elongation**

The primary physiological effects of auxin are cell division and cell elongation in the shoots. It is important in the secondary growth of stem and differentiation of xylem and phloem tissues.

#### **2. Apical dominance**

In many plants, if the terminal bud is intact and growing, the growth of lateral buds just below it remains suppressed. Removal of the apical bud results in the rapid growth of lateral buds. This phenomenon in which the apical bud dominates over the lateral buds and does not allow the lateral buds to grow is known as *apical dominance*.

Skoog and Thimann (1948) pointed out that the apical dominance might be under the control of auxin produced at the terminal bud and which is transported downward through the stem to the lateral buds and hinders the growth. They removed the apical bud and replaced it with agar block. This resulted in rapid growth of lateral buds. But when they replaced the apical bud with agar block containing auxin, the lateral buds remained suppressed and did not grow.



### 3. Root initiation

In contrast to stem, the higher concentration of auxin inhibits the elongation of roots but the number of lateral roots is considerably increased i.e., higher concentration of auxin induces more lateral branch roots. Application of IAA in lanolin paste (lanolin is a soft fat prepared from wool and is good solvent for auxin) to the cut end of a young stem results in an early and extensive rooting. This fact is of great practical importance and has been widely utilized to promote root formation in economically useful plants which are propagated by cuttings.

### 4. Prevention of abscission

Natural auxins prevent the formation of abscission layer which may otherwise result in the fall of leaves, flowers and fruits.

### 5. Parthenocarpy

Auxin can induce the formation of parthenocarpic fruits (fruit formation without pollination and fertilization). In parthenocarpic fruits, the concentration of auxin in the ovaries is higher than in the ovaries of plants which produce fruits only after fertilization. In

in the later cases, the concentration of the auxin in ovaries increases after pollination and fertilization.

## **6. Respiration**

Auxin stimulates respiration and there is a correlation between auxin induced growth and respiration. Auxin may increase the rate of respiration indirectly through increased supply of ADP by rapidly utilizing ATP in the expanding cells.

## **7. Callus formation**

Besides cell elongation, auxin may also be active in cell division. In many tissue cultures, where the callus growth is quite normal, the continued growth of such callus takes place only after the addition of auxin.

## **8. Eradication of weeds**

Some synthetic auxins especially 2, 4- D and 2, 4, 5-T are useful in eradication of weeds at higher concentrations.

## **9. Flowering and sex expression**

Auxins generally inhibit flowering but in pine apple and lettuce it promotes uniform flowering.

### **Distribution of auxin in plants**

In plants, auxin (IAA) is synthesized in growing tips or meristematic regions from where; it is transported to other plant parts. Hence, the highest concentration of IAA is found in growing shoot tips, young leaves and developing auxiliary shoots. In monocot seedling, the highest concentration of auxin is found in coleoptile tip which decreases progressively towards its base.

In dicot seedlings, the highest concentration is found in growing regions of shoot, young leaves and developing auxiliary shoots. Within the plants, auxin may present in two forms. i.e., *free auxins* and *bound auxins*. Free auxins are those which are easily extracted by various organic solvents such as diethyl ether. Bound auxins on the other hand, need more drastic methods such as hydrolysis, autolysis, enzymolysis etc. for extraction of auxin. Bound

auxins occur in plants as complexes with carbohydrates such as glucose, arabionse or sugar alcohols or proteins or amino acids such as aspartate, glutamate or with inositol.

### **Biosynthesis of auxin (IAA) in plants**

Thimann (1935) found that an amino acid, tryptophan is converted into Indole 3 acetic acid. Tryptophan is the primary precursor of IAA in plants. IAA can be formed from tryptophan by two different pathways.

1. By deamination of tryptophan to form indole-3-pyruvic acid followed by decarboxylation to from indole-3-acetaldehyde. The enzymes involved are tryptophan deamintion and indole pyruvate decarboxylase respectively.
2. By decarboxylation of tryptophan to form tryptamine followed by deamination to form indole-3-acetaldehyde and the enzymes involved are tryptophan decarboxylase and tryptamine oxidase respectively. Indole 3-acetaldehyde can readily be oxidized to indole 3-acetic acid (IAA) in the presence of indole 3-acetaldehyde dehydrogenase.


### **Transport of auxin in plant**

The transport of auxin is predominantly polar. In stems, polar transport of auxin is basipetal i.e., it takes place from apex towards base. Polar transport of auxin is inhibited by 2, 3, 5 Triiodobenzoic acid (TIBA) and Naphthyl thalamic acid (NPA). The substances are called as antiauxins.

### **Destruction / Inactivation of auxin in plants**

Auxin is destructed by the enzyme IAA oxidase in the presence of O<sub>2</sub> by oxidation.

IAA Oxidase



Rapid inactivation may also occur by irradiation with x-rays and gamma rays. UV light also reduces auxin levels in plants. Inactivation or decomposition of IAA by light has been called as photo oxidation.

### **Mechanism of Action**

IAA increases the plasticity of cell walls so that the cells stretch easily in response to turgor pressure. It has been suggested that IAA acts upon DNA to influence the production of mRNA. The mRNA codes for specific enzymes responsible for expansion of cell walls. Recent evidences indicate that IAA increases oxidative phosphorylation in respiration and enhanced oxygen uptake. The growth stimulation might be due to increased energy supply and it is also demonstrated that auxin induces production of ethylene in plants.

## **Gibberellins**

### **Discovery**

A Japanese scientist Kurosawa found that the rice seedlings infected by the fungus *Gibberella fujikuroi* grow taller and turned very thin and pale. An active substance was isolated from the infected seedlings and named as Gibberellin.

### Biosynthesis of gibberellins in plants

The primary precursor for the formation of gibberellins is acetate.

Acetate + COA → Acetyl COA → Mevalonic acid → MA pyrophosphate → Isopentanyl pyrophosphate → Geranyl pyrophosphate → GGPP → Kaurene → Gibberellins.

### **Physiological effects of gibberellins**

#### **1. Seed germination**

Certain light sensitive seeds eg. Lettuce and tobacco show poor germination in dark. Germination starts vigorously if these seeds are exposed to light or red light. This requirement of light is overcome if the seeds are treated with gibberellic acid in dark.

#### **2. Dormancy of buds**

In temperature regions the buds formed in autumn remain dormant until next spring due to severe cold. This dormancy of buds can be broken by gibberellin treatments. In potato also, there is a dormant period after harvest, but the application of gibberellin sprouts the refer vigorously.

### **3. Root growth**

Gibberellins have little or no effect on root growth. At higher concentration, some inhibition of root growth may occur. The initiation of roots is markedly inhibited by gibberellins in isolated cuttings.

### **4. Elongation of internodes**

The most pronounced effect of gibberellins on the plant growth is the elongation of the internodes. Therefore in many plants such as dwarf pea, dwarf maize etc gibberellins overcome the genetic dwarfism.

### **5. Bolting and flowering**

In many herbaceous plants, the early period of growth shows rosette habit with short stem and small leaves. Under short days, the rosette habit is retained while under long days bolting occurs i.e. the stem elongates rapidly and is converted into polar axis bearing flower primordia. This bolting can also be induced in such plants by the application of gibberellins even under non-inductive short days.

In *Hyoscyamus niger* (a long day plant) gibberellin treatment causes bolting and flowering under non-inductive short days. While in long day plants the gibberellin treatment usually results in early flowering. In short day plants, its effects are quite variable. It may either have no effect or inhibit or may activate flowering.

### **6. Parthenocarpy**

Germination of the pollen grains is stimulated by gibberellins; likewise, the growth of the fruit and the formation of parthenocarpic fruits can be induced by gibberellin treatment. In many cases, eg. pome and stone fruits where auxins have failed to induce parthenocarpy, the gibberellins have proven to be successful. Seedless and fleshly tomatoes and large sized seedless grapes are produced by gibberellin treatments on commercial scale.

## 7. Synthesis of the enzyme $\alpha$ - amylase

One important function of gibberellins is to cause the synthesis of the enzyme  $\alpha$  - amylase in the aleurone layer of the endosperm of cereal grains during germination. This enzyme brings about hydrolysis of starch to form simple sugars which are then translocated to growing embryo to provide energy source.

### Distribution of gibberellins in plant

Gibberellins are found in all parts of higher plants including shoots, roots, leaves, flower, petals, anthers and seeds. In general, reproductive parts contain much higher concentrations of gibberellins than the negative parts. Immature seeds are especially rich in gibberellins (10-100 mg per g fresh weight).

In plants, gibberellins occur in two forms free gibberellins and bound gibberellins. Bound gibberellins usually occur as gibberellin – glycosides.

## CYTOKININS (Kinetin)

Kinetin was discovered by Skoog and Miller (1950) from the tobacco pith callus and the chemical substance was identified as 6-furfuryl aminopurine. Because of its specific effect on *cytokinesis* (cell division), it was called as cytokinins or kinetin. The term, cytokinin was proposed by Letham (1963). Fairley and Kingour (1966) used the term, *phytokinins* for cytokinins because of their plant origin. Chemically cytokinins are kinins and they are purine derivatives.

Cytokinins, besides their main effect on cell division, also regulate growth and hence they are considered as natural plant growth hormones. Some of the very important and commonly known naturally occurring cytokinins are Coconut milk factor and Zeatin. It was also identified that cytokinin as a constituent of t-RNA.

## Naturally occurring cytokinins

Cytokinins can be extracted from coconut milk (liquid endosperm of coconut), tomato juice, flowers and fruits of *Pyrus malus*; fruits of *Pyrus communis* (Pear), *Prunus cerasiferae* (plum) and *Lycopersicum esculentum* (bhendi); Cambial tissues of *Pinus radiata*, *Eucalyptus regnans* and *Nicotiana tabacum*; immature fruits of *Zea mays*, *Juglans* sp. and *Musa* sp; female gametophytes of *Ginkgo biloba*; fruitlets, embryo and endosperms of *Prunus persica*; seedling of *Pisum sativum*; root exudates of *Helianthus annuus* and tumour tissues of tobacco. According to Skoog and Armstrong (1970), at least seven well established types of cytokinins have been reported from the plants.

## Biosynthesis

It is assumed that cytokinins are synthesised as in the case of purines in plants (nucleic acid synthesis). Root tip is an important site of its synthesis. However, developing seeds and cambial tissues are also the site of cytokinin biosynthesis. Kende (1965) reported that cytokinins move upwards perhaps in the xylem stream. However, basipetal movement in petiole and isolated stems are also observed. Seth *et al* (1966) found that auxin enhances kinetin movement (translocation) in bean stems.

## Physiological effects of cytokinins

### 1. Cell division

The most important biological effect of kinetin on plants is to induce cell division especially in tobacco pith callus, carrot root tissue, soybean cotyledon, pea callus etc.

### 2. Cell enlargement

Like auxins and gibberellins, the kinetin may also induce cell enlargement. Significant cell enlargement has been observed in the leaves of *Phaseolus vulgaris*, pumpkin cotyledons, tobacco pith culture, cortical cells of tobacco roots etc.

### 3. Concentration of apical dominance

External application of cytokinin promotes the growth of lateral buds and hence counteracts the effect of apical dominance

#### 4. Dormancy of seeds

Like gibberellins, the dormancy of certain light sensitive seeds such as lettuce and tobacco can also be broken by kinetin treatment.

#### 5. Delay of senescence ( Richmand - Lang effect)

The senescence of leaves usually accompanies with loss of chlorophyll and rapid breakdown of proteins. Senescence can be postponed to several days by kinetin treatment by improving RNA synthesis followed by protein synthesis.

Richmand and Lang (1957) while working on detached leaves of *Xanthium* found that kinetin was able to postpone the senescence for a number of days.

#### 6. Flower induction

Cytokinins can be employed successfully to induce flowering in short day plants.

#### 7. Morphogenesis

It has been shown that high auxin and low kinetin produced only roots whereas high kinetin and low auxin could promote formation of shoot buds.

#### 8. Accumulation and translocation of solutes

Plants accumulate solutes very actively with the help of Cytokinin and also help in solute translocation in phloem.

#### 9. Protein synthesis

Osborne (1962) demonstrated the increased rate of protein synthesis due to translocation by kinetin treatment.

#### 10. Other effects

Cytokinins provide resistance to high temperature, cold and diseases in some plants. They also help in flowering by substituting the photoperiodic requirements. In some cases, they stimulate synthesis of several enzymes involved in photosynthesis.

#### 11. Commercial applications

Cytokinins have been used for increasing shelf life of fruits, quickening of root induction and producing efficient root system, increasing yield and oil contents of oil seeds like ground nut.

### **Ethylene**

Ethylene is the only natural plant growth hormone exists in gaseous form.

#### **Important physiological elects**

1. The main role of ethylene is it hastens the ripening of fleshy fruits eg. Banana, apples, pears, tomatoes, citrus etc.
2. It stimulates senescence and abscission of leaves
3. It is effective in inducing flowering in pine apple
4. It causes inhibition of root growth
5. It stimulates the formation of adventitious roots
6. It stimulates fading of flowers
7. It stimulates epinasty of leaves.

### **Abscisic acid**

Addicott (1963) isolated a substance strongly antagonistic to growth from young cotton fruits and named Abscissin II. Later on this name was changed to Abscisic acid. This substance also induces dormancy of buds therefore it also named as Dormin.

Abscisic acid is a naturally occurring growth inhibitor.

### **Physiological effects**

The two main physiological effects are

1. Geotropism in roots
2. Stomatal closing
3. Besides other effects

### 1. Geotropism in roots

Geotropic curvature of root is mainly due to translocation of ABA in basipetal direction towards the root tip.

### 2. Stomatal closing

ABA is synthesized and stored in mesophyll chloroplast. In respond to water stress, the permeability of chloroplast membrane is lost which resulted in diffusion of ABA out of chloroplast into the cytoplasm of the mesophyll cells. From mesophyll cells it diffuses into guard cells where it causes closing of stomata.

### 3. Other effects

- i. Including bud dormancy and seed dormancy
- ii. Includes tuberisation
- iii. Induces senescence of leaves fruit ripening, abscission of leaves, flowers and fruits
- iv. Increasing the resistance of temperate zone plants to frost injury.

### **Growth retardants**

There is no. of synthesis compounds which prevent the gibberellins from exhibiting their usual responses in plants such as cell enlargement or stem elongation. So they are called as anti gibberellins or growth retardants. They are

1. Cycocel (2- chloroethyl trimethyl ammonium chloride (CCC)
2. Phosphon D – (2, 4 – dichlorobenzyl – tributyl phosphonium chloride)
3. AMO – 1618
4. Morphactins

5. Maleic hydrazide

## 21. PRACTICAL APPLICATION OF PLANT GROWTH REGULATORS IN CROP PRODUCTIVITY

### **Commercial uses of growth regulators**

#### **1. Rooting and plant propagation**

- a) Auxin compound like IBA NAA, 2,4-D, 2, 4,5-T
- b) IBA produces strong fibrous root system

#### **2. Germination and dormancy**

- a) Gibberellin is a potent germination promoter
- b) Abscisic acid – germination inhibitor (Anti – Gibberellin)
- c) Induce Dormancy - ABA
- d) Breaking of dormancy - Auxins and Gibberellin

#### **3. Fruit set and Development**

- a) Fruit setting → 2, 4, 5 – T
- b) Fruit size increment in grapes → Gibberellic acid
- c) Shelf life increment in fruits and flowers → Cytokinin
- d) Good fruit shape ----- Gibberellic acid + Cytokinin
- e) Parthenocarpic fruit – Gibberellins, IAA and PAA

#### **4. Sex expression**

Production of male flowers → Gibberellins (cucumber)

Production of female flowers → Auxins and Gibberellins  
 ↓                            ↓  
 Cucumber                 maize

#### **5. Abscission**

Control of abscission → NAA and IAA  
 Induce Abscission → Ethrel

### **6. Morphogenesis**

Auxin and Cytokinin

### **7. Weed control**

2, 4-D and 2, 4, 5-T

### **8. Plant organ size**

Increases plant height → GA  
 Shorten the plant height → TIBA  
    ↓  
    Tri iodo benzoic acid

Increases Tillering → Cytokinin  
 Ex: BAP (Benzyl amino purine) and TIBA

### **9. Antitranspirants → ABA and PMA**

↓  
 Phenyl mercury acetate

**10. Papaya Latex flow** → Ethephon

**11. Rubber latex flow** → 2, 4 – D and 2, 4,5 – T

**12. Fruit ripening** → Ethrel

**13. Sugarcane ripeners** → Glyphosphate and CCC


Hormone	Major Functions	Where Produced
Auxin	Stimulates cell elongation; involved in phototropism, gravitropism, apical dominance, and vascular differentiation; stimulates ethylene synthesis and induces adventitious roots on cuttings	Meristems of apical buds, embryo of seed, young leaves
Cytokinin	Stimulates cell division, reverses apical dominance, involved in shoot growth, delays leaf senescence	Synthesized in roots and transported to other organs
Ethylene	Stimulates fruit ripening, leaf and flower senescence, and abscission	Tissues of ripening fruits, nodes of stems, senescent leaves and flowers
Abscisic Acid	Inhibits growth, stimulates stomatal closure, maintains dormancy	Leaves, stems, green fruit
Gibberellin	Stimulates shoot elongation, stimulates bolting and flowering in biennials, regulates production of hydrolytic enzymes in grains	Meristems of apical buds and roots, young leaves, embryo

## 22. ENVIRONMENTAL STRESSES

The occurrence of unfavorable environmental factors such as moisture deficit / excess, high radiation, low and high temperature, salinity of water and soil, nutrient deficiency or toxicity and pollution of atmosphere, soil and water are likely to affect the crop growth in terms of morphology (plant size, architecture, malformation of plant organs, growth (height, volume, weight), physiological and metabolic processes and yield of crop plants.

### Stress and strain

Any environmental factor potentially unfavorable to plant is termed as **stress**. The effect of stress on plant condition is called **strain**. According to Newton's law of motion, a force is always accompanied by a counterforce, for an action there is always equal and opposite reaction. Stress is the action and whereas strain is the reaction.



### I. DROUGHT (Water stress)

*Drought is defined as the deficiency of water severe enough to check the plant growth.* Drought has been classified into two broad categories viz., soil drought and atmospheric drought. Soil drought leads to atmospheric drought. Atmospheric drought occurs due to low atmospheric humidity, high wind velocity and high temperature which cause a plant to lose most of its water.

### **Physiological changes occur due to drought**

#### 1. Functioning of stomata

In general, stomata lose their function and may die, because wilting after certain limit denatures the starch in the guard cells and also in the mesophyll cells.

#### 2. Carbohydrates metabolism in green leaves

The very first effect of drought on carbohydrates metabolism is that starch disappears from the wilted leaves and sugar accumulates simultaneously.

#### 3. Photosynthetic activity

CO<sub>2</sub> diffusion into the leaf is prevented due to decrease in stomatal opening and thereby reduces photosynthetic activity in green cells.

#### 4. Osmotic pressure

The reduced amount of water during drought causes an increase in the osmotic pressure of plant cell. This increase in osmotic pressure permits the plant to utilize better soil moisture.

#### 5. Permeability

The permeability to water and urea increases during drought.

#### 6. Biochemical effects

Water shortage alters the chemical composition. For example, starch is converted to sugar, besides this, there is a considerable increase in nitrate nitrogen and protein synthesis is adversely affected.

### **Adaptation to drought**

#### **Drought resistance**

*Drought resistance is defined as the capacity of plants to survive during the period of drought with little or no injury.* There are three important categories of plants growing in the areas facing drought. They are ephemerals, succulents and non-succulent perennials

### **1. Ephemerals**

These are short lived plants and they complete their life cycle within a short favourable period during rainy season. They pass dry periods in the form of seeds. They are called as *drought escaping plants*.

### **2. Succulent plants**


These plants accumulate large quantities of water and use it slowly during dry period. Thus, they pass dry periods or drought without facing it. Such plants develop several morphological adaptations for reducing transpiration such as thick cuticle, reduced leaf area, sunken stomata etc.

### **3. Non succulent plants**

These plants are in fact the real drought enduring (tolerant) plants. They tolerate drought without adapting any mechanism to ensure continuous supply of water. They develop many morphological adaptations which are collectively called *xeromorphy*. They develop, in general, greyish colour, reflecting surfaces, smaller leaves, extensive root system, leaf fall during dry season, sunken stomata and thick cuticle etc. They develop an elaborated conducting system. The stomata remain closed mostly in dry periods.

The plants develop several protoplasmic peculiarities such as cell size, cell structure, increased permeability, increased imbibition power, elasticity, small vacuoles, higher osmotic pressure etc.

### Osmotic adjustment



### Methods to overcome drought

- Selection of drought tolerant species
- Adjusting the time of sowing in such a way that the crop completes its lifecycle before the onset of drought
- Seed hardening with KCl, KH<sub>2</sub>PO<sub>4</sub>, CaCl<sub>2</sub> or Thiourea
- Thinning of poorly established plants
- Mulching to minimize the evaporative loss
- Foliar spray of antitranspirants such as Kaolin, PMA, Waxes and Silicone oils
- Foliar spray of KCl
- Foliar spray of growth retardants such as CCC and MC

## **HIGH MOISTURE STRESS - FLOODING / WATER LOGGING**

Water logging refers to a condition when water is present in excess amount than its optimum requirement. It creates an anaerobic situation in the rhizosphere due to which the plant experiences the stress ( $O_2$  deficient stress).

### **Nature of Water logging Stress**

In the water logged soils, water gets filled in the pores of the soil which are previously occupied by  $O_2$ . Such soils suffer  $O_2$  deficiency.

This  $O_2$  deficiency depresses growth and survival of plants growing in it.

Flood sensitive plants (eg. Tomato, soybean and sunflower) are killed in the water logged conditions, while the tolerant species (eg. Rice) withstand water logging for a considerable time. However, continuous submergence of rice for more than 10 days is also deleterious resulting in death and decay of the plants.

### **Plant Water Relations in Flooding Stress**

The flooding often induces stomatal closure mostly in  $C_3$  plants. This causes lower water flow in these plants. This also results in leaf dehydration because of reduced root permeability. Ultimately, wilting of leaves occurs due to the restricted water flow from the roots to the shoots.

Occurrence of these changes in leaves, shoots or roots is due to the transfer of toxic substances (acetaldehyde / alcohol) produced under anaerobic conditions in the roots as well as the levels of PGRs transported from the roots to shoots via transpiration stream.

### **Levels of Endogenous PGRs under Flooding Stress**

Endogenous levels of PGRs such as GA and cytokinins (CK) are reduced in the roots. This has enhanced levels of ABA and ethylene in the shoots causing stomatal closure and early onset of senescence respectively.

It is also reported that levels of auxins are reduced and that of Aminocyclopropane -1-Carboxylic Acid (ACC), precursor for the ethylene biosynthesis are increased under flooding stress.

Important roles played by these endogenous PGRs during high moisture (flooding) stress are summarized in the following table.

### **Effect of flooding stress on the endogenous levels of PGRs and their effect on plants**

<b>Sl. No.</b>	<b>Level of PGR in plants</b>	<b>Effects on plants under water logging</b>
01.	Reduced Auxins	Causes “Hypertrophy” (Swelling of stem base by collapse or enlargement of cells in cortex)
02.	Decreased GA	Causes reduction in cell enlargement and stem elongation
03.	Decreased CK	Results in early on-set of senescence and reduced rate of assimilate partitioning to the sinks
04.	Increased ABA	Cause stomatal closure with consequential decrease in the rate of gas exchanges during photosynthesis, respiration and transpiration; results in efflux of K <sup>+</sup> from the guard cells; decreases ion transport due to lower rate of transpiration; decrease the starch formation in the guard cells resulting in stomatal closure
05.	Increased Ethylene	Causes “Epinasty” of leaves (uneven growth of leaves due to more cell elongation on upper side than the lower side of the leaf); induces senescence and Hypertrophy in plants.

Thus, the O<sub>2</sub> stress in the roots under flooding produces signals, via transpiration stream, to the leaves affecting stomatal behaviour ultimately.

#### **Mitigation of High Moisture (Water logging) Stress**

1. Providing adequate drainage for draining excessive stagnating water around the root system.
2. Spray of growth retardant of 500 ppm cycocel for arresting apical dominance and thereby promoting growth of laterals
3. Foliar spray of 2% DAP + 1% KCl (MOP)
4. Nipping terminal buds for arresting apical dominance and thus promoting growth sympodial branches (as in cotton) for increasing productivity
5. Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods
6. Spray of 0.5 ppm brassinolide for increasing photosynthetic activity

7. Foliar spray of 100 ppm salicylic acid for increasing stem reserve utilization under high moisture stress
8. Foliar spray of 0.3 % Boric acid + 0.5 % ZnSO<sub>4</sub> + 0.5 % FeSO<sub>4</sub> + 1.0 % urea during critical stages of the stress.

## SALT STRESS

Salt stress occurs due to excess salt accumulation in the soil. As a result, water potential of soil solution decreases and therefore exosmosis occurs. This leads to physiological drought causing wilting of plants.

**Classification of saline soil:** 1. Saline soil 2. Alkaline soil

### 1. Saline soil

In saline soils, the electrical conductivity is greater than 4 dS/m, exchangeable sodium percentage is less than 15% and pH is less than 8.5. These soils are dominated by Cl<sup>-</sup> and SO<sub>4</sub><sup>2-</sup> ions.

### 2. Alkaline soil

Alkaline soils are also termed as sodic soils wherein, the electrical conductivity is less than 4 dS/m, exchangeable sodium percentage is greater than 15% and pH of the soil is greater than 8.5. These soils are dominated by CO<sub>3</sub><sup>2-</sup> and HCO<sub>3</sub><sup>-</sup> ions.

## Classification of plants

Plants are classified into two types based on the tolerance to salt stress. They are halophytes and glycophytes.

### 1. Halophytes

Halophytes are the plants that grow under high salt concentrations. They are again divided into two types based on extreme of tolerance.

*Euhalophytes*: can tolerate extreme salt stress

*Oligohalophytes*: can tolerate moderate salt stress

### 2. Glycophytes

Glycophytes are the plants that cannot grow under high salt concentration.

## **Effect of salt stress on plant growth and yield**

### 1. Seed germination

Salt stress delays seed germination due to the reduced activity of the enzyme,  $\alpha$ -amylase

### 2. Seedling growth

The early seedling growth is more sensitive. There is a significant reduction in root emergence, root growth and root length.

### 3. Vegetative growth

When plants attain vegetative stage, salt injury is more severe only at high temperature and low humidity. Because under these conditions, the transpiration rate will be very high as a result uptake of salt is also high.

### 4. Reproductive stage

Salinity affects panicle initiation, spikelet formation, fertilization and pollen grain germination.

### 5. Photosynthesis

Salinity drastically declines photosynthetic process. Thylakoid are damaged by high concentration of salt and chlorophyll *b* content is drastically reduced.

## **Mechanism of salt tolerance**

1. Some plants are able to maintain high water potential by reducing the transpiration rate.
2. Salts are accumulated in stem and older leaves in which metabolic processes take place in a slower rate.
3.  $\text{Na}^+$  (sodium ion) toxicity is avoided by accumulating high amount of  $\text{K}^+$  ions.
4. Accumulation of toxic ions in the vacuole but not in the cytoplasm.
5. Accumulation of proline and abscissic acid which are associated with tolerance of the plants to salt.

## **Relative salt tolerant crops**

Tolerant crops: Cotton, sugar cane, barley

Semi tolerant crops: Rice, maize, wheat, oats, sunflower, soybean

Sensitive crops: Cow pea, beans, groundnut and grams

### Mitigation of Salt Stress

1. Seed hardening with NaCl (10 mM concentration)
2. Application of gypsum @ 50% Gypsum Requirement (GR)
3. Incorporation of daincha (6.25 t/ha) in soil before planting
4. Foliar spray of 0.5 ppm brassinolide for increasing photosynthetic activity
5. Foliar spray of 2% DAP + 1% KCl (MOP) during critical stages
6. Spray of 100 ppm salicylic acid
7. Spray of 40 ppm of NAA for arresting pre-mature fall of flowers / buds / fruits
8. Extra dose of nitrogen (25%) in excess of the recommended
9. Split application of N and K fertilizers
10. Seed treatment + soil application + foliar spray of Pink Pigmented Facultative Methnaotrops (PPFM) @  $10^6$  as a source of cytokinins.

## TEMPERATURE STRESS

Temperature stress includes both high temperature stress and low temperature stress. Low temperature stress causes chilling injury and freezing injury.

### Low temperature stress

#### 1. Chilling injury

The tropical origin plants are injured when the temperature drops to some point close to 0°C. The injury which occurs due to low temperature but above zero degree centigrade is called chilling injury.

#### 2. Freezing injury

Freezing injury occurs when the temperature is 0°C or below.

### Effect of freezing and chilling injury plants

- The lipid molecules in cell membrane get solidified i.e. changed from liquid state to solid state. Hence, the semi-permeable nature of the membrane is changed and the membrane becomes leaky.
- Inactivation of mitochondria
- Streaming of protoplasm is stopped
- Accumulation of respiratory metabolites which become highly toxic
- Ice formation inside the cell occurs.

### **Prevention of cold injury**

- Some plants change the pattern of growth.
- The growth is completely arrested during this period.
- In cell membrane, unsaturated fatty acid content is increased.
- Intracellular ice formation is reduced.
- The quantity of free enzymes, sugars and proteins increases.

### **High temperature stress**

The effect of high temperature is heat Injury. Heat Injury occurs when plant temperature is higher than that of environment (exceeds 35°C).

#### **General effects of high temperature**

High temperature affects

1. Seedling growth and vigour
2. Water and nutrient uptake
3. Solute transport
4. Photosynthesis and respiration
5. General metabolic processes
6. Fertilization and maturation

#### **Cellular Changes during heat stress**

When plants are exposed to temperatures higher than 45°C it experiences heat stress.

The cellular changes due to heat stress are


1. Disruption of cytoskeleton and microtubules.
2. Fragmentation of golgi complex
3. Increase in number of lysosomes
4. Swelling of mitochondria thereby resulting in decreased respiration and oxidative phosphorylation
5. Disruption of normal protein synthesis
6. Disappearance of polysomes
7. Disruption of splicing of mRNA precursors
8. Cessation of pre-RNA processing
9. Decline in transcription by RNA polymerase I

10. Inhibition of chromatin assembly
11. Decline in DNA synthesis

### **Acclimation to high temperature**

#### **Morphological Adaptations**

- Reflective leaf hair
- Leaf waxes
- Leaf orientation
- Maximize conductive or convective loss of heat



Zeaxanthin decreases membrane fluidity and stabilises the membrane

#### **Heat Shock Proteins (HSPs)**

Plants have the capacity to interact with the environment in many different ways and to survive under extreme abiotic and perhaps also biotic stress conditions. The response to heat stress (hs) is highly conserved in organisms but owing to the sessile life style it is of utmost importance to plants. The hs-response is characterised by (i) a transient alteration of gene expression (synthesis of heat shock proteins: HSP) and (ii) by the acquisition of a higher level of stress tolerance (acclimation). The induction of HSP-expression is not restricted to high temperature stress, HSPs are also linked to a number of other abiotic stresses including cold, freezing, drought, dehydration, heavy metal, and oxidative stresses. HSPs are molecular chaperones, which either prevent complete denaturation (small HSP: sHSP) or are supporting proper folding (other HSP) of enzymes under or after protein denaturing conditions. Manipulation of the hs-response has the potential to improve common stress tolerance that may lead to a more efficient exploitation of the inherent genetic potential of agriculturally important plants.

HSPs are classified into different families and designated by molecular weight in kDa.

- HSP 100 k Da

- HSP 90
- HSP 70
- HSP 60
- 15 – 30 kDa low molecular mass HSPs or Small HSPs.

### **Functions**

- HSPs 60, 70 and 90 : act as molecular chaperons, involving ATP dependent stabilization and folding of proteins and assembly of oligomeric proteins.
- Some HSPs : assist in polypeptide transport across membranes into cellular compartments.
- Some HSPS : temporarily bind and stabilize an enzyme at a particular stage in cell development, later releasing the enzyme to become active.
- Binding of HSP with particular polypeptide within subcellular compartment avoid denaturation of many proteins at high temperatures.

### **Mitigation of Low Temperature Stress**

1. Seed hardening with 0.01% Ammonium molybdate and foliar spray of 0.1 % ammonium molybdate at critical stages of stress
2. Foliar spray of 2% calcium nitrate spray for membrane integrity
3. Foliar spray of 2% DAP + 1% KCl (MOP)
4. Foliar spray of 500 ppm cycocel for increasing root penetration in search of moisture for alleviation
5. Spray of 100 ppm salicylic acid
6. Brassinolide (0.5 ppm) for enhancing photosynthetic activity of plants
7. Seed treatment + soil application + foliar spray of Pink Pigmented Facultative Methnaotrops (PPFM) @  $10^6$  as a source of cytokinins.

### **Mitigation of High Temperature Stress**

1. Seed hardening with 0.5%  $\text{CaCl}_2$  solution for arresting membrane damage due to high temperature stress
2. Split application of N and K fertilizers

3. Foliar spray of 2% DAP + 1% KCl (MOP) (during the spray, sufficient moisture should be present in the soil for avoiding leaf scorching)
4. Foliar spray of 3% Kaoline
5. Foliar spray of 0.5% zinc sulphate + 0.3 % boric acid + 0.5 % Ferrous sulphate + 1% urea
6. Spray of 40 ppm NAA for controlling pre-mature flower / fruit drops due to high temperature stress
7. Foliar spray of 1% Urea + 2 % MgSO<sub>4</sub> + 0.5 % ZnSO<sub>4</sub> (for arresting chlorophyll degradation due to high temperature stress)
8. Foliar spray of 2% calcium nitrate spray for membrane integrity
9. Foliar spray of 0.5 ppm Brassinolide for increasing photosynthetic activity during stress
10. Spray of 100 ppm salicylic acid for increase stem reserve utilization and increasing Harvest Index of crops under stress
11. Seed treatment + soil application + foliar spray of Pink Pigmented Facultative Methnaotrops (PPFM) @ 10<sup>6</sup> as a source of cytokinins.

### **Low light and UV radiation stresses**

#### **Low Light Stress**

In some places (e.g. Thanjavur), the light intensity might be even up to 60000 lux in the first season but it would be low up to 30000 lux in the second season causing very poor productivity. Light quality is also very poor by showing about 400-440nm instead of the normal 600-640nm. The abnormal light intensity and quality causes reduced yield in any crops.

#### **UV-RADIATION STRESS**

UV radiation is divided into three categories

1. UV A – wavelength ranges from 320 to 400 nm and this is less lethal to the plants.
2. UV B – wavelength ranges from 280 to 320 nm and this is lethal to the plants.

3. UV C – wavelength is less than 280 nm and it is highly lethal to all biological systems.

The UV radiations cause environmental stress as the cell constituents like proteins and nucleic acids absorb UV radiation in the range of 250-400 nm (UV A and UV B) and cause death of the tissues. In general, on the outer atmosphere of the earth, CO<sub>2</sub>, ozone and water vapour form a layer and this layer prevent the entry of UV radiation. However, ozone depletion causes easy entry of UV radiation. In general, 1% reduction in ozone (O<sub>3</sub>) causes 2% increase in UV radiation.

### **UV radiation and plant response**

1. UV radiation slows down the growth of plants
2. Damage the process of photosynthesis
3. Prevent maturation and ripening process
4. Accelerate genetic mutation.

### **Mitigation of Low Light Stress:**

1. Foliar spray of 2% DAP + 1% KCl (MOP)
2. Spray of 2% coconut water
3. Spray of 40 ppm of NAA
4. 0.5 ppm Brassinolide spray
5. Spray of 100 ppm Salicylic acid
6. Spray of 500 ppm CC for arresting excessive vegetative growth
7. Split application of N and K fertilizers
8. Foliar spray of 0.3 % Boric acid + 0.5 % Zinc sulphate
9. Seed treatment + soil application + foliar spray of Pink Pigmented Facultative Methanotrophs (PPFM) @ 10<sup>6</sup> as a source of cytokinins.

## 23. SEED GERMINATION

The process of seed germination starts with the imbibition of water by seed coat and emergence of growing root tip of embryo. The process ends with the development of embryo into a seedling.

### **Physiological and biochemical changes during seed germination**

#### 1. Water uptake

Seed germination starts with the imbibition of water by dry seed coat. Due to imbibition of water, the seed coats become 1) More permeable to O<sub>2</sub> and water and 2) less resistant to outward growth of embryo.

#### 2. Respiration

Rapid increase in respiration rate of embryo occurs. Sucrose is probably the respiratory substrate at this stage which is provided by endosperm.

#### 3. Mobilization of reserve materials

As germination progresses, there is mobilization of reserve materials to provide.

1. building blocks for the development of embryo
2. energy for the biosynthetic process and
3. nucleic acids for control of protein synthesis and embryonic development

Changes in these components are as follows

##### i) Nucleic acids

In monocots, during imbibition, there is a rapid decrease of DNA and RNA content in the endosperm with a simultaneous increase in the embryonic axis probably due to their transportation as such. High concentration of RNA in the embryonic axis precedes cell division. Due to more cell division, DNA content is increased.

##### ii) Carbohydrates

Insoluble carbohydrates like starch are the important reserve food of cereals in the endosperm. During germination, starch is hydrolyzed first into maltose in the presence of α-

amylase and  $\beta$  - amylase and then maltose is converted into glucose by maltase. The glucose is further converted into soluble sucrose and transported to growing embryonic axis. During germination, the embryonic axis secretes gibberellic acid into the aleurone layer which causes synthesis of  $\alpha$ -amylase.

### 3. Lipids

Plants like castor bean, peanut etc., store large amount of neutral lipids or fats as reserve food in their seeds. During germination, the fats are hydrolyzed into fatty acids and glycerol by lipase enzyme. Fatty acids are further converted into acetyl – CoA by the process,  $\beta$  - oxidation. The acetyl CoA is further converted into sucrose via glyoxylate cycle and is transported to the growing embryonic axis.

### 4. Proteins

Some plants store proteins as reserve food in their seeds in the form of aleurone grains. Proteins are hydrolyzed into amino acids by peptidase enzyme. The amino acids may either provide energy by oxidation after deamination (removal of amino group) or may be utilized in the synthesis of new proteins.

### 5. Inorganic materials

A number of inorganic materials such as phosphate, calcium, magnesium and potassium are also stored in seeds in the form of phytin. These stored materials are liberated during germination due to the activity of various phosphatases including phytase.

#### **Emergence of seedling out of the seed coat**

All the changes described above gradually result in splitting of seed coat and emergence of the growing seedling. The radical comes out first and grows downward, and then plumule comes out and grows upward. Due to the continued growth of this seedling, the plumule comes out of the soil, exposed to light and develops its own photosynthetic apparatus.

Splitting of seed coat may take place either by imbibition pressure or by internal pressure created by the growing primary root or by hydrolytic enzymes which act on cell

wall contents of seed coat and digest it (e.g. cellulose and pectinase). Sometimes the seed coat may be extensively rotted by the activity of micro-organisms in the soil.

## DORMANCY OF SEEDS

All the viable seeds have capacity to germinate if placed under suitable conditions necessary for germination. But, some seeds fail to germinate sometimes even if placed under the condition favourable for germination. This may be due to some internal factors or due to specific requirement for some environmental factors. During this period, the growth of the seed remains suspended and they are said to be in rest stage or dormant stage and this phenomenon is called as dormancy of seeds.

### Factors causing dormancy of seeds

#### 1. Seed coats impermeable to water

The seeds of certain plants especially those belonging to the family's leguminaceae, solanaceae, malvaceae, etc. have very hard seed coats which are impermeable to water. The seeds remain dormant until the impermeable layer decay by the action of soil micro-organisms.

#### 2. Seeds coats impermeable to oxygen

In many plants such as cocklebur and many grasses, the seed dormancy is due to the impermeability of the seed coat to oxygen. However, during the period of dormancy, the seed coat gradually becomes more permeable to oxygen so that they may germinate.

#### 3. Immaturity of the Embryo

In certain orchids, the seed dormancy is due to the immaturity of the embryos which fail to develop fully by the time the seeds are shed. In such cases, the seeds germinate only after a period or rest during which the development of embryo inside the seed is completed.

#### 4. Germination Inhibitors

In certain seeds, the dormancy of the seeds is due to the presence of certain germination inhibitors like coumarin, ferulic acid, abscissic acid, etc. These may be present in endosperm, embryo, testa or juice or pulp of fruit.

#### 5. Chilling or low temperature requirement

In certain plants such as apple, rose, peach etc, the seeds remain dormant after harvest in the autumn as they have a low temperature or chilling requirement for germination. In nature, this requirement is fulfilled by the winter temperatures. In such case the seeds remain dormant throughout the winter season and germinate only in the following spring.

#### 6. Light sensitive seeds

In many species, the germination of the seeds is affected by light resulting in seed dormancy. Such light sensitive seeds are called *photo blastic*. Seeds of lettuce, tomato and tobacco are positively photo blastic and germinate only after they have been exposed to light. On the other hand, the seeds of certain plants are negatively photo blastic and their germination is inhibited by light.

#### Advantages of dormancy

1. In temperature zones, the dormancy of seeds helps the plants to tide over the severe colds which may be injurious for their vegetative and reproductive growth.
2. In tropical regions, the dormancy of seeds resulting from their impermeable seed coats ensures good chances of survival.
3. Dormancy of seeds in many cereals is of utmost importance to mankind. If these seeds germinate immediately after harvest in the field, they will become useless to man for consumption as food.

## 24. ABSCISSION AND SENESCENCE

Like human beings, plants also grow old and undergo aging and then they die. *Aging is the sum total of changes in the total plant or its organs.* During aging, the plants undergo chemical and structural changes. Aging leads to senescence and later phase of development that ultimately terminates to death.

### **Senescence**

*The deteriorative process which naturally terminates the functional life of an organ, organism or other life unit is collectively called senescence.* Senescence is a phase of the aging process. The major characteristic of senescence is that the metabolic processes are catabolic and eventually become irreversible and terminate to death.

Senescence is not confined only to whole plant. It may be limited to a particular plant organ such as leaf and flowers or cells or cell, organelles. Senescence is closely associated with the phenomenon of aging. Aging leads to senescence. Wheat plant dies after the development of fruit. This is the senescence of an entire plant. *Leaffall* in a coconut tree is an example of senescence.

### **Types of senescence**

*Leopold (1961)* has proposed types of senescence patterns in plants which are as follows.

#### **(a) Overall Senescence**

This type of senescence occurs in annuals where whole plant is affected. It is also called *whole plant senescence*. The entire plant dies after the development of fruit and seeds. E.g. Paddy, wheat, soybean etc.

#### **(b) Top Senescence**

In top senescence, the parts remaining above the ground or (shoot system) may die, but the root system and underground system remain viable. It is also called *shoot senescence*. E.g. Dock, perennial herbs.

#### **(c) Deciduous Senescence**

In deciduous woody plants, all the leaves die but the bulk of the stem and root system

remains viable. It is called *deciduous senescence* or *simultaneous* or *synchronous senescence*.  
E.g. Leaf fall in deciduous trees.

#### (d) Progressive Senescence

It is a gradual death of old leaves from the base to the top of the plants. It may occur at any time. It is also called *sequential senescence*. E.g. Leaf fall in a coconut tree.

### **Causes of Senescence**

1. Leaf senescence is accompanied by early loss in *chlorophyll*, RNA and enzymes.
2. Cellular constituents are decreased due to slower synthesis or faster break down.
3. Competition between vegetative and reproductive organs for nutrients.
4. A senescence factor (a hormone) is produced in soybean fruits that move to leaves where it causes senescence.
5. Short-day and long-night conditions induce flowering and leaf senescence.
6. Degradation of food reserves and loss of integrity in food storage cells of seeds.
7. Senescence is also hormonally controlled.

### **Physiology of Senescence**

The following physiological changes occur during senescence.

1. *Photosynthesis* stops.
2. *Chlorophyll* degradation: The colour of leaf changes from green to yellow.
3. *Anthocyanin* pigments accumulation in the leaves causing reddening in leaves.
4. The vacuoles function as *lysosomes* and digest the cellular materials.
5. The starch content decreased.
6. RNA and proteins are decreased.
7. DNA molecules are degraded by the enzyme DNase.
8. Growth promoting hormones such as cytokinin decrease.
9. The deteriorative hormones such as *ethylene* and *abscisic acid* (ABA) content are increased.

### **Senescence Promoters**

Senescence is promoted by hormones such as abscisic acid and ethylene. The

senescence accelerating ability of abscisic acid is well documented. The function of *ABA as a promoter of flower tissue senescence including initiation of colour fading or bleaching has been established.* The ABA content of aging leaves increases markedly as senescence is initiated. *Ethylene* plays a very important role in the senescence of certain plant parts, particularly fruit and petals and in the abscission process. It is an inducer in the senescence of flower tissue. Senescence Retardants: The primary plant hormones involved here are auxin, gibberellin and cytokinin.

### **Significance of Senescence**

1. The whole plant senescence occurs in monocarpic plants coinciding the seed setting and seed dispersal.
2. Due to the formation of abscission layer, the older leaves tend to fall down so that the nutrients will be diverted to the next young leaf.
3. The senescence process helps the mobilization of nutrients and of the vegetative parts of the plant into the fruits.
4. Plants escape the influence of seasonal adversity by undergoing senescence of its organs. Leaf fall in deciduous trees reduces the rate of transpiration to survive under adverse conditions.

### **Abscission**

*Shedding of leaves, flowers and fruits is called abscission.* Abscission is distinct in deciduous trees and shrubs. In autumn, all the leaves of deciduous plants fall, at about the same time giving the plants a naked appearance. In evergreen plants there is gradual abscission of leaves. The older leaves fall while new leaves are developed continuously throughout the year. In most of the herbaceous species, however the leaves are not shed even after they die. In many cases leaves are retained in withered dry condition even after the whole shoot is dead.

Abscission is a complex physiological process. During abscission, the colour of the leaves, flowers and fruits changes due to degradation of chlorophyll and the synthesis of *anthocyanin* pigment.

Leaf abscission takes place at the base of the petiole. The site of abscission is internally marked by a distinct zone called *abscission zone*. This zone is made up of one or more layers of cells arranged transversely across the petiole base. This is called *abscission layer*. The

abscission zone is pale or brown in colour. The cells of the abscission layer separate from each other due to the dissolution of middle lamellae and the primary cellulose walls under the influence of the activity of enzymes, *pectinase* and *cellulase*.

At this stage, the petiole remains attached to the stem by vascular elements only. But due to its own weight and the wind force, the leaf is detached from the stem. The broken vascular elements are soon plugged with *tyloses* or gums. Wound healing in cells proximal to the breaking point involves formation of a corky layer that protects the plant from pathogen invasion and excess water loss. *Suberin* and *lignin* are synthesized during healing.

Several environmental factors such as drought and N deficiency promote abscission. *Auxin* is synthesized in growing leaf blades and it strongly retards senescence and abscission. Abscission starts when the amount of auxin begins to decrease. Cytokinins and gibberellins arriving from the roots also delay senescence and abscission. Abscission is caused by the formation of cell wall degrading enzymes in the abscission zone, due to ethylene production.

### **Significance of Abscission**

1. It helps in diverting water and nutrients to the young leaves
2. It is a self pruning process through which fruits and injured organs are shed from the parent plant.
3. It helps in disseminating fruits and vegetative propagates.
4. Abscission serves as function in removing plant parts containing waste materials.

## 25. GLOBAL WARMING - PHYSIOLOGICAL EFFECTS ON CROP PRODUCTIVITY

### **GLOBAL WARMING (Green house Effect)**

In general, delicate plants which require protection from weather are grown in green house (glass house). In green house so many gases are produced like CO<sub>2</sub>, water vapour, methane, oxides of nitrogen and chloro fluoro carbon (CFC). These gases are produced from plants and accumulated inside the glass house; as a result glass house gets warming. In natural atmosphere also the same effect occurs i.e. global warming (due to the release of gases from plants).

But in glass house, glass roof is present to prevent the escape of gases from the glass house. In natural atmosphere, the gases such as ozone, water vapour, CO<sub>2</sub> methane etc. form a layer on the lower atmosphere and this layer prevents the heat escaping from the earth. If heat is released or escaped from earth, the temperature of earth would be below freezing point. The accumulation of heat or gases causes the warming of earth surface and leads to global warming.

#### **Global warming leads to the following effects:**

1. Rise in temperature
2. Average rise in the level of sea (about 6 cm/decade) due to melting of polar ice.
3. Steady increase (enrichment) in the CO<sub>2</sub>.

Atmosphere – (5.1 x 10<sup>18</sup>kg)

Lithosphere - (1.5 x 10<sup>22</sup> kg)                  are in a dynamic equilibrium

Hydrosphere – 1.4 x 10<sup>22</sup> kg)

Biosphere – 1.2 x 10<sup>15</sup> kg (dry wt.)

### **CO<sub>2</sub> fixation by green plants**

Total area of green plants 510 x 10<sup>6</sup> km<sup>2</sup>

CO<sub>2</sub> fixed – 1.39 x 10<sup>14</sup> kg year<sup>-1</sup> (0.27 kg m<sup>-2</sup> year<sup>-1</sup>)

### **Light utilization:**

A small portion of radiant energy (400 – 700 nm) is reaching the earth's surface

0.1 – 1.0 % utilized under natural vegetation or ordinary agriculture

- 2.0 – 2.5 % under intensive agriculture
- 6.0 – 10.0 % in some crop plants
- 20.0 – 25.0 % under laboratory condition

**Biosphere:**

O <sub>2</sub> in atmosphere	- $1.1 \times 10^{18}$ kg
O <sub>2</sub> released by photosynthesis	- $5.1 \times 10^{14}$ kg year <sup>-1</sup>
CO <sub>2</sub> content	- $5 \times 10^{16}$ kg (mostly dissolved in sea)
CO <sub>2</sub> consumed by photosynthesis	- $7 \times 10^{14}$ kg year <sup>-1</sup>

**Human activities disturbed the global ecosystem (MST- mesospheric,**

Stratospheric, Troposphere)

Landscape modification

Resource exploitation

Effluent flow

High temperature

Rainfall redistribution

Increased UV-B radiation due to stratospheric O<sub>3</sub> depletion

Increased level of atmospheric CO<sub>2</sub>

Other green house gases

Transparent to incoming short wave radiation

Absorb short wave and emit long wave radiation

Net emission of CO<sub>2</sub>

Bacterial fermentation in the anaerobic rice fields generate 120 million ton CH<sub>4</sub> every year

Ruminant gut bacteria produce 78 m tons of CH<sub>4</sub> every year and are released by Flatulence

**Green house gases**

CO<sub>2</sub>, CH<sub>4</sub>, NO, NO<sub>2</sub>, N<sub>2</sub>O, (N<sub>x</sub>O<sub>x</sub>), CFCl<sub>3</sub>, CF<sub>2</sub>Cl<sub>2</sub>, CFMs, O<sub>3</sub>, H<sub>2</sub>O

Fossil fuel reserves are large enough for climatic changes to occur, if these

reserves continue to be exploited at a higher rate in future

### **CO<sub>2</sub> enrichment and crop productivity**

1. CO<sub>2</sub> enrichment leads to increased photosynthesis and productivity
2. CO<sub>2</sub> enrichment also decreases stomatal conductance by closing the stomata, thereby decreasing the transpiration / unit area of the leaf.
3. In C<sub>3</sub> plant the efficiency of RuBP carboxylase enzyme is increased
4. Increased CO<sub>2</sub> concentration inhibits photorespiration in plants
5. CO<sub>2</sub> enrichment increases the yield and yield components.

### **Other green house gases**

1. Oxides of nitrogen (NO, NO<sub>2</sub>, N<sub>2</sub>O molecular N<sub>2</sub>) cause phototoxic, bleaching and necrosis (drying of tissues) in plants.
2. Ozone (O<sub>3</sub>) causes ozone injury to the plants.

### **Remedial measures for green house effect**

- Reduction in the use of fossil fuel
- Use of alternative sources of energy (renewable energy)
- Afforestation and community forestry
- Avoiding the use of CFCs and nuclear explosions
- Environmental awareness

### **Direct effect of CO<sub>2</sub> increase in the absence of climatic change**

- Doubling of CO<sub>2</sub> from 340 to 680 ppm increases 0 – 10 % increase in yield of C<sub>4</sub> plants (maize, sorghum, sugarcane) and 10 – 50 % increase in yield in C<sub>3</sub> plants (wheat, soybean, rice) depending upon specific crop and growing conditions
- Greater yield benefit accrue to the regions where the C<sub>3</sub> rather than C<sub>4</sub> crops dominate
- Higher CO<sub>2</sub> conc. reduces stomatal aperture, thereby reducing transpiration and WUE
- Doubling of CO<sub>2</sub> will cause about 40 % decrease in stomatal conductance in short term

### **Law of limiting factor:**

- When other environmental factors such as water, light, minerals & temperature limit yield, then higher conc. of CO<sub>2</sub> will have little or no effect.
- This generalizing concept has been challenged
- In certain stressful environments, the relative photosynthesis increased with increased in CO<sub>2</sub> conc.

C<sub>3</sub> – 95 % of world's biomass is of the C<sub>3</sub> category

- In C<sub>3</sub>, O<sub>2</sub> compete with CO<sub>2</sub> for the site of Rubisco – In C<sub>4</sub>, O<sub>2</sub> is not compete with CO<sub>2</sub> for the site of PEPCase
- At 340 ppm CO<sub>2</sub>, in the absence of O<sub>2</sub> Rubisco operating only at  $\frac{1}{2}$  to  $\frac{3}{4}$  of its substrate – saturated capacity
- PEPCase has high affinity for CO<sub>2</sub> than Rubisco – PEPCase is close to CO<sub>2</sub> saturation at the present atmospheric CO<sub>2</sub> conc., no significant enhance of C<sub>4</sub> crop growth from increased CO<sub>2</sub> so far as PEP Case is concerned

## Carbon Sequestration

Climate change is one of the most important global environmental challenges, with its implications on food production, water supply, healthy energy etc... are detrimental. Historically the responsibility of green house gas emission increase lies largely with the industrialized world and the developing countries are contributing very less amount only. (Jayant & Santhaye, 2006)

The increase in atmospheric concentration of CO<sub>2</sub> by 31% since 1950 from fossil fuel consumption and land use change indicates the threats of global warming since industrial revolution increases the global emission of carbon, estimated at  $270 \pm 30$  billion ton emission due to land use change include those by deforestation, due to agriculture and land use changes contributes  $78 \pm 12$  billion ton of carbon to the atmosphere. Well planned management practices enhances bio mass production, purifies surface and ground waters and reduces the rate of enrichment of atmospheric CO<sub>2</sub> due to fossil fuel (Lal, 2004)


The Kyoto protocol is a positive first step to prevent climatic change at global level and the responsibility going to each and every human being (Ravindranath, 2006).

The annual soil organic carbon (SOC) enrichment of atmospheric CO<sub>2</sub> has to be reduced. The annual SOC sequestration potential is only  $0.9 \pm 0.3$  Pg C/year. The CO<sub>2</sub> level

in atmosphere increase at the rate of 2.0-2.6 Pg C / year. So soil carbon sequestration methodologies plays significant role in reducing atmospheric CO<sub>2</sub> control (Lal, 2004)

Besides the artificial CO<sub>2</sub> capture and storage methods also play a significant role to cut down the increasing atmospheric CO<sub>2</sub> concentration. (IPCC, 2001)

Even though all the industrial and other carbon emissions cannot be controlled by soil carbon sequestration, it may be a viable method until a permanent solution found out.



**A carbon dioxide (CO<sub>2</sub>) sink** is a carbon dioxide reservoir that is increasing in size, and is the opposite of a carbon dioxide "source". The main natural sinks are (1) the oceans and (2) plants and other organisms that use photosynthesis to remove carbon from the atmosphere by incorporating it into biomass and release oxygen into the atmosphere. The process by which carbon dioxide sinks (natural and artificial) remove CO<sub>2</sub> from the atmosphere is known as carbon sequestration.

### Storage in vegetation and soils

Carbon stored in soils oxidizes rapidly; this, in addition to high rainfall levels, is the reason why tropical jungles have very thin organic soils. The forest eco-system may eventually become carbon neutral. Forest fires release absorbed carbon back into the atmosphere, as does deforestation due to rapidly increased oxidation of soil organic matter.

The dead trees, plants, and moss in peat bogs undergo slow anaerobic decomposition below the surface of the bog. This process is slow enough that in many cases the bog grows rapidly and fixes more carbon from the atmosphere than is released. Over time, the peat grows deeper. Peat bogs inter approximately one-quarter of the carbon stored in land plants and soils.

Under some conditions, forests and peat bogs may become sources of CO<sub>2</sub>, such as when a forest is flooded by the construction of a hydroelectric dam. Unless the forests and peat are harvested before flooding, the rotting vegetation is a source of CO<sub>2</sub> and methane comparable in magnitude to the amount of carbon released by a fossil-fuel powered plant of equivalent power.

## Oceans

Oceans are natural CO<sub>2</sub> sinks, and represent the largest active carbon sink on Earth. This role as a sink for CO<sub>2</sub> is driven by two processes, the solubility pump and the biological pump.<sup>[6]</sup> The former is primarily a function of differential CO<sub>2</sub> solubility in seawater and the thermohaline circulation, while the latter is the sum of a series of biological processes that transport carbon (in organic and inorganic forms) from the surface euphotic zone to the ocean's interior. A small fraction of the organic carbon transported by the biological pump to the seafloor is buried in anoxic conditions under sediments and ultimately forms fossil fuels such as oil and natural gas.

One way to increase the carbon sequestration efficiency of the oceans is to add micrometre-sized iron particles in the form of either hematite (iron oxide) or melanterite (iron sulfate) to certain regions of the ocean. This has the effect of stimulating growth of plankton. Iron is an important nutrient for phytoplankton, usually made available via upwelling along the continental shelves, inflows from rivers and streams, as well as deposition of dust suspended in the atmosphere. Natural sources of ocean iron have been

declining in recent decades, contributing to an overall decline in ocean productivity (NASA, 2003). Yet in the presence of iron nutrients plankton populations quickly grow, or 'bloom', expanding the base of biomass productivity throughout the region and removing significant quantities of CO<sub>2</sub> from the atmosphere via photosynthesis. A test in 2002 in the Southern Ocean around Antarctica suggests that between 10,000 and 100,000 carbon atoms are sunk for each iron atom added to the water. More recent work in Germany (2005) suggests that any biomass carbon in the oceans, whether exported to depth or recycled in the euphotic zone, represents long-term storage of carbon. This means that application of iron nutrients in select parts of the oceans, at appropriate scales, could have the combined effect of restoring ocean productivity while at the same time mitigating the effects of human caused emissions of carbon dioxide to the atmosphere.

Because the effect of periodic small scale phytoplankton blooms on ocean ecosystems is unclear, more studies would be helpful. Phytoplankton have a complex effect on cloud formation via the release of substances such as dimethyl sulfide (DMS) that are converted to sulfate aerosols in the atmosphere, providing cloud condensation nuclei, or CCN. But the effect of small scale plankton blooms on overall DMS production is unknown.

Other nutrients such as nitrates, phosphates, and silica as well as iron may cause ocean fertilization. There has been some speculation that using pulses of fertilization (around 20 days in length) may be more effective at getting carbon to ocean floor than sustained fertilization.

There is some controversy over seeding the oceans with iron however, due to the potential for increased toxic phytoplankton growth (e.g. "red tide"), declining water quality due to overgrowth, and increasing anoxia in areas harming other sea-life such as zooplankton, fish, coral, etc.

### Soils

Methods that significantly enhance carbon sequestration in soil include no-till farming, residue mulching, cover cropping, and crop rotation, all of which are more widely used in organic farming than in conventional farming. Because only 5% of US farmland currently uses no-till and residue mulching, there is a large potential for carbon sequestration.<sup>[26]</sup>

Conversion to pastureland, particularly with good management of grazing, can sequester even more carbon in the soil.

Terra preta, an anthropogenic, high-carbon soil, is also being investigated as a sequestration mechanism. By pyrolysing biomass, about half of its carbon can be reduced to charcoal, which can persist in the soil for centuries, and makes a useful soil amendment, especially in tropical soils (*biochar* or *agrichar*).

Savanna

### **Artificial sequestration**

For carbon to be sequestered artificially (i.e. not using the natural processes of the carbon cycle) it must first be captured, *or* it must be significantly delayed or prevented from being re-released into the atmosphere (by combustion, decay, etc.) from an existing carbon-rich material, by being incorporated into an enduring usage (such as in construction). Thereafter it can be passively stored *or* remain productively utilized over time in a variety of ways.

For example, upon harvesting, wood (as a carbon-rich material) can be immediately burned or otherwise serve as a fuel, returning its carbon to the atmosphere, *or* it can be incorporated into construction or a range of other durable products, thus sequestering its carbon over years or even centuries. One ton of dry wood is equivalent to 1.8 tons of Carbon dioxide.

### **Geological sequestration**

The method of *geo-sequestration* or *geological storage* involves injecting carbon dioxide directly into underground geological formations. Declining oil fields, saline aquifers, and unminable coal seams have been suggested as storage sites. Caverns and old mines that are commonly used to store natural gas are not considered, because of a lack of storage safety.

### **Mineral sequestration**

Mineral sequestration aims to trap carbon in the form of solid carbonate salts. This process occurs slowly in nature and is responsible for the deposition and accumulation of

limestone (calcium carbonate) over geologic time. Carbonic acid in groundwater slowly reacts with complex silicates to dissolve calcium, magnesium, alkalis and silica and leave a residue of clay minerals. The dissolved calcium and magnesium react with bicarbonate to precipitate calcium and magnesium carbonates, a process that organisms use to make shells. When the organisms die, their shells are deposited as sediment and eventually turn into limestone. Limestones have accumulated over billions of years of geologic time and contain much of Earth's carbon. Ongoing research aims to speed up similar reactions involving alkali carbonates.

## REFERENCES

1. Jain, J.K. 2007. Fundamentals of plant physiology, S.Chand & Company Ltd., New Delhi.
2. Pandey, S. N. and B. K.Sinha, 2006. Plant Physiology. Vikas Publishing House
3. Private Limited, New Delhi.
2. Purohit, S.S, 2005. Plant physiology, Student edition, Jodhpur.
3. Ray Noggle, G. and Fritz, G. J., 1991. Introductory Plant Physiology. Prentice Hall of India Pvt. Ltd., New Delhi.
5. Taiz. L. and Zeiger. E., 2006. Plant Physiology. Publishers: Sinauer Associates, Inc., Massachusetts, USA.

## ONLINE REFERENCE

1. <http://www.plantphys.org>
2. <http://www.Biologie.Uni-hamburg.de/b-online>
3. <http://4e.plantphys.net>
4. <http://3e.plantphys.net>
5. <http://www.botany.org>



This Book Download From e-course of ICAR  
**Visit for Other Agriculture books, News, Recruitment, Information, and Events at [WWW.AGRIMOON.COM](http://WWW.AGRIMOON.COM)**

Give FeedBack & Suggestion at [info@agrimoon.com](mailto:info@agrimoon.com)

**DISCLAIMER:**

The information on this website does not warrant or assume any legal liability or responsibility for the accuracy, completeness or usefulness of the courseware contents.

The contents are provided free for noncommercial purpose such as teaching, training, research, extension and self learning.

