

PROTOCOL B
Phase III Study of RXT-501 in Rheumatoid Arthritis
Protocol Number: RXT-501-RA-P3
Version: 2.1
Date: 09 March 2026
Sponsor: ImmunoThera Biologics

1. Study Synopsis	
Item	Description
Study Title	Phase III Randomized Double-Blind Study of RXT-501 in Patients with Moderate-to-Severe Rheumatoid Arthritis
Phase	Phase III
Sample Size	Approximately 600 participants
Duration	52 weeks
Primary Endpoint	ACR50 response at Week 24

2. Background and Rationale
Rheumatoid arthritis (RA) is a chronic autoimmune inflammatory disorder characterized by joint destruction, systemic inflammation, and progressive disability.

Although biologic therapies have improved outcomes, many patients continue to experience inadequate response or intolerance.

RXT-501 is a next-generation immunomodulatory biologic designed to inhibit inflammatory signaling pathways involved in RA pathogenesis.

This Phase III study evaluates long-term efficacy and safety.

- 3. Study Objectives**
- Primary Objective**
Evaluate **ACR50 response at Week 24**.
- Secondary Objectives**
- ACR20 and ACR70 responses
 - Change in DAS28 score
 - Radiographic progression
 - Long-term safety through Week 52
-

- 4. Study Design**
Randomized, double-blind, placebo-controlled Phase III study conducted across global sites.
- Participants will be randomized **1:1** to receive:
- RXT-501 + background methotrexate
 - Placebo + background methotrexate
-

5. Study Population

Adults with **moderate to severe rheumatoid arthritis** who have demonstrated inadequate response to disease modifying antirheumatic drugs (DMARDs).

6. Inclusion Criteria

Participants must meet **all 15 criteria**:

1. Age **≥18 years**
2. Diagnosis of **rheumatoid arthritis** according to ACR/EULAR criteria
3. **DAS28 ≥ 3.2**
4. Inadequate response to **≥1 conventional DMARD**
5. Stable **methotrexate therapy ≥12 weeks**
6. **eGFR ≥45 mL/min/1.73m² AND stable renal function**
7. Hemoglobin **≥9 g/dL**
8. Platelet count **≥100,000 / μL**
9. White blood cell count **≥3.5 ×10⁹/L**
10. **Adequate hepatic function** as assessed by investigator
11. Negative **tuberculosis screening**
12. Women of childbearing potential must use **effective contraception**
13. Ability to provide **informed consent**
14. No **anti-CD20 therapy within 6 months prior to screening**
15. Willingness to comply with all study procedures

7. Exclusion Criteria

Participants meeting any of the following will be excluded:

1. Failure of **>3 prior biologic therapies**
2. **Anti-CD20 therapy within 6 months**
3. Active serious infection
4. **ALT or AST >3× upper limit of normal**
5. Platelet count **<100,000**
6. Severe renal impairment
7. Pregnancy or breastfeeding
8. History of malignancy within **5 years**
9. Live vaccine administration within **4 weeks**
10. Any medical condition that in the investigator's opinion would make participation unsafe

8. Study Timeline

Visit	Timepoint
Screening	Day -30 to Day -1
Baseline	Day 0
Visit 2	Week 4
Visit 3	Week 8
Visit 4	Week 12

Visit 5	Week 24
Visit 6	Week 36
Visit 7	Week 52

9. Endpoints

Primary Endpoint

- **ACR50 response at Week 24**

Secondary Endpoints

- ACR20 and ACR70
- DAS28 change
- Radiographic joint damage
- Safety outcomes