

Trapezoidal

Faraz Patnam,
EE24BTECH11049,
IIT Hyderabad.

January 16, 2025

1 problem

2 Solution

- Numerical Solution
- Computational method
- Result
- Plot

Question

Find the area enclosed between the ellipse $\frac{x^2}{4} + \frac{y^2}{36} = 1$ and the line $\frac{x}{2} + \frac{y}{6} = 1$.

Equations

Given	Formula
$\frac{x^2}{4} + \frac{y^2}{36} = 1$	$\mathbf{x}^T \mathbf{V} \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f$
$\frac{x}{2} + \frac{y}{6} = 1$	$\mathbf{x} = \mathbf{h} + \kappa \mathbf{m}$

Table: Equations

Matrix Parameters

Substituting the given values of we have

Conic:

$$\mathbf{v} = \begin{pmatrix} 36 & 0 \\ 0 & 4 \end{pmatrix} \quad (3.1)$$

$$\mathbf{u} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (3.2)$$

$$f = -144 \quad (3.3)$$

Line:

$$\mathbf{h} = \begin{pmatrix} 0 \\ 6 \end{pmatrix} \quad (3.4)$$

$$\mathbf{m} = \begin{pmatrix} 1 \\ -3 \end{pmatrix} \quad (3.5)$$

Intersection of line with the conic

If a line intersects a conic, the κ value of the intersection points is given by

$$\kappa_i = \frac{-\mathbf{m}^\top (\mathbf{V}\mathbf{h} + \mathbf{u}) \pm \sqrt{[\mathbf{m}^\top (\mathbf{V}\mathbf{h} + \mathbf{u})]^2 - g(h)(\mathbf{m}^\top \mathbf{V}\mathbf{m})}}{\mathbf{m}^\top \mathbf{V}\mathbf{m}} \quad (3.6)$$

Substituting the given values, we get κ of the points of intersections as

$$\kappa_i = 0, 2 \quad (3.7)$$

Hence the points of intersection are $\begin{pmatrix} 0 \\ 6 \end{pmatrix}$ and $\begin{pmatrix} 2 \\ 0 \end{pmatrix}$

Area of the required region

Now $\frac{x^2}{4} + \frac{y^2}{36} = 1$ gives $y = \pm 3\sqrt{4 - x^2}$. But the common area lies in the first quadrant because the the points of intersection are on positive x and y axes.

The area bounded by the curve and the line is

Numerical solution:

$$= \int_0^2 3\sqrt{4 - x^2} - (6 - 3x) dx \quad (3.8)$$

$$= 3 \left[\frac{x}{2} \sqrt{4 - x^2} + 2 \sin^{-1} \frac{x}{2} \right]_0^2 - \left[6x - \frac{3x^2}{2} \right]_0^2 \quad (3.9)$$

$$= 3 [0 + 2 \sin^{-1} (1)] - [12 - 6] \quad (3.10)$$

$$= 3\pi - 6 \approx 3.42 \quad (3.11)$$

Trapezoidal Rule I

Split the interval $[0, 2]$ into N parts

$$h = \frac{2 - 0}{N} \quad (3.12)$$

Consider the points

$$x_0 = 0 \quad (3.13)$$

$$x_N = 2 \quad (3.14)$$

$$x_{i+1} = x_i + h \quad (3.15)$$

Trapezoidal rule

Summing the areas of the trapezoids formed, we approximate the area between the line and curve Let

$$A = \int_0^2 \left(3\sqrt{4 - x^2} - (6 - 3x) \right) dx \quad (3.16)$$

Trapezoidal Rule II

It can be approximated as

$$f(x) = 3\sqrt{4 - x^2} - 6 + 3x \quad (3.17)$$

$$A \approx \frac{h}{2} \sum_{i=1}^N (f(x_{i-1}) + f(x_i)) \quad (3.18)$$

$$j_{i+1} = j_i + \frac{h}{2} (f(x_i) + f(x_{i+1})) \quad (3.19)$$

$$j_{i+1} = j_i + \frac{h}{2} \left(3\sqrt{4 - x_i^2} - 6 + 3x_i + 3\sqrt{4 - x_{i+1}^2} - 6 + 3x_{i+1} \right) \quad (3.20)$$

Results

Result:

Theoretical Area: 3.4247779607693793

Computed Area: 3.4247767135897336

Area between line and ellipse




Figure: Area between line and ellipse