

Assignment 1

EE24BTECH11049

- 1) Consider three observations a , b and c such that $b = a + c$. If the standard deviation of $a + 2$, $b + 2$, $c + 2$ is d , then which of the following is true?


(2021-Mar)

- a) $b^2 = a^2 + c^2 + 3d^2$ c) $b^2 = 3(a^2 + c^2) + 9d^2$
 b) $b^2 = 3(a^2 + c^2) - 9d^2$ d) $b^2 = 3(a^2 + c^2 + d^2)$

- 2) Let a vector $\alpha\hat{\mathbf{i}} + \beta\hat{\mathbf{j}}$ be obtained by rotating the vector $\sqrt{3}\hat{\mathbf{i}} + \hat{\mathbf{j}}$ by an angle 45° about the origin in counterclockwise direction in the first quadrant. Then the area of triangle having vertices (α, β) , $(0, \beta)$ and $(0, 0)$ is equal to:

(2021-Mar)

- a) 1 b) $\frac{1}{2}$ c) $\frac{1}{\sqrt{2}}$ d) $2\sqrt{2}$



- 3) If for $a > 0$, the feet of perpendiculars from the points $\mathbf{A}(a, -2a, 3)$ and $\mathbf{B}(0, 4, 5)$ on the plane $lx + my + nz = 0$ are points $\mathbf{C}(0, -a, -1)$ and \mathbf{D} respectively, then the length of line segment \mathbf{CD} is equal to:

(2021-Mar)

a) $\frac{3}{4}$

b) $\frac{52}{867}$

c) $\frac{39}{50}$

d) $\frac{22}{425}$

- 8) If n is the number of irrational terms in the expansion of $\left[3^{\frac{1}{4}} + 5^{\frac{1}{8}}\right]^{60}$, then $(n - 1)$ is divisible by

(2021-Mar)

a) 8

b) 26

c) 7

d) 30

- 9) Let the position vectors of two points \mathbf{P} and \mathbf{Q} be $3\hat{\mathbf{i}} - \hat{\mathbf{j}} + 2\hat{\mathbf{k}}$ and $\hat{\mathbf{i}} + 2\hat{\mathbf{j}} - 4\hat{\mathbf{k}}$ respectively. Let \mathbf{R} and \mathbf{S} be two points such that the direction ratios of lines \mathbf{PR} and \mathbf{QS} are $(4, -1, 2)$ and $(-2, 1, -2)$ respectively. Let lines \mathbf{PR} and \mathbf{QS} intersect at \mathbf{T} . If the vector \mathbf{TA} is perpendicular to both \mathbf{PR} and \mathbf{QS} and the length of vector \mathbf{TA} is $\sqrt{5}$ units, then the modulus of a position vector of \mathbf{A} is:


(2021-Mar)

a) $\sqrt{5}$

b) $\sqrt{171}$

c) $\sqrt{227}$

d) $\sqrt{482}$



- 10) If the three normals drawn to the parabola, $y^2 = 2x$ pass through the point $(a, 0)$ $a \neq 0$, then ' a ' must be greater than

(2021-Mar)

a) 1

b) $\frac{1}{2}$

c) $-\frac{1}{2}$

d) -1

- 11) let

$$S_K = \sum_{r=1}^k \tan^{-1} \left[\frac{(6^r)}{(2^{r+1} + 3^{2r+1})} \right]. \text{ Then } \lim_{k \rightarrow \infty} S_k =$$

(2021-Mar)

- a) $\tan^{-1}\left(\frac{3}{2}\right)$ b) $\cot^{-1}\left(\frac{3}{2}\right)$ c) $\frac{\pi}{2}$ d) $\tan^{-1}(3)$

12) The number of roots of the equation, $(81)^{\sin^2 x} + (81)^{\cos^2 x} = 30$ in the interval $[0, \pi]$ is equal to :

(2021-Mar)

- a) 3 b) 2 c) 4 d) 8

13) If $y = y(x)$ is the solution of the differential equation,

$$\frac{dy}{dx} + 2y \tan x = \sin x,$$

$y\left(\frac{\pi}{3}\right) = 0$, then the maximum value of the function $y(x)$ over \mathbf{R} is equal to :

(2021-Mar)

- a) 8 b) $\frac{1}{2}$ c) $-\frac{15}{4}$ d) $\frac{1}{8}$

14) Which of the following Boolean expression is a tautology?

(2021-Mar)

- a) $(p \wedge q) \wedge (p \rightarrow q)$ c) $(p \wedge q) \vee (p \rightarrow q)$
 b) $(p \wedge q) \vee (p \vee q)$ d) $(p \wedge q) \rightarrow (p \rightarrow q)$

15) let $A = \begin{pmatrix} \iota & -\iota \\ -\iota & \iota \end{pmatrix}$. Then, the system of linear equations $A^8 \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 8 \\ 64 \end{pmatrix}$ has

(2021-Mar)

- a) No solution c) A unique solution
 b) Exactly two solutions d) Infinitely many solutions