

Assignment 3

2024-April

Session-04-05-2024-shift:1-16-30

EE24BTECH11049
Patnam Shariq Faraz Muhammed

MCQ

1) The integral

$$\int_0^{\frac{\pi}{4}} \frac{136 \sin x}{3 \sin x + 5 \cos x} dx$$

is equal to:

(2024-Apr)

- a) $3\pi - 10 \log_e (2\sqrt{2}) + 10 \log_e 5$ c) $3\pi - 30 \log_e 2 + 20 \log_e 5$
b) $3\pi - 50 \log_e 2 + 20 \log_e 5$ d) $3\pi - 25 \log_e 2 + 10 \log_e 5$

2) If $y = y(x)$ is the solution of the differential equation

$$\frac{dy}{dx} + 2y = \sin(2x), y(0) = \frac{3}{4}, \text{ then } y\left(\frac{\pi}{8}\right)$$

is equal to:

(2024-Apr)

- a) $e^{\frac{\pi}{8}}$ b) $e^{-\frac{\pi}{8}}$ c) $e^{\frac{\pi}{4}}$ d) $e^{-\frac{\pi}{4}}$

3) Let two straight line drawn from the origin **O** intersect the line $3x + 4y = 12$ at the points **P** and **Q** such that $\triangle OPQ$ is an isosceles triangle and $\angle POQ = 90^\circ$. If $l = \mathbf{OP}^2 + \mathbf{PQ}^2 + \mathbf{QO}^2$, then the greatest integer less than or equal to l is:

(2024-Apr)

- a) 44 b) 48 c) 42 d) 46

4) If the function

$$f(x) = \frac{\sin 3x + \alpha \sin x - \beta \cos 3x}{x^3}, x \in \mathbf{R}$$

is continuous at $x = 0$, then $f(0)$ is equal to:

(2024-Apr)

a) -4 b) 4 c) 2 d) -2

onsider the following statements:

Statement I :For any two complex numbers z_1, z_2 ,

$$(|z_1| + |z_2|) \left| \frac{z_1}{|z_1|} + \frac{z_2}{|z_2|} \right| \leq 2(|z_1| + |z_2|), \text{ and}$$

Statement II :If x, y, z and three distinct complex numbers and

a, b, c are three positive real numbers such that

$$\frac{a}{|y-z|} = \frac{b}{|z-x|} = \frac{c}{|x-y|}, \text{ then}$$

$$\frac{a^2}{y-z} + \frac{b^2}{z-x} + \frac{c^2}{x-y} = 1.$$

Between the above two statements:

(2024-Apr)

- a) statement I is correct but statement II is incorrect.
- b) both statement I and statement II are correct.
- c) statement I is incorrect but statement II is correct.
- d) both statement I and statement II are incorrect.

INTEGER

- 1) Let a_1, a_2, a_3, \dots be in arithmetic progression progression of positive terms. Let

$$A_k = a_1^2 - a_2^2 + a_3^2 - a_4^2 + \dots + a_{2k-1}^2 - a_{2k}^2.$$

If $A_3 = -153$, $A_5 = -435$ and $a_1^2 + a_2^2 + a_3^2 = 66$, then $a_{17} - A_7$ is equal to _____

(2024-Apr)

- 2) Suppose **AB** is focal chord of the parabola $y^2 = 12x$ of length l and slope $m < \sqrt{3}$,
If the distance of the chord **AB** from the origin is d , then ld^2 is equal to _____

(2024-Apr)

- 3) The number real roots of the equation $|x||x+2| - 5|x+1| - 1 = 0$ is _____

(2024-Apr)

- 4) If

$$S = \{a \in \mathbf{R} : |2a - 1| = 3[a] + 2\{a\}\}, A = 72 \sum_{a \in S} a,$$

where $[t]$ denotes the greatest integer less than or equal to t and $\{t\}$ represents the fractional part of t , then A is equal to _____

(2024-Apr)

- 5) Let $\bar{\mathbf{a}} = \hat{i} - 3\hat{j} + 7\hat{k}$, $\bar{\mathbf{b}} = 2\hat{i} - \hat{j} + \hat{k}$ and $\bar{\mathbf{c}}$ be a vector such that $(\bar{\mathbf{a}} + 2\bar{\mathbf{b}}) \times \bar{\mathbf{c}} = 3(\bar{\mathbf{c}} \times \bar{\mathbf{a}})$.
If $\bar{\mathbf{a}} \cdot \bar{\mathbf{c}} = 130$, then $\bar{\mathbf{b}} \cdot \bar{\mathbf{c}}$ is equal to _____

(2024-Apr)

- 6) The area of the region enclosed by the parabolas $y = x^2 - 5x$ and $y = 7x - x^2$ is _____

(2024-Apr)

7) let f be a differentiable function in the interval $(0, \infty)$ such that $f(1) = 1$ and

$$\lim_{t \rightarrow x} \frac{t^2 f(x) - x^2 f(t)}{t - x} = 1 \text{ for each } x > 0.$$

Then $2f(2) + 3f(3)$ is equal to _____

(2024-Apr)

8) If the constant term in the expansion of

$$\left(1 + 2x - 3x^3\right)\left(\frac{3}{2}x^2 - \frac{1}{3x}\right)^9 \text{ is } p,$$

then $108p$ is equal to _____

(2024-Apr)

9) From a lot of 10 items, which include 3 defective items, a sample of 5 items is drawn at random. Let the random variable X denote the numbers of defective items in the sample. If the variance of X is σ , then $96\sigma^2$ is equal to _____

(2024-Apr)

10) The number of ways of getting sum 16 on throwing a dice four times is _____

(2024-Apr)