Assignment 1

Chapter-11: Limits, Continuity and Differentiability

EE24BTECH11049

Patnam Shariq Faraz Muhammed

D (11 - 25): MCQs with One or More than One CORRECT

1) Let $g(x) = x \cdot f(x)$, where

$$f(x) = \begin{cases} x. \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$
. At $x = 0$

(1995)

- (a) g is differentiable but g' is not continuous
- (b) g is differentiable while f is not
- (c) both f and g are differentiable
- (d) g is differentiable and g' is continuous
- 2) The function $f(x) = \max\{(1-x), (1+x), 2\}$, $x \in (-\infty, \infty)$

(1995)

- (a) continuous at all points
- (b) differentiable at all points
- (c) differentiable at all points except at x = land x = -1
- (d) continuous at all points except at x = l and x = -1 where it is discontinuous
- 3) Let $h(x) = \min\{x, x^2\}$

(1998 - 2marks)

- (a) h is continuous for all x
- (b) h is differentiable for all x
- (c) h'(t) = 1, for all x > 1
- (d) h is not differentiable at two values of x

4)
$$\lim_{x\to 1} \frac{\sqrt{1-\cos 2(x-1)}}{x-1}$$

(1998 - 2marks)

- (a) exists and it equals $\sqrt{2}$
- (b) exists and it equals $-\sqrt{2}$
- (c) does not exist because $x 1 \mapsto 0$

(d) does not exist because the left-hand limit is not equal to the right-hand limit

5) If
$$f(x) = \min\{1, x^2, x^3\}$$

(2006, 5M, -1)

1

- (a) f(x) is continuous $\forall x \in R$
- (b) f(x) is continuous and differentiable everywhere
- (c) f(x) is not differentiable at two points
- (d) f(x) is not differentiable at one point

6) Let
$$L = \lim_{x\to 0} \frac{a - \sqrt{a^2 - x^2 - \frac{x^2}{4}}}{x^4}, a > 0.$$
 If L is finite, then

(2009)

- (a) a = 2
- (b) a = 1
- (c) $L = \frac{1}{64}$ (d) $L = \frac{1}{32}$
- 7) Let $f: R \to R$ be a function such that $f(x+y) = f(x) + f(y), \forall x, y \in R$. If f(x) is differentiable at x = 0, then

(2011)

- (a) f(x) is differentiable only in a finite interval containing zero
- (b) f(x) is continuous $\forall x \in R$
- (c) f'(x) is constant $\forall x \in R$
- (d) f(x) is differentiable except at finitely many points

8) If
$$f(x) = \begin{cases} -x - \frac{\pi}{2}, & x \le \frac{\pi}{2} \\ -\cos x, & \frac{\pi}{2} < x \le 0 \\ x - 1, & 0 < x \le 1 \\ \ln x, & x > 1 \end{cases}$$
 (2011)

- (a) f(x) is continuous at $x = \frac{\pi}{2}$
- (b) f(x) is not differentiable at x = 0
- (c) f'(x) is differentiable at x = 1
- (d) f(x) is differentiable at $x = \frac{3}{2}$
- 9) For every integer n, let a_n and b_n , be real numbers. Let function $f(x): IR \mapsto IR$ be given by $f(x) = \begin{cases} a_n + \sin \pi x, & for x \in [2n, 2n+1] \\ b_n + \cos \pi x, & for x \in (2n-1, 2n) \end{cases}$ for all integers n. If f is continuous, then which of the following hold(s) for all n

(2012)

- (a) $a_{n-1} b_{n-1} = 0$
- (b) $a_n b_n = 1$
- (c) $a_n b_{n+1} = 1$
- (d) $a_{n-1} b_n = -1$
- 10) For $a \in R$ (thesetofallrealnumbers), $a \neq -1 \lim_{n \to \infty} \frac{(1^a + 2^a + \dots + n^a)}{(n+1)^a \cdot [(na+1) + (na+2) + \dots + (na+n)]} = \frac{1}{60}$ Then $a = \frac{1}{60}$

(*JEE Adv.*2013)

- (a) 5
- (b) 7
- (c) $\frac{-15}{2}$
- (d) $\frac{2}{17}$
- 11) Let $f : [a, b] \mapsto [1, \infty)$ be a continuous function and let $g : R \mapsto R$ be defined as $f(x) = \begin{cases} 0, & if x < a, \\ \int_a^x f(t) dt, & if a \le x \le b; \\ \int_a^b f(t) dt, & if x > b \end{cases}$

(*JEE Adv.*2013)

- (a) g(x) is continuous but not differentiable at a
- (b) g(x) is differentiable on R
- (c) g(x) is continuous but not differentiable at b
- (d) g(x) is continuous and differentiable at either (a) or (b) but not both
- 12) For every pair of continuous functions f, g: $[0,1] \mapsto R$ such that max $\{f(x) : x \in [0,1]\}=$ max $\{g(x) : x \in [0,1]\}$, the correct statement(s) is(are):

(JEE Adv.2014)

- (a) $(f(c))^2 + 3 \cdot f(c) = (g(c))^2 + 3 \cdot g(c)$ for some $c \in [0, 1]$
- (b) $(f(c))^2 + f(c) = (g(c))^2 + 3 \cdot g(c)$ for some $c \in [0, 1]$
- (c) $(f(c))^2 + 3.f(c) = (g(c))^2 + g(c)$ for some $c \in [0, 1]$
- (d) $(f(c))^2 = (g(c))^2$ for some $c \in [0, 1]$

13) Let $g: R \mapsto R$ be a differentiable function with g(0) = 0, g'(0) = 0 and $g'(1) \neq 0$. Let $f(x) = \begin{cases} \frac{x}{|x|} \cdot g(x), & x \neq 0 \\ 0, & x = 0 \end{cases}$ and $h(x) = e^{|x|}$ for all $x \in R$. Let $(f \circ h)(x)$ denote f(h(x)) and $(h \circ f)(x)$ denote h(f(x)). Then which of the following is (are) true?

(JEE Adv.2015)

- (a) f is differentiable at x = 0
- (b) h is differentiable at x = 0
- (c) $f \circ h$ is differentiable at x = 0
- (d) $h \circ f$ is differentiable at x = 0
- 14) Let $a, b \in R$ and $f : R \mapsto R$ be defined by $f(x) = a \cdot \cos(|x^3 x|) + b \cdot |x| \cdot \sin(|x^3 + x|)$. Then f is

(*JEE Adv.*2016)

- (a) differentiable at x = 0 if a = 0 and b = 1
- (b) differentiable at x = 1 if a = 1 and b = 0
- (c) NOT differentiable at x = 0 if a = 1 and b = 0
- (d) NOT differentiable at x = 1 if a = 0 and b = 1
- 15) Let $f: \left[-\frac{1}{2}, 2\right] \mapsto R$ and $g: \left[-\frac{1}{2}, 2\right] \mapsto R$ be functions defined by $f(x) = \left[x^2 3\right]$ and $g(x) = |x| \cdot f(x) + |4x 7| \cdot f(x)$, where [y] denotes the greatest integer less than or equal to y for $y \in R$. Then

(JEE Adv. 2016)

- (a) f is discontinuous exactly at three points in $\left[-\frac{1}{2}, 2\right]$
- (b) f is discontinuous exactly at four points in $\left[-\frac{1}{2}, 2\right]$
- (c) g is NOT differentiable exactly at four points in $\left[-\frac{1}{2}, 2\right]$
- (d) g is NOT differentiable exactly at five points in $\left[-\frac{1}{2}, 2\right]$