

Crop Weather Calendar of Rabi Pointed gourd: Bogura region (Districts: Pabna, Sirajganj, Bogura, Joypurhat)

[illegible]

Crop Weather Calendar of *Kharif*season Pointed gourd: Bogura region (Districts: **Pabna, Sirajganj, Bogura, Joypurhat)**

Region: Bogura	Pointed Gourd- Kharif season									Duration: 100-120 days													
Month	March					April				May					June				July				
Std.Week/Normal	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	
Rainfall (mm)	5.0	2.0	4.0	3.0	8.0	8.0	12.0	26.0	28.0	37.0	42.0	45.0	58.0	59.0	74.0	70.0	92.0	76.0	93.0	80.0	91.0	76.0	
Max. Temp. (°C)	29.3	30.9	31.9	32.9	32.9	33.9	34.3	34.0	33.6	33.1	33.6	33.4	33.3	33.9	33.4	32.7	32.3	32.3	32.0	31.9	32.0	32.1	
Min. Temp. (°C)	16.4	17.4	18.3	20.0	21.1	22.3	23.0	23.0	23.1	23.3	23.9	24.3	24.6	25.0	25.9	26.0	26.1	26.1	26.0	26.1	26.0	26.1	
Mean Temp. (°C)	22.9	24.2	25.1	26.5	27.0	28.1	28.7	28.5	28.4	28.2	28.8	28.9	29.0	29.5	29.7	29.4	29.2	29.2	29.0	29.0	29.0	29.1	
RHmax (%)	88.7	89.1	86.4	86.3	87.7	88.0	88.3	90.6	90.9	92.4	92.4	92.7	93.0	93.7	94.1	95.1	95.1	95.6	95.9	95.9	95.9	95.9	
RHMin (%)	38.0	36.9	37.1	39.1	45.3	44.4	45.9	50.7	54.9	57.3	58.7	60.0	63.4	64.6	66.7	70.4	72.4	73.9	73.3	74.1	74.4	73.3	
RHmean (%)	63.4	63.0	61.8	62.7	66.5	66.2	67.1	70.7	72.9	74.9	75.6	76.4	78.2	79.2	80.4	82.8	83.8	84.8	85.6	85.0	85.2	84.6	
SShr (hrs)	60.0	59.0	61	58.0	56.0	56.0	56.0	55.0	52.0	48.0	50.0	49.0	46.0	44.0	39.0	33.0	28.0	26.0	28.0	31.0	29.0	28.0	
WD (deg)	235	230	225	205	176	173	164	149	134	127	126	126	121	135	134	134	134	132	141	135	138	143	
WS(Km/hr)	5.7	6.3	6.7	6.7	6.8	6.7	6.3	6.8	7.4	7.2	7.4	6.7	6.3	6.3	6.3	7.2	6.3	6.7	5.6	5.6	6.8	6.1	
																							
Stage	Spouting & Seedling					vegetative growth				Flowering & fruiting					Harvesting								
Favourable Environment																							
Temperature	Vine growth retarded below 15°C temperature, temperature 25-35°C (Vegetative growth, Flowering & fruiting)																						
Relative Humidity	60-80%																						
Soil Temperature	20-25°C																						
Sunshine Hour	Minimum 6 to 8 hrs. (Vine transplanting), 8-10 hrs. (Vegetative growth, Flowering & fruiting)																						
Favorable environmental conditions for insects and diseases																							
Fruit fly										Temperature: 24-30°C,RH: 55-75%													
Epilachna Beetle										Temperature: 25-30°C,RH: 55-65%													
Mealy bug										Temperature: 20-32°C,RH: 55-60%													
White fly										Temperature: 22-32°C,RH: 55-75%													
Red mite										Temperature: 15-26°C,RH: 60-70%													
Root knot (Nematode)										Temperature: 18-26°C,RH: 60-70%													
Powdery Mildew										Temperature: 22-27°C, Low humidity at day time and High humidity at night													
Weather warning																							
Rainfall (mm)						Early rainfall (10to 15mm)									More than 40mm rainfall								
Cloudy weather	Cloudy condition																						
Droughth	Drought due to excessive dry and warm weather(Pointed Gourd Drought Tolerant)																						
Duration of staying wet	Lands remain waterlogged for 2-3 days																						
Hailstorm	Early rainfall with hailstorm as well as seasonal rainfall																						
Windspeed (Km/hr.)	≥40 Km/hr																						
Temperature (°C)	Above 35°C																						

Crop Weather Calendar of *Rabi* season Pointed gourd: Cumilla region (Districts: Cumilla, Chandpur, Brahmanbaria)

Crop Weather Calendar of Rabi season Pointed Gourd: Cumilla Region (Districts: Cumilla, Chanchapur, Brahmanbaria)																		
Region: Cumilla	Pointed Gourd- Rabi season									Duration: 100-120 days								
Month	October				November					December				January				
Std.Week/Normal	40	41	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4	
Rainfall (mm)	51.0	46.5	30.5	16.0	11.5	18.0	8.5	4.0	2.5	0.00	4.5	1.0	1.5	0.0	0.0	0.0	1.5	
Max. Temp. (°C)	32.3	31.9	31.8	31.2	31.0	30.5	29.8	29.0	28.2	27.7	26.5	25.8	25.4	24.7	24.5	25.0	25.4	
Min. Temp. (°C)	25.4	24.7	24.3	22.8	21.8	21.0	19.3	18.1	16.9	15.8	15.0	14.2	13.4	13.0	12.7	12.7	12.6	
Mean Temp. (°C)	28.8	28.3	28.0	27.0	26.4	25.7	24.6	23.5	22.5	21.7	20.8	20.0	19.4	18.8	18.6	18.9	19.0	
RHmax (%)	95.6	95.7	95.5	95.2	95.3	95.6	95.3	95.4	95.8	95.8	96.2	96.0	96.0	96.0	95.5	95.3	94.4	
RHMin (%)	70.2	68.1	65.8	61.9	60.6	58.2	54.8	52.6	52.2	50.6	51.9	52.1	51.3	51.8	51.2	49.1	46.4	
RHmean (%)	82.9	81.9	80.6	78.5	77.9	76.9	75.0	74.0	74.0	73.2	74.1	74.1	73.6	73.9	73.4	72.2	70.4	
SShr (hrs)	39.0	43.0	44.5	50.5	47.0	49.5	50.0	50.5	48.5	47.5	42.5	40.5	40.0	40.0	37.0	43.5	47.0	
WD (deg)	203	214	201	187	202	183	184	190	195	206	192	211	213	205	209	218	212	
WS(Km/hr)	4.6	4.3	4.3	4.0	3.5	3.7	3.7	3.5	3.3	3.3	4.0	3.5	3.8	4.0	4.2	4.6	4.4	
																		
Stage	Spouting & Seedling				vegetative growth				Flowering & fruiting				Harvesting					
Favourable Environment																		
Temperature	Vine growth retarded below 15° C temperature, temperature 25-35° C (Vegetative growth, Flowering & fruiting)																	
Relative Humidity	60-80%																	
Soil Temperature	20-25° C																	
Sunshine Hour	Minimum 6 to 8 hrs. (Vine transplanting), 8-10 hrs. (Vegetative growth, Flowering & fruiting)																	
Favorable environmental conditions for insects and diseases																		
Fruit fly					Temperature: 24-30° C, RH: 55-75%													
Epilachna Beetle					Temperature: 25-30° C, RH: 55-65%													
Mealy bug					Temperature: 20-32° C, RH: 55-60%													
White fly					Temperature: 22-32° C, RH: 55-75%													
Red mite					Temperature: 15-26° C, RH: 60-70%													
Root knot (Nematode)					Temperature: 18-26° C, RH: 60-70%													
Powdery Mildew					Temperature: 22-27° C, Low humidity at day time and High humidity at night													
Weather warning																		
Rainfall (mm)	Late season rainfall (40mm or more)																	
Cloudy weather										Cloudy								
Droughth	Drought due to excessive dry and warm weather (Pointed Gourd Drought Tolerant)																	
Duration of staying wet	Lands remain waterlogged for 2-3 days																	
Hailstorm	Late season rainfall with hailstorm																	
Windspeed (Km/hr.)	≥ 40 Km/hr																	
Temperature (°C)	Below 15°C																	

Crop Weather Calendar of *Kharif* season Pointed gourd: Cumilla region (Districts: Cumilla, Chandpur, Brahmanbaria)

Region: Cumilla	Pointed Gourd- Kharif season									Duration: 100-120 days								
Month	February				March					April				May				
Std.Week/Normal	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
Rainfall (mm)	3.0	3.0	6.0	7.5	4.5	7.0	9.0	19.0	27.0	25.5	23.0	33.0	42.0	54.5	63.5	66.0	71.0	80.0
Max. Temp. (°C)	16.2	27.3	28.1	29.1	29.8	30.9	32.0	32.2	32.5	32.5	33.0	33.2	33.1	32.9	33.2	33.2	32.9	33.1
Min. Temp. (°C)	14.2	14.5	16.2	16.7	18.1	19.2	19.9	21.5	22.0	23.1	23.5	23.7	23.5	23.7	24.3	24.7	25.2	25.6
Mean Temp. (°C)	20.2	20.9	22.1	22.9	23.9	25.0	25.9	26.8	27.3	27.8	28.2	28.4	28.3	28.3	28.7	28.9	29.0	29.4
RHmax (%)	94.6	94.4	93.9	94.1	93.9	93.9	93.5	93.7	94.0	94.1	93.8	93.9	94.1	94.2	94.0	94.5	94.3	94.6
RHMin (%)	47.7	45.2	46.8	46.1	46.8	47.4	47.7	52.6	55.9	58.7	59.6	60.9	61.7	63.0	63.5	64.8	66.7	69.0
RHmean (%)	71.1	69.8	70.3	70.1	70.3	70.6	70.6	73.1	74.9	76.4	76.7	77.4	77.9	78.6	78.7	79.6	80.5	81.8
SShr (hrs)	47.0	50.0	49.5	50.5	52.5	52.5	52.5	51.0	48.5	51.0	52.0	52.5	51.0	47.5	47.5	46.5	41.0	38.0
WD (deg)	203	214	206	214	215	216	211	199	191	114	187	185	179	180	192	183	192	181
WS(Km/hr)	4.6	4.6	4.6	4.6	4.6	4.8	5.5	5.6	6.8	7.1	7.9	7.7	7.5	7.4	7.7	7.5	6.8	7.1
																		
Stage	Spouting & Seedling				vegetative growth				Flowering & fruiting					Harvesting				
Favourable Environment																		
Temperature	Vine growth retarded below 15° C temperature, temperature 25-35° C (Vegetative growth, Flowering & fruiting)																	
Relative Humidity	60-80%																	
Soil Temperature	20-25° C																	
Sunshine Hour	Minimum 6 to 8 hrs. (Vine transplanting), 8-10 hrs. (Vegetative growth, Flowering & fruiting)																	
Favorable environmental conditions for insects and diseases																		
Fruit fly									Temperature: 24-30° C, RH: 55-75%									
Epilachna Beetle									Temperature: 25-30° C, RH: 55-65%									
Mealy bug									Temperature: 20-32° C, RH: 55-60%									
White fly									Temperature: 22-32° C, RH: 55-75%									
Red mite									Temperature: 15-26° C, RH: 60-70%									
Root knot (Nematode)									Temperature: 18-26° C, RH: 60-70%									
Powdery Mildew									Temperature: 22-27° C, Low humidity at day time and High humidity at night									
Weather warning																		
Rainfall (mm)									Early rainfall (20 to 25mm)									
Cloudy weather									Cloudy condition									
Droughth	Drought due to excessive dry and warm weather (Pointed Gourd Drought Tolerant)																	
Duration of staying wet	Lands remain waterlogged for 2-3 days																	
Hailstorm									Early rainfall with hailstorm									
Windspeed (Km/hr.)	≥ 40 Km/hr																	
Temperature (°C)	Below 15°C Above 35°C																	

Crop Weather Calendar of *Kharif* Pointed gourd: Dhaka region(Districts: Dhaka, Narayanganja, Gazipur, Kishoreganj, Narsingdi, Munshiganj, Manikganj, Tangail)

Region: Dhaka	Pointed Gourd- Kharif Season													Duration: 100-120 days				
Month	February				March					April				May				
Std.Week/Normal	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
Rainfall (mm)	3.5	3.5	2.5	12.0	3.5	4.0	9.5	13.5	20.5	25.0	21.0	31.5	42.5	46.0	55.5	55.5	75.5	70.0
Max. Temp. (°C)	26.2	27.2	28.3	28.9	30.0	31.5	32.3	33.4	33.2	33.9	24.6	34.1	33.8	33.6	33.7	33.7	33.2	33.5
Min. Temp. (°C)	13.6	14.3	15.5	16.5	17.5	18.6	19.8	19.8	21.8	22.9	23.5	23.7	23.6	23.7	24.2	24.6	24.9	25.4
Mean Temp. (°C)	19.9	20.7	21.9	22.7	23.7	25.0	26.1	26.1	27.5	28.4	29.02	28.9	28.7	28.6	28.9	29.2	29.0	29.4
RHmax (%)	92.4	92.1	91.7	91.1	90.0	89.9	89.2	89.6	90.3	91.3	91.4	91.4	91.7	92.3	92.8	92.9	93.5	93.8
RHMin (%)	42.3	39.4	39.7	38.9	36.0	35.5	35.9	38.9	44.5	45.9	47.1	52.0	55.1	56.7	58.6	60.4	63.5	65.0
RHmean (%)	67.3	65.7	65.7	65.0	63.0	62.7	62.5	64.3	67.4	68.6	69.2	71.7	73.4	74.5	75.7	76.7	78.5	79.4
SShr (hrs)	49.5	51.5	53.0	54.5	56.0	55.0	58.0	55.5	52.5	53.5	54.0	55.0	54.5	51.5	48.0	48.0	42.5	42.0
WD (deg)	241	236	234	228	234	230	223	206	188	181	175	167	156	159	164	156	155	157
WS(Km/hr)	4.3	4.5	5.0	5.3	5.0	5.4	5.7	6.2	6.5	6.5	6.6	6.6	6.7	6.8	6.5	6.4	6.1	5.8
																		
Stage	Spouting & Seedling				vegetative growth				Flowering & fruiting					Harvesting				
Favourable Environment																		
Temperature	Vine growth retarded below 15°C temperature, temperature 25-35°C (Vegetative growth, Flowering & fruiting)																	
Relative Humidity	60-80%																	
Soil Temperature	20-25°C																	
Sunshine Hour	Minimum 6 to 8 hrs. (Vine transplanting), 8-10 hrs. (Vegetative growth, Flowering & fruiting)																	
Favorable environmental conditions for insects and diseases																		
Fruit fly					Temperature: 24-30°C,RH: 55-75%													
Epilachna Beetle					Temperature: 25-30°C,RH: 55-65%													
Mealy bug					Temperature: 20-32°C,RH: 55-60%													
White fly					Temperature: 22-32°C,RH: 55-75%													
Red mite					Temperature: 15-26°C,RH: 60-70%													
Root knot (Nematode)					Temperature: 18-26°C,RH: 60-70%													
Powdery Mildew					Temperature: 22-27°C, Low humidity at day time and High humidity at night													
Weather warning																		
Rainfall (mm)	Early rainfall (20 to 25mm)																	
Cloudy weather	Cloudy condition																	
Droughth	Drought due to excessive dry and warm weather(Pointed Gourd Drought Tolerant)																	
Duration of staying wet	Lands remain waterlogged for 2-3 days																	
Hailstorm	Early rainfall with hailstorm																	
Windspeed (Km/hr.)	≥40 Km/hr																	
Temperature (°C)	Below 15°C								Above 35°C									

Crop Weather Calendar of Rabi Pointed gourd: Dhaka region(Districts: Dhaka, Narayanganja, Gazipur, Kishoreganj,Narsingdi,Munshiganj, Manikganj, Tangail)

Region: Dhaka	Pointed Gourd- Rabi season													Duration: 100-120 days			
Month	November				December				January				February				
Std.Week/Normal	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8
Rainfall (mm)	12.5	10.0	6.0	1.5	4.5	1.5	4.5	1.5	3.5	1.0	0.0	0.0	0.5	3.5	3.5	2.5	12.0
Max. Temp. (°C)	30.9	30.6	29.8	29.0	28.3	27.6	26.2	25.5	24.8	24.0	23.9	24.5	25.0	26.2	27.2	28.3	28.9
Min. Temp. (°C)	21.5	20.5	18.7	17.7	16.11	15.5	14.7	13.5	12.9	12.4	12.0	12.0	12.1	13.6	14.3	15.5	16.5
Mean Temp. (°C)	26.2	25.5	24.3	23.3	22.2	21.5	20.5	19.5	28.8	18.2	17.9	18.3	18.6	19.9	20.7	21.9	22.7
RHmax (%)	94.5	95.0	94.7	94.6	94.3	94.8	95.3	95.1	95.0	94.6	94.3	93.8	93.4	92.4	92.1	91.7	91.1
RHMin (%)	54.6	51.5	49.3	47.5	46.3	47.6	49.4	49.8	49.1	50.3	48.8	46.2	43.3	42.3	39.4	39.7	38.9
RHmean (%)	74.6	73.2	72.0	71.0	70.3	71.2	72.3	72.5	72.1	72.4	71.6	70.0	68.3	67.3	65.7	65.7	65.0
SShr (hrs)	49.5	51.5	51.5	52.0	51.5	48.5	46.5	44.5	44.5	42.5	42.5	45.0	49.0	49.5	51.5	53.0	54.5
WD (deg)	190	189	199	212	211	219	224	225	233	238	238	232	242	241	236	234	228
WS(Km/hr)	3.7	3.7	3.7	3.7	3.6	3.7	3.7	3.7	3.7	4.1	4.0	4.1	4.3	4.3	4.5	5.0	5.3
																	
Stage	Spouting & Seedling				vegetative growth			Flowering & fruiting						Harvesting			
Favourable Environment																	
Temperature	Vine growth retarded below 15°C temperature, temperature 25-35°C (Vegetative growth, Flowering & fruiting)																
Relative Humidity	60-80%																
Soil Temperature	20-25°C																
Sunshine Hour	Minimum 6 to 8 hrs. (Vine transplanting), 8-10 hrs. (Vegetative growth, Flowering & fruiting)																
Favorable environmental conditions for insects and diseases																	
Fruit fly								Temperature: 24-30°C,RH: 55-75%									
Epilachna Beetle								Temperature: 25-30°C,RH: 55-65%									
Mealy bug								Temperature: 20-32°C,RH: 55-60%									
White fly								Temperature: 22-32°C,RH: 55-75%									
Red mite								Temperature: 15-26°C,RH: 60-70%									
Root knot (Nematode)								Temperature: 18-26°C,RH: 60-70%									
Powdery Mildew								Temperature: 22-27°C, Low humidity at day time and High humidity at night									
Weather warning																	
Rainfall (mm)	Early rainfall(15 to 20 mm)																
Cloudy weather	Cloudy condition																
Droughth	Drought due to excessive dry and warm weather(Pointed Gourd Drought Tolerant)																
Duration of staying wet	Land remain waterlogged for 2-3 days																
Hailstorm					Hailstorm with late seasonal rainfall												
Windspeed (Km/hr.)	≥40 Km/hr																
Temperature (°C)					Below 15°C									Above 35°C			

Crop Weather Calendar of kharif Pointed gourd:Dinaipur Region (Districts:Raishahi, Chapainawabgonj, Naoga, Natore) Bangladesh

Rongpur Region	kharif Pointed gourd										Life Span 140-150 day											
Months	May					June				July				August				September				
Std.Week	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	
Rainfall (mm)	31.0	36.0	57.0	55.0	73.0	57.0	77.0	111.0	96.0	109.0	105.0	81.0	84.0	70.0	60.0	93.0	103.0	71.0	66.0	61.0	73.0	
Max. Temp. (°C)	32.3	32.9	33.0	33.0	33.4	33.3	33.0	32.3	32.4	32.0	32.1	32.1	32.3	32.9	32.6	32.4	32.1	32.4	32.3	32.4	32.0	
Min. Temp. (°C)	22.3	22.9	23.6	24.0	24.4	25.0	25.1	25.6	26.0	26.0	26.0	26.0	26.0	26.3	26.7	26.1	26.0	26.0	26.0	25.7	25.1	
Mean Temp. (°C)	27.3	27.9	28.3	28.5	28.9	29.2	29.1	29.0	29.2	29.0	29.1	29.1	29.2	29.6	29.7	29.3	29.1	29.2	29.2	29.1	28.6	
RHmax (%)	92.3	92.0	92.1	92.4	92.7	93.4	93.9	94.7	94.4	95.1	94.6	94.4	94.7	94.1	94.0	95.0	95.7	95.6	95.6	96.1	96.0	
RHMin (%)	57.6	57.3	59.3	62.4	62.3	65.4	67.7	71.4	71.6	73.3	72.7	71.9	71.6	70.1	70.9	72.6	73.1	71.6	72.4	71.3	71.6	
RHmean (%)	75.0	74.7	75.7	77.4	77.5	79.4	80.8	83.1	83.0	84.2	83.7	83.2	83.2	82.1	82.5	83.8	84.4	83.6	84.0	83.7	83.8	
SShr (hrs)	49.0	51.0	48.0	44.0	42.0	40.0	35.0	27.0	31.0	27.0	29.0	31.0	36.0	36.0	37.0	34.0	34.0	37.0	37.0	36.0	33.0	
WD (deg)	118	115	119	108	117	111	118	120	121	127	121	126	134	132	132	134	139	141	133	143	146	
WS(Km/hr)	3.9	3.7	4.3	3.7	3.9	3.7	3.7	3.7	3.7	3.7	3.7	3.9	3.7	3.9	3.7	3.7	3.7	3.7	3.7	3.7	3.7	
																						
Stages	Vine Planting					Vegetative Growth				Flowering & Fruiting				Harvesting								
Favourable Environment																						
Temperature	Vine growth retarded below 15°C temperature, temperature 25-35°C (Vegetative growth, Flowering & fruiting)																					
Relative Humidity	60-80%																					
Soil Temperature	20-25°C																					
Sunshine Hour	Minimum 6 to 8 hrs. (Vine transplanting), 8-10 hrs.(Vegetative growth, Flowering & fruiting)																					
Favorable environmental conditions for insects and diseases																						
Fruit fly																Temperature: 24-30°C,RH: 55-75%						
Epilachna Beetle						Temperature: 25-30°C,RH: 55-65%																
Mealy bug						Temperature: 20-32°C,RH: 55-60%																
White fly						Temperature: 22-32°C,RH: 55-75%																
Red mite						Temperature: 15-26°C,RH: 60-70%																
Root knot (Nematode)						Temperature: 18-26°C,RH: 60-70%																
Powdery Mildew						Temperature: 22-27°C, Low humidity at day time and High humidity at night																
Weather warning																						
Rainfall (mm)	5-10mm (water logging)															Early rainfall (10-15mm)						
Cloudy weather						Cloudy condition																
Droughth						Drought due to excessive dry and warm weather(Pointed Gourd Drought Tolerant)																
Duration of staying wet						Land remain waterlogged for 2-3 days																
Hailstorm																Early rainfall with hailstorm						
Windspeed (Km/hr.)						≥40 Km/hr																
Temperature (°C)						Below 15°C										Above 35°C						

Crop Weather Calendar of Rabi Pointed Gourd: Jessore Region (Districts: Jessore, Chuadanga, Meherpur, Jinaidaha, magura, Kustia) Bangladesh

Region: Jessore					Rabi Pointed Gourd										Life cycle of crop, 140-250										
Months					October				November					December				January				February			
Std.Week					40	41	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8
Rainfall (mm)					44.5	33.0	26.5	20.0	9.0	8.0	7.5	1.0	2.0	1.5	4.0	0.0	3.5	0.5	1.5	2.5	3.0	6.5	5.0	4.5	8.0
Max. Temp. (°C)					33.2	33.0	32.7	31.9	31.1	31.1	30.3	29.6	28.5	27.7	26.6	25.7	25.2	24.5	24.5	25.1	25.6	26.6	27.8	28.9	30.0
Min. Temp. (°C)					25.1	24.4	23.4	21.6	20.7	19.4	18.0	16.7	15.1	14.0	13.4	12.3	11.4	10.9	10.7	11.0	10.8	12.0	13.3	14.7	15.6
Mean Temp. (°C)					29.1	28.7	28.0	26.7	25.9	25.3	24.2	23.1	21.8	20.9	20.0	19.0	18.3	17.7	17.6	18.0	18.2	19.3	20.5	21.8	22.8
RHmax (%)					96.5	96.5	96.5	96.1	95.9	95.9	95.7	95.7	95.4	95.2	95.8	95.5	95.7	95.4	95.3	95.3	94.9	94.9	94.2	94.0	93.7
RHMin (%)					67.0	64.4	61.5	57.1	56.2	52.0	49.8	48.1	46.3	47.2	49.2	49.4	47.9	49.6	47.9	47.0	44.7	44.2	42.2	41.0	39.3
RHmean (%)					81.7	80.4	79.0	76.6	76.0	74.0	72.8	71.9	70.8	71.2	72.5	72.5	71.8	72.5	71.6	71.2	69.8	69.5	68.2	67.5	66.5
SShr (hrs)					40.0	43.5	47.0	49.0	45.5	48.5	49.5	47.5	48.5	47.5	43.5	42.0	42.0	41.0	42.5	44.5	48.0	48.5	49.5	50.0	54.5
WD (deg)					195	198	187	182	153	175	177	179	178	184	183	177	174	158	156	188	192	186	191	203	214
WS(Km/hr)					3.1	3.0	2.9	2.5	2.5	2.3	2.4	2.3	2.4	2.3	2.4	2.5	2.5	3.0	3.0	3.0	3.0	3.2	3.1	3.3	3.7
																									
Stages					Vine Planting				Vegetative Growth				Flowering & Fruiting				Harvesting								
Favourable Environment																									
Temperature					Vine growth retarded below 15°C temperature, temperature 25-35°C (Vegetative growth, Flowering & fruiting)																				
Relative Humidity					60-80%																				
Soil Temperature					20-25°C																				
Sunshine Hour					Minimum 6 to 8 hrs. (Vine transplanting), 8-10 hrs.(Vegetative growth, Flowering & fruiting)																				
Favorable environmental conditions for insects and diseases																									
Fruit fly									Temperature: 24-30°C,RH: 55-75%																
Epilachna Beetle									Temperature: 25-30°C,RH: 55-65%																
Mealy bug									Temperature: 20-32°C,RH: 55-60%																
White fly									Temperature: 22-32°C,RH: 55-75%																
Red mite									Temperature: 15-26°C,RH: 60-70%																
Root knot (Nematode)									Temperature: 18-26°C,RH: 60-70%																
Powdery Mildew									Temperature: 22-27°C, Low humidity at day time and High humidity at night																
Weather warning																									
Rainfall (mm)					5-10mm (water logging)				Early rainfall (10-15mm)																
Cloudy weather									Cloudy condition																
Droughth									Drought due to excessive dry and warm weather(Pointed Gourd Drought Tolerant)																
Duration of staying wet									Land remain waterlogged for 2-3 days																
Hailstorm									Early rainfall with hailstorm																
Windspeed (Km/hr.)									≥40 Km/hr																
Temperature (°C)					Below 15°C				Above 35°C																

Crop Weather Calendar of Kharif Pointed Gourd: Jessore Region (Districts: Jessore, Chuadanga, Meherpur, Jinaidaha, magura, Kustia) Bangladesh

Region: Jessore	Kharif Pointed Gourd								Life cycle of crop, 140-250									
Months	March					April				May					June			
Std.Week	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26
Rainfall (mm)	4.0	2.5	4.5	11.5	12.0	11.5	7.0	12.5	21.5	28.5	30.5	36.0	44.5	38.5	49.5	54.5	81.0	72.5
Max. Temp. (°C)	31.0	32.4	33.6	34.7	34.6	35.5	36.1	36.5	36.3	36.0	36.0	35.9	35.5	35.6	35.3	34.0	33.3	33.0
Min. Temp. (°C)	16.4	17.7	19.0	20.8	21.8	22.8	23.5	24.1	24.2	24.2	25.2	25.0	25.5	26.0	26.2	26.2	26.1	26.2
Mean Temp. (°C)	23.7	25.0	26.3	27.7	28.2	29.2	29.8	30.3	30.3	30.1	30.6	30.4	30.5	30.8	30.7	30.1	29.7	29.6
RHmax (%)	93.2	92.5	92.1	92.1	92.4	92.4	92.9	92.4	92.3	92.8	92.7	93.1	93.6	94.0	94.1	94.8	95.7	96.0
RHMin (%)	36.5	35.4	34.9	36.7	42.0	41.5	41.6	44.3	46.9	49.9	51.7	54.8	58.7	60.0	62.1	67.5	71.7	73.5
RHmean (%)	64.9	63.9	63.9	64.4	67.2	66.9	67.2	68.3	69.6	71.3	72.2	74.0	76.1	77.0	78.1	81.1	83.7	84.7
SShr (hrs)	55.0	54.0	55.0	53.0	51.5	55.5	54.5	54.5	55.5	54.0	51.5	52.5	46.5	45.5	41.5	32.0	26.5	26.0
WD (deg)	219	213	230	212	208	209	204	193	188	186	186	180	181	180	183	174	175	178
WS(Km/hr)	3.5	3.9	4.1	4.4	4.7	5.1	5.2	5.7	5.6	5.6	5.8	5.2	5.2	5.0	4.6	4.9	4.6	4.6
																		
Stages	Germination and Seedling Planting				Vegetative Growth				Flowering & Fruiting					Harvesting				
Favourable Environment																		
Temperature	Vine growth retarded below 15°C temperature, temperature 25-35°C (Vegetative growth, Flowering & fruiting)																	
Relative Humidity	60-80%																	
Soil Temperature	20-25°C																	
Sunshine Hour	Minimum 6 to 8 hrs. (Vine transplanting), 8-10 hrs.(Vegetative growth, Flowering & fruiting)																	
Favorable environmental conditions for insects and diseases																		
Fruit fly									Temperature: 24-30°C,RH: 55-75%									
Epilachna Beetle									Temperature: 25-30°C,RH: 55-65%									
Mealy bug									Temperature: 20-32°C,RH: 55-60%									
White fly									Temperature: 22-32°C,RH: 55-75%									
Red mite									Temperature: 15-26°C,RH: 60-70%									
Root knot (Nematode)									Temperature: 18-26°C,RH: 60-70%									
Powdery Mildew					Temperature: 22-27°C, Low humidity at day time and High humidity at night													
Weather warning																		
Rainfall (mm)	5-10mm (water logging)								Early rainfall (10-15mm)									
Cloudy weather	Cloudy condition																	
Droughth					Drought due to excessive dry and warm weather(Pointed Gourd Drought Tolerant)													
Duration of staying wet	Land remain waterlogged for 2-3 days																	
Hailstorm														Early rainfall with hailstorm				
Windspeed (Km/hr.)	≥40 Km/hr																	
Temperature (°C)	Below 15°C								Above 35°C									

Crop Weather Calendar of *Kharif* season Pointed gourd:Mymensingh region (Districts: Mymensingh, Netrokona, Jamalpur, Sherpur)

Reg.: Mymensingh	Pointed gourd-Kharif season													Duration: 120-150 days							
Months	February				March					April				May					June		
Std.Week/Normal	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
Rainfall (mm)	2.0	1.0	5.0	7.0	1.0	4.0	6.0	11.0	17.0	20.0	30	40	36	55	62	69	97	81	96	103	98
Max. Temp. (°C)	25.9	26.6	27.4	28.1	29.1	30.3	31.0	31.9	31.3	32.1	32.7	32.0	31.7	31.6	31.9	32.3	32.0	32.0	32.0	31.7	31.4
Min. Temp. (°C)	13.0	13.7	15.0	15.9	16.6	17.4	18.6	19.9	20.9	21.9	22.4	22.9	22.9	22.9	23.6	24.0	24.3	25.0	25.6	25.7	26.0
Mean Temp. (°C)	19.5	20.2	21.2	22.0	22.9	23.9	24.8	25.9	26.1	27.0	27.6	27.5	27.3	27.3	27.8	28.2	28.2	28.5	28.8	28.7	28.7
RHmax (%)	92.6	92.1	91.4	90.6	90.6	90.6	90.1	89.7	90.4	89.9	90.9	91.1	91.3	92.4	92.6	92.7	92.7	94.0	94.1	94.7	95.0
RHMin (%)	47.9	45.3	45.0	45.1	44.3	43.6	44.6	48.3	54.7	55.0	56.6	61.1	62.6	64.6	66.0	66.4	69.4	71.9	73.0	75.4	76.6
RHmean (%)	70.3	68.7	68.2	67.9	67.6	67.1	67.4	69.0	72.6	72.5	73.8	76.1	77.0	78.5	79.3	79.6	81.1	83.0	83.6	85.1	85.8
SShr (hrs)	44.0	48.0	50.0	50.0	51.0	53.0	53.0	48.0	44.0	49.0	49.0	49.0	46.0	44.0	44.0	45.0	39.0	36.0	34.0	26.0	25.0
WD (deg)	228	218	222	210	214	202	208	179	165	149	146	143	128	126	134	131	126	130	142	138	143
WS(Km/hr)	3.7	3.9	4.4	5.4	5.0	5.4	5.6	6.3	6.1	7.2	6.8	7.2	7.6	7.4	6.8	7.4	7.4	6.8	6.1	7.4	6.8
																					
Stage	Spouting & Seedling				vegetative growth					Flowering & fruiting					Harvesting						
Favourable Environment																					
Temperature	Vine growth retarded below 15° C temperature, temperature, 25-35° C (Vegetative growth, Flowering & fruiting)																				
Relative Humidity	60-80%																				
Soil Temperature	20-25° C																				
Sunshine Hour	Minimum 6 to 8 hrs. (Vine transplanting), 8-10 hrs. (Vegetative growth, Flowering & fruiting)																				
Favorable environmental conditions for insects and diseases																					
Fruit fly										Temperature: 24-30° C, RH: 55-75%											
Epilachna Beetle										Temperature: 25-30° C, RH: 55-65%											
Mealy bug										Temperature: 20-32° C, RH: 55-60%											
White fly										Temperature: 22-32° C, RH: 55-75%											
Red mite										Temperature: 15-26° C, RH: 60-70%											
Root knot (Nematode)										Temperature: 18-26° C, RH: 60-70%											
Powdery Mildew										Temperature: 22-27° C, Low humidity at day time and High humidity at night											
Weather warning																					
Rainfall (mm)					Early rainfall (20 to 25 mm)																
Cloudy weather	Cloudy condition																				
Droughth	Drought due to excessive dry and warm weather (Pointed Gourd Drought Tolerant)																				
Duration of staying wet	Lands remain waterlogged for 2 to 3 days																				
Hailstorm					Early and seasonal rainfall with hailstorms																
Windspeed (Km/hr.)	≥ 40 Km/hr																				
Temperature (°C)	Below 15°C				Above 35°C																

Crop Weather Calendar of *Kharif* season Pointed gourd:Mymensingh region (Districts: Mymensingh, Netrokona, Jamalpur, Sherpur)

Reg.: Mymensingh	Pointed gourd-Kharif													Duration: 120-150 days			
Months	June				July				August					September			
Std.Week/Normal	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39
Rainfall (mm)	96.0	103.0	98.0	85.0	109.0	99.0	89.0	100.0	72.0	74.0	67.0	81.0	75.0	63.0	65.0	79.0	87.0
Max. Temp. (°C)	32.0	31.7	31.74	31.0	31.1	31.3	31.3	31.1	32.0	32.0	32.0	32.0	32.0	31.4	31.7	31.9	31.7
Min. Temp. (°C)	25.6	25.7	26.0	26.0	26.0	26.0	26.0	26.0	26.1	26.3	26.3	26.0	26.0	26.0	26.0	26.0	25.1
Mean Temp. (°C)	28.8	28.7	28.7	28.5	28.6	28.7	28.7	28.6	29.1	29.2	29.2	29.0	29.0	28.7	28.9	29.0	28.4
RHmax (%)	94.1	94.7	95.0	94.7	95.0	94.6	94.6	94.9	94.7	93.9	94.0	94.9	94.7	94.9	95.3	95.0	95.4
RHMin (%)	73.0	75.4	76.6	77.7	77.6	77.0	76.6	76.4	74.6	74.6	74.9	75.9	75.0	75.9	74.1	74.1	74.0
RHmean (%)	83.6	85.1	85.8	86.2	86.3	85.8	85.6	85.7	84.7	84.3	84.5	85.6	84.9	85.4	84.7	84.6	84.7
SShr (hrs)	34.0	26.0	25.0	23.0	24.0	24.0	25.0	30.0	33.0	31.0	30.0	26.0	31.0	28.0	31.0	31.0	31.0
WD (deg)	142.0	138	143	144	156	152	152	151	153	149	148	157	155	151	157	156	167
WS(Km/hr)	6.1	7.4	6.8	7.2	7.2	6.8	7.2	6.1	6.1	6.1	6.1	5.7	6.1	5.6	5.6	5.4	4.4
																	
Stage	Spouting & Seedling			vegetative growth			Flowering & fruiting					Harvesting					
Favourable Environment																	
Temperature	Vine growth retarded below 15° C temperature, temperature, 25-35° C (Vegetative growth, Flowering & fruiting)																
Relative Humidity	60-80%																
Soil Temperature	20-25° C																
Sunshine Hour	Minimum 6 to 8 hrs. (Vine transplanting), 8-10 hrs. (Vegetative growth, Flowering & fruiting)																
Favorable environmental conditions for insects and diseases																	
Fruit fly				Temperature: 24-30° C, RH: 55-75%													
Epilachna Beetle				Temperature: 25-30° C, RH: 55-65%													
Mealy bug				Temperature: 20-32° C, RH: 55-60%													
White fly				Temperature: 22-32° C, RH: 55-75%													
Red mite				Temperature: 15-26° C, RH: 60-70%													
Root knot (Nematode)				Temperature: 18-26° C, RH: 60-70%													
Powdery Mildew				Temperature: 22-27° C, Low humidity at day time and High humidity at night													
Weather warning																	
Rainfall (mm)	≥ 40 mm																
Cloudy weather	Cloudy condition																
Droughth	Drought due to lack of rainfall (Pointed Gourd Drought Tolerant)																
Duration of staying wet	Lands remain waterlogged for 2 to 3 days																
Hailstorm	Hailstorms with seasonal rainfall																
Windspeed (Km/hr.)	≥ 40 Km/hr																
Temperature (°C)	Above 35°C																

Crop Weather Calendar of *Rabi* season Pointed gourd: Rajshahi region (Districts: Rajshahi, Chapai Nawabganj, Naogaon, Natore)

Region: Rajshahi	Pointed Gourd- <i>Rabi</i> season													Duration: 100-120 days			
Months	September				October				November					December			
Std.Week/Normal	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52
Rainfall (mm)	54.0	59.0	52.0	83.0	44.0	35.0	23.0	12.0	7.0	5.0	1.0	0.0	2.0	1.0	3.0	0.0	4.0
Max. Temp. (°C)	32.9	33.0	33.0	32.9	32.7	32.3	31.7	32.3	31.7	31	29.6	28.9	28.3	27.0	26.0	25.4	24.4
Min. Temp. (°C)	26.0	26.0	26.0	25.3	25.0	24.1	23.1	24.1	23.1	21.4	17.6	16.6	15.0	14.0	13.1	12.4	11.6
Mean Temp. (°C)	29.5	29.5	29.5	29.1	28.9	28.2	27.4	28.2	27.4	26.2	23.6	22.8	21.7	20.5	19.6	18.9	18
RHmax (%)	96.3	96.7	96.3	96.7	96.3	96.4	96.1	96.4	96.1	96.0	95.7	95.3	94.7	94.7	95.9	95.3	95.7
RHMin (%)	71.3	71.0	69.9	68.9	67.4	64.3	60.9	64.3	60.9	57.4	49.4	47.4	45.6	46.9	50.1	50.0	50.0
RHmean (%)	83.8	83.9	83.0	82.8	81.9	80.4	78.5	80.4	78.5	76.7	72.6	71.4	70.2	70.8	73.0	72.7	72.9
SShr (hrs)	35.0	38.0	40.0	43.0	46.0	47.0	53.0	47.0	53.0	56.0	54.0	55.0	55.0	51.0	47.0	46.0	44.0
WD (deg)	142	143	143	153	145	154	146	154	146	146	151	146	145	143	146	144	167
WS(Km/hr)	5.6	5.6	5.6	5.4	4.8	3.9	3.9	3.9	3.9	4.3	3.7	3.7	3.7	3.7	3.7	3.7	3.7
																	
Stage	Spouting & Seedling			vegetative growth				Flowering & fruiting					Harvesting				
Favourable Environment																	
Temperature	Vine growth retarded below 15° C temperature, temperature 25-35° C (Vegetative growth, Flowering & fruiting)																
Relative Humidity	60-80%																
Soil Temperature	20-25° C																
Sunshine Hour	Minimum 6 to 8 hrs. (Vine transplanting), 8-10 hrs. (Vegetative growth, Flowering & fruiting)																
Favorable environmental conditions for insects and diseases																	
Fruit fly				Temperature: 24-30° C, RH: 55-75%													
Epilachna Beetle				Temperature: 25-30° C, RH: 55-65%													
Mealy bug				Temperature: 20-32° C, RH: 55-60%													
White fly				Temperature: 22-32° C, RH: 55-75%													
Red mite				Temperature: 15-26° C, RH: 60-70%													
Root knot (Nematode)				Temperature: 18-26° C, RH: 60-70%													
Powdery Mildew				Temperature: 22-27° C, Low humidity at day time and High humidity at night													
Weather warning																	
Rainfall (mm)	Rainfall 30 mm or above																
Cloudy weather	Cloudy condition																
Droughth								Drought due to excessive dry and warm weather (Pointed Gourd Drought Tolerant)									
Duration of staying wet	Lands remain waterlogged for 2 to 3 days																
Hailstorm	Hailstorms with seasonal rainfall																
Windspeed (Km/hr.)	≥ 40 Km/hr																
Temperature (°C)								Below 20 °C									

Crop Weather Calendar of *Kharif* season Pointed gourd:Rajshahi region (Districts: Rajshahi, Chapai Nawabganj, Naogaon, Natore)

[illegible]

Crop Weather Calendar of kharif Pointed gourd:Rongpur Region (Districts:Raishahi, Chapainawabgonj, Naoga, Natore) Bangladesh

Rongpur Region	kharif Pointed gourd										Life Span 140-150 day										
Months	October				November					December				January				February			
Std.Week	40	41	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8
Rainfall (mm)	58.0	53.0	35.5	10.0	4.0	3.0	2.0	0.0	1.0	0.0	2.5	0.0	2.5	1.0	2.5	2.0	1.0	1.0	1.5	3.5	1.5
Max. Temp. (°C)	32.1	31.5	31.2	30.6	30.0	29.8	29.0	28.3	27.7	26.7	25.1	24.7	24.1	22.7	22.4	23.2	23.4	24.7	25.9	26.7	27.5
Min. Temp. (°C)	24.9	24.0	22.5	21.0	20.0	19.0	17.5	16.4	15.0	14.2	13.5	12.5	11.5	10.9	10.4	10.8	10.8	11.8	12.7	13.7	14.4
Mean Temp. (°C)	28.5	27.8	26.8	25.8	25.0	24.4	23.3	22.3	21.3	20.4	19.3	18.6	17.8	16.8	16.4	17.0	17.0	18.3	19.3	20.2	21.0
RHmax (%)	95.5	95.4	95.2	95.0	94.5	94.4	94.6	94.9	94.9	95.0	95.4	95.8	96.1	96.0	96.0	95.8	95.7	95.5	94.3	93.7	92.9
RHMin (%)	69.3	67.9	64.5	60.8	58.7	55.9	54.0	50.9	50.6	53.5	56.2	56.2	55.2	58.5	57.8	56.2	54.2	51.6	48.0	47.1	45.3
RHmean (%)	82.4	81.7	79.9	77.9	76.6	75.1	74.3	72.9	72.7	74.2	75.8	76.0	75.6	77.3	76.9	76.0	74.9	73.5	71.2	70.4	69.1
SShr (hrs)	42.5	46.0	51.5	56.5	55.5	55.0	52.0	54.0	53.0	49.5	41.5	42.5	40.0	36.0	36.5	39.5	41.0	44.0	48.5	50.0	51.5
WD (deg)	147	143	134	127	115	106	126	134	150	129	158	160	187	199	214	199	210	208	192	201	201
WS(Km/hr)	6.0	6.4	6.3	6.1	6.6	6.6	6.3	6.4	5.6	6.0	5.7	5.6	5.6	5.7	5.7	5.7	5.6	6.2	6.6	7.5	7.2
																					
Stages	Vine Planting				Vegetative Growth				Flowering & Fruiting				Harvesting								
Favourable Environment																					
Temperature	Vine growth retarded below 15°C temperature, temperature 25-35°C (Vegetative growth, Flowering & fruiting)																				
Relative Humidity	60-80%																				
Soil Temperature	20-25°C																				
Sunshine Hour	Minimum 6 to 8 hrs. (Vine transplanting), 8-10 hrs.(Vegetative growth, Flowering & fruiting)																				
Favorable environmental conditions for insects and diseases																					
Fruit fly										Temperature: 24-30°C,RH: 55-75%											
Epilachna Beetle										Temperature: 25-30°C,RH: 55-65%											
Mealy bug										Temperature: 20-32°C,RH: 55-60%											
White fly										Temperature: 22-32°C,RH: 55-75%											
Red mite										Temperature: 15-26°C,RH: 60-70%											
Root knot (Nematode)										Temperature: 18-26°C,RH: 60-70%											
Powdery Mildew										Temperature: 22-27°C, Low humidity at day time and High humidity at night											
Weather warning																					
Rainfall (mm)	5-10mm (water logging)								Early rainfall (10-15mm)												
Cloudy weather	Cloudy condition																				
Droughth	Drought due to excessive dry and warm weather(Pointed Gourd Drought Tolerant)																				
Duration of staying wet	Land remain waterlogged for 2-3 days																				
Hailstorm										Early rainfall with hailstorm											
Windspeed (Km/hr.)	≥40 Km/hr																				
Temperature (°C)	Below 15°C									Above 35°C											