

Crop Weather Calendar of Rabi Tomato: Barisal Region (Districts: Patuakhali, Barisal, Jhalokhati, Pirozpur, Borghuna, Bhola) Bangladesh

Region: Barisal	Rabi Tomato												Life cycle of crop, 150-180							
Months	December				January				February				March				April			
Std.Week	৪৯	৫০	৫১	৫২	১	২	৩	৪	৫	৬	৭	৮	৯	১০	১১	১২	১৩	১৪	১৫	১৬
Rainfall (mm)	১.৭	৪.০	১.৩	০.৭	১.৩	১.০	১.০	১.৭	২.৭	৩.৭	৫.০	৮.০	৫.৭	৫.৩	৬.০	১৪.৩	১৮.০	১৮.০	২১.৭	২৫.০
Max. Temp. (°C)	২৭.৮	২৭.০	২৬.২	২৫.৭	২৫.১	২৫.২	২৫.৮	২৬.২	২৭.০	২৭.৮	২৮.৯	২৯.৭	৩০.৭	৩১.৭	৩২.৭	৩৩.০	৩২.৯	৩৩.৩	৩৩.৭	৩৩.৫
Min. Temp. (°C)	১৫.৬	১৪.৯	১৪.০	১৩.১	১২.৬	১২.৫	১২.৭	১২.৫	১৪.০	১৪.৬	১৬.২	১৬.৯	১৮.৩	১৯.৫	২০.৫	২২.১	২২.৮	২৩.৭	২৪.০	২৪.৬
Mean Temp. (°C)	২১.৭	২১.০	২০.১	১৯.৪	১৮.৯	১৮.৯	১৯.২	১৯.৪	২০.৫	২১.২	২২.৫	২৩.৩	২৪.৫	২৫.৬	২৬.৬	২৭.৬	২৭.৯	২৮.৫	২৮.৮	২৯.১
RHmax (%)	৯৮.০	৯৭.৯	৯৮.২	৯৮.২	৯৮.০	৯৭.৮	৯৭.৬	৯৭.৬	৯৭.৩	৯৭.১	৯৭.০	৯৭.০	৯৬.৭	৯৭.১	৯৬.৭	৯৬.৫	৯৬.৫	৯৬.৬	৯৬.৪	৯৬.০
RHMin (%)	৫০.৬	৫০.৯	৫০.৭	৫০.৩	৫০.১	৪৯.৩	৪৮.৫	৪৫.২	৪৬.২	৪৪.৮	৪৬.২	৪৫.১	৪৪.৯	৪৫.২	৪৫.৩	৫০.৭	৫৪.৪	৫৮.০	৫৯.০	৬১.৯
RHmean (%)	৭৪.৩	৭৪.৪	৭৪.৪	৭৪.৩	৭৪.১	৭৩.৬	৭৩.১	৭১.৪	৭১.৭	৭০.৯	৭১.৬	৭১.০	৭০.৮	৭১.২	৭১.০	৭৩.৬	৭৫.৪	৭৭.৩	৭৮.১	৭৮.৯
SShr (hrs)	৪৮.৭	৪৬.৩	৪৪.৩	৪৩.৭	৪৬.৭	৪৪.০	৪৭.৩	৫২.৭	৫০.৩	৫২.৩	৫১.৭	৫২.০	৫৩.৭	৫৪.০	৫৪.৩	৫২.৩	৫০.৭	৫৩.০	৫১.৩	৫৪.৭
WD (deg)	২২৫	২২৩	২১২	২১২	২০৯	২১১	২০৪	২১৬	২১৭	২১১	২১৪	২১১	২১৪	২২৬	২১৮	২১০	১৯৮	২০০০	১৯৮	১৯৭
WS(Km/hr)	৩.১	৩.৪	৩.৩	৩.৯	৪.৫	৪.৪	৪.৫	৪.৩	৫.০	৪.৬	৪.৯	৪.৯	৫.১	৫.২	৫.৪	৬.৩	৬.৭	৬.৯	৭.০	৭.৫
																				
Stages	Germination and Seedling Planting				Vegetative Growth				Flowering& Fruiting				Harvesting							
Favourable Weather																				
Temperature									18-25 °C											
Relative humidity									50-70%											
Soil temperature									20-25 °C											
Sunshine hour									6-8 hr											
Favourable weather for disease and pest infestation																				
White fly									15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly											
Tomato fruit borer									19.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %											
Leaf miner									Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C; Rainfall had positive and significant effect on leaf miner population											
Early blight									Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall											
Late blight									Temperature: 15-27°C, prolonged hot dry days increased disease infestation; relative humidity 75 to 90% or greater											
Fungal Wilting									Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%											
Weather warning																				
Rainfall (mm)									≥20 mm rainfall per day at the end of the rainy season (September-November)											
Cloudy weather									Cloudy condition											
Drought									Drought (moisture stress) due to excessive dry weather											
Duration of staying wet									Land remain waterlogged for 2 to 4 days											


Hailstorm	Late season rainfall with hailstorm	
Windspeed	≥ 15 km/hr	
Temperature (°C)	30 to 32°C	

Crop Weather Calendar of Robi Tomato: Dinajpur Region (Districts: Dinajpur, Thakurgaon, Panchagaor) Bangladesh

Dinajpur Region		Robi Tomato					Life span 120-130											
Months	November					December				January				February				
Std.Week	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	
Rainfall (mm)	5.0	3.0	0.0	0.0	0.0	0.0	3.0	0.0	4.0	2.0	2.0	3.0	2.0	1.0	2.0	3.0	1.0	
Max. Temp. (°C)	30.3	29.7	29.0	28.7	28.0	27.0	25.4	24.9	23.9	22.6	22.4	23.0	23.7	25.0	25.9	26.7	27.7	
Min. Temp. (°C)	19.4	18.3	16.9	15.6	14.0	13.3	12.9	12.0	10.9	10.7	10.3	10.0	10.6	11.1	12.1	13.6	14.1	
Mean Temp. (°C)	24.9	24.0	23.0	22.2	21.0	20.2	19.2	18.5	17.4	16.7	16.4	16.5	17.2	18.1	19.0	20.2	20.9	
RHmax (%)	96.0	95.9	95.9	95.3	95.6	96.1	96.4	96.3	96.6	96.7	96.3	96.4	95.9	94.7	94.0	93.4	92.0	
RHMin (%)	55.4	53.0	51.6	48.1	47.6	49.0	52.9	52.7	54.0	55.1	53.7	52.4	50.4	47.7	43.1	42.4	40.9	
RHmean (%)	75.7	74.5	73.7	71.7	71.6	72.5	74.6	74.5	75.3	75.9	75.0	74.4	73.1	71.2	68.5	67.9	66.4	
SShr (hrs)	55	56	52.0	54.0	53.0	48.0	42.0	41.0	37.0	34.0	34.0	37.0	41.0	44.0	46.0	52.0	54.0	
WD (deg)	149	144	154	164	177	183	193	200	213	225	238	243	227	241	225	230	225	
WS(Km/hr)	3.5	3.7	3.7	3.7	3.5	2.6	3.7	2.9	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	
																		
Stages	Germination and Seedling Planting					Vegetative Growth				Flowering & Fruiting				Harvesting				
Favourable Weather																		
Temperature	18-25 °C																	
Relative humidity	50-70%																	
Soil temperature	20-25 °C																	
Sunshine hour	6-8 hr																	
Favourable weather for disease and pest infestation																		
White fly	15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																	
Tomato fruit borer						1 9.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %												
Leaf miner						Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C ; Rainfall had positive and significant effect on leaf miner population												
Early blight						Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall												
Late blight						Temperature: 15-27°C, prolonged hot dry days increased disease infestation; relative humidity 75 to 90% or greater												
Fungal Wilting	Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%																	
Weather warning																		
Rainfall (mm)	≥20 mm rainfall per day at the end of the rainy season (September-November)																	
Cloudy weather	Cloudy condition																	
Drought										Drought (moisture stress) due to excessive dry weather								
Duration of staying wet	Land remain waterlogged for 2 to 4 days																	
Hailstorm	Late season rainfall with hailstorm																	
Windspeed	≥15 km/hr																	
Temperature (°C)	30 to 32°C																	

Dinajpur Region	KharifTomato					Life span 120-130											
Months	March					April				May				June			
Std.Week	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
Rainfall (mm)	1.0	1.0	4.0	2.0	7.0	8.0	8.0	25.0	23.0	31.0	36.0	57.0	55.0	73.0	57.0	77.0	111.0
Max. Temp. (°C)	28.6	30.1	31.1	32.0	32.1	33.0	33.4	33.0	32.4	32.3	32.9	33.0	33.0	33.4	33.3	33.0	32.3
Min. Temp. (°C)	15.0	16.1	17.1	18.6	19.7	20.4	21.1	22.0	22.0	22.3	22.9	23.6	24.0	24.4	25.0	25.1	25.6
Mean Temp. (°C)	21.8	23.1	24.1	25.3	25.9	26.7	17.3	27.5	27.2	27.3	27.9	28.3	28.5	28.9	29.2	29.1	29.0
RHmax (%)	90.6	88.9	87.6	86.0	87.1	86.6	87.4	89.4	91.1	92.3	92.0	92.1	92.4	92.7	93.4	93.9	94.7
RHMin (%)	38.4	35.3	35.1	36.4	43.3	40.9	43.4	50.9	53.7	57.6	57.3	59.3	62.4	62.3	65.4	67.7	71.4
RHmean (%)	64.5	62.1	61.4	61.2	65.2	63.8	65.4	70.2	72.4	75.0	74.7	75.7	77.4	77.5	79.4	80.8	83.1
SShr (hrs)	57.0	59.0	59.0	55.0	49.0	52.0	55.0	49.0	50.0	49.0	51.0	48.0	44.0	42.0	40.0	35.0	27.0
WD (deg)	228	230	212	192	170	171	150	142	123	118	115	119	108	117	111	118	120
WS(Km/hr)	3.8	4.2	5.4	5.6	5.0	5.6	4.8	4.8	5.4	3.9	3.7	4.3	3.7	3.9	3.7	3.7	3.7
Stages	Germination and Seedling Planting				Vegetative Growth				Flowering & Fruiting			Harvesting					
Favourable Weather																	
Temperature	26-32°C																
Relative humidity	50-80%																
Soil temperature	22-26 °C																
Sunshine hour	8-10 hr																
Favourable weather for disease and pest infestation																	
White fly	15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																
Tomato fruit borer										1 9.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %							
Leaf miner						Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C; Rainfall had positive and significant effect on leaf miner population											
Early blight						Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall											
Late blight						Temperature: 15-27°C, prolonged hot dry days increased disease infestation; relative humidity 75 to 90% or greater											
Fungal Wilting	Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%																
Weather warning																	
Rainfall (mm)	≥20 mm rainfall per day at the end of the rainy season (September-November)																
Cloudy weather	Cloudy condition																
Drought										Drought (moisture stress) due to excessive dry weather							
Duration of staying wet	Land remain waterlogged for 2 to 4 days																
Hailstorm	Late season rainfall with hailstorm																
Windspeed	≥15 km/hr																
Temperature (°C)	30 to 32°C																

Crop Weather Calendar of Rabi Tomato: Jessore Region (Districts: Jessore, Chuadanga, Meherpur, Jinaidaha, magura, Kustia) Bangladesh

Region: Jessore	Rabi Tomato										Life cycle of crop, 140-150											
Months	November					December				January				February				March				
Std.Week	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13
Rainfall (mm)	9.0	8.0	7.5	1.0	2.0	1.5	4.0	0.0	3.5	0.5	1.5	2.5	3.0	6.5	5.0	4.5	8.0	4.0	2.5	4.5	11.5	12.0
Max. Temp. (°C)	31.1	31.1	30.3	29.6	28.5	27.7	26.6	25.7	25.2	24.5	24.5	25.1	25.6	26.6	27.8	28.9	30.0	31.0	32.4	33.6	34.7	34.6
Min. Temp. (°C)	20.7	19.4	18.0	16.7	15.1	14.0	13.4	12.3	11.4	10.9	10.7	11.0	10.8	12.0	13.3	14.7	15.6	16.4	17.7	19.0	20.8	21.8
Mean Temp. (°C)	25.9	25.3	24.2	23.1	21.8	20.9	20.0	19.0	18.3	17.7	17.6	18.0	18.2	19.3	20.5	21.8	22.8	23.7	25.0	26.3	27.7	28.2
RHmax (%)	95.9	95.9	95.7	95.7	95.4	95.2	95.8	95.5	95.7	95.4	95.3	95.3	94.9	94.9	94.2	94.0	93.7	93.2	92.5	92.1	92.1	92.4
RHMin (%)	56.2	52.0	49.8	48.1	46.3	47.2	49.2	49.4	47.9	49.6	47.9	47.0	44.7	44.2	42.2	41.0	39.3	36.5	35.4	34.9	36.7	42.0
RHmean (%)	76.0	74.0	72.8	71.9	70.8	71.2	72.5	72.5	71.8	72.5	71.6	71.2	69.8	69.5	68.2	67.5	66.5	64.9	63.9	63.9	64.4	67.2
SShr (hrs)	45.5	48.5	49.5	47.5	48.5	47.5	43.5	42.0	42.0	41.0	42.5	44.5	48.0	48.5	49.5	50.0	54.5	55.0	54.0	55.0	53.0	51.5
WD (deg)	153	175	177	179	178	184	183	177	174	158	156	188	192	186	191	203	214	219	213	230	212	208
WS(Km/hr)	2.5	2.3	2.4	2.3	2.4	2.3	2.4	2.5	2.5	3.0	3.0	3.0	3.0	3.2	3.1	3.3	3.7	3.5	3.9	4.1	4.4	4.7
																						
Stages	Germination and Seedling Planting					Vegetative Growth					Flowering & Fruiting					Harvesting						
Favorable environment																						
Temperature											18 to 25 °C											
Relative Humidity											50 to 70%											
Soil Temperature											20 to 25 °C											
Sunshine Hour											6 to 8 hr											
Favourable weather for disease and pest infestation																						
White fly	15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																					
Tomato fruit borer											19.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %											
Leaf miner						Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C; Rainfall had positive and significant effect on leaf miner population																
Early blight						Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall																
Late blight						Temperature: 15-27°C, prolonged hot dry days increased disease infestation; relative humidity 75 to 90% or greater																
Fungal Wilting	Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%																					
Weather warning																						
Rainfall (mm)	≥20 mm rainfall per day at the end of the rainy season (September-November)																					
Cloudy weather	Cloudy condition																					
Drought											Drought (moisture stress) due to excessive dry weather											
Duration of staying wet	Land remain waterlogged for 2 to 4 days																					
Hailstorm	Late season rainfall with hailstorm																					
Windspeed	≥15 km/hr																					
Temperature (°C)	30 to 32°C																					

Crop Weather Calendar of Kharif Tomato: Jessore Region (Districts: Jessore, Chuadanga, Meherpur, Jinaidaha, magura, Kustia) Bangladesh

Region: Jessore		Kharif Tomato								Life cycle of crop, 140-150									
Months		March					April				May				June				
Std.Week		9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26
Rainfall (mm)		4.0	2.5	4.5	11.5	12.0	11.5	7.0	12.5	21.5	28.5	30.5	36.0	44.5	38.5	49.5	54.5	81.0	72.5
Max. Temp. (°C)		31.0	32.4	33.6	34.7	34.6	35.5	36.1	36.5	36.3	36.0	36.0	35.9	35.5	35.6	35.3	34.0	33.3	33.0
Min. Temp. (°C)		16.4	17.7	19.0	20.8	21.8	22.8	23.5	24.1	24.2	24.2	25.2	25.0	25.5	26.0	26.2	26.2	26.1	26.2
Mean Temp. (°C)		23.7	25.0	26.3	27.7	28.2	29.2	29.8	30.3	30.3	30.1	30.6	30.4	30.5	30.8	30.7	30.1	29.7	29.6
RHmax (%)		93.2	92.5	92.1	92.1	92.4	92.4	92.9	92.4	92.3	92.8	92.7	93.1	93.6	94.0	94.1	94.8	95.7	96.0
RHMin (%)		36.5	35.4	34.9	36.7	42.0	41.5	41.6	44.3	46.9	49.9	51.7	54.8	58.7	60.0	62.1	67.5	71.7	73.5
RHmean (%)		64.9	63.9	63.9	64.4	67.2	66.9	67.2	68.3	69.6	71.3	72.2	74.0	76.1	77.0	78.1	81.1	83.7	84.7
SShr (hrs)		55.0	54.0	55.0	53.0	51.5	55.5	54.5	54.5	55.5	54.0	51.5	52.5	46.5	45.5	41.5	32.0	26.5	26.0
WD (deg)		219	213	230	212	208	209	204	193	188	186	186	180	181	180	183	174	175	178
WS(Km/hr)		3.5	3.9	4.1	4.4	4.7	5.1	5.2	5.7	5.6	5.6	5.8	5.2	5.2	5.0	4.6	4.9	4.6	4.6
																			
Stages		Germination and Seedling Planting					Vegetative Growth				Flowering & Fruiting				Harvesting				
Favourable Weather																			
Temperature		18-25 °C																	
Relative humidity		50-70%																	
Soil temperature		20-25 °C																	
Sunshine hour		6-8 hr																	
Favourable weather for disease and pest infestation																			
White fly		15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																	
Tomato fruit borer											19.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %								
Leaf miner							Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C; Rainfall had positive and significant effect on leaf miner population												
Early blight							Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall												
Late blight							Temperature: 15-27°C, prolonged hot dry days increased disease infestation; relative humidity 75 to 90% or greater												
Fungal Wilting		Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%																	
Weather warning																			
Rainfall (mm)		≥20 mm rainfall per day at the end of the rainy season (September-November)																	
Cloudy weather		Cloudy condition																	
Drought											Drought (moisture stress) due to excessive dry weather								
Duration of staying wet		Land remain waterlogged for 2 to 4 days																	
Hailstorm		Late season rainfall with hailstorm																	
Windspeed		≥15 km/hr																	

Temperature (°C)

30 to 32°C

Crop Weather Calendar of Rabi Tomato: Khulna Region (Districts: Khulna, Norail, Bagerhat & sathkhira) Bangladesh

Crop Weather Calendar of RabiTomato, Panchajanya Region (District: Panchajanya, Nepal; Elevation: 1000m; Soil: Reddish Brown)					Life cycle of crop, 140-150																	
RabiTomato					Life cycle of crop, 140-150																	
October					November					December				January				February				
Months	40	41	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	
Std.Week	40	41	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	
Rainfall (mm)	41.0	38.5	31.5	26.5	9.5	13.0	8.0	3.0	3.5	1.0	4.0	0.0	1.0	3.0	1.5	0.0	3.5	5.5	6.5	5.0	13.5	
Max. Temp. (°C)	32.7	32.7	32.4	31.7	31.0	31.0	30.1	29.5	28.5	27.7	27.0	26.0	25.6	25.0	25.0	25.8	26.2	27.0	27.7	29.0	30.1	
Min. Temp. (°C)	25.5	25.0	24.2	22.5	21.5	20.6	19.3	18.0	16.3	15.3	14.5	13.4	12.6	12.2	12.2	12.7	12.4	13.9	14.5	16.2	17.0	
Mean Temp. (°C)	29.1	28.9	28.3	27.1	26.3	25.8	24.7	23.7	22.4	21.5	20.7	19.7	19.1	18.6	18.6	19.2	19.3	20.4	21.1	22.6	23.6	
RHmax (%)	96.8	96.5	96.2	96.3	96.0	95.6	95.9	95.8	95.5	94.7	95.7	95.8	95.7	95.4	95.2	94.9	94.9	95.1	94.5	94.3	94.3	
RHMin (%)	69.2	65.6	63.3	59.2	57.4	53.9	51.3	49.0	46.7	47.3	48.8	49.2	47.8	48.7	46.9	46.8	44.3	44.9	43.4	43.9	41.8	
RHmean (%)	83.0	81.1	79.8	77.8	76.7	74.7	73.6	72.4	71.1	71.0	72.2	72.5	71.7	72.0	71.0	70.8	69.6	70.0	69.0	69.1	68.0	
SShr (hrs)	41.5	47.5	49.5	52.5	48.5	53.5	52.5	53.5	53.0	52.0	49.5	48.0	48.5	47.0	46.5	49.5	53.0	52.5	53.5	55.5	55.5	
WD (deg)	189	203	193	184	173	176	189	197	197	194	189	181	177	175	182	198	198	190	193	209	218	
WS(Km/hr)	4.3	3.7	4.1	3.8	3.4	4.1	3.8	4.3	4.3	4.0	4.5	4.1	.6	4.6	4.7	4.3	4.6	4.9	5.0	4.8	5.0	
																						
Stages	Germination and Seedling Planting				Vegetative Growth					Flowering & Fruiting				Harvesting								
Favourable Weather																						
Temperature											18-25 °C											
Relative humidity											50-70%											
Soil temperature											20-25 °C											
Sunshine hour											6-8 hr											
Favourable weather for disease and pest infestation																						
White fly	15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																					
Tomato fruit borer											19.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %											
Leaf miner	Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C ; Rainfall had positive and significant effect on leaf miner population																					
Early blight	Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall																					
Late blight						Temperature: 15-27°C, prolonged hot dry days increased disease infestation; relative humidity 75 to 90% or greater																
Fungal Wilting	Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%																					
Weather warning																						
Rainfall (mm)	≥20 mm rainfall per day at the end of the rainy season (September-November)																					
Cloudy weather	Cloudy condition																					
Drought											Drought (moisture stress) due to excessive dry weather											
Duration of staying wet	Land remain waterlogged for 2 to 4 days																					
Hailstorm	Late season rainfall with hailstorm																					
Windspeed											≥15 km/hr											
Temperature (°C)											30 to 32°C											

Khulna Region	RabiTomato					V												
Months	March					April				May					June			
Std. Week	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26
Rainfall (mm)	6.5	2.5	6.5	13.5	16.0	12.5	12.5	17.5	25.0	31.5	32.5	37.0	47.5	50.0	52.0	77.5	79.0	93.5
Max. Temp. (°C)	31.0	32.4	33.4	34.1	34.0	35.0	35.2	35.2	35.2	35.1	35.2	35.3	35.1	35.0	34.7	33.4	32.7	32.3
Min. Temp. (°C)	18.1	19.4	20.4	22.0	22.7	24.0	24.3	24.9	25.0	25.0	25.3	25.8	26.0	26.4	26.8	26.2	26.3	26.1
Mean Temp. (°C)	24.5	25.9	26.9	28.1	28.4	29.5	29.8	30.1	30.1	30.0	30.2	30.6	30.5	30.7	30.7	29.8	29.3	29.2
RHmax (%)	93.7	93.7	93.5	94.5	94.1	94.3	93.7	93.3	93.2	93.0	93.1	93.4	93.7	93.9	94.2	95.1	95.7	95.8
RHMin (%)	40.7	40.7	40.2	44.2	47.3	48.8	49.2	52.4	53.7	55.3	57.4	57.9	60.7	62.0	64.9	70.2	73.3	75.1
RHmean (%)	67.2	67.2	66.8	69.3	70.7	71.5	71.4	72.8	73.4	74.1	75.2	75.7	77.2	77.9	79.5	82.6	84.5	85.4
SShr (hrs)	57.5	57.5	59.0	57.0	56.5	59.0	60.0	61.5	58.0	59.5	56.5	55.0	47.0	50.0	41.5	29.5	23.5	25.0
WD (deg)	222	227	233	204	203	202	196	192	186	186	183	177	180	184	178	170	175	179
WS(Km/hr)	4.9	5.2	5.6	6.3	7.2	7.1	7.6	7.6	8.0	7.5	7.5	7.1	7.2	7.3	6.7	6.8	6.8	6.8
																		
Stages	Germination and Seedling Planting				Vegetative Growth				Flowering & Fruiting				Harvesting					
Favourable Weather																		
Temperature	26-32°C																	
Relative humidity	50-80%																	
Soil temperature	22-26 °C																	
Sunshine hour	8-10 hr																	
Favourable weather for disease and pest infestation																		
White fly	15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																	
Tomato fruit borer											1 9.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %							
Leaf miner						Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C; Rainfall had positive and significant effect on leaf miner population												
Early blight						Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall												
Late blight						Temperature: 15-27°C, prolonged hot dry days increased disease infestation; relative humidity 75 to 90% or greater												
Fungal Wilting	Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%																	
Weather warning																		
Rainfall (mm)	≥20 mm rainfall per day at the end of the rainy season (September-November)																	
Cloudy weather	Cloudy condition																	
Drought											Drought (moisture stress) due to excessive dry weather							
Duration of staying wet	Land remain waterlogged for 2 to 4 days																	
Hailstorm	Late season rainfall with hailstorm																	

Windspeed	≥ 15 km/hr
Temperature (°C)	30 to 32°C

Crop Weather Calendar of *Rabi* seasonTomato: Chittagong region (Districts: Cox's Bazar, Chittagong, Feni, Noakhali, Laksmipur)

Region: Chittagong	Tomato-Rabi season													Duration: 120-150 days							
Month	September				October				November					December				January			
Std.Week/Normal	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	01	02	03	04
Rainfall (mm)	92.7	73.7	73.7	60.3	66.7	75.7	43.7	26	19.3	21.7	14.3	9.0	11.7	10	4	2.3	0.2	0.3	0.7	0.7	1.3
Max. Temp. (°C)	31.4	31.9	31.9	32.1	32.2	31.1	32.2	31.9	31.5	31.2	30.5	29.9	29	28.6	27.8	26.9	26.6	26.5	26.1	26.6	26.9
Min. Temp. (°C)	25.2	25.1	25.2	25.1	25	24.8	24.5	23.3	22.8	21.6	20.2	19.2	18	17	16.3	15.3	14.9	14.1	13.9	14.1	14.1
Mean. Temp. (°C)	28.3	28.5	28.5	28.6	28.6	28.3	28.3	27.6	27.1	26.4	25.3	24.6	23.5	22.8	22.1	21.1	20.8	20.3	20	20.4	20.5
RHmax (%)	95.6	95.7	95.7	95.7	96	96.1	95.8	95	94.9	94.7	94.4	94.1	94.5	94.4	94.2	94.4	94.1	93.8	93.1	92.9	91.6
RHmin (%)	73.4	72.2	71.5	70.2	69.9	68.8	66.7	62.2	62.2	58.1	55.4	52	50.8	49.5	50	50.3	49.1	48.1	47.1	44.9	42.3
RHmean (%)	84.5	83.9	83.6	82.9	82.9	82.4	81.2	78.6	78.6	76.9	74.9	73.1	72.6	71.9	72.1	72.3	71.6	71	70.1	68.9	67
SShr (hrs)	35.4	39.3	39.5	42.5	42.8	45.8	49.5	53.3	52.3	54	55.5	56.5	55.5	54.8	52.8	50.8	51.3	53.3	50.8	53.8	56.6
WD (degree)	185	190	194	202	209	203	215	201	196	195	199	203	196	204	196	199	207	200	193	203	203
WS(Km/hr)	8.6	7.6	7.6	7.4	6.8	6.7	6.9	6	6	5.9	5.6	6.1	5.9	5.1	5.7	5.8	6.2	6.6	6.9	6.8	7
																					
Stages	Spouting & Seedling				Vegetative growth				Flowering and Fruiting					Harvesting							
Favourable Weather																					
Temperature	18-25 °C																				
Relative humidity	50-70%																				
Soil temperature	20-25 °C																				
Sunshine hour	6-8 hr																				
Favourable weather for disease and pest infestation																					
White fly	15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																				
Tomato fruit borer									1 9.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %												
Leaf miner									Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C; Rainfall had positive and significant effect on leaf miner population												
Early blight					Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall																
Late blight					Temperature: 15-27°C, prolonged hot dry days increased disease infestation; relative humidity 75 to 90% or greater																
Fungal Wilting	Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%																				
Weather warning																					
Rainfall (mm)	≥20 mm rainfall per day at the end of the rainy season (September-November)																				
Cloudy weather	Cloudy condition																				
Drought									Drought (moisture stress) due to excessive dry weather												
Duration of staying wet	Land remain waterlogged for 2 to 4 days																				
Hailstorm	Late season rainfall with hailstorm																				
Windspeed	≥15 km/hr																				
Temperature (°C)	30 to 32°C																				

Crop Weather Calendar of *Rabi* season Tomato: Cumilla region (Districts: Cumilla, Chandpur, Brahmanbaria)

Region: Cumilla	Tomato-Rabi sesaon													Duration: 120-150 days							
Month	September				October				November					December				January			
Std.Week/Normal	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	01	02	03	04
Rainfall (mm)	53	66.5	68	53	51	46.5	30.5	16	11.5	18	8.5	4	2.5	0	4.5	1	1.5	0	0	0	1.5
Max. Temp. (°C)	32.8	32	32	32	32.3	31.9	31.8	31.2	31	30.5	29.8	29	28.2	27.7	26.5	25.8	25.4	24.7	24.5	25	25.4
Min. Temp. (°C)	25.7	25.6	25.6	25.6	25.4	24.7	24.3	22.8	21.8	21	19.3	18.1	16.9	15.8	15	14.2	13.8	13	12.7	12.7	12.6
Mean. Temp. (°C)	28.8	28.8	28.8	28.8	28.8	28.3	28	27	26.4	25.7	24.6	23.5	22.5	21.7	20.8	20	19.9	18.8	18.6	18.9	19
RHmax (%)	95.2	95.4	95.4	95.6	95.6	95.5	95.5	95.2	95.3	95.6	95.3	95.4	95.8	95.8	96.2	96	96	96	95.5	95.3	94.4
RHmin (%)	73.5	73.1	72.2	71.1	70.2	68.1	65.8	61.9	60.6	58.2	54.8	5.6	52.2	50.6	51.9	52.1	51.3	51.8	51.2	49.1	46.4
RHmean (%)	84.3	84.3	83.8	83.3	82.9	81.9	80.6	78.6	77.9	76.9	75	74	74	73.2	74.1	74.1	73.6	73.9	73.4	72.2	70.4
SShr (hrs)	32	33	33.5	35	39	43	44.5	50.5	47	49.5	50	50.5	48.5	47.5	42.5	40.5	40	40	37	43.5	47
WD (degree)	186	191	197	201	203	214	201	187	202	183	184	190	195	206	192	211	213	205	209	218	212
WS(Km/hr)	5.3	5.7	4.7	4.7	4.6	4.3	4.3	4	3.5	3.7	3.7	3.5	3.3	3.3	4	3.5	3.8	4	4.2	4.6	4.4
																					
Stages	Spouting & Seedling				Vegetative growth				Flowering and Fruting					Harvesting							
Favourable Weather																					
Temperature									18-25 °C												
Relative humidity									50-70%												
Soil temperature									20-25 °C												
Sunshine hour									6-8 hr												
Favourable weather for disease and pest infestation																					
White fly									15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly												
Tomato fruit borer									1 9.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %												
Leaf miner									Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C ; Rainfall had positive and significant effect on leaf miner population												
Early blight									Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall												
Late blight									Temperature: 15-27°C, prolonged hot dry days increased disease infestation; relative humidity 75 to 90% or greater												
Fungal Wilting									Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%												
Weather warning																					
Rainfall (mm)									≥20 mm rainfall at the end of the rainy season (September- January)												
Cloudy weather									Cloudy condition												
Drought									Drought (moisture stress) due to excessive dry weather												
Duration of staying wet									Land remain waterlogged for 2 to 4 days												
Hailstorm									Late season rainfall with hailstorm												
Windspeed									≥15 km/hr												
Temperature (°C)									30-32°C												

Crop Weather Calendar of *Rabi* season Tomato: Dhaka region (Districts: Narayanganj, Gazipur, Narsinghdi, Munsiganj, Manikganj, Tangail)

Region: Dhaka					Tomato-Rabi season														Duration: 120-150 days							
Month	September				October				November					December					January				February			
Std. Week/Normal	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	01	02	03	04	05	06	07	08	
Rainfall (mm)	60.5	68.5	70	69	47	42	40.5	26	12.5	10	6	1.5	4.5	1.5	4.5	1.5	3.5	1	0	0	0.5	3.5	3.5	2.5	12	
Max. Temp. (°C)	32.3	32.2	32.3	32.3	32.5	32.3	32	31.5	30.9	30.6	29.8	29	28.3	27.6	26.2	25.5	24.8	24	23.9	24.5	25	26.2	27.2	28.3	28.9	
Min. Temp. (°C)	26	26	26	25.7	25.3	24.8	23.8	22.3	21.5	20.5	18.7	17.7	16.1	15.5	14.7	13.5	12.9	12.4	12	12	12.1	13.6	14.3	15.5	16.5	
Mean. Temp. (°C)	29.2	29.1	29.1	29	28.9	28.5	27.9	26.9	26.2	25.5	24.3	23.3	22.2	21.5	20.5	19.5	18.8	18.2	17.9	18.3	18.6	19.9	20.7	21.9	22.7	
RHmax (%)	94.8	95.2	95	95.5	95.4	95.4	95	94.8	94.5	95	94.7	94.6	94.3	94.8	95.3	95.1	95	94.6	94.3	93.8	93.4	92.3	92.1	91.7	91.1	
RHmin (%)	70.2	69.4	69.7	68	65.7	63	60.2	55.5	54.6	51.5	49.3	47.5	46.3	47.6	49.4	49.1	49.1	50.3	48.8	46.2	43.3	42.3	39.4	39.7	38.9	
RHmean (%)	82.5	82.3	82.3	81.8	80.6	79.6	77.6	75.2	74.4	73.2	72	71	70.3	71.2	72.3	72.5	72.1	72.4	71.6	70	68.3	67.3	65.7	65.7	65	
SShr (hrs)	33.5	32	32	34	40	44	47.5	54	49.5	51.5	51.5	52	51.5	48.6	46.5	44.5	44.4	42.5	42.5	45	49	49.6	51.5	53	54.5	
WD (degree)	163	177	176	195	194	১৯৮	192	190	190	189	199	212	211	219	224	225	233	238	238	232	242	241	236	234	228	
WS(Km/hr)	5.5	5.3	4.3	4.6	4	3.9	4.1	3.8	3.7	3.7	3.7	3.7	3.6	3.7	3.7	3.7	3.7	4.1	4	4.1	4.3	4.3	4.5	5	5.3	
																										
Stages	Spouting & Seedling				Vegetative growth				Flowering and Fruting								Harvesting									
Favourable Weather																										
Temperature									18-25 °C																	
Relative humidity									50-70%																	
Soil temperature									20-25 °C																	
Sunshine hour									6-8 hr																	
Favourable weather for disease and pest infestation																										
White fly	15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																									
Tomato fruit borer									1 9.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %																	
Leaf miner					Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C ; Rainfall had positive and significant effect on leaf miner population																					
Early blight									Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall																	
Late blight									Temperature: 15-27°C, prolonged hot dry days increased disease infestation; relative humidity 75 to 90% or greater																	
Wilting	Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%																									
Weather warning																										
Rainfall (mm)					≥20 mm at the end of the rainy season (September- February)																					
Cloudy weather									Cloudy condition																	
Drought									Drought (moisture stress) due to excessive dry weather																	
Duration of staying wet					Land remain waterlogged for 2 to 4 days																					
Hailstorm					Late season rainfall with hailstorm																					
Windspeed									≥15 km/hr																	
Temperature (°C)									30 to 32°C																	

Crop Weather Calendar of *Kharif* season Tomato: Dhaka region (Districts: Narayanganj, Gazipur, Narsinghdi, Munsiganj, Manikganj, Tangail)

Region: Mymensingh	Tomato-Kharif season													Duration: 120-150 days							
Month	February				March				April				May					June			
Std.Week/Normal	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
Rainfall (mm)	3.5	3.5	2.5	12.0	3.5	4.0	9.5	13.5	20.5	25.0	21.0	31.5	42.5	46.0	55.5	55.5	75.5	70.0	72.5	86.5	81.0
Max. Temp. (°C)	26.2	27.2	28.3	28.9	30.0	31.5	32.3	33.4	33.2	33.9	24.6	34.1	33.8	33.6	33.7	33.7	33.2	33.5	33.4	32.8	32.2
Min. Temp. (°C)	13.6	14.3	15.5	16.5	17.5	18.6	19.8	19.8	21.8	22.9	23.5	23.7	23.6	23.7	24.2	24.6	24.9	25.4	25.8	25.9	26.2
Mean. Temp. (°C)	19.9	20.7	21.9	22.7	23.7	25.0	26.1	26.1	27.5	28.4	29.0	28.9	28.7	28.6	28.9	29.2	29.0	29.4	29.6	29.3	29.2
RHmax (%)	92.4	92.1	91.7	91.1	90.0	89.9	89.2	89.6	90.3	91.3	91.4	91.4	91.7	92.3	92.8	92.9	93.5	93.8	94.0	94.5	94.4
RHmin (%)	42.3	39.4	39.7	38.9	36.0	35.5	35.9	38.9	44.5	45.9	47.1	52.0	55.1	56.7	58.6	60.4	63.5	65.0	66.7	69.9	71.5
RHmean (%)	67.3	65.7	65.7	65.0	63.0	62.7	62.5	64.3	67.4	68.6	69.2	71.7	73.4	74.5	75.7	76.7	78.5	79.4	80.3	82.2	83.0
SShr (hrs)	49.5	51.5	53.0	54.5	56.0	55.0	58.0	55.5	52.5	53.5	54.0	55.0	54.5	51.5	48.0	48.0	42.5	42.0	38.5	31.0	26.0
WD (degree)	241	236	234	228	234	230	223	206	188	181	175	167	156	159	164	156	155	157	161	157	157
WS(Km/hr)	4.3	4.5	5.0	5.3	5.0	5.4	5.7	6.2	6.5	6.5	6.6	6.6	6.7	6.8	6.5	6.4	6.1	5.8	5.6	6.3	6.5
																					
Stages	Spouting & Seedling				Vegetative growth				Flowering and Fruting						Harvesting						
Favourable Weather																					
Temperature	26-32 °C																				
Relative humidity	50-80%																				
Soil temperature	22-26 °C																				
Sunshine hour	8-10 hr																				
Favourable weather for disease and pest infestation																					
White fly	15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																				
Tomato fruit borer					19.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %																
Leaf miner					Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C ; Rainfall had positive and significant effect on leaf miner population																
Early blight					Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall																
Fungal Wilting	Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%																				
Weather warning																					
Rainfall (mm)	≥20 mm rainfall per day at the end of the rainy season (February - June)																				
Cloudy weather	Cloudy condition																				
Drought											Drought (moisture stress) due to excessive dry weather										
Duration of staying wet	Land remain waterlogged for 2 to 4 days																				
Hailstorm	Late season rainfall with hailstorm																				
Windspeed											≥15 km/hr										
Temperature (°C)											30 to 32°C										

Crop Weather Calendar of *Rabi* season Tomato: Mymensingh region (Districts: Mymensingh, Netrokona, Jamalpur, Sherpur)

Region: Mymensingh	Tomato-Rabi season													Duration: 120-150 days							
Month	September				October				November					December				January			
Std.Week/Normal	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	01	02	03	04
Rainfall (mm)	63.0	65.0	79.0	87.0	71.0	52.0	45.0	15.0	7.0	5.0	5.0	0.0	4.0	1.0	3.0	0.0	2.0	0.0	3.0	1.0	0.0
Max. Temp. (°C)	31.4	31.7	31.9	31.7	32.0	31.7	31.6	31.1	30.9	30.3	30.0	28.9	28.3	27.6	26.3	25.4	25.0	24.4	24.0	24.0	25.1
Min. Temp. (°C)	26.0	26.0	26.0	25.1	25.0	24.3	23.4	21.7	20.9	19.7	28.3	17.0	15.4	15.0	14.0	13.3	12.4	12.0	11.6	11.9	12.0
Mean. Temp. (°C)	28.7	28.9	29.0	28.4	28.5	28.0	27.5	26.4	25.9	25.0	24.2	23.0	21.9	21.3	20.2	19.4	18.7	18.2	17.8	18.0	18.6
RHmax (%)	94.9	95.3	95.0	95.4	95.6	95.3	95.3	95.4	95.0	95.0	94.6	95.0	95.1	95.3	95.6	95.6	95.3	95.1	94.4	94.4	93.4
RHmin (%)	75.9	74.1	74.1	74.0	72.6	70.3	66.7	62.7	61.0	58.4	55.9	53.0	51.6	52.3	54.1	53.6	53.7	53.6	53.1	51.4	47.0
RHmean (%)	85.4	84.7	84.6	84.7	84.1	82.8	81.0	79.1	78.1	76.7	75.3	74.0	73.4	73.8	74.9	74.9	74.5	74.4	73.8	72.9	70.2
SShr (hrs)	28.0	30.0	31.0	31.0	40.0	41.0	47.0	53.0	51.0	53.0	52.0	53.0	50.0	49.0	43.0	40.0	42.0	41.0	39.0	41.0	45.0
WD (degree)	151	157	156	167	181	174	193	197	186	181	186	186	193	191	206	199	221	230	236	216	238
WS(Km/hr)	5.6	5.6	5.4	4.4	4.3	3.9	3.7	3.7	3.7	3.9	3.7	3.7	3.7	3.0	3.5	3.7	3.7	3.7	3.7	3.7	3.7
																					
Stages	Spouting & Seedling				Vegetative growth				Flowering and Fruting					Harvesting							
Favourable Weather																					
Temperature	18-25 °C																				
Relative humidity	50-70%																				
Soil temperature	20-25 °C																				
Sunshine hour	6-8 hr																				
Favourable weather for disease and pest infestation																					
White fly	15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																				
Tomato fruit borer									19.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %												
Leaf miner					Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C ; Rainfall had positive and significant effect on leaf miner population																
Early blight					Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall																
late blight					Temperature: 15-27°C, prolonged hot dry days increased disease infestation; relative humidity 75 to 90% or greater																
Fungal Wilting	Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%																				
Weather warning																					
Rainfall (mm)	≥20 mm rainfall at the end of the rainy season (September-January)																				
Cloudy weather	Cloudy condition																				
Drought									Drought due to excessive dry weather												
Duration of staying wet	Land remain waterlogged for 2 to 4 days																				
Hailstorm	Late season rainfall with hailstorm																				
Windspeed	≥15 km/hr																				
Temperature (°C)	30 to 32°C																				

Crop Weather Calendar of *Kharif*season Tomato: Mymensingh region (Districts: Mymensingh, Netrokona, Jamalpur, Sherpur)

Region: Mymensingh	Tomato-Kharif season													Duration: 120-150 days							
Month	February				March					April				May					June		
Std.Week/Normal	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
Rainfall (mm)	2.0	1.0	5.0	7.0	1.0	4.0	6.0	11.0	17.0	20.0	30.0	40.0	36.0	55.0	62.0	69.0	97.0	81.0	96.0	103	98.0
Max. Temp. (°C)	25.9	26.6	27.4	28.1	29.1	30.3	31.0	31.9	31.3	32.1	32.7	32.0	31.7	31.6	31.9	32.3	32.0	32.0	32.0	31.7	31.4
Min. Temp. (°C)	13.0	13.7	15.0	15.9	16.6	17.4	18.6	19.9	20.9	21.9	22.4	22.9	22.9	22.9	23.6	24.0	24.3	25.0	25.6	25.7	26.0
Mean. Temp. (°C)	19.5	20.2	21.2	22.0	22.9	23.9	24.8	25.9	26.1	27.0	27.6	27.5	27.3	27.3	27.8	28.2	28.2	28.5	28.8	28.7	28.7
RHmax (%)	92.6	92.1	91.4	90.6	90.6	90.6	90.1	89.7	90.4	89.9	90.9	91.1	91.3	92.4	92.6	92.7	92.7	94.0	94.1	94.7	95.0
RHmin (%)	47.9	45.3	45.0	45.1	44.3	43.6	44.6	48.3	54.7	55.0	56.0	61.1	62.6	64.6	66.0	66.4	69.4	71.9	73.0	75.4	76.6
RHmean (%)	70.3	68.7	68.2	67.9	67.6	67.1	67.4	69.0	72.6	72.5	73.8	76.1	77.0	78.5	79.3	79.6	81.1	83.3	83.6	85.1	85.8
SShr (hrs)	44.0	48.0	50.0	50.0	51.0	53.0	53.0	48.0	44.0	49.0	49.0	49.0	46.0	44.0	44.0	45.0	39.0	36.0	44.0	26.0	25.0
WD (degree)	228	218	222	210	214	202	208	179	165	149	146	143	128	126	134	131	126	130	142	138	143
WS(Km/hr)	3.7	3.9	4.4	5.4	5.0	5.4	5.6	6.3	6.1	7.2	6.8	7.2	7.6	7.4	6.8	7.4	7.4	6.8	6.1	7.4	6.8
																					
Stages	Spouting & Seedling				Vegetative growth					Flowering and Fruiting					Harvesting						
Favourable Weather																					
Temperature		18-25 °C																			
Relative humidity		50-70%																			
Soil temperature		20-25 °C																			
Sunshine hour		6-8 hr																			
Favourable weather for disease and pest infestation																					
White fly		15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																			
Tomato fruit borer												19.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %									
Leaf miner												Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C ; Rainfall had positive and significant effect on leaf miner population									
Early blight												Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall									
late blight												Temperature: 15-27°C, prolonged hot dry days increased disease infestation; relative humidity 75 to 90% or greater									
Fungal Wilting		Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%																			
Weather warning																					
Rainfall (mm)		≥20 mm rainfall at the end of the rainy season (September-June)																			
Cloudy weather		Cloudy condition																			
Drought												Drought (moisture stress) due to excessive dry weather									
Duration of staying wet		Land remain waterlogged for 2 to 4 days																			

Hailstorm	Late season rainfall with hailstorm			
Windspeed		≥15 km/hr		
Temperature (°C)		30 to 32°C		

Region: Mymensingh	Tomato-Kharif season													Duration: 120-150 days								
Month	February				March					April				May					June			
Std.Week/Normal	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	
Rainfall (mm)	2.0	1.0	5.0	7.0	1.0	4.0	6.0	11.0	17.0	20.0	30.0	40.0	36.0	55.0	62.0	69.0	97.0	81.0	96.0	103	98.0	
Max. Temp. (°C)	25.9	26.6	27.4	28.1	29.1	30.3	31.0	31.9	31.3	32.1	32.7	32.0	31.7	31.6	31.9	32.3	32.0	32.0	32.0	31.7	31.4	
Min. Temp. (°C)	13.0	13.7	15.0	15.9	16.6	17.4	18.6	19.9	20.9	21.9	22.4	22.9	22.9	22.9	23.6	24.0	24.3	25.0	25.6	25.7	26.0	
Mean. Temp. (°C)	19.5	20.2	21.2	22.0	22.9	23.9	24.8	25.9	26.1	27.0	27.6	27.5	27.3	27.3	27.8	28.2	28.2	28.5	28.8	28.7	28.7	
RHmax (%)	92.6	92.1	91.4	90.6	90.6	90.6	90.1	89.7	90.4	89.9	90.9	91.1	91.3	92.4	92.6	92.7	92.7	94.0	94.1	94.7	95.0	
RHmin (%)	47.9	45.3	45.0	45.1	44.3	43.6	44.6	48.3	54.7	55.0	56.0	61.1	62.6	64.6	66.0	66.4	69.4	71.9	73.0	75.4	76.6	
RHmean (%)	70.3	68.7	68.2	67.9	67.6	67.1	67.4	69.0	72.6	72.5	73.8	76.1	77.0	78.5	79.3	79.6	81.1	83.3	83.6	85.1	85.8	
SShr (hrs)	44.0	48.0	50.0	50.0	51.0	53.0	53.0	48.0	44.0	49.0	49.0	49.0	46.0	44.0	44.0	45.0	39.0	36.0	44.0	26.0	25.0	
WD (degree)	228	218	222	210	214	202	208	179	165	149	146	143	128	126	134	131	126	130	142	138	143	
WS(Km/hr)	3.7	3.9	4.4	5.4	5.0	5.4	5.6	6.3	6.1	7.2	6.8	7.2	7.6	7.4	6.8	7.4	7.4	6.8	6.1	7.4	6.8	
																						
Stages	Spouting & Seedling				Vegetative growth				Flowering and Fruting					Harvesting								
Favourable Weather																						
Temperature	26-32 °C																					
Relative humidity	50-80%																					
Soil temperature	22-26°C																					
Sunshine hour	8-10 hr																					
Favourable weather for disease and pest infestation																						
White fly	15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																					
Tomato fruit borer					1 9.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %																	
Leaf miner					Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C; Rainfall had positive and significant effect on leaf miner population																	
Early blight					Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall																	
Wilting	Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%																					
Weather warning																						
Rainfall (mm)	Early rainfall (15-20 mm)													More than 40 mm rainfall in the season								
Cloudy weather	Cloudy condition																					
Drought	Drought due to excessive dry weather																					
Duration of staying wet									Land remain waterlogged for 2 to 5 days													
Hailstorm	Early rainfall with hailstorm																					
Windspeed	≥40 km/hr																					
Temperature (°C)	≥35 °C																					

District: Rajshahi						Tomato-Rabi season								Duration: 120-150 days														
Month	November					December				January				February				March					April					
Std.Week/Normal	44	45	46	47	48	49	50	51	52	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16			
Rainfall (mm)	7.0	5.0	1.0	0.0	2.0	1.0	3.0	0.0	4.0	1.0	1.0	2.0	1.0	4.0	3.0	2.0	3.0	5.0	4.0	3.0	4.0	14.0	13.0	10.0	19.0			
Max. Temp. (°C)	31.7	31.0	29.6	28.9	28.3	27.0	26.0	25.4	24.4	24.0	23.0	24.0	24.9	25.6	26.7	28.1	29.3	30.4	32.1	33.1	34.6	35.0	36.1	36.7	36.4			
Min. Temp. (°C)	23.1	21.4	17.6	16.6	15.0	14.0	13.1	12.4	11.6	11.0	10.9	10.9	10.9	12.0	12.3	13.6	14.1	15.1	16.4	17.6	19.4	20.6	22.1	23.0	23.4			
Mean. Temp. (°C)	27.4	26.2	23.6	22.8	21.7	20.5	196	18.9	18.0	17.5	17.4	17.5	17.9	18.8	19.5	20.9	21.7	22.8	24.3	25.4	27.0	27.8	29.1	29.9	29.9			
RHmax (%)	96.1	96.0	95.7	95.3	94.7	94.7	95.9	95.3	95.7	95.6	95.1	95.3	95.3	95.3	94.0	93.9	92.0	91.6	90.3	88.7	88.3	89.1	88.9	89.0	90.1			
RHmin (%)	60.9	57.4	49.4	47.4	45.6	46.9	50.1	50.0	50.0	50.1	48.4	47.9	45.7	44.7	42.0	40.3	36.6	34.6	32.7	30.4	30.4	35.9	34.3	34.6	41.4			
RHmean (%)	78.5	76.7	72.6	71.4	70.2	70.8	73.0	72.7	72.9	72.9	71.6	71.6	70.5	70.0	68.0	67.1	64.3	63.1	61.5	59.3	59.4	62.5	61.6	61.8	65.8			
SShr (hrs)	53.0	56.0	54.0	55.0	55.0	51.0	47.0	46.0	44.0	42.0	42.0	46.0	52.0	51.0	55.0	56.0	58.0	59.0	59.0	59.0	59.0	52.0	57.0	57.0	57.0			
WD (degree)	146	146	151	146	145	143	146	144	167	146	161	174	196	198	195	197	215	209	220	214	208	186	185	185	267			
WS(Km/hr)	3.9	4.3	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.9	3.7	3.7	4.4	3.9	5.0	3.9	5.0	5.0	5.4	5.4	5.6	5.6	6.1	6.1	8.2			
Stages	Spouting & Seedling					Vegetative growth				Flowering and Fruiting								Harvesting										
Favourable Weather																												
Temperature	18-25 °C																											
Relative humidity	50-70%																											
Soil temperature	20-25 °C																											
Sunshine hour	6-8 hr																											
Favourable weather for disease and pest infestation																												
White fly	15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																											
Tomato fruit borer														19.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %														
Leaf miner						Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C ; Rainfall had positive and significant effect on leaf miner population																						
Early blight						Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall																						
late blight						Temperature: 15-27°C, prolonged hot dry days increased disease infestation; relative humidity 75 to 90% or greater																						
Fungal Wilting	Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%																											
Weather warning																												
Rainfall (mm)	≥20 mm rainfall at the end of the rainy season (September-November)																											
Cloudy weather	Cloudy condition																											
Drought														Drought (moisture stress) due to excessive dry weather														
Duration of staying wet	Land remain waterlogged for 2 to 4 days																											
Hailstorm	Late season rainfall with hailstorm																											
Windspeed																	≥15 km/hr											
Temperature (°C)																	30 to 32°C											

Crop Weather Calendar of Kharif Tomato: Rajshahi region (Districts: Rajshahi, Chapai Nawabganj, Naogaon, Natore)

Region: Rajshahi	Tomato- Kharif season										Duration: 120-150 days									
Month	May					June					July				August					
Std.Week/Normal	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35		
Rainfall (mm)	24.0	29.0	33.0	36.30	34.0	48.0	51.0	78.0	72.0	66.0	71.0	75.0	72.0	52.0	57.0	64.0	53.0	57.10		
Max. Temp. (°C)	35.4	35.3	35.1	35.0	35.3	35.0	33.9	33.1	33.0	32.7	32.9	32.3	32.6	32.9	32.9	32.7	33.0	33.0		
Min. Temp. (°C)	23.7	24.3	24.6	25.0	25.4	25.9	26.0	26.0	26.0	26.0	26.0	26.0	26.0	26.0	26.1	26.0	26.1	26.0		
Mean. Temp. (°C)	29.6	29.8	29.9	30.0	30.4	30.5	30.0	29.6	29.5	29.4	29.5	29.2	29.3	29.5	29.5	29.4	29.6	29.5		
RHmax (%)	92.1	92.7	93.3	93.1	93.7	94.6	95.4	96.1	96.6	96.6	96.9	97.0	96.9	96.7	96.6	96.7	96.1	96.4		
RHmin (%)	50.0	51.9	55.6	58.3	58.9	62.6	67.9	71.1	73.4	73.7	74.3	75.3	73.9	72.6	72.6	74.1	72.0	70.9		
RHmean (%)	71.1	72.3	74.5	75.7	76.3	78.6	81.7	83.6	85.0	85.2	85.6	86.2	85.4	84.7	84.6	85.4	84.1	83.7		
SShr (hrs)	52.0	53.0	54.0	48.0	51.0	48.0	35.0	31.0	31.0	30.0	30.0	30.0	34.0	34.0	36.0	33.0	34.0	40.0		
WD (degree)	136	136	135	126	133	139	129	127	132	141	134	138	140	131	131	141	143	133		
WS(Km/hr)	7.4	7.2	7.2	7.2	7.2	6.7	6.7	6.8	6.8	5.7	6.1	6.3	6.7	6.3	5.6	5.7	5.6	5.6		
																				
Stages	Spouting & Seedling					Vegetative growth					Flowering and Fruting					Harvesting				
Favourable Weather																				
Temperature	18-25 °C																			
Relative humidity	50-70%																			
Soil temperature	20-25 °C																			
Sunshine hour	6-8 hr																			
Favourable weather for disease and pest infestation																				
White fly	15°C to 32°C, the optimum temperature for white fly population buildup is 20 °C to 24°C; Hot weather (high temperature) and low humidity with little or no rainfall are conducive for the multiplication and infestation of white fly																			
Tomato fruit borer								19.73°C to 31.44 °C; Average relative humidity ranging from 86.18% to 93.61 %												
Leaf miner						Grow best at 29.5°C to 32.0 °C and maximum temperature range for it to resist is at 34.18°C to 40.5 °C ; Rainfall had positive and significant effect on leaf miner population														
Early blight						Disease develops at moderate to warm (15°C to 30°C) temperatures; relative humidity 90% or greater, moderately to heavy rainfall														
late blight						Temperature: 15-27°C, prolonged hot dry days increased disease infestation; relative humidity 75 to 90% or greater														
Fungal Wilting	Dry atmospheric conditions, Max. Soil temperature 35°C to 43.5°C; Mini. Soil temperature 25°C to 30 °C; RH: 30%-50%																			
Weather warning																				
Rainfall (mm)	≥20 mm rainfall at the end of the rainy season (September-November)																			
Cloudy weather	Cloudy condition																			
Drought															Drought (moisture stress) due to excessive dry weather					
Duration of staying wet	Land remain waterlogged for 2 to 4 days																			
Hailstorm	Late season rainfall with hailstorm																			
Windspeed												≥15 km/hr								
Temperature (°C)												30 to 32°C								