

Crop Weather Calendar of *Rabi* season Sweet gourd: Barisal region (Districts: Patuakhali, Barisal, Jhalokati, Barguna, Bhola)

Region: Barisal		Sweet gourd– <i>Rabi</i> season														Duration:120-150 days				
Months		October		November				December				January				February				
Std.Week/Normal	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	
Rainfall (mm)	41.0	25.0	19.3	25.3	9.0	6.0	7.0	1.7	4.0	1.3	0.7	1.3	1.0	1.0	1.7	2.7	3.7	5.0	8.0	
Max. Temp. (°C)	32.0	31.7	31.0	30.5	29.9	29.3	28.3	27.8	27.0	26.2	25.7	25.1	25.2	25.8	26.2	27.0	27.8	28.9	29.7	
Min. Temp. (°C)	24.3	22.9	22.1	21.0	19.6	28.4	16.7	15.6	14.9	14.0	13.1	12.6	12.5	12.7	12.5	14.0	14.6	16.2	16.9	
Mean Temp. (°C)	28.1	27.3	26.5	25.7	24.8	28.9	22.5	21.7	21.0	20.1	19.4	18.9	18.9	19.2	19.4	20.5	21.2	22.5	23.3	
RHmax (%)	97.8	97.8	97.8	97.7	97.8	97.8	97.8	98.0	97.9	98.2	98.2	98.0	97.8	97.6	97.6	97.3	97.1	97.0	97.0	
RHMin (%)	68.0	63.7	63.2	60.7	56.6	53.7	51.8	50.6	50.9	50.7	50.3	50.1	49.3	48.5	45.2	46.2	44.8	46.2	45.1	
RHmean (%)	82.9	80.7	80.5	79.2	77.2	75.8	74.8	74.3	74.4	74.4	74.3	74.1	73.6	73.1	71.4	71.7	70.9	71.6	71.0	
SShr (hrs)	44.7	48.7	47.0	47.3	49.0	50.0	51.3	48.7	46.3	44.3	43.7	46.7	44.0	47.3	52.7	50.3	52.3	51.7	52.0	
WD (deg)	228	212	214	214	232	217	212	225	223	212	212	209	211	204	216	217	211	214	211	
WS(Km/hr)	3.6	3.1	3.5	3.1	2.7	3.3	3.6	3.1	3.4	3.3	3.9	4.5	4.4	4.5	4.3	5.0	4.6	4.9	4.9	
																				
Stage	Germination & Seedling				vegetative growth				Flowering & fruiting						Harvesting					
Favourable condition																				
Temperature	21 to 32 °C				20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)															
Relative Humidity	55-70%																			
Soil Temperature	18 to 25 °C																			
Sunshine Hour	6 to 8 hrs. (at least)																			
Favorable environmental conditions for insects and diseases																				
Fruit fly					Temperature: 25-30°C,RH: 60-91%															
Pumpkin beetle					Temperature: 25-32°C,RH: 65-80%															
Aphid					Temperature: 22-30°C,RH:60-85%															
Powdery mildew					Temperature: 22-27°C,RH: Lower RH during day with higher RH during night															
Downy mildew					Temperature: 25-32°C,RH: 75-95%															
Wilting	Temperature: 22-27°C (Atmosphere), >24°C (Soil) RH: 70-80% (Soil)																			
Anthraxnose/Fruit rot					Temperature: 25-32°C,RH: High (89-85%), Higher infestation with continuous rainfall															
Weather warning																				
Rainfall (mm)	20 to 30 mm rainfall (creates waterlogging condition)														Early rainfall (15 mm or more)					
Cloudy weather	Cloudy condition																			
Droughth					Drought due to excessive dry weather															
Duration of staying wet	Land remain waterlogged for 3 to 7 days																			
Hailstorm	Seasonal rainfall with hailstorm														Early rainfall with hailstorm					
Windspeed (Km/hr.)	≥40 Km/hr																			
Temperature (°C)	>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																			

Crop Weather Calendar of *Kharif*season Sweet gourd:Barisal region (Districts: Patuakhali, Barisal, Jhalokati, Barguna, Bhola)

Region:Barisal	Sweet gourd – Kharif season																Duration:120-150 days					
Months	March				April				May				June				July					
Std.Week/Normal	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	
Rainfall (mm)	5.3	6.0	14.3	18.0	18.0	21.7	25.0	34.7	40.7	52.7	49.0	69.3	66.7	85.3	105.3	129.7	126.3	129.0	91.7	111.7	115.0	
Max. Temp. (°C)	31.7	32.7	33.0	32.9	33.3	33.7	33.5	33.5	33.6	33.4	33.4	33.2	33.3	32.8	31.9	31.3	31.0	31.0	31.1	30.8	31.0	
Min. Temp. (°C)	19.5	20.5	22.1	22.8	23.7	24.0	24.6	24.4	24.4	25.1	25.2	25.7	26.0	26.1	26.0	26.0	26.0	26.0	26.0	26.0	25.9	
Mean Temp. (°C)	25.6	26.6	27.6	27.9	28.5	28.8	29.1	29.0	29.0	29.3	29.3	29.5	29.6	29.4	28.9	28.6	28.5	28.5	28.5	28.6	28.5	
RHmax (%)	97.1	96.7	96.5	96.5	96.6	96.4	96.0	96.0	96.1	96.2	96.1	96.2	96.5	96.4	96.6	97.0	97.3	97.4	97.4	97.4	97.6	
RHMin (%)	45.2	45.3	50.7	54.4	58.0	59.30	61.9	62.5	64.2	65.9	67.5	69.6	70.9	73.2	76.9	79.2	80.6	80.7	79.4	81.0	80.0	
RHmean (%)	71.2	71.0	73.6	75.4	77.3	78.1	78.9	79.3	80.1	81.1	81.8	82.9	83.7	84.8	86.8	88.1	80.0	89.0	88.4	89.2	88.8	
SShr (hrs)	54.0	54.3	52.3	50.7	53.0	51.3	54.7	51.7	49.0	48.0	46.0	37.3	37.3	30.7	30.0	21.3	20.3	20.3	23.3	21.0	25.3	
WD (deg)	226	218	210	198	200	198	197	191	190	192	189	186	18/7	185	177	179	175	180	175	177	175	
WS(Km/hr)	5.2	5.4	6.3	6.7	6.9	7.0	7.5	7.2	7.0	7.3	7.1	7.0	7.0	6.3	6.5	6.8	6.5	6.5	6.5	6.5	6.5	
																						
Stage	Germination & Seedling				vegetative growth				Flowering & fruiting				Harvesting									
Favourable condition																						
Temperature	21 to 32 °C				20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																	
Relative Humidity	55-70%																					
Soil Temperature	18 to 25 °C																					
Sunshine Hour	6 to 8 hrs. (at least)																					
Favorable environmental conditions for insects and diseases																						
Fruit fly					Temperature: 25-30°C,RH: 60-91%																	
Pumpkin beetle					Temperature: 25-32°C,RH: 65-80%																	
Aphid					Temperature: 22-30°C,RH:60-85%																	
Powdery mildew					Temperature: 22-27°C,RH: Lower RH during day with higher RH during night																	
Downy mildew					Temperature: 25-32°C,RH: 75-95%																	
Wilting	Temperature: 22-27°C (Atmosphere), >24°C (Soil) RH: 70-80% (Soil)																					
Anthracnose/Fruit rot									Temperature: 25-32°C,RH: High (89-85%), Higher infestation with continuous rainfall													
Weather warning																						
Rainfall (mm)	Early rainfall (15 mm or more)										40 mm or more (creates waterlogging condition)											
Cloudy weather	Cloudy condition																					
Droughth	Drought due to lack of rainfall																					
Duration of staying wet	Land remain waterlogged for 3 to 7 days																					
Hailstorm	Early rainfall and seasonal rainfall with hailstorm																					
Windspeed (Km/hr.)	≥40 Km/hr																					
Temperature (°C)	>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																					

Reg.: Chhattogram		Sweet gourd – Rabi season											Duration:110-120 days					
Months		September				October				November					December			
Std.Week/Normal	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	
Rainfall (mm)	118.3	92.7	73.3	73.7	60.3	66.7	75.7	26.0	19.3	21.7	14.3	9.0	11.7	1.0	4.0	2.3	2.0	
Max. Temp. (°C)	31.3	31.4	31.9	31.9	32.1	32.2	31.9	31.9	31.5	31.2	30.5	29.9	29.0	28.6	27.8	26.9	26.6	
Min. Temp. (°C)	25.3	25.2	25.1	25.2	25.1	25.0	24.8	23.3	22.8	21.6	20.2	19.2	18.0	17.0	16.3	15.3	14.9	
Mean Temp. (°C)	28.3	28.3	28.5	28.5	28.6	28.6	28.3	27.6	27.1	26.4	25.3	24.6	23.5	22.8	22.1	21.1	20.8	
RHmax (%)	95.9	95.6	95.7	95.7	95.7	96.0	96.1	95.0	94.9	94.7	94.4	94.1	94.5	94.4	94.2	94.4	94.1	
RHMin (%)	74.2	73.4	72.2	71.5	70.2	69.9	68.8	62.2	62.2	58.1	55.4	52.0	50.8	49.5	50.0	50.3	49.1	
RHmean (%)	85.0	84.5	83.9	83.6	82.9	82.9	82.4	78.6	78.6	76.4	74.9	73.1	72.6	71.9	72.2	72.3	71.6	
SShr (hrs)	36.8	35.8	39.3	39.5	42.5	42.8	45.8	53.3	52.3	54.0	55.5	56.5	55.5	54.8	52.8	50.8	51.3	
WD (deg)	179	185	190	194	202	209	203	201	196	195	199	203	196	204	196	199	207	
WS(Km/hr)	9.1	8.6	8.3	7.6	7.4	6.8	6.7	6.2	6.0	5.9	5.6	6.1	5.9	5.1	5.7	5.8	6.2	
Stage	Germination and seedling				vegetative growth				Flowering & fruiting					Harvesting				
Favourable condition																		
Temperature	21 to 32 °C				20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)													
Relative Humidity	55-70%																	
Soil Temperature	18 to 25 °C																	
Sunshine Hour	6 to 8 hrs. (at least)																	
Favorable environmental conditions for insects and diseases																		
Fruit fly					Temperature: 25-30° C, RH: 60-91%													
Pumpkin beetle					Temperature: 25-32° C, RH: 65-80%													
Aphid					Temperature: 22-30° C, RH:60-85%													
Powdery mildew					Temperature: 22-27° C, RH: Lower RH during day with higher RH during night													
Downy mildew					Temperature: 25-32° C, RH: 75-95%													
Wilting	Temperature: 22-27° C (Atmosphere), >24° C (Soil) RH: 70-80% (Soil)																	
Anthrachnose/Fruit rot									Temperature: 25-32° C, RH: High (89-85%), Higher infestation with continuous rainfall									
Weather warning																		
Rainfall (mm)	20 to 30 mm rainfall (creates waterlogging condition)																	
Cloudy weather	Cloudy condition																	
Droughth	Drought due to lack of rainfall and excessive dry weather																	
Duration of staying wet	Land remain waterlogged for 3 to 7 days																	
Hailstorm	Seasonal rainfall with hailstorm																	
Windspeed (Km/hr.)	≥ 40 Km/hr																	
Temperature (°C)	>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																	

Crop Weather Calendar of *Kharif* season Sweet gourd: Chattogram region (Districts: Cox's bazar, Chattogram, Feni, Noakhali, Lakshmipur)

Region: Chattogram	Sweet gourd – Kharif season												Duration:120-125 days				
Months	April				May					June			July				
Std.Week/Normal	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Rainfall (mm)	21.3	27.0	34.3	36.0	62.7	61.3	70.3	88.7	88.0	119.3	161.0	180.7	197.3	201.7	156.0	189.0	173.3
Max. Temp. (°C)	32.1	32.6	32.7	32.9	32.8	32.9	32.8	32.5	32.3	32.1	31.5	31.3	30.9	30.7	31.1	30.7	30.9
Min. Temp. (°C)	23.5	24.0	24.4	24.5	24.5	24.8	25.2	25.2	25.6	25.6	25.4	25.4	25.2	25.4	25.3	25.2	25.3
Mean Temp. (°C)	27.8	28.3	28.6	28.7	28.7	28.9	29.0	28.9	29.0	28.8	28.5	28.3	28.1	28.0	28.2	27.9	28.1
RHmax (%)	93.3	92.9	92.6	92.8	92.9	93.3	93.5	93.9	94.0	94.9	95.3	95.1	95.9	96.0	95.7	96.1	95.8
RHMin (%)	60.1	60.5	61.6	62.0	62.3	64.1	66.1	68.9	70.2	71.9	74.8	75.3	77.3	77.7	76.4	77.6	76.3
RHmean (%)	76.7	76.7	77.1	77.4	77.6	78.7	79.8	81.4	82.1	83.4	85.1	85.2	86.6	86.9	86.0	86.9	86.1
SShr (hrs)	56.0	56.0	54.8	55.3	52.8	55.0	48.0	42.8	40.3	36.8	28.0	28.0	25.3	24.0	29.0	25.5	30.0
WD (deg)	209	205	205	202	199	203	199	186	185	182	177	174	173	173	173	174	174
WS(Km/hr)	10.0	11.1	10.6	10.6	10.1	10.0	10.1	10.0	10.2	9.8	10.5	11.0	10.9	10.7	10.8	10.9	10.3
																	
Stage	Germination and seedling				Vegetative growth			Flowering & fruiting					Harvesting				
Favourable condition																	
Temperature	21-32 °C			20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)													
Relative Humidity	55-70%																
Soil Temperature	18 to 25 °C																
Sunshine Hour	6 to 8 hrs. (at least)																
Favorable environmental conditions for insects and diseases																	
Fruit fly					Temperature: 25-30° C, RH: 60-91%												
Pumpkin beetle					Temperature: 25-32° C, RH: 65-80%												
Aphid					Temperature: 22-30° C, RH:60-85%												
Powdery mildew					Temperature: 22-27° C, RH: Lower RH during day with higher RH during night												
Downy mildew					Temperature: 25-32° C, RH: 75-95%												
Wilting	Temperature: 22-27° C (Atmosphere), >24° C (Soil) RH: 70-80% (Soil)																
Anthrachnose/Fruit rot					Temperature: 25-32° C, RH: High (89-85%), Higher infestation with continuous rainfall												
Weather warning																	
Rainfall (mm)	Early rainfall (15 mm or more)								40 mm or more (creates waterlogging condition)								
Cloudy weather	Cloudy condition																
Droughth	Drought due to lack of rainfall																
Duration of staying wet	land remain waterlogged for 3 to 7 days																
Hailstorm	Early rainfall and seasonal rainfall with hailstorm																
Windspeed (Km/hr.)	≥ 40 Km/hr																
Temperature (°C)	>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																

Crop Weather Calendar of *Rabi* season Sweet gourd: Dhaka region (Districts: Dhaka, Narayanganj, Gazipur, Kishoreganj, Narsingdi, Munshiganj, Manikganj, Tangail)

Region: Dhaka		Sweet gourd – Rabi season										Duration:120-150 days									
Months	October		November				December				January				February				March		
Std.Week/Normal	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10
Rainfall (mm)	40.5	26.0	12.5	10.0	6.0	1.5	4.5	1.5	4.5	1.5	3.5	1.0	0.0	0.0	0.5	3.5	3.5	2.5	12.0	3.5	4.0
Max. Temp. (°C)	32.0	31.5	30.9	30.6	29.8	29.0	28.3	27.6	26.2	25.5	24.8	24.0	23.9	24.5	25.0	26.2	27.2	28.3	28.9	30.0	31.5
Min. Temp. (°C)	23.8	22.3	21.5	20.5	28.7	17.7	16.1	15.5	14.7	13.5	12.9	12.4	12.0	12.0	12.1	13.6	14.3	15.5	16.5	17.5	18.6
Mean Temp. (°C)	27.9	26.9	26.2	25.5	24.3	23.3	22.2	21.5	20.5	19.5	18.8	18.2	17.9	18.3	18.6	19.9	20.7	21.9	22.7	23.7	25.1
RHmax (%)	95.0	94.8	94.5	95.0	94.7	94.6	94.3	94.8	95.3	95.1	95.0	94.6	94.3	93.8	93.4	92.4	92.1	91.7	91.1	90.0	89.9
RHMin (%)	60.2	55.5	54.6	51.5	49.3	47.5	46.3	47.6	49.4	49.8	49.1	50.3	48.8	46.2	43.3	42.3	39.4	39.7	38.9	36.0	35.5
RHmean (%)	77.6	75.2	74.6	73.2	72.0	71.0	70.3	71.2	72.3	72.5	72.1	72.4	71.6	70.0	68.3	67.3	65.7	65.7	65.0	63.0	62.7
SShr (hrs)	47.5	54.0	49.5	51.5	51.5	52.0	51.5	48.5	46.5	44.5	44.5	42.5	42.5	45.0	49.0	49.5	51.5	53.0	54.5	56.0	55.0
WD (deg)	192	190	190	189	199	212	211	219	224	225	233	238	238	232	242	241	236	234	228	234	230
WS(Km/hr)	4.1	3.8	3.7	3.7	3.7	3.7	3.6	3.7	3.7	3.7	3.7	4.1	4.0	4.1	4.3	4.3	4.5	5.0	5.3	5.0	5.4
																					
Stage	Germination & Seedling				vegetative growth				Flowering & fruiting				Harvesting								
Favourable condition																					
Temperature	21-32 °C				20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																
Relative Humidity					55-70%																
Soil Temperature					18 to 25 °C																
Sunshine Hour					6 to 8 hrs. (at least)																
Favorable environmental conditions for insects and diseases																					
Fruit fly					Temperature: 25-30° C, RH: 60-91%																
Pumpkin beetle					Temperature: 25-32° C, RH: 65-80%																
Aphid					Temperature: 22-30° C, RH:60-85%																
Powdery mildew					Temperature: 22-27° C, RH: Lower RH during day with higher RH during night																
Downy mildew					Temperature: 25-32° C, RH: 75-95%																
Wilting	Temperature: 22-27° C (Atmosphere), >24° C (Soil) RH: 70-80% (Soil)																				
Anthracnose/Fruit rot									Temperature: 25-32° C, RH: High (89-85%), Higher infestation with continuous rainfall												
Weather warning																					
Rainfall (mm)	20 to 30 mm rainfall (creates waterlogging condition)								Early rainfall (10 to 15 mm or more)												
Cloudy weather					Cloudy condition																
Droughth					Drought due to excessive dry weather																
Duration of staying wet					Land remain waterlogged for 3 to 7 days																
Hailstorm					Seasonal rainfall with hailstorm								Early rainfall with hailstorm								
Windspeed (Km/hr.)					≥ 40 Km/hr																
Temperature (°C)					>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																

Crop Weather Calendar of *Kharif* season Sweet gourd: Dhaka region (Districts: Dhaka, Narayanganj, Gazipur, Kishoreganj, Narsingdi, Munshiganj, Manikganj, Tangail)

Region: Dhaka	Sweet gourd – Kharif season															Duration:130-150 days						
Months	March					April				May				June						July		
Std.Week/Normal	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Rainfall (mm)	3.5	4.0	9.5	13.5	20.5	25.0	21.0	31.5	42.5	46.0	55.5	55.5	75.5	70.0	72.5	86.5	81.0	77.5	82.5	73.5	91.0	81.5
Max. Temp. (°C)	30.0	31.5	32.3	33.4	33.2	33.9	24.6	34.1	33.8	33.6	33.7	33.7	33.2	33.5	33.4	32.8	32.2	32.2	32.1	32.0	31.7	32.2
Min. Temp. (°C)	17.5	18.6	19.8	19.8	21.8	22.9	23.5	23.7	23.6	23.7	24.2	24.6	24.9	25.4	25.8	25.9	26.2	26.2	26.1	26.2	26.0	26.0
Mean Temp. (°C)	23.7	25.0	26.1	26.1	27.5	28.4	29.0	28.9	28.7	28.6	28.9	29.2	29.0	29.4	29.6	29.3	29.2	29.2	29.1	29.1	28.8	29.1
RHmax (%)	90.0	89.9	89.2	89.6	90.3	91.3	91.4	91.4	91.7	92.3	92.8	92.9	93.5	93.8	94.0	94.5	94.4	94.7	94.7	94.1	94.5	94.3
RHMin (%)	36.0	35.5	35.9	38.9	44.5	45.9	47.1	52.0	55.1	56.7	58.6	60.4	63.5	65.0	66.7	69.9	71.5	72.9	73.5	73.1	73.5	71.1
RHmean (%)	63.0	62.7	62.5	64.3	67.4	68.6	69.2	71.7	73.4	74.5	75.7	76.7	78.5	79.4	80.3	82.2	83.0	83.8	84.1	83.6	84.0	82.7
SShr (hrs)	56.0	55.0	58.0	55.5	52.5	53.5	54.0	55.0	54.5	51.5	48.0	48.0	42.5	42.0	38.5	31.0	26.0	26.0	27.0	27.5	28.5	32.0
WD (deg)	234	230	223	206	188	181	175	167	156	159	164	156	155	157	161	157	157	158	162	157	161	159
WS(Km/hr)	5.0	5.4	5.7	6.2	6.5	6.5	6.6	6.6	6.7	6.8	6.5	6.4	6.1	5.8	5.6	6.3	6.5	6.5	5.8	5.9	6.1	5.9
																						
Stage	Germination & Seedling					vegetative growth				Flowering & fruiting				Harvesting								
Favourable condition																						
Temperature	21-32 °C					20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																
Relative Humidity	55-70%																					
Soil Temperature	18 to 25 °C																					
Sunshine Hour	6 to 8 hrs. (at least)																					
Favorable environmental conditions for insects and diseases																						
Fruit fly						Temperature: 25-30° C, RH: 60-91%																
Pumpkin beetle						Temperature: 25-32° C, RH: 65-80%																
Aphid						Temperature: 22-30° C, RH:60-85%																
Powdery mildew						Temperature: 22-27° C, RH: Lower RH during day with higher RH during night																
Downy mildew						Temperature: 25-32° C, RH: 75-95%																
Wilting	Temperature: 22-27° C (Atmosphere), >24° C (Soil) RH: 70-80% (Soil)																					
Anthraco nose/Fruit rot										Temperature: 25-32° C, RH: High (89-85%), Higher infestation with continuous rainfall												
Weather warning																						
Rainfall (mm)	Early rainfall (15 mm or more)									40 mm or more (creates waterlogging condition)												
Cloudy weather	Cloudy condition																					
Droughth	Drought due to lack of rainfall and excessive dry weather																					
Duration of staying wet	Land remain waterlogged for 3 to 7 days																					
Hailstorm	Early rainfall and seasonal rainfall with hailstorm																					
Windspeed (Km/hr.)	≥ 40 Km/hr																					
Temperature (°C)	>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																					

Reg.: Mymensingh	Sweet gourd – <i>Rabi</i> season													Duration:130-150 days							
Months	October				November					December				January				February			
Std.Week/Normal	40	41	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8
Rainfall (mm)	71.0	52.0	45.0	15.0	7.0	5.0	5.0	0.0	4.0	1.0	3.0	0.0	2.0	0.0	3.0	1.0	0.0	2.0	1.0	5.0	7.0
Max. Temp. (°C)	32.0	31.7	31.6	31.1	30.9	30.3	30.0	28.9	28.3	27.6	26.3	25.4	25.0	24.4	24.0	24.0	25.1	25.9	26.6	27.4	28.1
Min. Temp. (°C)	25.0	24.3	23.4	21.7	20.9	19.7	18.3	17.0	15.4	15.0	14.0	13.3	12.4	12.0	11.6	11.9	12.0	13.0	13.7	15.0	15.9
Mean Temp. (°C)	28.5	28.0	27.5	26.4	25.9	25.0	24.2	23.0	21.9	21.3	21.2	19.4	18.7	18.2	17.8	18.0	18.6	19.5	20.2	21.2	22.0
RHmax (%)	95.6	95.3	95.3	95.4	95.0	95.0	94.6	95.0	95.1	95.3	95.6	95.6	95.3	95.1	94.4	94.4	93.4	92.6	92.1	91.4	90.6
RHMin (%)	72.6	70.3	66.7	62.7	61.0	58.4	55.9	53.0	51.6	52.3	54.1	53.6	53.7	53.6	53.1	51.4	47.0	47.9	45.3	45.0	45.1
RHmean (%)	84.1	82.8	81.0	79.1	78.0	76.7	75.3	74.0	73.4	73.8	74.9	74.9	74.5	74.4	73.8	72.9	70.2	70.3	68.7	68.2	67.9
SShr (hrs)	40.0	41.0	47.0	53.0	51.0	53.0	52.0	53.0	50.0	49.0	43.0	40.0	42.0	41.0	39.0	41.0	45.0	44.0	48.0	50.0	50.0
WD (deg)	181	174	193	197	186	181	186	186	193	191	206	199	221	230	236	216	238	228	218	222	210
WS(Km/hr)	4.3	3.9	3.7	3.7	3.7	3.9	3.7	3.7	3.7	3.0	3.5	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.9	4.4	5.4
Stage	Germination & Seedling				vegetative growth				Flowering & fruiting					Harvesting							
Favourable condition	Temperature 21 to 32 °C				20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)				55-70%					18 to 25 °C							
Relative Humidity									6 to 8 hrs. (at least)												
Soil Temperature																					
Sunshine Hour																					
Favorable environmental conditions for insects and diseases																					

Reg.: Mymensingh	Sweet gourd – Kharif season													Duration:120-150 days							
Months	April				May					June				July				August			
Std.Week/Normal	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34
Rainfall (mm)	20.0	30.0	40.0	36.0	55.0	62.0	69.0	97.0	81.0	96.0	103.0	98.0	85.0	109.0	99.0	89.0	100.0	72.0	74.0	67.0	81.0
Max. Temp. (°C)	32.1	32.7	32.0	31.7	31.6	31.9	32.3	32.0	32.0	32.0	31.7	31.4	31.0	31.1	31.3	31.3	31.1	32.0	32.0	32.0	32.0
Min. Temp. (°C)	21.9	22.4	22.9	22.9	22.9	23.6	24.0	24.3	25.0	25.6	25.7	26.0	26.0	26.0	26.0	26.0	26.0	26.0	26.3	26.3	26.0
Mean Temp. (°C)	27.0	27.6	27.5	27.3	27.3	27.8	28.2	28.2	28.5	28.8	28.7	28.7	28.5	28.6	28.7	28.7	28.6	29.1	29.2	29.2	29.0
RHmax (%)	89.9	90.9	91.1	91.3	92.4	92.6	92.7	92.7	94.0	94.1	94.7	95.0	94.7	95.0	94.6	94.6	94.9	94.7	93.9	94.0	94.9
RHMin (%)	55.0	56.6	61.1	62.6	64.6	66.0	66.4	69.4	71.9	73.0	75.4	76.6	77.7	77.6	77.0	76.6	76.4	74.6	74.6	74.9	75.9
RHmean (%)	72.5	73.8	76.1	77.0	78.5	79.3	79.6	81.1	83.0	83.6	85.1	85.8	86.2	86.3	85.8	85.6	85.7	84.7	84.3	84.5	85.4
SShr (hrs)	49.0	49.0	49.0	46.0	44.0	44.0	45.0	39.0	36.0	34.0	26.0	25.0	23.0	24.0	24.0	25.0	30.0	33.0	31.0	30.0	26.0
WD (deg)	149	146	143	128	126	134	131	126	130	142	138	143	144	156	152	152	151	153	149	148	157
WS(Km/hr)	7.2	6.8	7.2	7.6	7.4	6.8	7.4	7.4	6.8	6.1	7.4	6.8	7.2	7.2	6.8	7.2	6.1	6.1	6.1	6.1	5.7
Stage	Germination & Seedling				vegetative growth					Flowering & fruiting				Harvesting							
Favourable condition	21 to 32 °C				20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)					55-70%				18 to 25 °C							
Temperature																					
Relative Humidity																					
Soil Temperature																					
Sunshine Hour																					

Region:Sylhet	Sweet gourd – Rabi season													Duration:110-120 days			
Months	October				November					December				January			
Std.Week/Normal	40	41	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4
Rainfall (mm)	63.0	60.0	36.5	24.5	12.0	14.5	8.5	0.5	7.0	0.5	4.5	1.0	4.0	0.0	0.5	0.5	1.0
Max. Temp. (°C)	31.9	31.7	31.6	31.4	30.9	30.1	29.6	29.0	28.2	27.8	26.9	26.1	25.7	25.3	24.7	25.1	25.8
Min. Temp. (°C)	24.0	23.4	22.7	21.2	20.4	19.1	17.3	16.6	15.3	14.3	13.7	12.6	12.0	11.4	11.3	11.3	11.1
Mean Temp. (°C)	27.9	27.5	27.2	26.3	25.7	24.6	23.5	22.8	21.7	21.1	20.3	19.3	18.8	18.3	18.0	18.0	18.4
RHmax (%)	96.4	96.1	96.0	95.6	95.4	95.2	95.2	95.3	95.5	94.7	95.7	95.3	96.1	95.8	96.1	95.5	94.9
RHMin (%)	96.4	96.1	66.5	63.1	61.9	59.2	56.2	54.4	53.5	51.4	53.6	53.0	52.7	53.5	53.9	50.7	48.1
RHmean (%)	83.8	84.4	81.3	79.3	78.6	77.2	75.7	74.9	74.5	73.0	74.6	74.2	74.4	74.6	75.0	73.1	71.5
SShr (hrs)	71.2	68.7	49.5	54.5	52.5	55.5	55.5	56.5	56.0	55.5	50.5	51.5	50.5	51.0	45.0	48.0	51.0
WD (deg)	190	191	187	191	173	179	180	184	190	177	186	191	203	200	211	213	219
WS(Km/hr)	83.8	82.4	4.6	4.5	4.7	4.5	4.3	4.6	4.3	4.2	4.5	4.5	4.3	4.6	4.9	5.1	4.8
Stage	Germination & Seedling				vegetative growth			Flowering & fruiting				Harvesting					
Favourable condition	Temperature 21 to 32 °C				20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)												
Relative Humidity	55-70%																
Soil Temperature	18 to 25 °C																
Sunshine Hour	6 to 8 hrs. (at least)																
Favorable environmental conditions for insects and diseases	Temperature: 25-30° C, RH: 60-91%																
Fruit fly					Temperature: 25-32° C, RH: 65-80%												
Pumpkin beetle					Temperature: 22-30° C, RH:60-85%												
Aphid					Temperature: 22-27° C, RH: Lower RH during day with higher RH during night												
Powdery mildew					Temperature: 25-32° C, RH: 75-95%												
Downy mildew																	
Wilting	Temperature: 22-27° C (Atmosphere), >24° C (Soil) RH: 70-80% (Soil)																
Anthracoise/Fruit rot					Temperature: 25-32° C, RH: High (89-85%), Higher infestation with continuous rainfall												
Weather warning																	
Rainfall (mm)	20 to 30 mm rainfall (creates waterlogging condition)																
Cloudy weather	Cloudy condition																
Droughth	Drought due to lack of rainfall																
Duration of staying wet	Land remain waterlogged for 3 to 7 days																
Hailstorm	Seasonal rainfall with hailstorm																
Windspeed (Km/hr.)	≥ 40 Km/hr																
Temperature (°C)	>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																

Region:Sylhet	Sweet gourd – Kharif season																Duration:120-150 days					
Months	April				May					June				July				August				
Std.Week/Normal	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	
Rainfall (mm)	20.0	30.0	40.0	36.0	55.0	62.0	69.0	97.0	81.0	96.0	103.0	98.0	85.0	109.0	99.0	89.0	100.0	72.0	74.0	67.0	81.0	
Max. Temp. (°C)	32.1	32.7	32.0	31.7	31.6	31.9	32.3	32.0	32.0	32.0	31.7	31.4	31.0	31.1	31.3	31.3	31.1	32.0	32.0	32.0	32.0	
Min. Temp. (°C)	21.9	22.4	22.9	22.9	22.9	23.6	24.0	24.3	25.0	25.6	25.7	26.0	26.0	26.0	26.0	26.0	26.0	26.1	26.3	26.3	26.0	
Mean Temp. (°C)	27.0	27.6	27.5	27.3	27.3	27.8	28.2	28.2	28.5	28.8	28.7	28.7	28.5	28.6	28.7	28.7	28.6	29.1	29.2	29.2	29.0	
RHmax (%)	89.9	90.9	91.1	91.3	92.4	92.6	92.7	92.7	94.0	94.1	94.7	95.0	94.7	95.0	94.6	94.6	94.9	94.7	939.9	94.0	94.9	
RHMin (%)	55.0	56.6	61.1	62.6	64.6	66.0	66.4	69.4	71.9	73.0	75.4	76.6	77.7	77.6	77.0	76.6	76.4	74.6	74.6	74.9	75.9	
RHmean (%)	72.5	73.8	76.1	77.0	78.5	79.3	79.6	81.1	83.0	83.6	85.1	85.8	86.2	86.3	85.8	85.6	85.7	84.7	84.3	84.5	85.4	
SShr (hrs)	49.0	49.0	49.0	46.0	44.0	44.0	45.0	39.0	36.0	34.0	26.0	25.0	23.0	24.0	24.0	25.0	30.0	33.0	31.0	30.0	26.0	
WD (deg)	149	146	143	128	126	134	131	126	130	142	138	143	144	156	152	152	151	153	149	148	157	
WS(Km/hr)	7.2	6.8	7.2	7.6	7.4	6.8	7.4	7.4	6.8	6.1	7.4	6.8	7.2	7.2	6.8	7.2	6.1	6.1	6.1	6.1	5.7	
Stage	Germination & Seedling				vegetative growth					Flowering & fruiting					Harvesting							
Favourable condition																						
Temperature	21 to 32 °C				20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																	
Relative Humidity	55-70%																					
Soil Temperature	18 to 25 °C																					
Sunshine Hour	6 to 8 hrs. (at least)																					
Favorable environmental conditions for insects and diseases																						
Fruit fly					Temperature: 25-30° C, RH: 60-91%																	
Pumpkin beetle					Temperature: 25-32° C, RH: 65-80%																	
Aphid					Temperature: 22-30° C, RH:60-85%																	
Powdery mildew					Temperature: 22-27° C, RH: Lower RH during day with higher RH during night																	
Downy mildew					Temperature: 25-32° C, RH: 75-95%																	
Wilting	Temperature: 22-27° C (Atmosphere), >24° C (Soil) RH: 70-80% (Soil)																					
Anthracoese/Fruit rot										Temperature: 25-32° C, RH: High (89-85%), Higher infestation with continuous rainfall												
Weather warning																						
Rainfall (mm)	Early rainfall (15 mm or more)									40 mm or more (creates waterlogging condition)												
Cloudy weather	Cloudy condition																					
Droughth	Drought due to lack of rainfall and excessive dry weather																					
Duration of staying wet	Land remain waterlogged for 3 to 7 days																					
Hailstorm	Early rainfall and Seasonal rainfall with hailstorm																					
Windspeed (Km/hr.)	≥ 40 Km/hr																					
Temperature (°C)	>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																					

Crop Weather Calendar of Rabi Sweet Gourd: Bogra Region (Districts: Bogra, Pabna, Sirajgonj, Joypurhat) Bangladesh

Bogra Region				Rabi Sweet Gourd: Bogra Region (Districts: Bogra, Pabna, Sirajgonj, Jesspura / Bangladesh)												Life Span 140-150 day				
Months	October		November				December				January				February					
Std.Week/Normal	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	
Rainfall (mm)	25.0	18.0	5.0	5.0	0.0	0.0	3.0	0.0	3.0	0.0	4.0	0.0	2.0	2.0	1.0	1.0	1.0	4.0	3.0	
Max. Temp. (°C)	32.0	31.6	31.0	30.7	30.1	29.7	28.9	27.7	26.3	25.0	25.7	24.0	24.0	24.3	25.4	25.9	26.9	27.7	28.6	
Min. Temp. (°C)	23.4	21.7	21.0	19.9	18.3	17.3	15.7	15.0	14.1	12.0	13.4	12.0	11.4	11.4	12.0	12.9	13.6	14.6	15.4	
Mean Temp. (°C)	27.7	26.7	26.0	25.3	24.2	23.5	22.3	21.4	20.2	18.7	19.6	18.1	17.7	17.9	18.7	19.4	20.3	21.2	22.0	
RHmax (%)	95.7	95.1	94.3	94.6	94.1	93.9	93.9	93.7	94.6	94.3	94.6	94.1	94.3	94.3	93.9	93.1	92.1	91.4	90.4	
RHMin (%)	62.1	57.9	55.6	52.4	49.9	48.3	47.0	48.1	50.6	50.6	50.0	50.9	49.0	47.7	46.0	44.9	41.4	40.3	39.1	
RHmean (%)	78.9	76.5	75.0	73.5	72.0	71.1	70.5	70.9	72.6	72.5	72.3	72.5	71.7	71.0	70.0	69.0	66.8	65.9	64.8	
SShr (hrs)	54.0	57.0	52.0	55.0	55.0	56	56	54	48	49.0	46.0	44.0	44.0	46.0	50.0	50.0	52.0	53.0	56.0	
WD (deg)	199	198	186	200	202	205	217	222	232	249	248	258	247	239	252	249	233	234	229	
WS(Km/hr)	5.0	4.4	4.4	3.7	3.7	3.7	3.7	3.7	3.9	3.7	3.7	4.3	5.4	4.8	5.0	5.4	5.6	5.6	6.7	
																				
Stage	Germination and Seedling Planting				Vegetative Growth				Flowering & Fruiting				Harvesting							
Favourable condition																				
Temperature	21 to 32 °C				20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)															
Relative Humidity	55-70%																			
Soil Temperature	18 to 25 °C																			
Sunshine Hour	6 to 8 hrs. (at least)																			
Favorable environmental conditions for insects and diseases																				
Fruit fly					Temperature: 25-30°C,RH: 60-91%															
Pumpkin beetle					Temperature: 25-32°C,RH: 65-80%															
Aphid					Temperature: 22-30°C,RH:60-85%															
Powdery mildew					Temperature: 22-27°C,RH: Lower RH during day with higher RH during night															
Downy mildew					Temperature: 25-32°C,RH: 75-95%															
Wilting	Temperature: 22-27°C (Atmosphere), >24°C (Soil) RH: 70-80% (Soil)																			
Anthrachnose/Fruit rot									Temperature: 25-32°C,RH: High (89-85%), Higher infestation with continuous rainfall											
Weather warning																				
Rainfall (mm)	20 to 30 mm rainfall (creates waterlogging condition)												Early rainfall (15 mm or more)							
Cloudy weather	Cloudy condition																			
Droughth					Drought due to excessive dry weather															
Duration of staying wet	Land remain waterlogged for 3 to 7 days																			
Hailstorm	Seasonal rainfall with hailstorm												Early rainfall with hailstorm							
Windspeed (Km/hr.)	≥40 Km/hr																			
Temperature (°C)	>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																			

Crop Weather Calendar of Kharif Sweet Gourd: Bogra Region (Districts: Bogra, Pabna, Sirajgonj, Joypurhat) Bangladesh

Bogra Region	Rabi Sweet Gourd												Life Span 120-130 day				
Months	April				May					June				July			
Std.Week/Normal	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
Rainfall (mm)	8.0	12.0	26.0	28.0	37.0	42.0	45.0	58.0	59.0	74.0	70.0	92.0	76.0	93.0	80.0	91.0	76.0
Max. Temp. (°C)	33.9	34.3	34.0	33.6	33.1	33.6	33.4	33.3	33.9	33.4	32.7	32.3	32.3	32.0	31.9	32.0	32.1
Min. Temp. (°C)	22.3	23.0	23.0	23.1	23.3	23.9	24.3	24.6	25.0	25.9	26.0	26.1	26.1	26.0	26.1	26.0	26.1
Mean Temp. (°C)	28.1	28.7	28.5	28.4	28.2	28.8	28.9	29.0	29.5	29.7	29.4	29.2	29.2	29.0	29.0	29.0	29.1
RHmax (%)	88.0	88.3	90.6	90.9	92.4	92.4	92.7	93.0	93.7	94.1	95.1	95.1	95.6	95.9	95.9	95.9	95.9
RHMin (%)	44.4	45.9	50.7	54.9	57.3	58.7	60.0	63.4	64.6	66.7	70.4	72.4	73.9	73.3	74.1	74.4	73.3
RHmean (%)	66.2	67.1	70.7	72.9	74.9	75.6	76.4	78.2	79.2	80.4	82.8	83.8	84.8	84.6	85.0	85.2	84.6
SShr (hrs)	56.0	56.0	55.0	52.0	48.0	50.0	49.0	46.0	44.0	39.0	33.0	28.0	26.0	28.0	31.0	29.0	28.0
WD (deg)	173	164	149	134	127	126	126	121	135	134	134	134	132	141	135	138	143
WS(Km/hr)	6.7	6.3	6.8	7.4	7.2	7.4	6.7	6.3	6.3	6.3	7.2	6.3	6.7	5.6	5.6	6.8	6.1
																	
Stage	Germination and Seedling Planting			Vegetative Growth				Flowering & Fruiting					Harvesting				

Favourable condition			
Temperature	21 to 32 °C	20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)	
Relative Humidity	55-70%		
Soil Temperature	18 to 25 °C		
Sunshine Hour	6 to 8 hrs. (at least)		
Favorable environmental conditions for insects and diseases			
Fruit fly		Temperature: 25-30°C,RH: 60-91%	
Pumpkin beetle		Temperature: 25-32°C,RH: 65-80%	
Aphid		Temperature: 22-30°C,RH:60-85%	
Powdery mildew		Temperature: 22-27°C,RH: Lower RH during day with higher RH during night	
Downy mildew		Temperature: 25-32°C,RH: 75-95%	
Wilting	Temperature: 22-27°C (Atmosphere), >24°C (Soil) RH: 70-80% (Soil)		
Anthraxnose/Fruit rot		Temperature: 25-32°C,RH: High (89-85%), Higher infestation with continuous rainfall	
Weather warning			
Rainfall (mm)	20 to 30 mm rainfall (creates waterlogging condition)		Early rainfall (15 mm or more)
Cloudy weather	Cloudy condition		
Droughth	Drought due to excessive dry weather		
Duration of staying wet	Land remain waterlogged for 3 to 7 days		
Hailstorm	Seasonal rainfall with hailstorm		Early rainfall with hailstorm
Windspeed (Km/hr.)	≥40 Km/hr		
Temperature (°C)	>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)		

Crop Weather Calendar of **Rabi Sweet Gourd**: Cumilla Region (Districts: Cumilla, Chandpur, B.Barua) Bangladesh

Cumilla Region	Rabi Sweet Gourd											Life Span 110-120 day					
Months	October		November					December				January				February	
Std. Week/Normal	42	43	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6
Rainfall (mm)	30.5	16.0	11.5	18.0	8.5	4.0	2.5	0.0	4.5	1.0	1.5	0.0	0.0	0.0	1.5	3.0	3.0
Max. Temp. (°C)	31.8	31.2	31.0	30.5	29.8	29.0	28.2	27.7	26.5	25.8	25.4	24.7	24.5	25.0	25.4	16.2	27.3
Min. Temp. (°C)	24.3	22.8	21.8	21.0	19.3	18.1	16.9	15.8	15.0	14.2	13.4	13.0	12.7	12.7	12.6	14.2	14.5
Mean Temp. (°C)	28.0	27.0	26.4	25.7	24.6	23.5	22.5	21.7	20.8	20.0	19.4	18.8	18.6	18.9	19.0	20.2	20.9
RHmax (%)	95.5	95.2	95.3	95.6	95.3	95.4	95.8	95.8	96.2	96.0	96.0	96.0	95.5	95.3	94.4	94.6	94.4
RHMin (%)	65.8	61.9	60.6	58.2	54.8	52.6	52.2	50.6	51.9	52.1	51.3	51.8	51.2	49.1	46.4	47.7	45.2
RHmean (%)	80.6	78.5	77.9	76.9	75.0	74.0	74.0	73.2	74.1	74.1	73.6	73.9	73.4	72.2	70.4	71.1	69.8
SShr (hrs)	44.5	50.5	47.0	49.5	50.0	50.5	48.5	47.5	42.5	40.5	40.0	40.0	37.0	43.5	47.0	47.0	50.0
WD (deg)	201	187	202	183	184	190	195	206	192	211	213	205	209	218	212	203	214
WS(Km/hr)	4.3	4.0	3.5	3.7	3.7	3.5	3.3	3.3	4.0	3.5	3.8	4.0	4.2	4.6	4.4	4.6	4.6
																	
Stage	Seedling raising & planting		vegetative growth					Flowering & fruiting				Harvesting					

Favourable condition			
Temperature	21 to 32 °C	20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)	
Relative Humidity		55-70%	
Soil Temperature		18 to 25 °C	
Sunshine Hour		6 to 8 hrs. (at least)	
Favorable environmental conditions for insects and diseases			
Fruit fly		Temperature: 25-30°C,RH: 60-91%	
Pumpkin beetle		Temperature: 25-32°C,RH: 65-80%	
Aphid		Temperature: 22-30°C,RH:60-85%	
Powdery mildew		Temperature: 22-27°C,RH: Lower RH during day with higher RH during night	
Downy mildew		Temperature: 25-32°C,RH: 75-95%	
Wilting	Temperature: 22-27°C (Atmosphere), >24°C (Soil) RH: 70-80% (Soil)		
Anthraxnose/Fruit rot		Temperature: 25-32°C,RH: High (89-85%), Higher infestation with continuous rainfall	
Weather warning			
Rainfall (mm)	20 to 30 mm rainfall (creates waterlogging condition)		Early rainfall (15 mm or more)
Cloudy weather	Cloudy condition		
Droughth		Drought due to excessive dry weather	
Duration of staying wet	Land remain waterlogged for 3 to 7 days		
Hailstorm	Seasonal rainfall with hailstorm		Early rainfall with hailstorm
Windspeed (Km/hr.)	≥40 Km/hr		
Temperature (°C)	>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)		

Crop Weather Calendar of **Kharif Sweet Gourd**: Cumilla Region (Districts: Cumilla, Chandpur, B.Barua) Bangladesh

Cumilla Region	Rabi Sweet Gourd									Life Span 110-120 day							
Months	February				March					April				May			
Std. Week/Normal	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Rainfall (mm)	3.0	3.0	6.0	7.5	4.5	7.0	9.0	19.0	27.0	25.5	23.0	33.0	42.0	54.5	63.5	66.0	71.0
Max. Temp. (°C)	16.2	27.3	28.1	29.1	29.8	30.9	32.0	32.2	32.5	32.5	33.0	33.2	33.1	32.9	33.2	33.2	32.9
Min. Temp. (°C)	14.2	14.5	16.2	16.7	18.1	19.2	19.9	21.5	22.0	23.1	23.5	23.7	23.5	23.7	24.3	24.7	25.2
Mean Temp. (°C)	20.2	20.9	22.1	22.9	23.9	25.0	25.9	26.8	27.3	27.8	28.2	28.4	28.3	28.3	28.7	28.9	29.0
RHmax (%)	94.6	94.4	93.9	94.1	93.9	93.9	93.5	93.7	94.0	94.1	93.8	93.9	94.1	94.2	94.0	94.5	94.3
RHMin (%)	47.7	45.2	46.8	46.1	46.8	47.4	47.7	52.6	55.9	58.7	59.6	60.9	61.7	63.0	63.5	64.8	66.7
RHmean (%)	71.1	69.8	70.3	70.1	70.3	70.6	70.6	73.1	74.9	76.4	76.7	77.4	77.9	78.6	78.7	79.6	80.5
SShr (hrs)	47.0	50.0	49.5	50.5	52.5	52.5	52.5	51.0	48.5	51.0	52.0	52.5	51.0	47.5	47.5	46.5	41.0
WD (deg)	203	214	206	214	215	216	211	199	191	194	187	185	179	180	192	183	192
WS(Km/hr)	4.6	4.6	4.6	4.6	4.6	4.8	5.5	5.6	6.8	7.1	7.9	7.7	7.5	7.4	7.7	7.5	6.8
																	
Stage	Seedling raising & planting				vegetative growth				Flowering & fruiting				Harvesting				

Favourable condition			
Temperature	21 to 32 °C	20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)	
Relative Humidity	55-70%		
Soil Temperature	18 to 25 °C		
Sunshine Hour	6 to 8 hrs. (at least)		
Favorable environmental conditions for insects and diseases			
Fruit fly		Temperature: 25-30°C,RH: 60-91%	
Pumpkin beetle		Temperature: 25-32°C,RH: 65-80%	
Aphid		Temperature: 22-30°C,RH:60-85%	
Powdery mildew		Temperature: 22-27°C,RH: Lower RH during day with higher RH during night	
Downy mildew		Temperature: 25-32°C,RH: 75-95%	
Wilting	Temperature: 22-27°C (Atmosphere), >24°C (Soil) RH: 70-80% (Soil)		
Anthracoese/Fruit rot		Temperature: 25-32°C,RH: High (89-85%), Higher infestation with continuous rainfall	
Weather warning			
Rainfall (mm)	20 to 30 mm rainfall (creates waterlogging condition)		Early rainfall (15 mm or more)
Cloudy weather	Cloudy condition		
Droughth		Drought due to excessive dry weather	
Duration of staying wet	Land remain waterlogged for 3 to 7 days		
Hailstorm	Seasonal rainfall with hailstorm		Early rainfall with hailstorm
Windspeed (Km/hr.)	≥40 Km/hr		
Temperature (°C)	>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)		

Crop Weather Calendar of Rabi Sweet Gourd:JashoreRegion (Districts: Jashore, Kushtia, Chuadanga, Jhenaidah, Meherpur, Magura) **Bangladesh**

Region: Jesshore		Rabi Sweet Gourd																Life Span 140-150 day					
Months		November					December				January				February				March				
Std.Week/Normal		44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13
Rainfall (mm)		9.0	8.0	7.5	1.0	2.0	1.5	4.0	0.0	3.5	0.5	1.5	2.5	3.0	6.5	5.0	4.5	8.0	4.0	2.5	4.5	11.5	12.0
Max. Temp. (°C)		31.1	31.1	30.3	29.6	28.5	27.7	26.6	25.7	25.2	24.5	24.5	25.1	25.6	26.6	27.8	28.9	30.0	31.0	32.4	33.6	34.7	34.6
Min. Temp. (°C)		20.7	19.4	18.0	16.7	15.1	14.0	13.4	12.3	11.4	10.9	10.7	11.0	10.8	12.0	13.3	14.7	15.6	16.4	17.7	19.0	20.8	21.8
Mean Temp. (°C)		25.9	25.3	24.2	23.1	21.8	20.9	20.0	19.0	18.3	17.7	17.6	18.0	18.2	19.3	20.5	21.8	22.8	23.7	25.0	26.3	27.7	28.2
RHmax (%)		95.9	95.9	95.7	95.7	95.4	95.2	95.8	95.5	95.7	95.4	95.3	95.3	94.9	94.9	94.2	94.0	93.7	93.2	92.5	92.1	92.1	92.4
RHMin (%)		56.2	52.0	49.8	48.1	46.3	47.2	49.2	49.4	47.9	49.6	47.9	47.0	44.7	44.2	42.2	41.0	39.3	36.5	35.4	34.9	36.7	42.0
RHmean (%)		76.0	74.0	72.8	71.9	70.8	71.2	72.5	72.5	71.8	72.5	71.6	71.2	69.8	69.5	68.2	67.5	66.5	64.9	63.9	63.9	64.4	67.2
SShr (hrs)		45.5	48.5	49.5	47.5	48.5	47.5	43.5	42.0	42.0	41.0	42.5	44.5	48.0	48.5	49.5	50.0	54.5	55.0	54.0	55.0	53.0	51.5
WD (deg)		153	175	177	179	178	184	183	177	174	158	156	188	192	186	191	203	214	219	213	230	212	208
WS(Km/hr)		2.5	2.3	2.4	2.3	2.4	2.3	2.4	2.5	2.5	3.0	3.0	3.0	3.0	3.2	3.1	3.3	3.7	3.5	3.9	4.1	4.4	4.7
																							
Stage		Seedling raising & planting					vegetative growth					Flowering & fruiting					Harvesting						
Favourable condition																							
Temperature		21 to 32 °C					20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																
Relative Humidity		55-70%																					
Soil Temperature		18 to 25 °C																					
Sunshine Hour		6 to 8 hrs. (at least)																					
Favorable environmental conditions for insects and diseases																							
Fruit fly							Temperature: 25-30°C,RH: 60-91%																
Pumpkin beetle							Temperature: 25-32°C,RH: 65-80%																
Aphid							Temperature: 22-30°C,RH:60-85%																
Powdery mildew							Temperature: 22-27°C,RH: Lower RH during day with higher RH during night																
Downy mildew							Temperature: 25-32°C,RH: 75-95%																
Wilting		Temperature: 22-27°C (Atmosphere), >24°C (Soil) RH: 70-80% (Soil)																					
Anthracnose/Fruit rot							Temperature: 25-32°C,RH: High (89-85%), Higher infestation with continuous rainfall																
Weather warning																							
Rainfall (mm)		Early rainfall (15 mm or more)										40 mm or more (creates waterlogging condition)											
Cloudy weather		Cloudy condition																					
Droughth		Drought due to lack of rainfall																					
Duration of staying wet		Land remain waterlogged for 3 to 7 days																					
Hailstorm		Early rainfall and seasonal rainfall with hailstorm																					
Windspeed (Km/hr.)		≥40 Km/hr																					
Temperature (°C)		>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																					

Crop Weather Calendar of Kharif Sweet Gourd:JashoreRegion (Districts: Jashore, Kushtia, Chuadanga, Jhenaidah, Meherpur, Magura) Bangladesh

Region: Jesshore	Kharif Sweet Gourd									Life Span 140-150 day												
Months	April				May					June				July				August				
Std.Week/Normal	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35
Rainfall (mm)	11.5	7.0	12.5	21.5	28.5	30.5	36.0	44.5	38.5	49.5	54.5	81.0	72.5	82.0	72.0	83.5	73.0	63.5	56.0	68.0	48.5	50.5
Max. Temp. (°C)	35.5	36.1	36.5	36.3	36.0	36.0	35.9	35.5	35.6	35.3	34.0	33.3	33.0	32.7	32.9	32.7	32.7	33.0	32.8	32.7	33.1	33.3
Min. Temp. (°C)	22.8	23.5	24.1	24.2	24.2	25.2	25.0	25.5	26.0	26.2	26.2	26.1	26.2	26.0	26.1	26.2	26.2	26.0	26.2	26.0	26.1	26.0
Mean Temp. (°C)	29.2	29.8	30.3	30.3	30.1	30.6	30.4	30.5	30.8	30.7	30.1	29.7	29.6	29.4	29.5	29.5	29.4	29.5	29.5	29.3	29.6	29.6
RHmax (%)	92.4	92.9	92.4	92.3	92.8	92.7	93.1	93.6	94.0	94.1	94.8	95.7	96.0	96.1	96.1	96.1	96.1	96.0	96.2	96.5	96.2	96.2
RHMin (%)	41.5	41.6	44.3	46.9	49.9	51.7	54.8	58.7	60.0	62.1	67.5	71.7	73.5	74.4	73.4	74.6	73.4	72.6	73.0	73.0	71.7	71.0
RHmean (%)	66.9	67.2	68.3	69.6	71.3	72.2	74.0	76.1	77.0	78.1	81.1	83.7	84.7	85.2	84.7	85.3	84.7	84.3	84.6	84.7	83.9	83.6
SShr (hrs)	55.5	54.5	54.5	55.5	54.0	51.5	52.5	46.5	45.5	41.5	32.0	26.5	26.0	24.5	29.0	25.0	29.5	32.5	30.0	28.0	30.5	32.5
WD (deg)	209	204	193	188	186	186	180	181	180	183	174	175	178	186	178	180	177	172	175	175	185	171
WS(Km/hr)	5.1	5.2	5.7	5.6	5.6	5.8	5.2	5.2	5.0	4.6	4.9	4.6	4.6	4.4	4.3	4.4	4.7	4.3	4.4	4.3	3.9	4.1
																						
Stage	Seedling raising & planting				vegetative growth					Flowering & fruiting							Harvesting					
Favourable condition																						
Temperature	21 to 32 °C				20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																	
Relative Humidity					55-70%																	
Soil Temperature					18 to 25 °C																	
Sunshine Hour					6 to 8 hrs. (at least)																	
Favorable environmental conditions for insects and diseases																						
Fruit fly					Temperature: 25-30°C,RH: 60-91%																	
Pumpkin beetle					Temperature: 25-32°C,RH: 65-80%																	
Aphid					Temperature: 22-30°C,RH:60-85%																	
Powdery mildew					Temperature: 22-27°C,RH: Lower RH during day with higher RH during night																	
Downy mildew					Temperature: 25-32°C,RH: 75-95%																	
Wilting					Temperature: 22-27°C (Atmosphere), >24°C (Soil) RH: 70-80% (Soil)																	
Anthrachnose/Fruit rot					Temperature: 25-32°C,RH: High (89-85%), Higher infestation with continuous rainfall																	
Weather warning																						
Rainfall (mm)					Early rainfall (15 mm or more)					40 mm or more (creates waterlogging condition)												
Cloudy weather					Cloudy condition																	
Droughth					Drought due to lack of rainfall																	
Duration of staying wet					Land remain waterlogged for 3 to 7 days																	
Hailstorm					Early rainfall and seasonal rainfall with hailstorm																	
Windspeed (Km/hr.)					≥40 Km/hr																	
Temperature (°C)					>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																	

Crop Weather Calendar of Rabi Sweet Gourd: Khulna Region (Districts: Khulna, Norail, Bagerhat&sathkhira) Bangladesh

Khulna Region			Rabi Sweet Gourd										Life Span 110-120 day						
Months			October		November				December				January				February		
Std.Week/Normal			42	43	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6
Rainfall (mm)			31.5	26.5	9.5	13.0	8.0	3.0	3.5	1.0	4.0	0.0	1.0	3.0	1.5	0.0	3.5	5.5	6.5
Max. Temp. (°C)			32.4	31.7	31.0	31.0	30.1	29.5	28.5	27.7	27.0	26.0	25.6	25.0	25.0	25.8	26.2	27.0	27.7
Min. Temp. (°C)			24.2	22.5	21.5	20.6	19.3	18.0	16.3	15.3	14.5	13.4	12.6	12.2	12.2	12.7	12.4	13.9	14.5
Mean Temp. (°C)			28.3	27.1	26.3	25.8	24.7	23.7	22.4	21.5	20.7	19.7	19.1	18.6	18.6	19.2	19.3	20.4	21.1
RHmax (%)			96.2	96.3	96.0	95.6	95.9	95.8	95.5	94.7	95.7	95.8	95.7	95.4	95.2	94.9	94.9	95.1	94.5
RHMin (%)			63.3	59.2	57.4	53.9	51.3	49.0	46.7	47.3	48.8	49.2	47.8	48.7	46.9	46.8	44.3	44.9	43.4
RHmean (%)			79.8	77.8	76.7	74.7	73.6	72.4	71.1	71.0	72.2	72.5	71.7	72.0	71.0	70.8	69.6	70.0	69.0
SShr (hrs)			49.5	52.5	48.5	53.5	52.5	53.5	53.0	52.0	49.5	48.0	48.5	47.0	46.5	49.5	53.0	52.5	53.5
WD (deg)			193	184	173	176	189	197	197	194	189	181	177	175	182	198	198	190	193
WS(Km/hr)			4.1	3.8	3.4	4.1	3.8	4.3	4.3	4.0	4.5	4.1	.6	4.6	4.7	4.3	4.6	4.9	5.0
																			
Stage			Seedling raising & planting		vegetative growth				Flowering & fruiting				Harvesting						
Favourable condition																			
Temperature		21 to 32 °C				20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)													
Relative Humidity		55-70%																	
Soil Temperature		18 to 25 °C																	
Sunshine Hour		6 to 8 hrs. (at least)																	
Favorable environmental conditions for insects and diseases																			
Fruit fly						Temperature: 25-30°C,RH: 60-91%													
Pumpkin beetle						Temperature: 25-32°C,RH: 65-80%													
Aphid						Temperature: 22-30°C,RH:60-85%													
Powdery mildew						Temperature: 22-27°C,RH: Lower RH during day with higher RH during night													
Downy mildew						Temperature: 25-32°C,RH: 75-95%													
Wilting		Temperature: 22-27°C (Atmosphere), >24°C (Soil) RH: 70-80% (Soil)																	
Anthraxnose/Fruit rot											Temperature: 25-32°C,RH: High (89-85%), Higher infestation with continuous rainfall								
Weather warning																			
Rainfall (mm)		Early rainfall (15 mm or more)										40 mm or more (creates waterlogging condition)							
Cloudy weather		Cloudy condition																	
Droughth		Drought due to lack of rainfall																	
Duration of staying wet		Land remain waterlogged for 3 to 7 days																	
Hailstorm		Early rainfall and seasonal rainfall with hailstorm																	
Windspeed (Km/hr.)		≥40 Km/hr																	
Temperature (°C)		>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																	

Crop Weather Calendar of Kharif Sweet Gourd: Khulna Region (Districts: Khulna, Norail, Bagerhat&sathkhira) Bangladesh

Khulna Region		Kharif Sweet Gourd									Life Span 110-120 day							
Months	February				March					April				May				
Std. Week/Normal	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	
Rainfall (mm)	5.5	6.5	5.0	13.5	6.5	2.5	6.5	13.5	16.0	12.5	12.5	17.5	25.0	31.5	32.5	37.0	47.5	
Max. Temp. (°C)	27.0	27.7	29.0	30.1	31.0	32.4	33.4	34.1	34.0	35.0	35.2	35.2	35.2	35.1	35.2	35.3	35.1	
Min. Temp. (°C)	13.9	14.5	16.2	17.0	18.1	19.4	20.4	22.0	22.7	24.0	24.3	24.9	25.0	25.0	25.3	25.8	26.0	
Mean Temp. (°C)	20.4	21.1	22.6	23.6	24.5	25.9	26.9	28.1	28.4	29.5	29.8	30.1	30.1	30.0	30.2	30.6	30.5	
RHmax (%)	95.1	94.5	94.3	94.3	93.7	93.7	93.5	94.5	94.1	94.3	93.7	93.3	93.2	93.0	93.1	93.4	93.7	
RHMin (%)	44.9	43.4	43.9	41.8	40.7	40.7	40.2	44.2	47.3	48.8	49.2	52.4	53.7	55.3	57.4	57.9	60.7	
RHmean (%)	70.0	69.0	69.1	68.0	67.2	67.2	66.8	69.3	70.7	71.5	71.4	72.8	73.4	74.1	75.2	75.7	77.2	
SShr (hrs)	52.5	53.5	55.5	55.5	57.5	57.5	59.0	57.0	56.5	59.0	60.0	61.5	58.0	59.5	56.5	55.0	47.0	
WD (deg)	190	193	209	218	222	227	233	204	203	202	196	192	186	186	183	177	180	
WS(Km/hr)	4.9	5.0	4.8	5.0	4.9	5.2	5.6	6.3	7.2	7.1	7.6	7.6	8.0	7.5	7.5	7.1	7.2	
																		
Stage	Seedling raising & planting				vegetative growth				Flowering & fruiting				Harvesting					
Favourable condition																		
Temperature	21 to 32 °C				20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)													
Relative Humidity	55-70%																	
Soil Temperature	18 to 25 °C																	
Sunshine Hour	6 to 8 hrs. (at least)																	
Favorable environmental conditions for insects and diseases																		
Fruit fly					Temperature: 25-30°C,RH: 60-91%													
Pumpkin beetle					Temperature: 25-32°C,RH: 65-80%													
Aphid					Temperature: 22-30°C,RH:60-85%													
Powdery mildew					Temperature: 22-27°C,RH: Lower RH during day with higher RH during night													
Downy mildew					Temperature: 25-32°C,RH: 75-95%													
Wilting	Temperature: 22-27°C (Atmosphere), >24°C (Soil) RH: 70-80% (Soil)																	
Anthracnose/Fruit rot					Temperature: 25-32°C,RH: High (89-85%), Higher infestation with continuous rainfall													
Weather warning																		
Rainfall (mm)	Early rainfall (15 mm or more)									40 mm or more (creates waterlogging condition)								
Cloudy weather	Cloudy condition																	
Droughth	Drought due to lack of rainfall																	
Duration of staying wet	Land remain waterlogged for 3 to 7 days																	
Hailstorm	Early rainfall and seasonal rainfall with hailstorm																	
Windspeed (Km/hr.)	≥40 Km/hr																	
Temperature (°C)	>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)																	

Crop Weather Calendar of Rabi Sweet Gourd:RajshahiRegion (Districts:Rajshahi, Chapainawabgonj, Naoga, Natore) Bangladesh

RajshahiRegion	Rabi Sweet Gourd									Life Span 140-150 day													
Months	November					December				January				February				March					APRIL
Std. Week/Normal	৪৪	৪৫	৪৬	৪৭	৪৮	৪৯	৫০	৫১	৫২	১	২	৩	৪	৫	৬	৭	৮	৯	১০	১১	১২	১৩	১৪
Rainfall (mm)	৭.০	৫.০	১.০	০.০	২.০	১.০	৩.০	০.০	৪.০	১.০	১.০	২.০	১.০	৪.০	৩.০	২.০	৩.০	৫.০	৪.০	৩.০	৪.০	১৪.০	১৩.০
Max. Temp. (°C)	৩১.৭	৩১.০	২৯.৬	২৮.৯	২৮.৩	২৭.০	২৬.০	২৫.৪	২৪.৪	২৪.০	২৩.০	২৪.০	২৪.৯	২৫.৬	২৬.৭	২৮.১	২৯.৩	৩০.৪	৩২.১	৩৩.১	৩৪.৬	৩৫.০	৩৬.১
Min. Temp. (°C)	২৩.১	২১.৪	১৭.৬	১৬.৬	১৫.০	১৪.০	১৩.১	১২.৪	১১.৬	১১.০	১০.৯	১০.৯	১০.৯	১২.০	১২.৩	১৩.৬	১৪.১	১৫.১	১৬.৪	১৭.৬	১৯.৪	২০.৬	২২.১
Mean Temp. (°C)	২৭.৪	২৬.২	২৩.৬	২২.৮	২১.৭	২০.৫	১৯.৬	১৮.৯	১৮.০	১৭.৫	১৭.৪	১৭.৫	১৭.৯	১৮.৮	১৯.৫	২০.৯	২১.৭	২২.৮	২৪.৩	২৫.৪	২৭.০	২৭.৮	২৯.১
RHmax (%)	৯৬.১	৯৬.০	৯৫.৭	৯৫.৩	৯৪.৭	৯৪.৭	৯৫.৯	৯৫.৩	৯৫.৭	৯৫.৬	৯৫.১	৯৫.৩	৯৫.৩	৯৫.৩	৯৪.০	৯৩.৯	৯২.০	৯১.৬	৯০.৩	৮৮.৭	৮৮.৩	৮৯.১	৮৮.৯
RHMin (%)	৬০.৯	৫৭.৪	৪৯.৪	৪৭.৪	৪৫.৬	৪৬.৯	৫০.১	৫০.০	৫০.০	৫০.১	৪৮.৪	৪৭.৯	৪৫.৭	৪৪.৭	৪২.০	৪০.৩	৩৬.৬	৩৪.৬	৩২.৭	৩০.৪	৩০.৪	৩৫.৯	৩৪.৩
RHmean (%)	৭৮.৫	৭৬.৭	৭২.৬	৭১.৪	৭০.২	৭০.৮	৭৩.০	৭২.৭	৭২.৯	৭২.৯	৭১.৬	৭১.৬	৭০.৫	৭০.০	৬৮.০	৬৭.১	৬৪.৩	৬৩.১	৬১.৫	৫৯.৪	৫৯.৪	৬২.৫	৬১.৬
SShr (hrs)	৫৩.০	৫৬.০	৫৪.০	৫৫.০	৫৫.০	৫১.০	৪৭.০	৪৬.০	৪৪.০	৪২.০	৪২.০	৪৬.০	৫২.০	৫১.০	৫৫.০	৫৬.০	৫৮.০	৫৯.০	৫৯.০	৫৯.০	৫৯.০	৫২.০	৫৭.০
WD (deg)	১৪৬	১৪৬	১৫১	১৪৬	১৪৫	১৪৩	১৪৬	১৪৪	১৬৭	১৪৬	১৬১	১৭৪	১৯৬	১৯৮	১৯৫	১৯৭	২১৫	২০৯	২২০	২১৪	২০৮	১৮৬	১৮৫
WS(Km/hr)	৩.৯	৪.৩	৩.৭	৩.৭	৩.৭	৩.৭	৩.৭	৩.৭	৩.৭	৩.৯	৩.৭	৩.৭	৪.৪	৩.৯	৫.০	৩.৯	৫.০	৫.০	৫.৪	৫.৪	৫.৬	৫.৬	৬.১
																							
Stage	অঙ্কুরোদগম ও চারা রোপণ				অঙ্গজ বৃদ্ধি				ফুল আসা ও ফল ধারণ				ফসল উত্তোলন										

Favourable condition		
Temperature	21 to 32 °C	20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)
Relative Humidity	55-70%	
Soil Temperature	18 to 25 °C	
Sunshine Hour	6 to 8 hrs. (at least)	
Favorable environmental conditions for insects and diseases		
Fruit fly		Temperature: 25-30°C,RH: 60-91%
Pumpkin beetle		Temperature: 25-32°C,RH: 65-80%
Aphid		Temperature: 22-30°C,RH:60-85%
Powdery mildew		Temperature: 22-27°C,RH: Lower RH during day with higher RH during night
Downy mildew		Temperature: 25-32°C,RH: 75-95%
Wilting	Temperature: 22-27°C (Atmosphere), >24°C (Soil) RH: 70-80% (Soil)	
Anthraxnose/Fruit rot		Temperature: 25-32°C,RH: High (89-85%), Higher infestation with continuous rainfall
Weather warning		
Rainfall (mm)	Early rainfall (15 mm or more)	40 mm or more (creates waterlogging condition)
Cloudy weather	Cloudy condition	
Droughth	Drought due to lack of rainfall	
Duration of staying wet	Land remain waterlogged for 3 to 7 days	
Hailstorm	Early rainfall and seasonal rainfall with hailstorm	
Windspeed	>40 Km/hr	

(Km/hr.)	
Temperature (°C)	>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)

Crop Weather Calendar of Rabi Sweet Gourd: Rajshahi Region (Districts:Rajshahi, Chapainawabgonj, Naoga, Natore) Bangladesh

RajshahiRegion	Rabi Sweet Gourd													Life Span 140-150 day							
Months	এপ্রিল (চৈত্র-বৈশাখ)				মে (বৈশাখ-জ্যৈষ্ঠ)					জুন (জ্যৈষ্ঠ-আষাঢ়)				জুলাই (আষাঢ়-শ্রাবণ)				আগস্ট (শ্রাবণ-ভাদ্র)			
Std.Week/Nom al	১৪	১৫	১৬	১৭	১৮	১৯	২০	২১	২২	২৩	২৪	২৫	২৬	২৭	২৮	২৯	৩০	৩১	৩২	৩৩	৩৪
Rainfall (mm)	১৩.০	১০.০	১৯.০	১৭.০	২৪.০	২৯.০	৩৩.০	৩৬.০	৩৪.০	৪৮.০	৫১.০	৭৮.০	৭২.০	৬৬.০	৭১.০	৭৫.০	৭২.০	৫২.০	৫৭.০	৬৪.০	৫৩.০
Max. Temp. (°C)	৩৬.১	৩৬.৭	৩৬.৪	৩৫.৯	৩৫.৪	৩৫.৩	৩৫.১	৩৫.০	৩৫.৩	৩৫.০	৩৩.৯	৩৩.১	৩৩.০	৩২.৭	৩২.৯	৩২.৩	৩২.৬	৩২.৯	৩২.৯	৩২.৭	৩৩.০
Min. Temp. (°C)	২২.১	২৩.০	২৩.৪	২৩.৯	২৩.৭	২৪.৩	২৪.৬	২৫.০	২৫.৪	২৫.৯	২৬.০	২৬.০	২৬.০	২৬.০	২৬.০	২৬.০	২৬.০	২৬.০	২৬.১	২৬.০	২৬.১
Mean Temp. (°C)	২৯.১	২৯.৯	২৯.৯	২৯.৯	২৯.৬	২৯.৮	২৯.৯	৩০.০	৩০.৪	৩০.৫	৩০.০	২৯.৬	২৯.৫	২৯.৪	২৯.৫	২৯.২	২৯.৩	২৯.৫	২৯.৫	২৯.৪	২৯.৬
RHmax (%)	৮৮.৯	৮৯.০	৯০.১	৯০.৭	৯২.১	৯২.৭	৯৩.৩	৯৩.১	৯৩.৭	৯৪.৬	৯৫.৪	৯৬.১	৯৬.৬	৯৬.৬	৯৬.৯	৯৭.০	৯৬.৯	৯৬.৭	৯৬.৬	৯৬.৭	৯৬.১
RHMin (%)	৩৪.৩	৩৪.৬	৪১.৪	৪৫.১	৫০.০	৫১.৯	৫৫.৬	৫৮.৩	৫৮.৯	৬২.৬	৬৭.৯	৭১.১	৭৩.৪	৭৩.৭	৭৪.৩	৭৫.৩	৭৩.৯	৭২.৬	৭২.৬	৭৪.১	৭২.০
RHmean (%)	৬১.৬	৬১.৮	৬৫.৮	৬৭.৯	৭১.১	৭২.৩	৭৪.৫	৭৫.৭	৭৬.৩	৭৮.৬	৮১.৭	৮৩.৬	৮৫.০	৮৫.২	৮৫.৬	৮৬.২	৮৫.৪	৮৪.৭	৮৪.৬	৮৫.৪	৮৪.১
SShr (hrs)	৫৭.০	৫৭.০	৫৭.০	৫৬.০	৫২.০	৫৩.০	৫৪.০	৪৮.০	৫১.০	৪৮.০	৩৫.০	৩১.০	৩১.০	৩০.০	৩০.০	৩০.০	৩৪.০	৩৪.০	৩৬.০	৩৩.০	৩৪.০
WD (deg)	১৮.৫	১৮.৫	২৬.৭	১৪.৭	১৩.৬	১৩.৬	১৩.৫	১২.৬	১৩.৩	১৩.৯	১২.৯	১২.৭	১৩.২	১৪.১	১৩.৪	১৩.৮	১৪.০	১৩.১	১৩.১	১৪.১	১৪.৩
WS(Km/hr)	৬.১	৬.১	৮.২	৭.৪	৭.৪	৭.২	৭.২	৭.২	৭.২	৬.৭	৬.৭	৬.৮	৬.৮	৫.৭	৬.১	৬.৩	৬.৭	৬.৩	৫.৬	৫.৭	৫.৬
																					
Stage	অঙ্কুরোদগম ও চারা রোপণ				অঙ্গজ বৃদ্ধি					ফুল আসা ও ফল ধারণ				ফসল উত্তোলন							

Favourable condition			
Temperature	21 to 32 °C	20 to 25°C (During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)	
Relative Humidity	55-70%		
Soil Temperature	18 to 25 °C		
Sunshine Hour	6 to 8 hrs. (at least)		
Favorable environmental conditions for insects and diseases			
Fruit fly		Temperature: 25-30°C,RH: 60-91%	
Pumpkin beetle		Temperature: 25-32°C,RH: 65-80%	
Aphid		Temperature: 22-30°C,RH:60-85%	
Powdery mildew		Temperature: 22-27°C,RH: Lower RH during day with higher RH during night	
Downy mildew		Temperature: 25-32°C,RH: 75-95%	
Wilting	Temperature: 22-27°C (Atmosphere), >24°C (Soil) RH: 70-80% (Soil)		
Anthracnose/Fruit rot			Temperature: 25-32°C,RH: High (89-85%), Higher infestation with continuous rainfall
Weather warning			
Rainfall (mm)	Early rainfall (15 mm or more)		40 mm or more (creates waterlogging condition)
Cloudy weather	Cloudy condition		
Droughth	Drought due to lack of rainfall		
Duration of staying wet	Land remain waterlogged for 3 to 7 days		
Hailstorm	Early rainfall and seasonal rainfall with hailstorm		
Windspeed (Km/hr.)	≥40 Km/hr		
Temperature (°C)	>35° C(During cultivation period high temperature and long day length increase no. of male flowers and reduce no. of female flowers)		