

Crop Weather Calendar of Rabi: Onion, Bogra Region (Districts: Bogra, Pabna, Sirajgonj, Joypurhat) Bangladesh

Bogra Region	Rabi:Onion																	
	Life span 100-120 days																	
Months	August				September					October				November				
Std.Week	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48
Rainfall (mm)	63.0	61.0	72.0	67.0	56.0	60.0	60.0	76.0	74.0	51.0	42.0	25.0	18.0	5.0	5.0	0.0	0.0	3.0
Max. Temp. (°C)	32.3	32.6	32.1	32.3	32.6	32.3	32.0	32.1	32.0	32.1	32.3	32.0	31.6	31.0	30.7	30.1	29.7	28.9
Min. Temp. (°C)	26.3	26.9	26.3	26.4	26.1	26.0	26.0	26.0	25.1	25.0	24.0	23.4	21.7	21.0	19.9	18.3	17.3	15.7
Mean Temp. (°C)	29.3	29.8	29.2	29.4	29.4	29.2	29.0	29.1	28.6	28.6	28.2	27.7	26.7	26.0	25.3	24.2	23.5	22.3
RHmax (%)	95.1	95.3	95.6	95.9	95.7	95.9	96.3	96.1	96.4	96.0	96.0	95.7	95.1	94.3	94.6	94.1	93.9	93.9
RHMin (%)	72.0	71.9	73.3	72.7	71.7	72.0	72.3	70.7	71.0	68.7	66.0	62.1	57.9	55.6	52.4	49.9	48.3	47.0
RHmean (%)	83.6	83.6	84.5	84.3	83.7	84.0	84.3	83.4	83.7	82.4	81.0	78.9	76.5	75.0	73.5	72.0	71.1	70.5
SShr (hrs)	35.0	37.0	36.0	31.0	35.0	35.0	35.0	37.0	37.0	44.0	48.0	54.0	57.0	52.0	55.0	55.0	56	56
WD (deg)	137	136	141	147	142	147	152	160	174	170	184	199	198	186	200	202	205	217
WS(Km/hr)	6.3	5.7	6.1	5.6	5.6	5.6	5.6	5.6	5.4	5.0	5.0	5.0	4.4	4.4	3.7	3.7	3.7	3.7
																		
Stages	Germination and Seedling Planting				Vegetative Growth					Bulb formation				Harvesting				
Favorable environment																		
Temperature	10°C to 30°C				The optimum temperature for vegetative phase is 13- 24°C					The optimum temperature for bulb development is15-25°C				Dry and clear weather is more suitable				
Relative Humidity	Requires about 70% relative humidity for good growth																	
Soil Temperature	A wide range of soil temperatures from 20°C to 35 °C, with the optimum at 24°C																	
Sunshine Hour	9 to 10 hours									10 to 12 hours								
Favorable environmental conditions for insects and diseases																		
Thrips	Hot and dry weather increases Trips population; Heavy rainfall reduces population; RH:47.6% to 78.62%																	
Common cut worm	Favored by low temperature and high relative humidity; The upper development threshold temperature for all stages is 37 °C; RH: 50% to 90%																	
Purple blotch	Warm temperature (21°C to 30 °C); High humidity (80% to 90%); Moderate to heavy frequent rainfall and heavy dew during growing stage causes prolonged leaf wetness (12 hours or more) favored disease development.																	
Basal bulb rot	Atmospheric temperature: 25°C to 32°C; Soil temperature: 25°C to 28 °C along with high soil moisture; relative humidity above 70% and Dry atmospheric conditions																	
Weather warning																		
Rainfall (mm)	Heavy-rainfall events in short spells result in surface water stagnation (5 to 10 mm per day)																	
Cloudy weather										Cloudy weather								
Drought										7 to 10 days								
Duration of stay wet (day)	3 to 5 days									2 to 3 days								
Hailstorms	Hailstorms with seasonal rainfall																	
Wind speed (Km/hr.)	≥ 16 km/hr																	
Temperature (°C)	Above 35°C																	


Crop Weather Calendar of Rabi Onion: Chottogram Region (Districts: Chottogram, Cox's Bazar, Noakhali, Feni, Laxmipur) Bangladesh

Chottogram Region	Rabi Onion												Life span 100-120 day					
Months	December				January				February				March					
Std.Week/Normal	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13	
Rainfall (mm)	1.0	4.0	2.3	2.0	0.3	0.7	0.7	1.3	2.0	1.3	2.3	8.3	4.7	5.3	6.3	13.3	21.3	
Max. Temp. (°C)	28.6	27.8	26.9	26.6	26.5	26.1	26.6	26.9	27.7	28.7	29.3	29.8	30.6	31.2	32.0	32.0	31.8	
Min. Temp. (°C)	17.0	16.3	15.3	14.9	14.1	13.9	14.1	14.1	14.9	15.8	16.9	17.6	18.8	19.8	20.7	21.8	22.7	
Mean Temp. (°C)	22.8	22.1	21.1	20.8	20.3	20.0	20.4	20.5	21.3	22.2	23.1	23.7	24.7	25.5	26.3	26.9	27.2	
RHmax (%)	94.4	94.2	94.4	94.1	93.8	93.1	92.9	91.6	91.0	90.6	90.4	91.0	90.8	91.1	91.3	93.0	93.0	
RHMin (%)	49.5	50.0	50.3	49.1	48.2	47.1	44.9	42.3	43.6	41.2	43.3	43.9	44.8	46.6	47.4	54.0	58.2	
RHmean (%)	71.9	72.1	72.3	71.6	71.0	70.1	68.9	67.0	67.3	65.9	66.8	67.5	67.8	68.8	69.4	73.5	75.6	
SShr (hrs)	54.8	52.8	50.8	51.3	53.3	50.8	53.8	56.5	56.3	57.8	57.5	58.5	60.3	59.3	56.8	57.0	52.8	
WD (deg)	204	196	199	207	200	193	203	203	201	203	208	214	221	215	223	218	207	
WS(Km/hr)	5.1	5.7	5.8	6.2	6.6	6.9	6.8	7.0	7.1	7.3	7.7	7.9	8.6	8.7	8.9	9.4	9.7	
																		
Stage	Seedling raising & Transplanting				vegetative growth				Flowering & fruiting				Harvesting					
Favorable environment																		
Temperature	10°C to 30°C				The optimum temperature for vegetative phase is 13- 24°C				The optimum temperature for bulb development is15-25°C				Dry and clear weather is more suitable					
Relative Humidity	Requires about 70% relative humidity for good growth																	
Soil Temperature	A wide range of soil temperatures from 20°C to 35 °C, with the optimum at 24°C																	
Sunshine Hour	9 to 10 hours				9 to 10 hours				10 to 12 hours									
Favorable environmental conditions for insects and diseases																		
Thrips	Hot and dry weather increases Trips population; Heavy rainfall reduces population; RH:47.6% to 78.62%																	
Common cut worm	Favored by low temperature and high relative humidity; The upper development threshold temperature for all stages is 37 °C; RH: 50% to 90%																	
Purple blotch	Warm temperature (21°C to 30 °C); High humidity (80% to 90%); Moderate to heavy frequent rainfall and heavy dew during growing stage causes prolonged leaf wetness (12 hours or more) favored disease development.																	
Basal bulb rot	Atmospheric temperature: 25°C to 32°C; Soil temperature: 25°C to 28 °C along with high soil moisture; relative humidity above 70% and Dry atmospheric conditions																	
Weather warning																		
Rainfall (mm)	Heavy-rainfall events in short spells result in surface water stagnation (5 to 10 mm per day)																	
Cloudy weather													Cloudy weather					
Drought									7 to 10 days									
Duration of stay wet (day)	3 to 5 days								2 to 3 days									
Hailstorms	Hailstorms with seasonal rainfall																	
Wind speed (Km/hr.)	≥ 16 km/hr																	
Temperature (°C)	Above 35°C																	


Crop Weather Calendar of Rabi Onion: CumillaRegion (Districts: Cumilla, Chandpur, B.Baria) Bangladesh

[illegible]


Crop Weather Calendar of Kharif Onion: Cumilla Region (Districts: Cumilla, Chandpur, B.Baria) Bangladesh

Crop Weather Calendar of Kharif Onion (District: Cumilla, Chandrapur, D.Dhaka, Bangladesh)																	
Cumilla Region	Kharif Onion						Life span: 100-110 days										
Months	July				August			September				October					
Std.Week	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43
Rainfall (mm)	96.0	88.0	107.0	95.0	84.0	66.5	82.5	73.0	66.0	53.0	66.5	68.0	53.0	51.0	46.5	30.5	16.0
Max. Temp. (°C)	31.2	31.5	31.3	31.5	32.0	32.0	31.9	32.0	32.0	31.8	32.0	32.0	32.0	32.3	31.9	31.8	31.2
Min. Temp. (°C)	26.0	26.0	25.7	26.0	25.9	26.0	26.0	26.0	25.9	25.7	25.6	25.6	25.6	25.4	24.7	24.3	22.8
Mean Temp. (°C)	28.6	28.7	28.5	28.7	28.9	29.0	28.9	29.0	28.9	28.8	28.8	28.8	28.8	28.8	28.3	28.0	27.0
RHmax (%)	95.4	95.0	95.5	95.3	94.9	94.8	94.9	95.3	95.0	95.2	95.5	95.4	95.6	95.6	95.7	95.5	95.2
RHMin (%)	76.8	75.9	76.6	75.2	74.2	73.9	74.8	74.0	73.4	73.5	73.1	72.2	71.1	70.2	68.1	65.8	61.9
RHmean (%)	86.1	85.4	86.0	85.2	84.5	84.3	84.9	84.6	84.2	84.3	84.3	83.8	83.3	82.9	81.9	80.6	78.5
SShr (hrs)	23.5	30.0	26.5	31.0	30.0	31.5	32.0	30.5	34.5	32.0	33.0	33.5	35.0	39.0	43.0	44.5	50.5
WD (deg)	179	174	174	179	178	182	177	182	179	186	191	197	201	203	214	201	187
WS(Km/hr)	7.3	7.0	7.5	6.7	6.7	6.2	6.6	5.8	5.6	5.3	5.7	4.7	4.7	4.6	4.3	4.3	4.0
																	
Stages	Germination and Seedling Planting				Vegetative Growth			Bulb formation					Harvesting				
Favorable environment																	
Temperature	10°C to 30°C				The optimum temperature for vegetative phase is 13- 24°C			The optimum temperature for bulb development is 15-25°C					Dry and clear weather is more suitable				
Relative Humidity	Requires about 70% relative humidity for good growth																
Soil Temperature	A wide range of soil temperatures from 20°C to 35 °C, with the optimum at 24°C																
Sunshine Hour	9 to 10 hours				9 to 10 hours			10 to 12 hours									
Favorable environmental conditions for insects and diseases																	
Thrips	Hot and dry weather increases Trips population; Heavy rainfall reduces population; RH:47.6% to 78.62%																
Common cut worm	Favored by low temperature and high relative humidity; The upper development threshold temperature for all stages is 37 °C; RH: 50% to 90%																
Purple blotch	Warm temperature (21°C to 30 °C); High humidity (80% to 90%); Moderate to heavy frequent rainfall and heavy dew during growing stage causes prolonged leaf wetness (12 hours or more) favored disease development.																
Basal bulb rot	Atmospheric temperature: 25°C to 32°C; Soil temperature: 25°C to 28 °C along with high soil moisture; relative humidity above 70% and Dry atmospheric conditions																
Weather warning																	
Rainfall (mm)	Heavy-rainfall events in short spells result in surface water stagnation (5 to 10 mm per day)																
Cloudy weather	Cloudy weather																
Drought	7 to 10 days																
Duration of stay wet (day)	3 to 5 days																
Hailstorms	Hailstorms with seasonal rainfall																
Wind speed (Km/hr.)	≥ 16 km/hr																
Temperature (°C)	Above 35°C																


Crop Weather Calendar of Rabi Onion: **Dhaka Region** (Districts: Dhaka, Tangail, Gazipur, Narsingdi, Narayanganj, Munshiganj, Manikgonj) **Bangladesh**

Dhaka Region				Rabi Onion					Life span: 100-110 days															
Months	November				December				January				February								March			
Std.Week	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13			
Rainfall (mm)	10.0	6.0	1.5	4.5	1.5	4.5	1.0	3.5	1.0	0.0	0.0	0.5	3.5	3.5	2.5	12.0	3.5	4.0	9.5	13.5	20.5			
Max. Temp. (°C)	30.6	29.8	29.0	28.3	27.6	26.2	25.5	24.8	24.0	23.9	24.5	25.0	26.2	27.2	28.3	28.9	30.0	31.5	32.3	33.4	33.2			
Min. Temp. (°C)	20.5	18.7	17.7	16.1	15.5	14.7	13.5	12.9	12.4	12.0	12.0	12.1	13.6	14.3	15.5	16.5	17.5	18.6	19.8	19.8	21.8			
Mean Temp. (°C)	25.5	24.3	23.3	22.2	21.5	20.5	19.5	18.8	18.2	17.9	18.3	18.6	19.9	20.7	21.9	22.7	23.7	25.0	26.1	26.1	27.5			
RHmax (%)	95.0	94.7	94.6	94.3	94.8	95.3	95.1	95.0	94.6	94.3	93.8	93.4	92.4	92.1	91.7	91.1	90.0	89.9	89.2	89.6	90.3			
RHMin (%)	51.5	49.3	47.5	46.3	47.6	49.4	49.8	49.1	50.3	48.8	46.2	43.3	42.3	39.4	39.7	38.9	36.0	35.5	35.9	38.9	44.5			
RHmean (%)	73.2	72.0	71.0	70.3	71.2	72.3	72.5	72.1	72.4	71.6	70.0	68.3	67.3	65.7	65.7	65.0	63.0	62.7	62.5	64.3	67.4			
SShr (hrs)	51.5	51.5	52.0	51.5	48.5	46.5	44.5	44.5	42.5	42.5	45.0	49.0	49.5	51.5	53.0	54.5	56.0	55.0	58.0	55.5	52.5			
WD (deg)	189	199	212	211	219	224	225	233	238	238	232	242	241	236	234	228	234	230	223	206	188			
WS(Km/hr)	3.7	3.7	3.7	3.6	3.7	3.7	3.7	3.7	4.1	4.0	4.1	4.3	4.3	4.5	5.0	5.3	5.0	5.4	5.7	6.2	6.5			
																								
Stages	Germination and Seedling Planting				Vegetative Growth				Bulb formation								Harvesting							
Favorable environment																								
Temperature	10°C to 30°C				The optimum temperature for vegetative phase is 13- 24°C				The optimum temperature for bulb development is15- 25°C								Dry and clear weather is more suitable							
Relative Humidity	Requires about 70% relative humidity for good growth																							
Soil Temperature	A wide range of soil temperatures from 20°C to 35 °C, with the optimum at 24°C																							
Sunshine Hour	9 to 10 hours								10 to 12 hours															
Favorable environmental conditions for insects and diseases																								
Thrips	Hot and dry weather increases Trips population; Heavy rainfall reduces population; RH:47.6% to 78.62%																							
Common cut worm	Favored by low temperature and high relative humidity; The upper development threshold temperature for all stages is 37 °C; RH: 50% to 90%																							
Purple blotch	Warm temperature (21°C to 30 °C); High humidity (80% to 90%); Moderate to heavy frequent rainfall and heavy dew during growing stage causes prolonged leaf wetness (12 hours or more) favored disease development.																							
Basal bulb rot	Atmospheric temperature: 25°C to 32°C; Soil temperature: 25°C to 28 °C along with high soil moisture; relative humidity above 70% and Dry atmospheric conditions																							
Weather warning																								
Rainfall (mm)	Heavy-rainfall events in short spells result in surface water stagnation (5 to 10 mm per day)																							
Cloudy weather									Cloudy weather															
Drought	7 to 10 days																							
Duration of stay wet (day)	3 to 5 days								2 to 3 days															
Hailstorms	Hailstorms with seasonal rainfall																							
Wind speed (Km/hr.)	≥ 16 km/hr																							
Temperature (°C)	Above 35°C																							


Crop Weather Calendar of Kharip Onion: **Dhaka Region** (Districts: Dhaka, Tangail, Gazipur, Narsingdi, Narayanganj, Munshiganj, Manikgonj) **Bangladesh**

Dhaka Region	Kharif Onion								Life span: 100-110 days																
Months	April				May				June				July			August									
Std.Week	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34				
Rainfall (mm)	25.0	21.0	31.5	42.5	46.0	55.5	55.0	75.5	70.0	72.5	86.5	81.0	77.5	82.5	73.5	91.0	81.5	67.0	59.5	77.0	62.5				
Max. Temp. (°C)	33.9	34.6	34.1	33.8	33.6	33.7	33.7	33.2	33.5	33.4	32.8	32.2	32.2	32.1	32.0	31.7	32.2	32.2	32.2	31.9	32.3				
Min. Temp. (°C)	22.9	23.5	23.7	23.6	23.7	24.2	24.6	24.9	25.4	25.8	25.9	26.2	26.2	26.1	26.2	26.0	26.0	26.2	26.7	26.2	26.2				
Mean Temp. (°C)	28.4	29.0	28.9	28.7	28.6	28.9	29.2	29.0	29.4	29.6	29.3	29.2	29.2	29.1	29.1	28.9	29.1	29.2	29.4	29.0	29.3				
RHmax (%)	91.3	91.4	91.4	91.7	92.3	92.8	92.9	93.5	93.8	94.0	94.5	94.4	94.7	94.7	94.1	94.5	94.3	94.0	94.1	94.4	94.8				
RHMin (%)	45.9	47.1	52.0	55.1	56.7	58.6	60.4	63.5	65.0	66.7	69.9	71.5	72.9	73.5	73.1	73.5	71.1	71.0	71.6	72.2	71.2				
RHmean (%)	68.6	69.2	71.7	73.4	74.5	75.7	76.7	78.5	79.4	80.3	82.2	83.0	83.8	84.1	83.6	84.0	82.7	82.5	82.8	83.3	83.0				
SShr (hrs)	53.5	54.0	55.0	54.5	51.5	48.0	48.0	42.5	42.0	38.5	31.0	26.0	26.0	27.0	27.5	28.5	32.0	33.5	32.5	31.5	31.5				
WD (deg)	181	175	167	156	159	164	156	155	157	161	157	157	158	162	157	161	159	156	165	157	165				
WS(Km/hr)	6.5	6.6	6.6	6.7	6.8	6.5	6.4	6.1	5.8	5.6	6.3	6.5	6.5	5.8	5.9	6.1	5.9	6.1	5.8	5.6	5.4				
																									
Stages	Germination and Seedling Planting				Vegetative Growth				Bulb formation							Harvesting									
Favorable environment																									
Temperature		10°C to 30°C				The optimum temperature for vegetative phase is 13- 24°C							The optimum temperature for bulb development is15-25°C							Dry and clear weather is more suitable					
Relative Humidity		Requires about 70% relative humidity for good growth																							
Soil Temperature		A wide range of soil temperatures from 20°C to 35 °C, with the optimum at 24°C																							
Sunshine Hour		9 to 10 hours											10 to 12 hours												
Favorable environmental conditions for insects and diseases																									
Thrips		Hot and dry weather increases Trips population; Heavy rainfall reduces population; RH:47.6% to 78.62%																							
Common cut worm		Favored by low temperature and high relative humidity; The upper development threshold temperature for all stages is 37 °C; RH: 50% to 90%																							
Purple blotch		Warm temperature (21°C to 30 °C); High humidity (80% to 90%); Moderate to heavy frequent rainfall and heavy dew during growing stage causes prolonged leaf wetness (12 hours or more) favored disease development.																							
Basal bulb rot		Atmospheric temperature: 25°C to 32°C; Soil temperature: 25°C to 28 °C along with high soil moisture; relative humidity above 70% and Dry atmospheric conditions																							
Weather warning																									
Rainfall (mm)		Heavy-rainfall events in short spells result in surface water stagnation (5 to 10 mm per day)																							
Cloudy weather																	Cloudy weather								
Drought													7 to 10 days												
Duration of stay wet (day)		3 to 5 days																				2 to 3 days			
Hailstorms		Hailstorms with seasonal rainfall																							
Wind speed (Km/hr.)		≥ 16 km/hr																							
Temperature (°C)		Above 35°C																							


Crop Weather Calendar of Rabi Onion: **Dinajpur Region** (Districts:Dinajpur, Thakurgaon, Panchagaor) **Bangladesh**

Dinajpur Region	Rabi Onion								Life span: 100-110 days															
Months	November				December				January				February				March							
Std.Week	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13			
Rainfall (mm)	3.0	0.0	0.0	0.0	0.0	3.0	0.0	4.0	2.0	2.0	3.0	2.0	1.0	2.0	3.0	1.0	1.0	1.0	4.0	2.0	7.0			
Max. Temp. (°C)	29.7	29.0	28.7	28.0	27.0	25.4	24.9	23.9	22.6	22.4	23.0	23.7	25.0	25.9	26.7	27.7	28.6	30.1	31.1	32.0	32.1			
Min. Temp. (°C)	18.3	16.9	15.6	14.0	13.3	12.9	12.0	10.9	10.7	10.3	10.0	10.6	11.1	12.1	13.6	14.1	15.0	16.1	17.1	18.6	19.7			
Mean Temp. (°C)	24.0	23.0	22.2	21.0	20.2	19.2	18.5	17.4	16.7	16.4	16.5	17.2	18.1	19.0	20.2	20.9	21.8	23.1	24.1	25.3	25.9			
RHmax (%)	95.9	95.9	95.3	95.6	96.1	96.4	96.3	96.6	96.7	96.3	96.4	95.9	94.7	94.0	93.4	92.0	90.6	88.9	87.6	86.0	87.1			
RHMin (%)	53.0	51.6	48.1	47.6	49.0	52.9	52.7	54.0	55.1	53.7	52.4	50.4	47.7	43.1	42.4	40.9	38.4	35.3	35.1	36.4	43.3			
RHmean (%)	74.5	73.7	71.7	71.6	72.5	74.6	74.5	75.3	75.9	75.0	74.4	73.1	71.2	68.5	67.9	66.4	64.5	62.1	61.4	61.2	65.2			
SShr (hrs)	56	52.0	54.0	53.0	48.0	42.0	41.0	37.0	34.0	34.0	37.0	41.0	44.0	46.0	52.0	54.0	57.0	59.0	59.0	55.0	49.0			
WD (deg)	144	154	164	177	183	193	200	213	225	238	243	227	241	225	230	225	228	230	212	192	170			
WS(Km/hr)	3.7	3.7	3.7	3.5	2.6	3.7	2.9	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.7	3.8	4.2	5.4	5.6	5.0			
																								
Stages	Germination and Seedling Planting				Vegetative Growth				Bulb formation								Harvesting							
Favorable environment																								
Temperature	10°C to 30°C				The optimum temperature for vegetative phase is 13- 24°C				The optimum temperature for bulb development is15-25°C								Dry and clear weather is more suitable							
Relative Humidity	Requires about 70% relative humidity for good growth																							
Soil Temperature	A wide range of soil temperatures from 20°C to 35 °C, with the optimum at 24°C																							
Sunshine Hour	9 to 10 hours								10 to 12 hours															
Favorable environmental conditions for insects and diseases																								
Thrips	Hot and dry weather increases Trips population; Heavy rainfall reduces population; RH:47.6% to 78.62%																							
Common cut worm	Favored by low temperature and high relative humidity; The upper development threshold temperature for all stages is 37 °C; RH: 50% to 90%																							
Purple blotch	Warm temperature (21°C to 30 °C); High humidity (80% to 90%); Moderate to heavy frequent rainfall and heavy dew during growing stage causes prolonged leaf wetness (12 hours or more) favored disease development.																							
Basal bulb rot	Atmospheric temperature: 25°C to 32°C; Soil temperature: 25°C to 28 °C along with high soil moisture; relative humidity above 70% and Dry atmospheric conditions																							
Weather warning																								
Rainfall (mm)	Heavy-rainfall events in short spells result in surface water stagnation (5 to 10 mm per day)																							
Cloudy weather													Cloudy weather											
Drought	7 to 10 days																							
Duration of stay wet (day)	3 to 5 days												2 to 3 days											
Hailstorms	Hailstorms with seasonal rainfall																							
Wind speed (Km/hr.)	≥ 16 km/hr																							
Temperature (°C)	Above 35°C																							


Crop Weather Calendar of Kharif Onion: **Dinajpur Region** (Districts:Dinajpur, Thakurgaon, Panchagaor) **Bangladesh**

Dinajpur Region		Kharif Onion							Life span: 100-110 days													
Months	April				May				June				July			August						
Std.Week	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	
Rainfall (mm)	8.0	8.0	25.0	23.0	31.0	36.0	57.0	55.0	73.0	57.0	77.0	111.0	96.0	109.0	105.0	81.0	84.0	70.0	60.0	93.0	103.0	
Max. Temp. (°C)	33.0	33.4	33.0	32.4	32.3	32.9	33.0	33.0	33.4	33.3	33.0	32.3	32.4	32.0	32.1	32.1	32.3	32.9	32.6	32.4	32.1	
Min. Temp. (°C)	20.4	21.1	22.0	22.0	22.3	22.9	23.6	24.0	24.4	25.0	25.1	25.6	26.0	26.0	26.0	26.0	26.0	26.3	26.7	26.1	26.0	
Mean Temp. (°C)	26.7	17.3	27.5	27.2	27.3	27.9	28.3	28.5	28.9	29.2	29.1	29.0	29.2	29.0	29.1	29.1	29.2	29.6	29.7	29.3	29.1	
RHmax (%)	86.6	87.4	89.4	91.1	92.3	92.0	92.1	92.4	92.7	93.4	93.9	94.7	94.4	95.1	94.6	94.4	94.7	94.1	94.0	95.0	95.7	
RHMin (%)	40.9	43.4	50.9	53.7	57.6	57.3	59.3	62.4	62.3	65.4	67.7	71.4	71.6	73.3	72.7	71.9	71.6	70.1	70.9	72.6	73.1	
RHmean (%)	63.8	65.4	70.2	72.4	75.0	74.7	75.7	77.4	77.5	79.4	80.8	83.1	83.0	84.2	83.7	83.2	83.2	82.1	82.5	83.8	84.4	
SShr (hrs)	52.0	55.0	49.0	50.0	49.0	51.0	48.0	44.0	42.0	40.0	35.0	27.0	31.0	27.0	29.0	31.0	36.0	36.0	37.0	34.0	34.0	
WD (deg)	171	150	142	123	118	115	119	108	117	111	118	120	121	127	121	126	134	132	132	134	139	
WS(Km/hr)	5.6	4.8	4.8	5.4	3.9	3.7	4.3	3.7	3.9	3.7	3.7	3.7	3.7	3.7	3.7	3.9	3.7	3.9	3.7	3.7	3.7	
																						
Stages	Germination and Seedling Planting				Vegetative Growth				Bulb formation							Harvesting						
Favorable environment																						
Temperature	10°C to 30°C				The optimum temperature for vegetative phase is 13- 24°C				The optimum temperature for bulb development is15-25°C							Dry and clear weather is more suitable						
Relative Humidity	Requires about 70% relative humidity for good growth																					
Soil Temperature	A wide range of soil temperatures from 20˚C to 35 ˚C, with the optimum at 24˚C																					
Sunshine Hour	9 to 10 hours								10 to 12 hours													
Favorable environmental conditions for insects and diseases																						
Thrips	Hot and dry weather increases Trips population; Heavy <i>rainfall reduces population</i> ; RH:47.6% to 78.62%																					
Common cut worm	Favored by low temperature and high relative humidity; The upper development threshold temperature for all stages is 37 ˚C; RH: 50% to 90%																					
Purple blotch	Warm temperature (21°C to 30 ˚C); High humidity (80% to 90%); Moderate to heavy frequent rainfall and heavy dew during growing stage causes prolonged leaf wetness (12 hours or more) favored disease development.																					
Basal bulb rot	Atmospheric temperature: 25°C to 32°C; Soil temperature: 25°C to 28 ˚C along with high soil moisture; relative humidity above 70% and Dry atmospheric conditions																					
Weather warning																						
Rainfall (mm)	Heavy-rainfall events in short spells result in surface water stagnation (5 to 10 mm per day)																					
Cloudy weather									Cloudy weather													
Drought	7 to 10 days																					
Duration of stay wet (day)	3 to 5 days								2 to 3 days													
Hailstorms	Hailstorms with seasonal rainfall																					
Wind speed (Km/hr.)	≥ 16 km/hr																					
Temperature (°C)	Above 35˚C																					

Crop Weather Calendar of Rabi:Onion, Faridpur Region (Districts: Faridpur, Madaripur, Sariatpur, Gopalganj, Rajbari) Bangladesh


Faridpur Region				Rabi:Onion														Life span 100-120 days						
Months	November				December				January				February						March					
Std.Week	44	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13		
Rainfall (mm)	11.5	11.0	8.0	3.0	4.5	0.5	3.5	0.5	2.0	0.5	0.5	0.5	1.5	3.5	5.5	5.0	7.5	5.0	6.5	7.0	12.5	18.5		
Max. Temp. (°C)	30.9	30.4	29.7	29.2	28.0	27.5	26.5	25.5	25.1	24.3	24.4	24.8	25.4	26.5	27.4	28.5	29.5	30.7	31.8	32.9	33.8	33.4		
Min. Temp. (°C)	22.0	20.7	19.1	18.2	16.4	15.2	14.7	13.7	12.9	12.2	12.0	12.1	12.1	13.1	13.9	15.5	16.0	17.3	18.5	19.5	21.2	21.8		
Mean Temp. (°C)	26.4	25.5	24.4	23.7	22.2	21.3	20.6	19.6	19.0	18.3	18.2	18.4	18.7	19.8	20.7	22.0	22.7	24.0	25.1	26.2	27.5	27.6		
RHmax (%)	95.2	95.4	95.5	95.8	95.6	95.6	96.1	96.2	96.0	95.7	95.4	95.5	95.2	94.7	94.7	64.1	93.8	93.4	92.7	92.4	92.9	93.2		
RHMin (%)	57.6	55.5	53.2	51.2	49.9	50.5	51.4	51.3	51.0	52.2	50.9	48.9	46.3	46.2	43.3	43.3	41.5	39.5	38.8	38.4	42.1	46.7		
RHmean (%)	76.4	75.4	74.3	73.5	72.7	73.0	73.7	73.7	73.5	73.9	73.1	72.2	70.7	70.4	69.0	68.7	67.7	66.4	65.7	65.4	67.5	69.9		
SShr (hrs)	48.5	51.0	49.5	50.5	51.0	47.0	44.5	40.5	43.0	41.5	39.5	44.5	50.0	49.0	52.0	52.0	53.0	54.5	53.0	54.5	51.5	50.5		
WD (deg)	202	189	202	201	199	201	197	196	183	191	191	210	204	205	203	214	217	222	230	230	216	200		
WS(Km/hr)	2.8	2.8	2.8	2.8	2.8	2.8	2.8	2.9	3.1	3.4	3.6	3.3	3.4	3.4	3.1	4.0	4.3	4.5	4.5	4.5	5.5	5.6		
																								
Stages	Germination and Seedling Planting				Vegetative Growth				Bulb formation								Harvesting							
Favorable environment																								
Temperature		10°C to 30°C					The optimum temperature for vegetative phase is 13-24°C							The optimum temperature for bulb development is 15-25°C							Dry and clear weather is more suitable			
Relative Humidity		Requires about 70% relative humidity for good growth																						
Soil Temperature		A wide range of soil temperatures from 20°C to 35 °C, with the optimum at 24°C																						
Sunshine Hour		9 to 10 hours										10 to 12 hours												
Favorable environmental conditions for insects and diseases																								
Thrips		Hot and dry weather increases Trips population; Heavy rainfall reduces population; RH:47.6% to 78.62%																						
Common cut worm		Favored by low temperature and high relative humidity; The upper development threshold temperature for all stages is 37 °C; RH: 50% to 90%																						
Purple blotch		Warm temperature (21°C to 30 °C); High humidity (80% to 90%); Moderate to heavy frequent rainfall and heavy dew during growing stage causes prolonged leaf wetness (12 hours or more) favored disease development.																						
Basal bulb rot		Atmospheric temperature: 25°C to 32°C; Soil temperature: 25°C to 28 °C along with high soil moisture; relative humidity above 70% and Dry atmospheric conditions																						
Weather warning																								
Rainfall (mm)		Heavy-rainfall events in short spells result in surface water stagnation (5 to 10 mm per day)																						
Cloudy weather																			Cloudy weather					
Drought		7 to 10 days																						
Duration of stay wet (day)		3 to 5 days																	2 to 3 days					
Hailstorms		Hailstorms with seasonal rainfall																						
Wind speed (Km/hr.)		≥ 16 km/hr																						
Temperature (°C)		Above 35°C																						

Crop Weather Calendar of Rabi Onion: Jessore Region (Districts: Jessore, Chuadanga, Meherpur, Jinaidaha, Magura, Kustia) Bangladesh


Region: Jessore	Rabi Onion										Life cycle of crop, 140-150							
Months	February				March					April				May				
Std.Week	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
Rainfall (mm)	6.5	5.0	4.5	8.0	4.0	2.5	4.5	11.5	12.0	11.5	7.0	12.5	21.5	28.5	30.5	36.0	44.5	38.5
Max. Temp. (°C)	26.6	27.8	28.9	30.0	31.0	32.4	33.6	34.7	34.6	35.5	36.1	36.5	36.3	36.0	36.0	35.9	35.5	35.6
Min. Temp. (°C)	12.0	13.3	14.7	15.6	16.4	17.7	19.0	20.8	21.8	22.8	23.5	24.1	24.2	24.2	25.2	25.0	25.5	26.0
Mean Temp. (°C)	19.3	20.5	21.8	22.8	23.7	25.0	26.3	27.7	28.2	29.2	29.8	30.3	30.3	30.1	30.6	30.4	30.5	30.8
RHmax (%)	94.9	94.2	94.0	93.7	93.2	92.5	92.1	92.1	92.4	92.4	92.9	92.4	92.3	92.8	92.7	93.1	93.6	94.0
RHMin (%)	44.2	42.2	41.0	39.3	36.5	35.4	34.9	36.7	42.0	41.5	41.6	44.3	46.9	49.9	51.7	54.8	58.7	60.0
RHmean (%)	69.5	68.2	67.5	66.5	64.9	63.9	63.9	64.4	67.2	66.9	67.2	68.3	69.6	71.3	72.2	74.0	76.1	77.0
SShr (hrs)	48.5	49.5	50.0	54.5	55.0	54.0	55.0	53.0	51.5	55.5	54.5	54.5	55.5	54.0	51.5	52.5	46.5	45.5
WD (deg)	186	191	203	214	219	213	230	212	208	209	204	193	188	186	186	180	181	180
WS(Km/hr)	3.2	3.1	3.3	3.7	3.5	3.9	4.1	4.4	4.7	5.1	5.2	5.7	5.6	5.6	5.8	5.2	5.2	5.0
																		
Stages	Germination and Seedling Planting				Vegetative Growth					Flowering & Fruiting				Harvesting				
Favorable environment																		
Temperature	10°C to 30°C				The optimum temperature for vegetative phase is 13- 24°C					The optimum temperature for bulb development is15-25°C				Dry and clear weather is more suitable				
Relative Humidity	Requires about 70% relative humidity for good growth																	
Soil Temperature	A wide range of soil temperatures from 20°C to 35 °C, with the optimum at 24°C																	
Sunshine Hour	9 to 10 hours									10 to 12 hours								
Favorable environmental conditions for insects and diseases																		
Thrips	Hot and dry weather increases Trips population; Heavy rainfall reduces population; RH:47.6% to 78.62%																	
Common cut worm	Favored by low temperature and high relative humidity; The upper development threshold temperature for all stages is 37 °C; RH: 50% to 90%																	
Purple blotch	Warm temperature (21°C to 30 °C); High humidity (80% to 90%); Moderate to heavy frequent rainfall and heavy dew during growing stage causes prolonged leaf wetness (12 hours or more) favored disease development.																	
Basal bulb rot	Atmospheric temperature: 25°C to 32°C; Soil temperature: 25°C to 28 °C along with high soil moisture; relative humidity above 70% and Dry atmospheric conditions																	
Weather warning																		
Rainfall (mm)	Heavy-rainfall events in short spells result in surface water stagnation (5 to 10 mm per day)																	
Cloudy weather										Cloudy weather								
Drought										7 to 10 days								
Duration of stay wet (day)	3 to 5 days									2 to 3 days								
Hailstorms	Hailstorms with seasonal rainfall																	
Wind speed (Km/hr.)	≥ 16 km/hr																	
Temperature (°C)	Above 35°C																	

Cumilla Region					Rabi Onion				Life span: 100-110 days												
Months	November				December				January				February				March				
Std.Week	45	46	47	48	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13
Rainfall (mm)	5.0	5.0	0.0	4.0	1.0	3.0	0.0	2.0	0.0	3.0	1.0	0.0	3.0	3.0	6.0	7.5	4.5	7.0	9.0	19.0	27.0
Max. Temp. (°C)	30.3	30.0	28.9	28.3	27.6	26.3	25.4	25.0	24.4	24.0	24.0	25.1	16.2	27.3	28.1	29.1	29.8	30.9	32.0	32.2	32.5
Min. Temp. (°C)	19.7	18.3	17.0	15.4	15.0	14.0	13.3	12.4	12.0	11.6	11.9	12.0	14.2	14.5	16.2	16.7	18.1	19.2	19.9	21.5	22.0
Mean Temp. (°C)	25.0	24.2	23.0	21.9	21.3	20.2	19.4	18.7	18.2	17.8	18.0	18.6	20.2	20.9	22.1	22.9	23.9	25.0	25.9	26.8	27.3
RHmax (%)	95.0	94.6	95.0	95.1	95.3	95.6	95.6	95.3	95.1	94.4	94.4	93.4	94.6	94.4	93.9	94.1	93.9	93.9	93.5	93.7	94.0
RHMin (%)	58.4	55.9	53.0	51.6	52.3	54.1	53.6	53.7	53.6	53.1	51.4	47.0	47.7	45.2	46.8	46.1	46.8	47.4	47.7	52.6	55.9
RHmean (%)	76.7	75.3	74.0	73.4	73.8	74.9	74.9	74.5	74.4	73.8	72.9	70.2	71.1	69.8	70.3	70.1	70.3	70.6	70.6	73.1	74.9
SShr (hrs)	53.0	52.0	53.0	50.0	49.0	43.0	40.0	42.0	41.0	39.0	41.0	45.0	47.0	50.0	49.5	50.5	52.5	52.5	52.5	51.0	48.5
WD (deg)	181	186	186	193	191	206	199	221	230	236	216	238	203	214	206	214	215	216	211	199	191
WS(Km/hr)	3.9	3.7	3.7	3.7	3.0	3.5	3.7	3.7	3.7	3.7	3.7	3.7	4.6	4.6	4.6	4.6	4.6	4.8	5.5	5.6	6.8
Stages	Germination and Seedling Planting				Vegetative Growth				Bulb formation				Harvesting								
Favorable environment																					
Temperature	10°C to 30°C				The optimum temperature for vegetative phase is 13- 24°C				The optimum temperature for bulb development is 15- 25°C				Dry and clear weather is more suitable								
Relative Humidity	Requires about 70% relative humidity for good growth																				
Soil Temperature	A wide range of soil temperatures from 20°C to 35 °C, with the optimum at 24°C																				
Sunshine Hour	9 to 10 hours								10 to 12 hours												
Favorable environmental conditions for insects and diseases																					
Thrips	Hot and dry weather increases Trips population; Heavy rainfall reduces population; RH:47.6% to 78.62%																				
Common cut worm	Favored by low temperature and high relative humidity; The upper development threshold temperature for all stages is 37 °C; RH: 50% to 90%																				
Purple blotch	Warm temperature (21°C to 30 °C); High humidity (80% to 90%); Moderate to heavy frequent rainfall and heavy dew during growing stage causes prolonged leaf wetness (12 hours or more) favored disease development.																				
Basal bulb rot	Atmospheric temperature: 25°C to 32°C; Soil temperature: 25°C to 28 °C along with high soil moisture; relative humidity above 70% and Dry atmospheric conditions																				
Weather warning																					
Rainfall (mm)	Heavy-rainfall events in short spells result in surface water stagnation (5 to 10 mm per day)																				
Cloudy weather									Cloudy weather												
Drought	7 to 10 days																				
Duration of stay wet (day)	3 to 5 days								2 to 3 days												
Hailstorms	Hailstorms with seasonal rainfall																				
Wind speed (Km/hr.)	≥ 16 km/hr																				
Temperature (°C)	Above 35°C																				

Crop Weather Calendar of Rabi Onion: Region Raishahi(Districts:Raishahi, Chapainawabgonj, Naoga, Natore) Bangladesh

Region Rajshahi	Rabi Onion				Life span 100-120 day												
Months	December				January				February				March				
Std.Week/Normal	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13
Rainfall (mm)	1.0	3.0	0.0	4.0	1.0	1.0	2.0	1.0	4.0	3.0	2.0	3.0	5.0	4.0	3.0	4.0	14.0
Max. Temp. (°C)	27.0	26.0	25.4	24.4	24.0	23.9	24.0	24.9	25.6	26.7	28.1	29.3	30.4	32.1	33.1	34.6	35.0
Min. Temp. (°C)	14.0	13.1	12.4	11.6	11.0	10.9	10.9	10.9	12.0	12.3	13.6	14.1	15.1	16.4	17.6	19.4	20.6
Mean Temp. (°C)	20.5	19.6	18.9	18.0	17.5	17.4	17.5	17.9	18.8	19.5	20.9	21.7	22.8	24.3	25.4	27.0	27.8
RHmax (%)	94.7	95.9	95.3	95.7	95.6	95.1	95.3	95.3	95.3	94.0	93.9	92.0	91.6	90.3	88.7	88.3	89.1
RHMin (%)	46.9	50.1	50.0	50.0	50.1	48.4	47.9	45.7	44.7	42.0	40.3	36.6	34.6	32.7	30.4	30.4	35.9
RHmean (%)	70.8	73.0	72.7	72.9	72.9	71.8	71.6	70.5	70.0	68.0	67.1	64.3	63.1	61.5	59.4	59.4	62.5
SShr (hrs)	51.0	47.0	46.0	44.0	42.0	42.0	46.0	52.0	51.0	55.0	56.0	58.0	59.0	59.0	59.0	59.0	52.0
WD (deg)	143	146	144	167	146	161	174	196	198	195	197	215	209	220	214	208	186
WS(Km/hr)	3.7	3.7	3.7	3.7	3.9	3.7	3.7	4.4	3.9	5.0	3.9	5.0	5.0	5.4	5.4	5.6	5.6
																	
Stage	Seedling raising & Transplanting				vegetative growth				Flowering & fruiting				Harvesting				
Favorable environment																	
Temperature	10°C to 30°C				The optimum temperature for vegetative phase is 13- 24°C				The optimum temperature for bulb development is15-25°C				Dry and clear weather is more suitable				
Relative Humidity	Requires about 70% relative humidity for good growth																
Soil Temperature	A wide range of soil temperatures from 20°C to 35 °C, with the optimum at 24°C																
Sunshine Hour	9 to 10 hours								10 to 12 hours								
Favorable environmental conditions for insects and diseases																	
Thrips	Hot and dry weather increases Trips population; Heavy rainfall reduces population; RH:47.6% to 78.62%																
Common cut worm	Favored by low temperature and high relative humidity; The upper development threshold temperature for all stages is 37 °C; RH: 50% to 90%																
Purple blotch	Warm temperature (21°C to 30 °C); High humidity (80% to 90%); Moderate to heavy frequent rainfall and heavy dew during growing stage causes prolonged leaf wetness (12 hours or more) favored disease development.																
Basal bulb rot	Atmospheric temperature: 25°C to 32°C; Soil temperature: 25°C to 28 °C along with high soil moisture; relative humidity above 70% and Dry atmospheric conditions																
Weather warning																	
Rainfall (mm)	Heavy-rainfall events in short spells result in surface water stagnation (5 to 10 mm per day)																
Cloudy weather									Cloudy weather								
Drought									7 to 10 days								
Duration of stay wet (day)	3 to 5 days								2 to 3 days								
Hailstorms	Hailstorms with seasonal rainfall																
Wind speed (Km/hr.)	≥ 16 km/hr																
Temperature (°C)	Above 35°C																

Crop Weather Calendar of Rabi Onion: Rangpur Region (Districts: Rangpur, Lalmonirhat, Nilphamari, Kurigram, Gaibandha) Bangladesh

Rangpur Region	Rabi Onion								Life span 100-120 day									
Months	December				January				February				March					
Std.Week/Normal	49	50	51	52	1	2	3	4	5	6	7	8	9	10	11	12	13	
Rainfall (mm)	0.0	2.5	0.0	2.5	1.0	2.5	2.0	1.0	1.0	1.5	3.5	1.5	2.5	1.0	2.5	5.5	18.5	
Max. Temp. (°C)	26.7	25.1	24.7	24.1	22.7	22.4	23.2	23.4	24.7	25.9	26.7	27.5	28.5	29.9	30.8	31.6	31.4	
Min. Temp. (°C)	14.2	13.5	12.5	11.5	10.9	10.4	10.8	10.8	11.8	12.7	13.7	14.4	15.3	16.1	17.3	18.4	19.5	
Mean Temp. (°C)	20.4	19.3	18.6	17.8	16.8	16.4	17.0	17.0	18.3	19.3	20.2	21.0	21.9	23.0	24.0	25.0	25.4	
RHmax (%)	95.0	95.4	95.8	96.1	96.0	96.0	95.8	95.7	95.5	94.3	93.7	92.9	91.4	89.8	88.7	88.1	87.7	
RHMin (%)	53.5	56.2	56.2	55.2	58.5	57.8	56.2	54.2	51.6	48.0	47.1	45.3	42.8	39.6	40.8	42.5	48.8	
RHmean (%)	74.2	75.8	76.0	75.6	77.3	76.9	76.0	74.9	73.5	71.2	70.4	69.1	67.1	64.7	64.7	65.3	68.3	
SShr (hrs)	49.5	41.5	42.5	40.0	36.0	36.5	39.5	41.0	44.0	48.5	50.0	51.5	56.5	57.5	57.0	54.0	49.0	
WD (deg)	129	158	160	187	199	214	199	210	208	192	201	201	201	207	196	179	152	
WS(Km/hr)	6.0	5.7	5.6	5.6	5.7	5.7	5.7	5.6	6.2	6.6	7.5	7.2	7.7	7.8	8.2	8.7	8.7	
																		
Stage	Seedling raising & Transplanting				vegetative growth				Flowering & fruiting				Harvesting					
Favorable environment																		
Temperature	10°C to 30°C				The optimum temperature for vegetative phase is 13- 24°C				The optimum temperature for bulb development is15-25°C				Dry and clear weather is more suitable					
Relative Humidity	Requires about 70% relative humidity for good growth																	
Soil Temperature	A wide range of soil temperatures from 20°C to 35 °C, with the optimum at 24°C																	
Sunshine Hour	9 to 10 hours								10 to 12 hours									
Favorable environmental conditions for insects and diseases																		
Thrips	Hot and dry weather increases Trips population; Heavy rainfall reduces population; RH:47.6% to 78.62%																	
Common cut worm	Favored by low temperature and high relative humidity; The upper development threshold temperature for all stages is 37 °C; RH: 50% to 90%																	
Purple blotch	Warm temperature (21°C to 30 °C); High humidity (80% to 90%); Moderate to heavy frequent rainfall and heavy dew during growing stage causes prolonged leaf wetness (12 hours or more) favored disease development.																	
Basal bulb rot	Atmospheric temperature: 25°C to 32°C; Soil temperature: 25°C to 28 °C along with high soil moisture; relative humidity above 70% and Dry atmospheric conditions																	
Weather warning																		
Rainfall (mm)	Heavy-rainfall events in short spells result in surface water stagnation (5 to 10 mm per day)																	
Cloudy weather									Cloudy weather									
Drought									7 to 10 days									
Duration of stay wet (day)	3 to 5 days								2 to 3 days									
Hailstorms	Hailstorms with seasonal rainfall																	
Wind speed (Km/hr.)	≥ 16 km/hr																	
Temperature (°C)	Above 35°C																	