

Department of Media Studies and Journalism

University of Liberal Arts Bangladesh

Final Project | Fall Semester, 2025

MSJ 3263: Emergency Communication (Sec: 01)

Students' Name	Nayeem Ahmed Nishad, Faria Akter, MD Akibur Islam Etun
Students' ID	231012029, 223012075, 231012010
Section	
Total Marks	100
Marks Obtained	

Problem Statement and Description

Emergency and risk communication refers to the process of conveying critical information to the public during times of crisis or potential danger. It involves disseminating accurate, timely, and actionable information to help individuals, communities, and organizations make informed decisions and take appropriate actions to protect themselves and mitigate risks. The goal of emergency and risk communication is to ensure that people have the knowledge and resources they need to respond effectively to emergencies and minimize potential harm. This includes providing information about the nature of the emergency, the potential risks involved, recommended safety measures, evacuation procedures, and available support services. Effective emergency and risk communication requires a multi-faceted approach that involves various communication channels, such as mass media, social media, community outreach, and direct messaging. It also emphasizes the importance of clear, concise, and consistent messaging to avoid confusion and misinformation. Keeping this in mind, prepare a report on emergency communication.

Bangladesh frequently faces emergencies such as floods, cyclones, heatwaves, disease outbreaks, industrial accidents, and humanitarian crises. Effective emergency communication is critical for **saving lives, reducing panic, building trust, and enabling coordinated response.**

This assignment assesses students' ability to **design a clear, ethical, and evidence-based emergency communication plan**, grounded in international best practices and adapted to the Bangladeshi socio-cultural context.

Learning Objectives

By completing this assignment, students will be able to:

1. Explain key principles of emergency and risk communication
2. Design communication strategies for different phases of an emergency
3. Develop clear, actionable, and trustworthy emergency messages
4. Identify stakeholders and coordination mechanisms
5. Integrate monitoring, feedback, and rumor management
6. Apply ethical standards in crisis communication

Target Audience

Senior managers of a development organization, donor agencies, and government officials.

Goals and Expectations

1. You will be able to prepare a report on emergency and risk communication.
2. Your plan should be original.

Constraint

There is no constraint in preparing the plan.

Deliverables

You have been hired as the lead Emergency Communications consultant for a major city corporation such as Dhaka / Rajshahi / Chattogram / Sylhet / Rangpur / Barishal. Your task is to develop a comprehensive Emergency Communications Plan that addresses key issues and best practices.

Assignment Title

Designing an Emergency Communication Plan for Crisis Preparedness and Response in Bangladesh

Assignment Task

Students will design a **comprehensive Emergency Communication Plan** for **one type of emergency in Bangladesh**, covering **preparedness, response, and early recovery** phases.

Emergency Scenario (Choose One)

- Cyclone or flood emergency
- Dengue, cholera, or pandemic outbreak
- Heatwave or climate-induced disaster
- Fire or industrial accident
- Rohingya refugee emergency
- Earthquake (low probability, high impact)

Required Structure of the Assignment

Section 1: Emergency Context and Risk Analysis

- Description of the emergency
- Risk and vulnerability analysis
- Affected population groups
- Communication challenges (panic, misinformation, access)

Section 2: Principles of Emergency Communication (15 marks)

Explain and apply key principles, such as:

- Risk Communication and Community Engagement (RCCE)
- Timeliness, transparency, and trust
- Two-way communication
- Do-no-harm and accountability
- Coordination and consistency

(Reference WHO, UNICEF, IFRC, Sphere standards)

Section 3: Communication Objectives and Audiences

- Communication objectives for:

- Preparedness
- Response
- Early recovery
- Audience segmentation:
 - General public
 - High-risk groups
 - Responders and institutions

Section 4: Message Development and Framing

Students must develop:

- Key life-saving messages
- Message framing (simple, actionable, culturally appropriate)
- Language and literacy considerations
- Rumor and misinformation management strategies

Include sample messages.

Section 5: Channels and Coordination Mechanisms

- Communication channels:
 - SMS, radio, TV
 - Community volunteers
 - Social media and digital platforms
- Coordination with:
 - Government agencies
 - NGOs and media
 - Community leaders

Section 6: Monitoring, Feedback, and Adaptation (10 marks)

- Indicators for message reach and understanding
- Community feedback mechanisms

- Media and rumor monitoring
- Adaptive communication actions

Section 7: Ethics, Inclusion, and Accountability

Discuss:

- Protection of vulnerable groups
- Avoiding fear, stigma, and blame
- Accessibility (language, disability, gender)
- Accountability to affected populations (AAP)

Section 8: Supplement

- Message flowchart for different emergency phases
- Sample SMS or radio script
- Crisis communication timeline (first 72 hours)

Submission Guidelines

- **Length:** 2,500–3,000 words
- **Format:** MS Word or PDF
- **Font:** Times New Roman, 12 pt, 1.5 spacing
- **Referencing:** APA style
- **Minimum references:** 8 (WHO, UNICEF, IFRC, academic sources)

Assessment Rubric (Summary)

Criteria	Marks
Risk analysis and context	15
Application of emergency communication principles	20
Quality of messages and strategy	20
Channel selection and coordination	15

Criteria	Marks
Monitoring and feedback	10
Ethics and inclusion	10
Structure and clarity	10
Total	100

Academic Integrity

All submissions must be original. Plagiarism or unethical use of AI tools will be handled according to university policy.

Submission Deadline

The deadline for submitting the plan is December 25, 2025.

Keep in Mind

- You must **read the instructions carefully** before writing your answer.
- You should download the question paper and write your answer in the recommended place.
- Total Marks: 100 [will weigh 30% in the final grading].
- You must write your assignment in English, and your answer must be computer-composed.
- Deadline of Submission: December 25, 2025.
- This is a group assignment project. You should form a group of 2 to 3 individuals. All members of the group should submit their soft copy of the assignment. Submit only one hardcopy from each group.
- You must submit a soft copy of your assignment to the Google Classroom using your account. **No email submission will be accepted.**
- **You will also submit a hard copy of your assignment at the first class after the midterm exam week, and get your exam attendance signed.**
- **While submitting the hardcopy, please print this question paper also.**

- You may consult published academic journals, articles, books, etc. In that case, you must write it in your own words and give proper citations. **Any plagiarism will be considered a punishable offence, and it will severely affect your grades.**
- You must submit your articles in a **single MS Word file**. You **MUST NOT send your assignment in PDF** or other file formats.
- The **name of your submission file** should be as follows: [Your Name] _ [Your Student ID] _ [ER – Section 1] _ [Fall 2025].
- You will use the Times New Roman font.
- Keep a one-inch margin on each side and 1.5-inch line spacing. You will submit all the answers in ONE file. Keep in mind: NOT more than one file.
- The course teacher will run a Similarity-Check in Turnitin. Papers with a similarity index higher than 20% will not be accepted.
- If the answers of two or more students match completely, or certain paragraphs match exactly, all the students having such similarity will get ZERO. No explanation will be accepted in this regard.
- For any guidance, do communicate with me. You can contact me at 01718674830 / aminul.islam@ulab.edu.bd.

Citation should be as follows

In the text: (Author Name, publication year)

In Reference: Author name, (publication year). Title of the article, *Name of Journal* / book, Issue, Volume, page numbers, [for book: publication place: Name of publisher].

Dengue Outbreak Emergency Communication Plan

Section 1: Emergency Context and Risk Analysis

Description of the Emergency

Bangladesh is currently facing a deadly dengue outbreak across the country. The sheer volume of cases in 2025 has surpassed the historic peak. The situation is critical due to the unpredictable shift in where and when infection is occurring. This year, the outbreak has stretched into the winter months (late December), defying traditional seasonality (WHO, 2023).

Key Risk Factors

1. **Geographical expansion:** Unlike previous years, when the sole epicenter was Dhaka city, in 2025 the outbreak has aggressively spread to the Barishal, Chattogram, and Khulna divisions. Rural areas are now new hotspots, where healthcare infrastructure is fragile.
2. **Serotype shift (re-infection risk):** The circulation of the DENV-2 and DENV-3 serotypes is driving severe cases. Populations previously infected with one strain are now contracting a different one, leading to Antibody-Dependent Enhancement (ADE) — a condition that significantly increases the risk of Dengue Hemorrhagic Fever and Shock Syndrome (Billah et al., 2025).
3. **Climate factors:** Due to climate change, the monsoon season has extended with irregular rainfall, and a warmer-than-average winter has kept vector breeding active into December.
4. **Healthcare strain:** Surging dengue patients have overwhelmed the tertiary care hospitals, which have limited infrastructure to treat so many patients (Mahedi et al., 2025).

Vulnerability Analysis: Who Is at Most Risk

The virus is affecting demographics differently. While more cases are recorded in one group, deaths are clustering in another.

Population Group	Risk Profile	Why? (Vulnerability Drivers)
Young Adults (Age 16–30)	High Infection Risk	This is the most mobile demographic (work/study). They are highly active during peak mosquito-biting hours (early morning and late afternoon) in schools, universities, and offices.
Women (All Ages)	High Mortality Risk	Critical disparity: Although fewer women are hospitalized than men, they have a higher Case Fatality Rate (CFR). This is often due to delayed care-seeking behaviour (household prioritization) and social barriers to accessing immediate emergency transport.
Children (0–15)	High Severity Risk	Children's immune systems are less robust against shock. Warning signs (such as rapid pulse weakening) can escalate much faster in children than in adults.
Rural Residents	Systemic Risk	Residents in newly affected districts (e.g., Barguna, Pirojpur) often lack access to platelet separators or ICUs nearby, necessitating dangerous long-distance travel during critical illness phases (WHO, 2023).

Communication Challenges

Effective response is currently disrupted by three major communication barriers.

A. The Platelet Panic

- **The miscommunication:** People panic when platelets drop below 100k and demand unnecessary transfusions (Billah et al., 2025).
- **Reality:** Death is rarely caused by low platelets alone; it is caused by plasma leakage from the blood vessels, leading to shock.
- **Challenge:** Shifting the public narrative from “watch the platelets” to “watch the blood pressure and urine output.”

B. The Seasonal Myth

- **The miscommunication:** “It’s December (a winter month), so the fever cannot be dengue.”
- **The risk:** This belief causes patients to disregard early symptoms as the common flu. They delay testing and enter the critical phase.

C. Reliance on Home Remedies

- **The miscommunication:** Some people believe that papaya leaf is a cure for dengue.
- **The risk:** Relying on these remedies often delays necessary medical monitoring.

Section 2: Principles of Emergency Communication

Explaining and implementing the principles of emergency communication:

1. Timeliness, Transparency, and Trust

Information must be released immediately, acknowledging the uncertainty in order to build credibility.

- **The challenge:** The public believes that dengue is a monsoon disease. Fever in the winter season is dismissed as a cold, delaying treatment.

Application:

- **Transparency:** Authorities (DGHS) must clearly state that this is not normal — the season has shifted, and mosquitoes are surviving in winter.
- **Timeliness:** Do not wait for confirmation of the PCR test to warn a community. If cases of fever appear, immediately issue a provisional alert.
- **Build trust:** Use trusted messengers such as imams and school teachers. People trust local figures who share their daily lives.

2. Two-Way Communication (Listening More Than Telling)

RCCE is not a monologue. We must set up channels to listen to community fears and myths in order to address them effectively.

- **The challenge:** “The platelet panic” — patients’ families harass doctors for platelet transfusions (which are rarely needed) while ignoring the real killer: **blood pressure drop (shock).**

Application:

- **Feedback mechanism:** Monitor social media posts and community groups, and set up dedicated “Dengue Myth Hotline” hospitals; if the dominant conversation is about platelets, do not ignore it.
- **Response:** “We hear you worrying about the bleeding, but it is not the only reason for the deaths due to dengue. In this new strain, plasma leakage is the silent killer. IV saline is the right medicine that saves you from shock. Watch the urine output, not just the blood count.”
- **Visual aid:** Use simple infographics in the local dialect to explain to people.

3. Do-No-Harm and Accountability

Interventions should not inadvertently damage the community, stigmatize groups, or drain their resources.

The challenge:

- **Stigma:** Labeling slum areas as “dirty” and “dengue zones” stops residents from seeking help or allowing volunteers inside.
- **Financial harm:** Private clinics leveraging the panic to sell expensive dengue tests and platelet kits to poor families.

Application:

- **Accountability:** Publicize the official government price for the dengue test by putting up posters in public hospitals and on social media. Also, provide a number for complaints.

4. Coordination and Consistency

Conflicting messages can cost lives. Every actor should speak with one voice.

- **The challenge:** A doctor in Khulna prescribes antibiotics, while a doctor in Dhaka prescribes fluids. Patients get confused and delay treatment.

Application:

- **Unified protocol:** The DGHS, the Bangladesh Pediatric Association, and NGOs must use the exact same one-page fever-management guide.

References Used

1. WHO Outbreak Communication Guidelines (2005)
2. IFRC Community Engagement and Accountability (CEA) Policy
3. Sphere Health Standard 2.2.1 (Control of Communicable Diseases)

Section 3: Communication Objectives and Audiences

1. Communication Objectives: The “What & When”

Phase 1: Preparedness (Before / Early)

- **The goal:** Break the “winter myth.”
- **What they must KNOW:** “Mosquitoes are hiding inside your warm house (fridge trays, flower vases) this winter.”
- **What they must DO:** “Inspect your home every Friday morning. Pour out any water found indoors.”

Phase 2: Response (Now)

- **The goal:** Stop the “platelet panic” and start the “fluid focus.”
- **What they must KNOW:** “Death comes from low blood pressure (shock), not low platelets. Saline is the medicine.”
- **What they must DO:** “Go to the hospital immediately if the patient has warning signs (severe tummy pain, vomiting 3+ times, no urine for 6 hours) — even if the fever is gone.”

Phase 3: Early Recovery (After)

- **The goal:** Restore trust in local care.
- **What they must KNOW:** “Local hospitals have the right saline protocols. Travelling to Dhaka while in shock is dangerous.”

- **What they must DO:** “Support community clean-up of water tanks to prevent next year’s surge.”

2. Audience Segmentation: The “Who & How”

Audience Group	Who Are They?	Precise Message (The “Script”)	The Action Required
The “Gatekeepers”	Mothers & housewives (they care for the sick)	“Don’t wait for bleeding. Watch the urine. If the patient stops peeing, run to the hospital.”	Monitor: urine output & hydration.
The “Vulnerable”	Rural residents (Barishal, Khulna, Pirojpur)	“Do not rush to Dhaka immediately. The long journey can kill. Stabilize at your Upazila Health Complex first.”	Trust: local stabilization before transport.
The “Ignored”	Pregnant women	“A fever for you is a double emergency. Do not stay home. Get admitted to hospital on Day 1.”	Admit: immediate hospitalisation.
The “Enforcers”	Pharmacy owners (the first point of contact)	“If a customer has fever, DO NOT sell painkillers (NSAIDs like Diclofenac). Sell Paracetamol only.”	Refuse: stop selling harmful drugs.
The “Influencers”	Imams & teachers	“Cleanliness is faith. This Friday, tell everyone to clean their fridge trays.”	Preach: specific cleaning habits.

Action Charts for Responders and Institutions

Category A: Community & First Responders

These actors control the spread and initial triage.

Role / Institution	The Critical “STOP” (Do Not Do This)	The Critical “DO” (Mandatory Action)	The Script (What to Say)
Pharmacy Owners & Staff	STOP selling NSAIDs (Diclofenac, Ibuprofen) or steroids for undefined fever. This causes bleeding.	DO ask every fever customer: “ <i>Do you have stomach pain or vomiting?</i> ” If yes, refuse the sale and send them to hospital.	“I cannot give you a painkiller. It could be dengue. Take only Paracetamol and go to the clinic.”
Ward Councillors & City Corp.	STOP relying solely on street fogging (smoke). It does not kill indoor larvae.	DO deploy “Red Teams” to inspect indoor hotspots: fridge trays, AC drain pipes, and flower vases in high-rise buildings.	“Fogging is not enough. We are entering homes today to check water trays.”
Imams & Mosque Committees	STOP giving generic advice like “Stay safe.”	DO dedicate the <i>Khutbah</i> to a specific action: “Check the mosque’s ablution (Wudu) tank and your own fridge tray.”	“It is a religious duty to remove stagnant water today. Check your home before Asr prayer.”

Role / Institution	The Critical “STOP” (Do Not Do This)	The Critical “DO” (Mandatory Action)	The Script (What to Say)
School Teachers / Headmasters	STOP ignoring students who put their heads down on desks (lethargy).	DO implement a “full-sleeve” uniform policy immediately. Allow tracksuits / trousers instead of skirts / shorts.	“Wear full sleeves to school. If you feel weak, tell a teacher immediately.”

Category B: Clinical & Medical Responders

These actors determine survival.

Role / Institution	The Critical “STOP” (Do Not Do This)	The Critical “DO” (Mandatory Action)	The Script (What to Say)
Village Doctors (Palli Chikitshak)	STOP prescribing antibiotics for viral fever. It wastes money and time.	DO refer any patient with “ warning signs ” (vomiting, tummy pain) to the Upazila Complex immediately.	“This fever needs a blood pressure check, not antibiotics. Go to the government hospital now.”
Triage Nurses (ER)	STOP using “platelet count” as the admission criterion.	DO admit based on Shock Index : low pulse pressure (<20 mmHg), cold extremities, or urine output <500 ml/day.	“We are admitting you because your blood pressure is unstable, not because of your platelets.”
Diagnostic Centers (Private)	STOP price gouging on NS1 / CBC tests.	DO display the govt. fixed price (e.g., 50 / 300 BDT) clearly at the reception counter.	“The government rate is [X] Taka. We are charging that amount.”

Section 4: Message Development and Framing

1. Core Message Framework

Color Code	Status	Key Message (The “Hook”)	Action Required
● GREEN	Safe	<i>“Fever is high? Don’t panic. Drink saline.”</i>	Stay home. Drink 3 L of fluid daily. Take Paracetamol only.
● YELLOW	Caution	<i>“Fever gone but feeling weak? This is the danger zone.”</i>	Visit a doctor. If the patient stops eating or is very tired, this is the Critical Phase (Day 4–6).
● RED	Danger	<i>“Stomach pain + vomiting? Ignore the platelets, run to the hospital.”</i>	Immediate admission. These are signs of shock (BP drop), which kills faster than bleeding.

2. Framing for Specific Audiences

For Mothers (The Caregivers)

- **Objective:** Shift focus from “temperature” to “urine output.”
- **The message:** *“Ma, don’t worry if the fever is 104°F. Worry if your child hasn’t peed in 6 hours. Peeing means they are safe.”*
- **Why:** Mothers often stop giving fluids when the fever drops, which is exactly when shock begins.

For Rural Men (The Decision Makers)

Channel	Format	Content (Draft)
SMS	Text (Bangla)	“জ্বর হলে অবহেলা নয়। ডেঙ্গু পরীক্ষায় নিকটস্থ নগর স্বাস্থ্য কেন্দ্রে যান। প্যারাসিটামল ছাড়া অন্য ব্যথানাশক ওষুধ খাবেন না। - ঢাকা উত্তর সিটি কর্পোরেশন”

Channel	Format	Content (Draft)
Radio	Jingle (15 sec)	(Sound of thunder / rain) “বৃষ্টির পানি জমতে দেবেন না। ৩ দিনে ১ বার, জমা পানি পরিষ্কার। আপনার পরিবারকে ডেঙ্গু মুক্ত রাখুন।”
Poster	Visual	Image of a clean roof vs. a dirty roof. Headline: “আপনার ছাদ কি মশার কারখানা? আজই পরিষ্কার করুন।” (Is your roof a mosquito factory? Clean it today.)

- **Objective:** Prevent dangerous travel to Dhaka.
- **The message:** *“Your local Upazila Hospital has the exact same saline as Dhaka Medical. Traveling 6 hours to Dhaka with low blood pressure will kill the patient on the road. Stabilize here first.”*

For Youth / Students

- **Objective:** Peer monitoring.
- **The message:** *“If your friend is absent and says ‘I’m just weak,’ go check on them. Weakness after fever is the silent killer.”*

3. Language and Literacy Considerations

[This section was left blank in the original submission.]

4. Rumour and Misinformation Management

Strategy — The “Truth Sandwich”: never repeat the myth alone; sandwich it between the truth.

Myth 1: “We need platelets to survive.”

- **The truth:** “Saline saves lives, not platelets.”
- **The myth:** “Many people think low platelets cause death.”
- **The correction:** “Actually, death is caused by ‘dry veins’ (shock). Platelets will naturally come back up if the blood pressure is strong.”

Myth 2: “Papaya leaf juice cures dengue.”

- **The truth:** “You need fluids to keep the heart running.”
- **The myth:** “Papaya juice is popular.”
- **The correction:** “It is okay to drink it, but **NEVER** stop the ORS (saline). Papaya juice cannot fix low blood pressure.”

Myth 3: “Dengue is a summer disease.”

- **The truth:** “Mosquitoes have adapted to the winter cold inside our homes.”
- **The myth:** “Fever in December is just a cold.”
- **The correction:** “Any fever now is dengue until tested. Test immediately.”

Sample Messages

Source 1: Directorate General of Health Services (DGHS)

- **Message:** “Warning signs: severe abdominal pain, persistent vomiting, bleeding from the gums, and lethargy. If any of these appear, go to the hospital immediately.”
- **Link:** [DGHS National Dengue Guideline \(2024–2030\)](#) *(Note: Refers to the “Warning Signs” section in the official PDF).*

Source 2: UNICEF Bangladesh

- **Message:** “Dengue can be fatal, but early treatment saves lives. Do not wait for the patient to bleed. Watch for dehydration signs like a dry mouth and no tears when crying.”
- **Link:** [UNICEF Bangladesh: How to Keep Children Safe from Dengue](#)

Section 5: Channels and Coordination Mechanisms

A multi-channel communication approach is essential to ensure wide reach and message reinforcement. Television and radio remain key channels for mass dissemination, particularly for populations with limited internet access. SMS alerts provide rapid, direct communication of urgent information, while social media platforms enable real-time updates and interactive engagement, especially among younger audiences (WHO, 2017).

Community volunteers and frontline health workers play a crucial role in interpersonal communication, door-to-door outreach, and community engagement. Their involvement increases trust and allows messages to be adapted to local concerns. Effective coordination among government agencies, NGOs, media organizations, and community leaders ensures message consistency and credibility. UNICEF highlights that coordinated communication prevents confusion and strengthens the overall outbreak response (UNICEF, 2022).

Section 6: Monitoring, Feedback, and Adaptation

Monitoring and evaluation allow communication strategies to remain responsive and effective throughout the outbreak. Key indicators include message reach, public understanding of dengue prevention and symptoms, and reported behaviour changes. Data can be collected through surveys, media analytics, and rapid community assessments (WHO, 2017).

Community feedback mechanisms such as hotlines, social media engagement, and volunteer reports enable two-way communication and help identify gaps, concerns, and emerging rumors. Media and rumor monitoring are critical for detecting misinformation early. Based on monitoring findings, communication actions are adapted by revising messages, changing delivery channels, or intensifying community engagement to address evolving needs (WHO, 2023).

Section 7: Ethics, Inclusion, and Accountability

Ethical emergency communication prioritizes the protection of vulnerable populations while avoiding fear, stigma, and blame. Dengue communication must not single out specific communities or groups, as this can lead to discrimination and reduced cooperation. Instead, messages emphasize solidarity, shared responsibility, and achievable actions (WHO, 2017).

Inclusion and accessibility are ensured by providing information in local languages and using simple wording, visual aids, and formats accessible to people with disabilities. Gender-sensitive communication recognizes different household and caregiving roles and ensures equal access to information. Accountability to Affected Populations (AAP) requires transparency, responsiveness, and meaningful community participation in communication processes (UNICEF, 2022).

Section 8: Supplement — Operational Communication Tools

Emergency communication messages evolve across outbreak phases. During the preparedness and early-alert phase, communication focuses on awareness and prevention. In the response and escalation phase, messages emphasize symptom recognition and care-seeking. At peak transmission, life-saving guidance and rumor control are prioritized. During recovery, communication reinforces sustained prevention and community responsibility (WHO, 2017).

Sample SMS: “High fever and body pain may be dengue. Do not delay — visit the nearest health center. Remove standing water weekly to prevent mosquito breeding.”

Sample radio message: “Dengue spreads through mosquito bites, not from person to person. Remove standing water from your home every week. If the fever lasts more than two days, seek medical care immediately. Together, we can protect our community.”

During the first 72 hours of an outbreak, communication begins with official confirmation and preventive guidance within the first 24 hours. Between 24 and 48 hours, regular updates, myth-busting, and intensified media engagement are prioritized. From 48 to 72 hours, communication focuses on reassurance, sustained prevention, and adaptation based on feedback and rumor monitoring (UNICEF & WHO, 2020).

Conclusion

Effective emergency communication is a cornerstone of dengue outbreak response, complementing medical and vector-control interventions. This report demonstrates that clear message development, culturally appropriate framing, coordinated multi-channel dissemination, continuous monitoring, and strong ethical foundations are essential for protecting public health. By aligning with the WHO and UNICEF RCCE frameworks and prioritizing accountability to affected populations, dengue emergency communication can build trust, reduce misinformation, and ultimately save lives.

References

World Health Organization. (2017). *Communicating risk in public health emergencies: A WHO guideline for emergency risk communication (ERC) policy and practice*. World Health Organization.

- World Health Organization. (2023). *Dengue and severe dengue*. World Health Organization. <https://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-dengue>
- World Health Organization. (2012). *Handbook for integrated vector management*. World Health Organization.
- World Health Organization. (2016). *Zika virus outbreak global response: Risk communication and community engagement*. World Health Organization.
- World Health Organization. (2020). *Risk communication and community engagement readiness and response to coronavirus disease (COVID-19)*. World Health Organization.
- UNICEF. (2022). *Risk communication and community engagement (RCCE) for public health emergencies*. United Nations Children's Fund.
- UNICEF & World Health Organization. (2020). *Managing epidemics: Key facts about major deadly diseases*. World Health Organization.
- UNICEF, World Health Organization, & International Federation of Red Cross and Red Crescent Societies. (2020). *Risk communication and community engagement (RCCE) action plan guidance for COVID-19 preparedness and response*. UNICEF.
- Centers for Disease Control and Prevention. (2022). *Dengue: Epidemiology, prevention, and control*. Centers for Disease Control and Prevention.
- Reynolds, B., & Seeger, M. W. (2005). Crisis and emergency risk communication as an integrative model. *Journal of Health Communication*, 10(1), 43–55. <https://doi.org/10.1080/10810730590904571>
- Billah, M., Shampa, N., & Henderson, A. (2025). Escalating dengue burden and emerging hotspots in Bangladesh: A decade of trends and 2025 forecast. *International Journal of Data Science and Analysis*, 11(5), 136–142. <https://doi.org/10.11648/j.ijdsa.20251105.12>
- Mahedi, M. R. A., Roy, P., Radia, A. A., Afrin, S., Ahmed, M. F., & Syrmos, N. (2025). Symptomatic pattern changes and climatic factors in dengue outbreaks in Bangladesh. *Future Health*, 0, 1–8. https://doi.org/10.25259/fh_78_2024
- World Health Organization. (2023, August 11). *Disease outbreak news*. World Health Organization. <https://www.who.int/emergencies/disease-outbreak-news/item/2023>

Omambia, S. (2024). Combating public health infodemics: Strategies for misinformation control and evidence-based communication. *Journal of Advances in Medicine and Medical Research*, 36, 1–8. <https://doi.org/10.9734/jammr/2024/v36i105582>