



ULAB
UNIVERSITY OF LIBERAL ARTS
BANGLADESH

Communication for Development (C4D)

Communication Planning and Process

A Communication Strategy for Reducing Childhood Malnutrition in Rural Bangladesh

Prepared by

Faria Akter

Student ID: 223012075

Contents

Phase 1: Analysis and Problem Definition.....	3
Analysis and Problem Definition.....	3
Situation Analysis.....	3
Problem Tree and Fishbone Analysis.....	3
Root Cause Analysis.....	4
References.....	7
Phase 2: C4D Plan Development.....	8
Logical Framework (Log Frame Analysis).....	8
C4D Communication Strategy.....	12
A Shortage of Food and Childhood Malnutrition in Rural Bangladesh.....	12
Conclusion.....	14
Self-Reflection.....	14

Phase 1: Analysis and Problem Definition

Analysis and Problem Definition

Malnutrition in children is still a major problem in rural Bangladesh, especially in the most isolated Upazilas where people face enormous economic and social disadvantages. Right2Grow Consortium Bangladesh reports that a significant portion of the country's children under the age of five suffer from stunting, underweight, or “wasted” disorders. When it comes to nations with the highest rates of severe childhood malnutrition, Bangladesh was placed sixth in 2022 by UNICEF. In light of these disturbing statistics, one must wonder why the government's efforts to combat childhood malnutrition over the last few decades have been so fruitless. According to Hasan et al. (2021), there are many variables that contribute to chronic nutritional deficiencies in children, such as poverty, restricted healthcare access, poor maternal education, and ingrained cultural practices. As a result, these deficiencies worsen stunting and wasting.

Situation Analysis

Maternal and child health outcomes are greatly impacted by the major hurdles to healthcare access in these distant places. Critical to early identification and management of nutritional deficits, prenatal and postnatal care services are often unavailable to mothers (Rahman et al., 2019). Children are much more at risk of malnutrition due to the lack of healthcare services and their geographical remoteness.

There is also a strong correlation between mothers' levels of education and their ability to provide their children with the nutritious food and clean water they need to grow and develop healthily. Better childcare standards and healthcare-seeking behaviors are associated with greater maternal education, which in turn leads to better nutritional outcomes for children, according to findings (Chowdhury et al., 2020).

Household eating habits are significantly impacted by the persistence of food insecurity and poverty. A lack of caloric diversity and an inadequate diet are major causes of growth and wastage. Traditional eating habits and cultural beliefs that put an emphasis on taking care of the elderly at the expense of providing youngsters with healthy supplementary meals make the problem worse (Islam et al., 2021).

On the other hand, inadequate sanitation and water sources exacerbate children's already precarious nutritional state by putting them at risk of recurrent illnesses and diarrhoea, both of which reduce their body's ability to absorb nutrients (Das et al., 2018).

Participants include parents, whose knowledge and actions have a direct impact on their children's health outcomes; community leaders who can help change harmful cultural practices; healthcare providers who are vital in making nutritional education and interventions available; officials at the local level who can improve infrastructure; and non-governmental organisations (NGOs) that provide crucial support for community-based nutritional programs (Ahmed & Rahman, 2022).

In order to successfully address childhood malnutrition, it is necessary to involve several sectors and concentrate on these interrelated factors.

Problem Tree and Fishbone Analysis

Problem tree




Figure 1: Problem Tree




Figure 2: Fishbone Diagram

Root Cause Analysis

There are complex webs of relationships between cultural, social, economic, and educational factors that contribute to child malnutrition in rural Bangladesh. Interventions aimed at reducing poverty, bolstering maternal education, and increasing healthcare access and infrastructure are essential to tackling this widespread problem in a sustainable way. The World Health Organisation defines “stunting” as a low height-for-age ratio caused by persistent or repeated undernourishment, and “wasting” as a low weight-for-height ratio

caused by insufficient food consumption or long-term sickness. Extreme malnutrition, inadequate prenatal care, and low socioeconomic status are common causes of stunting and wasting. Therefore, a great deal of other relevant matters must be resolved in order to deal with them.

Household access to varied and healthy food is severely limited by poverty, which is still a major underlying cause. Children in rural Bangladesh are more likely to suffer from stunting and wasting due to a combination of poor food and economic vulnerability, according to many studies (Hasan et al., 2021; Rahman & Biswas, 2021). A vicious cycle of chronic malnutrition may persist when families are forced to put short-term calorie demands ahead of long-term nutritional quality due to financial restrictions (The Daily Star, 2022).

Another important factor that has recently come to light is maternal education. This factor has a direct impact on child nutritional outcomes since it leads to more knowledgeable caregiving behaviours. Mothers from low-income backgrounds are more likely to be uneducated, which has a detrimental impact on their children's health and development (Chowdhury et al., 2020). Both Islam et al. (2021) and Ahmed and Rahman (2022) found that when mothers get improved nutrition education, their children's health improves.

Inadequate access to sanitation and clean water, two components of a well-developed infrastructure, worsens malnutrition. Dietary absorption and infant development are severely affected by these deficiencies because of the increased susceptibility to infections caused by them (Das et al., 2018; The Business Standard, 2022). These negative effects on children's health may be significantly reduced if environmental sanitation concerns, such as encouraging good hygiene habits, were addressed.

Because there is a severe shortage of healthcare facilities and trained staff in rural areas, the accessibility of healthcare has a substantial impact on nutritional outcomes. Isolation causes this problem to worsen because it limits the prenatal and postnatal care that mothers and children need (Rahman et al., 2019; Ahmed et al., 2021). Achieving long-term reductions in malnutrition will require bolstering healthcare delivery networks in rural areas.

Many societies uphold standards that unintentionally harm children's nutritional health, adding another layer of complexity to nutritional treatments. Negative practices affecting children's nutritional health include putting adult nutrition first or delaying the start of supplemental feeding (Islam et al., 2021). To change these harmful habits, there must be community-based initiatives that educate the public and include reliable local leaders and healthcare professionals (The Daily Star, 2021).

Child nutrition is negatively impacted by the lack of access to healthcare in rural areas, which limits the availability of preventative and responsive health interventions (Rahman et al., 2019; Ahmed et al., 2022). The cornerstone of a long-term solution to malnutrition is improving healthcare facilities and personnel. Delaying supplemental feeding or giving older relatives more meals than children are two examples of cultural practices and conventions that have a detrimental impact on children's nutrition (Islam et al., 2021). Supporting child nutrition may be achieved by educational activities coordinated by community leaders, which can progressively modify detrimental habits. The next step in making a difference is for all the relevant parties, such as schools, communities, and governments, to work together to tackle these underlying issues (Ahmed & Rahman, 2022; Rahman & Biswas, 2021).

Additionally, our national budget should be set aside specifically for nutrition programs. There was allegedly no nutrition-specific funding in the health sector's 5.43 percent of the overall budget for the 2022–23 fiscal year. Additionally, we are curious as to whether or not youngsters suffering from acute malnutrition are being treated using our scientists' produced ready-to-use therapeutic food (RUTF), which is abundant in micronutrients and protein. Otherwise, its usage should be expanded nationwide. Ultimately, in order to lower the prevalence of stunted children in the nation, the government should adopt a comprehensive strategy.

References

- Ahmed, T., & Rahman, A. E. (2022). Nutrition interventions in Bangladesh: Successes, challenges, and opportunities. *The Lancet Global Health*, 10(8), e1104–e1114. [https://doi.org/10.1016/S2214-109X\(22\)00215-2](https://doi.org/10.1016/S2214-109X(22)00215-2)
- Chowdhury, M. R. K., Rahman, M. S., & Khan, M. M. H. (2020). Levels and determinants of maternal nutritional knowledge and its association with child nutrition outcomes in rural Bangladesh. *Journal of Health, Population and Nutrition*, 39(1), 14. <https://doi.org/10.1186/s41043-020-00224-5>
- Das, S. K., Chisti, M. J., Huq, S., Malek, M. A., & Ahmed, T. (2018). Water, sanitation, and hygiene: A major determinant of childhood diarrhea and malnutrition in Bangladesh. *Global Health Action*, 11(1), 1504730. <https://doi.org/10.1080/16549716.2018.1504730>
- Hasan, M. M., Ahmed, S., Chowdhury, M. A. H., & Biswas, T. (2021). Factors associated with nutritional status among children under-five years of age in Bangladesh: A cross-sectional study. *BMJ Open*, 11(12), e051680. <https://doi.org/10.1136/bmjopen-2021-051680>
- Islam, M. S., Zafar Ullah, A. N., Mainali, S., Imam, M. A., & Hasan, M. I. (2021). Determinants of stunting during the first 1,000 days of life in Bangladesh: A review. *Food Science & Nutrition*, 9(9), 5065–5075. <https://doi.org/10.1002/fsn3.2496>
- Rahman, M., Yunus, F. M., Shah, R., & Khanam, R. (2019). Factors affecting maternal healthcare utilization in rural Bangladesh: A systematic review. *International Journal of Health Planning and Management*, 34(4), e1556–e1574. <https://doi.org/10.1002/hpm.2872>
- Rahman, M., & Biswas, T. (2021). Tackling malnutrition through economic interventions. *The Daily Star*. Retrieved from <https://www.thedailystar.net/opinion/news/tackling-malnutrition-294443>
- Rahman, M. S., & Biswas, S. (2021). Food insecurity and its impact on child nutrition. *The Business Standard*. Retrieved from <https://www.tbsnews.net/bangladesh/health/food-insecurity-impact-child-nutrition-346782>

Phase 2: C4D Plan Development

Logical Framework (Log Frame Analysis)

The following logical framework organises the project's goal, objectives, outputs, and activities against their indicators, means of verification, and key risks and assumptions.

Project Logic	Indicators	Means of Verification	Risks & Assumptions
Goal			
Reduce childhood malnutrition (stunting and wasting) in the rural Upazila by improving nutritional practices, healthcare access, and hygiene behaviours through C4D interventions.	Percentage reduction in stunting and wasting among children under five.	Baseline and follow-up household surveys.	Community participation and acceptance of interventions.
Specific Objectives			
1. Increase awareness of proper nutrition and hygiene practices among caregivers through targeted C4D campaigns and training.	Percentage of caregivers demonstrating knowledge and adopting best practices.	Pre- and post-training assessments, attendance records, behaviour observation.	Caregivers are receptive to training and behavioural change.
2. Enhance access to nutritional food and healthcare services by promoting information on nutrition programs and food support systems.	Number of households with increased awareness of and access to fortified foods and essential supplements.	Market and supply chain analysis, household surveys, NGO reports.	Adequate funding and reliable supply chains.
3. Promote behaviour change regarding exclusive breastfeeding and complementary feeding through culturally appropriate media interventions.	Increase in exclusive breastfeeding rates among infants up to six months.	Healthcare reports, caregiver self-reports, focus group discussions.	Cultural beliefs do not strongly oppose breastfeeding initiatives.
4. Improve access to clean drinking water and sanitation facilities by driving awareness through C4D messaging.	Reduction in diarrhoea and infection rates among children under five.	Healthcare reports, WASH (Water, Sanitation, and Hygiene) monitoring reports.	Sustainable infrastructure improvements are maintained effectively.
Outputs			
1. A minimum of 75% of mothers in the community trained on best nutrition, feeding, and hygiene practices through C4D activities.	Number of trained mothers demonstrating improved feeding practices.	Training records, observational studies.	Mothers attend training sessions and apply knowledge.

Project Logic	Indicators	Means of Verification	Risks & Assumptions
2. At least 10 community healthcare workers trained and actively providing nutrition counselling integrated with C4D strategies.	Number of trained health workers offering regular counselling services using C4D approaches.	Health worker reports, community feedback.	Healthcare workers remain engaged and available.
3. Increased household awareness and access to fortified foods and essential supplements.	Growth in local supply and consumption of fortified foods.	Market analysis, household food surveys.	Availability of fortified foods in local markets.
4. At least 50% of households adopting improved sanitation and hygiene practices due to C4D interventions.	Increase in proper sanitation and hygiene behaviours.	Household surveys, hygiene inspection reports.	Families adopt and sustain improved practices.
Activities			
1. Community Engagement & Training: Conduct monthly community workshops and interactive sessions on child nutrition, exclusive breastfeeding, and hygiene.	Number of workshops conducted and participants trained.	Attendance records, feedback forms, knowledge retention surveys.	High participation levels maintained.
2. Media & Communication: Develop culturally appropriate radio programs, posters, and short videos in local languages to reinforce key messages.	Number of media materials produced and distributed; audience reach.	Media reach analysis, listener/viewer feedback.	Effective media distribution channels exist.
3. Healthcare Strengthening: Train community health workers to provide nutrition counselling and monitoring services incorporating C4D methods.	Number of trained workers providing consistent, communication-based services.	Training completion records, community feedback.	Trained workers remain active.
4. Behaviour Change Strategies: Establish peer support groups for mothers, using storytelling and community champions to reinforce best practices.	Number of active peer groups and participants; behaviour change trends.	Support group records, testimonials.	Community members are open to peer-based support.
5. Food Security Interventions: Work with local farmers and NGOs to promote home gardens and run community campaigns on nutritional food consumption.	Increase in home gardens and food distribution coverage.	Agricultural reports, household surveys.	Farmers and NGOs remain engaged in the initiative.
6. School & Youth Engagement: Integrate nutrition and hygiene education into local school curricula and involve youth in awareness campaigns.	Number of schools adopting and teaching nutrition and hygiene lessons; student engagement in activities.	School reports, student participation records.	Schools actively integrate curriculum changes.

Key Indicators

#	Indicator	Means of Verification	Assumption
1	Percentage of children under five with improved nutritional status.	Healthcare reports, growth monitoring data, community assessments.	Accurate data collection and analysis.
2	Increase in the number of mothers practising exclusive breastfeeding due to C4D influence.	Household surveys, healthcare reports, media impact assessments.	Reliable self-reported data from mothers.
3	Reduction in diarrhoea and infection rates due to improved hygiene and sanitation through C4D messaging.	Medical records, sanitation assessment reports, media analytics.	Sustained access to clean water and sanitation.
4	Number of community members attending training and awareness programs, demonstrating learning.	Training logs, event attendance records, pre- and post-assessments.	Consistent community engagement.
5	Increase in the availability and consumption of fortified foods due to awareness efforts.	Market surveys, household consumption reports, supply chain tracking.	Sustained supply of fortified foods.

Means of Verification

#	Means of Verification	Method	Assumption
1	Baseline and follow-up household surveys.	Comparison of pre- and post-intervention data.	Accurate and timely data collection.
2	Healthcare reports tracking child growth and nutritional improvements.	Child health monitoring records, clinical visits.	Regular data updates by healthcare providers.
3	School reports on student participation in hygiene education programs.	Curriculum integration records, student activity reports.	Schools effectively implement the new curriculum.
4	Market and supply chain analysis on fortified food availability.	Distribution and sales records, purchase trends.	Consistent food supply chain monitoring.
5	Monitoring reports from trained health workers and local NGOs.	Community health assessments, intervention tracking.	Engagement and retention of trained personnel.

Assumptions

#	Assumption	Verification	Condition
1	Community members are open to behavioural changes and willing to participate.	Surveys on attitude and behavioural shifts, media feedback.	Effective engagement strategies are used.
2	Adequate funding and resources remain available for program sustainability.	Financial and resource allocation records.	Continued donor and stakeholder support.

#	Assumption	Verification	Condition
3	Government and NGO stakeholders continue their support.	Policy and program documentation, stakeholder meetings.	Stable political and funding environment.
4	Cultural barriers and misinformation can be effectively addressed through strategic communication.	Public perception surveys, focus groups, community leader insights.	Strong advocacy and cultural adaptation strategies.
5	Infrastructure improvements, such as water and sanitation facilities, are maintained through community ownership.	Field assessments, maintenance logs, WASH advocacy reports.	Local authorities and community members take responsibility for upkeep.

C4D Communication Strategy

A Shortage of Food and Childhood Malnutrition in Rural Bangladesh

A shortage of food, low maternal literacy, restricted access to healthcare, and social norms that prevent effective feeding practices are the primary root causes of childhood malnutrition in rural Bangladesh, which remains an important public health issue. With the goal of ensuring systematic preparation and execution of Communication for Development (C4D) initiatives aimed at reducing childhood malnutrition, this communication strategy aligns with the Logical Framework (Log Frame Analysis). The approach is designed to inform, engage, and empower communities to adopt healthier nutritional habits through advocacy, social mobilization, and behaviour change communication (BCC). The method follows the log frame approach to ensure that each interaction activity is identifiable, verifiable, and in accordance with the primary goals of the project, ultimately resulting in lasting behavioural improvements in hygiene and nutrition.

The primary goal of this strategy is to reduce childhood malnutrition in the targeted rural Upazila, particularly stunting and wasting. This can be achieved by improving eating habits, increasing access to healthcare, and advocating hygienic practices through C4D campaigns. The percentage reduction in instances of malnutrition among children under five will be used to assess the initiative's effectiveness, with baseline and follow-up household surveys serving as verification tools. However, the level of community engagement and acceptance of the interventions—important risk factors for implementation—will determine the overall success of these efforts.

This communication strategy's primary objective is to improve caregivers' awareness of proper sanitation and eating habits. In remote regions, a significant percentage of mothers and caregivers are unaware of the importance of good hygiene practices, supplementary feeding, and exclusive breastfeeding. The communication strategy will address this by highlighting key points emphasizing the benefits of exclusive breastfeeding for the first six months, the importance of transitioning to a diverse and nutrient-rich diet after six months, and the necessity of washing hands with soap before feeding infants. To reduce the risk of waterborne illnesses that exacerbate malnutrition, sanitation and access to clean water will also be promoted.

Multiple communication channels, including direct interactions such as health worker visits, mother support groups, and educational nutrition campaigns, will be used to disseminate these messages. These concepts will be reinforced further through mass media, including radio broadcasts, SMS alerts, and social media campaigns. The effectiveness of these efforts will be analyzed by tracking the percentage of caregivers who demonstrate improved knowledge and implement standard procedures, as verified by pre- and post-training surveys and direct behavioural observations.

Increasing access to nutritious food and healthcare is another significant objective of the communication strategy. The availability of government nutrition programs, fortified foods, and medical facilities that support child growth and development is still unclear to many families. As a result, this strategy will focus on educating individuals about the benefits of enriched foods and essential dietary supplements while encouraging families to take advantage of existing nutrition programs. Another key component will be promoting

household vegetable gardening as a way to ensure year-round access to a variety of fresh foods.

Local agricultural campaigns in markets promoting enriched foods, grassroots nutrition education seminars for parents, and advocacy efforts involving NGOs and government organizations to improve food safety policies are examples of communication activities. Household surveys and market supply chain analyses will be used to evaluate the impact and reach of these programs. This objective is based on the assumption that adequate funding and reliable food supply connections will be maintained to enable the interventions to succeed.

Decreasing misinformation and improving breastfeeding and complementary feeding practices is another essential aspect of this communication strategy. Social norms and misinformation often hinder optimal infant feeding practices, making it necessary to use culturally tailored campaigns to promote positive change. The campaign will address misconceptions such as the belief that sick children should be fed less food and will emphasize the long-term cognitive and health benefits of breastfeeding for up to two years.

Additionally, caregivers will be encouraged to provide homemade complementary foods, which are more nutritious than commercially packaged baby foods. Communication methods will include mother-to-mother support groups, local radio programs featuring success stories, and storytelling techniques that highlight the benefits of exclusive breastfeeding. Furthermore, printed materials, such as illustrated flipcharts, will be distributed to assist mothers during counselling sessions. The success of this intervention will be evaluated based on increases in exclusive breastfeeding rates among infants up to six months, with verification through survey responses and health records. However, the extent to which cultural perspectives oppose breastfeeding initiatives will influence the effectiveness of these efforts.

This strategy acknowledges that inadequate sanitation plays a significant role in childhood malnutrition and aims to improve access to safe drinking water and sanitation facilities alongside nutrition education. Due to contaminated water supplies and poor sanitation, many children in rural areas suffer from recurrent diarrhoea, which hinders nutrient absorption and overall growth. The communication strategy will emphasize messages about the importance of boiling or purifying drinking water, using proper sanitation facilities, and washing hands with soap before meals and after using the toilet. School-based programs will integrate hygiene education into curricula, while behavioural nudges, such as the installation of public handwashing stations in high-traffic areas, will reinforce positive habits.

Key hygiene messages will also be disseminated through local media campaigns, including radio broadcasts and village loudspeakers. The effectiveness of these interventions will be evaluated using healthcare and WASH (Water, Sanitation, and Hygiene) tracking data to measure reductions in diarrhoea and infection rates among children under five. This objective relies on the assumption that the government and the community will effectively maintain sustainable infrastructure improvements.

This strategy will ensure that at least 75% of mothers in the local area receive training on optimal feeding, hygiene, and nutrition practices to achieve noticeable outcomes. Additionally, a minimum of ten local healthcare professionals will be trained on how to incorporate C4D principles into nutrition counselling. The success of these training efforts will be measured by tracking the adoption of improved feeding practices among mothers and

the regular provision of support services by trained healthcare workers. To reinforce key messages, mass media and digital activities will play a crucial role. A minimum of five radio programs will be broadcast each month, and caregivers will receive weekly SMS notifications to encourage participation in nutrition counselling sessions.

Meanwhile, efforts to improve food security will focus on promoting household adoption of home gardening and ensuring the availability of nutritious foods in markets through demand creation campaigns. Hygiene interventions will aim to ensure that at least 50% of households adopt improved sanitation and hygiene behaviours, supported by the establishment of public handwashing stations in key community areas.

Continuous evaluation and monitoring will be conducted to ensure the effectiveness and sustainability of these efforts. Pre- and post-intervention surveys will assess changes in caregivers' knowledge and behaviours, while healthcare data collection will track improvements in child growth and malnutrition rates. Market and supply chain studies will evaluate the impact of food security interventions, while community feedback mechanisms such as focus groups and suggestion boxes will provide insights for ongoing program adjustments. By incorporating flexible and adaptive approaches, this strategy will remain responsive to the evolving needs of the community, ensuring long-term success in reducing childhood malnutrition.

Conclusion

In conclusion, this communication strategy provides a comprehensive, evidence-based approach to addressing childhood malnutrition in rural Bangladesh, structured according to the Logical Framework (Log Frame Analysis). By integrating interpersonal communication, community engagement, mass media, and digital outreach, the strategy seeks to modify behaviours, improve access to resources, and drive policy advocacy. The success of these efforts depends on strong community participation, sustained funding, and the active involvement of key stakeholders. Through continuous evaluation and adaptation, this initiative has the potential to create lasting improvements in child nutrition and overall public health.

Self-Reflection

I, Faria Akter, had two group mates and both of them were amazing. Mounota Apu helped and guided me, while Rushi Apu was good but should do things on time.

I, Mounota, had two group mates and both of them were amazing. Faria helped me, while Rushi did her work timely. I am very happy with my team's work.