Unclassified Acidobacteria Phylum

non−ED vs. T1 adjusted p = 8.4e−05
non−ED vs. T2 adjusted p = 1e−04
T1 vs. T2 adjusted p = 0.82



Unclassified Candidatus Saccharibacte

non−ED vs. T1 adjusted p = 0.00038
non−ED vs. T2 adjusted p = 1e−04
T1 vs. T2 adjusted p = 0.97



Unclassified Desulfobacterales Order

non−ED vs. T1 adjusted p = 0.00017
non−ED vs. T2 adjusted p = 1e−04
T1 vs. T2 adjusted p = 0.72



Coriobacteriaceae

non−ED vs. T1 adjusted p = 0.00017
non−ED vs. T2 adjusted p = 0.00019
T1 vs. T2 adjusted p = 0.91



Gottschalkiaceae

non−ED vs. T1 adjusted p = 0.0043
non−ED vs. T2 adjusted p = 0.00038
T1 vs. T2 adjusted p = 0.72



Methanothermaceae

non−ED vs. T1 adjusted p = 0.072
non−ED vs. T2 adjusted p = 0.00038
T1 vs. T2 adjusted p = 0.14



Unclassified Synechococcales Order

non−ED vs. T1 adjusted p = 2.8e−05
non−ED vs. T2 adjusted p = 0.00049
T1 vs. T2 adjusted p = 0.93



Coprothermobacteraceae

non−ED vs. T1 adjusted p = 0.00017
non−ED vs. T2 adjusted p = 0.00093
T1 vs. T2 adjusted p = 0.93



Unclassified Tissierellia Class

non−ED vs. T1 adjusted p = 0.074
non−ED vs. T2 adjusted p = 0.00093
T1 vs. T2 adjusted p = 0.72



Unclassified Chroococcales Order

non−ED vs. T1 adjusted p = 0.00017
non−ED vs. T2 adjusted p = 0.0011
T1 vs. T2 adjusted p = 1



Kangiellaceae

non−ED vs. T1 adjusted p = 0.029
non−ED vs. T2 adjusted p = 0.0013
T1 vs. T2 adjusted p = 0.72



Acholeplasmataceae

non−ED vs. T1 adjusted p = 0.0072
non−ED vs. T2 adjusted p = 0.0014
T1 vs. T2 adjusted p = 0.74



Unclassified Coriobacteriales Order

non−ED vs. T1 adjusted p = 8e−04
non−ED vs. T2 adjusted p = 0.0014
T1 vs. T2 adjusted p = 0.72



Anaplasmataceae

non−ED vs. T1 adjusted p = 0.0078
non−ED vs. T2 adjusted p = 0.0018
T1 vs. T2 adjusted p = 0.83



Holosporaceae

non−ED vs. T1 adjusted p = 0.034
non−ED vs. T2 adjusted p = 0.0018
T1 vs. T2 adjusted p = 0.72



Blattabacteriaceae

non−ED vs. T1 adjusted p = 0.011
non−ED vs. T2 adjusted p = 0.0018
T1 vs. T2 adjusted p = 0.72



Ferroplasmaceae

non−ED vs. T1 adjusted p = 0.0055
non−ED vs. T2 adjusted p = 0.0021
T1 vs. T2 adjusted p = 0.98



Unclassified Candidatus Gracilibacteria P

non−ED vs. T1 adjusted p = 0.015
non−ED vs. T2 adjusted p = 0.0022
T1 vs. T2 adjusted p = 0.72



Unclassified Saccharomycetales Order

non−ED vs. T1 adjusted p = 4.4e−06
non−ED vs. T2 adjusted p = 0.0022
T1 vs. T2 adjusted p = 0.46



Unclassified Nostocales Order

non−ED vs. T1 adjusted p = 0.074
non−ED vs. T2 adjusted p = 0.0025
T1 vs. T2 adjusted p = 0.72



Unclassified Coriobacteriia Class

non−ED vs. T1 adjusted p = 0.0081
non−ED vs. T2 adjusted p = 0.0031
T1 vs. T2 adjusted p = 0.76



Acidimicrobiaceae

non−ED vs. T1 adjusted p = 0.031
non−ED vs. T2 adjusted p = 0.0037
T1 vs. T2 adjusted p = 0.83



Borreliaceae

non−ED vs. T1 adjusted p = 0.04
non−ED vs. T2 adjusted p = 0.0043
T1 vs. T2 adjusted p = 0.72



Unclassified Fusobacteriales Order

non−ED vs. T1 adjusted p = 0.17
non−ED vs. T2 adjusted p = 0.0057
T1 vs. T2 adjusted p = 0.72



Succinivibrionaceae

non−ED vs. T1 adjusted p = 0.00034
non−ED vs. T2 adjusted p = 0.0087
T1 vs. T2 adjusted p = 0.89



Akkermansiaceae

non−ED vs. T1 adjusted p = 0.16
non−ED vs. T2 adjusted p = 0.012
T1 vs. T2 adjusted p = 0.72



Sordariaceae

non−ED vs. T1 adjusted p = 0.056
non−ED vs. T2 adjusted p = 0.012
T1 vs. T2 adjusted p = 0.72



Aerococcaceae

non−ED vs. T1 adjusted p = 0.012
non−ED vs. T2 adjusted p = 0.013
T1 vs. T2 adjusted p = 0.9



Dictyoglomaceae

non−ED vs. T1 adjusted p = 0.07
non−ED vs. T2 adjusted p = 0.015
T1 vs. T2 adjusted p = 0.72



Salinivirgaceae

non−ED vs. T1 adjusted p = 0.074
non−ED vs. T2 adjusted p = 0.015
T1 vs. T2 adjusted p = 0.72



Spiroplasmataceae

non−ED vs. T1 adjusted p = 0.074
non−ED vs. T2 adjusted p = 0.015
T1 vs. T2 adjusted p = 0.72



Unclassified Silvanigrellales Order

non−ED vs. T1 adjusted p = 0.029
non−ED vs. T2 adjusted p = 0.015
T1 vs. T2 adjusted p = 0.82



Silvanigrellaceae

non−ED vs. T1 adjusted p = 0.15
non−ED vs. T2 adjusted p = 0.016
T1 vs. T2 adjusted p = 0.72



Egibacteraceae

non−ED vs. T1 adjusted p = 0.027
non−ED vs. T2 adjusted p = 0.018
T1 vs. T2 adjusted p = 0.83



Aphanizomenonaceae

non−ED vs. T1 adjusted p = 0.076
non−ED vs. T2 adjusted p = 0.019
T1 vs. T2 adjusted p = 0.72



Methanotrichaceae

non−ED vs. T1 adjusted p = 0.0078
non−ED vs. T2 adjusted p = 0.019
T1 vs. T2 adjusted p = 0.72



Thermodesulfobiaceae

non−ED vs. T1 adjusted p = 0.2
non−ED vs. T2 adjusted p = 0.019
T1 vs. T2 adjusted p = 0.72



Candidatus Midichloriaceae

non−ED vs. T1 adjusted p = 0.041
non−ED vs. T2 adjusted p = 0.025
T1 vs. T2 adjusted p = 0.72



Unclassified Actinobacteria Phylum

non−ED vs. T1 adjusted p = 0.0044
non−ED vs. T2 adjusted p = 0.025
T1 vs. T2 adjusted p = 0.72



Ignavibacteriaceae

non−ED vs. T1 adjusted p = 0.0025
non−ED vs. T2 adjusted p = 0.025
T1 vs. T2 adjusted p = 0.82



Nautiliaceae

non−ED vs. T1 adjusted p = 0.051
non−ED vs. T2 adjusted p = 0.025
T1 vs. T2 adjusted p = 0.72



Fibrobacteraceae

non−ED vs. T1 adjusted p = 0.0053
non−ED vs. T2 adjusted p = 0.026
T1 vs. T2 adjusted p = 0.72



Mycoplasmataceae

non−ED vs. T1 adjusted p = 0.073
non−ED vs. T2 adjusted p = 0.026
T1 vs. T2 adjusted p = 0.72



Sanguibacteraceae

non−ED vs. T1 adjusted p = 0.01
non−ED vs. T2 adjusted p = 0.027
T1 vs. T2 adjusted p = 0.72



Candidatus Deianiraeaceae

non−ED vs. T1 adjusted p = 0.55
non−ED vs. T2 adjusted p = 0.028
T1 vs. T2 adjusted p = 0.72



Lacipirellulaceae

non−ED vs. T1 adjusted p = 0.048
non−ED vs. T2 adjusted p = 0.028
T1 vs. T2 adjusted p = 1



Lichenihabitantaceae

non−ED vs. T1 adjusted p = 0.055
non−ED vs. T2 adjusted p = 0.028
T1 vs. T2 adjusted p = 1



Unclassified Flavobacteriales Order

non−ED vs. T1 adjusted p = 0.19
non−ED vs. T2 adjusted p = 0.028
T1 vs. T2 adjusted p = 0.72



Chamaesiphonaceae

non−ED vs. T1 adjusted p = 0.063
non−ED vs. T2 adjusted p = 0.032
T1 vs. T2 adjusted p = 0.72



Cohaesibacteraceae

non−ED vs. T1 adjusted p = 0.011
non−ED vs. T2 adjusted p = 0.032
T1 vs. T2 adjusted p = 0.74



Unclassified Entomoplasmatales Order

non−ED vs. T1 adjusted p = 0.21
non−ED vs. T2 adjusted p = 0.032
T1 vs. T2 adjusted p = 0.72



Unclassified Thermoplasmatales Order

non−ED vs. T1 adjusted p = 0.01
non−ED vs. T2 adjusted p = 0.032
T1 vs. T2 adjusted p = 0.72



Clavicipitaceae

non−ED vs. T1 adjusted p = 0.01
non−ED vs. T2 adjusted p = 0.033
T1 vs. T2 adjusted p = 0.89



Fusobacteriaceae

non−ED vs. T1 adjusted p = 0.71
non−ED vs. T2 adjusted p = 0.033
T1 vs. T2 adjusted p = 0.71



Brachyspiraceae

non−ED vs. T1 adjusted p = 0.11
non−ED vs. T2 adjusted p = 0.037
T1 vs. T2 adjusted p = 0.72



Glomerellaceae

non−ED vs. T1 adjusted p = 0.018
non−ED vs. T2 adjusted p = 0.037
T1 vs. T2 adjusted p = 0.72



Unclassified Enterobacterales Order

non−ED vs. T1 adjusted p = 0.5
non−ED vs. T2 adjusted p = 0.037
T1 vs. T2 adjusted p = 0.72



Ktedonosporobacteraceae

non−ED vs. T1 adjusted p = 0.018
non−ED vs. T2 adjusted p = 0.039
T1 vs. T2 adjusted p = 0.72



Unclassified Bacillales Order

non−ED vs. T1 adjusted p = 0.055
non−ED vs. T2 adjusted p = 0.039
T1 vs. T2 adjusted p = 0.97



Unclassified Burkholderiales Order

non−ED vs. T1 adjusted p = 0.029
non−ED vs. T2 adjusted p = 0.039
T1 vs. T2 adjusted p = 0.74



Brevibacteriaceae

non−ED vs. T1 adjusted p = 0.027
non−ED vs. T2 adjusted p = 0.039
T1 vs. T2 adjusted p = 0.82



Cryomorphaceae

non−ED vs. T1 adjusted p = 0.074
non−ED vs. T2 adjusted p = 0.039
T1 vs. T2 adjusted p = 0.72



Erwiniaceae

non−ED vs. T1 adjusted p = 0.022
non−ED vs. T2 adjusted p = 0.039
T1 vs. T2 adjusted p = 0.72



Kosmotogaceae

non−ED vs. T1 adjusted p = 0.04
non−ED vs. T2 adjusted p = 0.041
T1 vs. T2 adjusted p = 0.96



Euzebyaceae

non−ED vs. T1 adjusted p = 0.066
non−ED vs. T2 adjusted p = 0.041
T1 vs. T2 adjusted p = 0.97



Francisellaceae

non−ED vs. T1 adjusted p = 0.056
non−ED vs. T2 adjusted p = 0.041
T1 vs. T2 adjusted p = 0.82



Methanocaldococcaceae

non−ED vs. T1 adjusted p = 0.07
non−ED vs. T2 adjusted p = 0.041
T1 vs. T2 adjusted p = 0.77



Unclassified Campylobacterales Order

non−ED vs. T1 adjusted p = 0.04
non−ED vs. T2 adjusted p = 0.042
T1 vs. T2 adjusted p = 0.89



Halothiobacillaceae

non−ED vs. T1 adjusted p = 0.074
non−ED vs. T2 adjusted p = 0.044
T1 vs. T2 adjusted p = 0.96



Chroococcidiopsidaceae

non−ED vs. T1 adjusted p = 0.21
non−ED vs. T2 adjusted p = 0.044
T1 vs. T2 adjusted p = 0.72



Piscirickettsiaceae

non−ED vs. T1 adjusted p = 0.048
non−ED vs. T2 adjusted p = 0.046
T1 vs. T2 adjusted p = 0.87



Unclassified Verrucomicrobiales Order

non−ED vs. T1 adjusted p = 0.95
non−ED vs. T2 adjusted p = 0.046
T1 vs. T2 adjusted p = 0.24



Bernardetiaceae

non−ED vs. T1 adjusted p = 0.12
non−ED vs. T2 adjusted p = 0.049
T1 vs. T2 adjusted p = 0.72



Methanococcaceae

non−ED vs. T1 adjusted p = 0.034
non−ED vs. T2 adjusted p = 0.051
T1 vs. T2 adjusted p = 0.92



Unclassified Oceanospirillales Order

non−ED vs. T1 adjusted p = 0.21
non−ED vs. T2 adjusted p = 0.051
T1 vs. T2 adjusted p = 0.72



Thermoplasmataceae

non−ED vs. T1 adjusted p = 0.4
non−ED vs. T2 adjusted p = 0.052
T1 vs. T2 adjusted p = 0.72



Oxalobacteraceae

non−ED vs. T1 adjusted p = 0.034
non−ED vs. T2 adjusted p = 0.054
T1 vs. T2 adjusted p = 0.73



Bifidobacteriaceae

non−ED vs. T1 adjusted p = 0.011
non−ED vs. T2 adjusted p = 0.054
T1 vs. T2 adjusted p = 0.72



Ardenticatenaceae

non−ED vs. T1 adjusted p = 0.047
non−ED vs. T2 adjusted p = 0.054
T1 vs. T2 adjusted p = 0.72



Hungateiclostridiaceae

non−ED vs. T1 adjusted p = 0.045
non−ED vs. T2 adjusted p = 0.054
T1 vs. T2 adjusted p = 0.91



Ichthyobacteriaceae

non−ED vs. T1 adjusted p = 0.18
non−ED vs. T2 adjusted p = 0.054
T1 vs. T2 adjusted p = 0.72



Nostocaceae

non−ED vs. T1 adjusted p = 0.059
non−ED vs. T2 adjusted p = 0.055
T1 vs. T2 adjusted p = 0.96



Endomicrobiaceae

non−ED vs. T1 adjusted p = 0.19
non−ED vs. T2 adjusted p = 0.056
T1 vs. T2 adjusted p = 0.72



Natranaerobiaceae

non−ED vs. T1 adjusted p = 0.16
non−ED vs. T2 adjusted p = 0.056
T1 vs. T2 adjusted p = 0.72



Vulgatibacteraceae

non−ED vs. T1 adjusted p = 0.027
non−ED vs. T2 adjusted p = 0.058
T1 vs. T2 adjusted p = 0.72



Corynebacteriaceae

non−ED vs. T1 adjusted p = 0.024
non−ED vs. T2 adjusted p = 0.062
T1 vs. T2 adjusted p = 0.72



Methylacidiphilaceae

non−ED vs. T1 adjusted p = 0.011
non−ED vs. T2 adjusted p = 0.062
T1 vs. T2 adjusted p = 0.72



Conexibacteraceae

non−ED vs. T1 adjusted p = 0.01
non−ED vs. T2 adjusted p = 0.062
T1 vs. T2 adjusted p = 0.72



Ilumatobacteraceae

non−ED vs. T1 adjusted p = 0.012
non−ED vs. T2 adjusted p = 0.062
T1 vs. T2 adjusted p = 0.72



Nitrosopumilaceae

non−ED vs. T1 adjusted p = 0.096
non−ED vs. T2 adjusted p = 0.062
T1 vs. T2 adjusted p = 0.72



Ruminococcaceae

non−ED vs. T1 adjusted p = 0.00017
non−ED vs. T2 adjusted p = 0.062
T1 vs. T2 adjusted p = 0.72



Unclassified Streptosporangiales Order

non−ED vs. T1 adjusted p = 0.055
non−ED vs. T2 adjusted p = 0.062
T1 vs. T2 adjusted p = 0.72



Immundisolibacteraceae

non−ED vs. T1 adjusted p = 0.032
non−ED vs. T2 adjusted p = 0.065
T1 vs. T2 adjusted p = 0.72



Isosphaeraceae

non−ED vs. T1 adjusted p = 0.034
non−ED vs. T2 adjusted p = 0.065
T1 vs. T2 adjusted p = 0.72



Ruaniaceae

non−ED vs. T1 adjusted p = 0.013
non−ED vs. T2 adjusted p = 0.065
T1 vs. T2 adjusted p = 0.72



Thiobacillaceae

non−ED vs. T1 adjusted p = 0.029
non−ED vs. T2 adjusted p = 0.065
T1 vs. T2 adjusted p = 0.73



Unclassified Chromatiales Order

non−ED vs. T1 adjusted p = 0.029
non−ED vs. T2 adjusted p = 0.065
T1 vs. T2 adjusted p = 0.72



Crocinitomicaceae

non−ED vs. T1 adjusted p = 0.085
non−ED vs. T2 adjusted p = 0.068
T1 vs. T2 adjusted p = 0.82



Mesoaciditogaceae

non−ED vs. T1 adjusted p = 0.074
non−ED vs. T2 adjusted p = 0.068
T1 vs. T2 adjusted p = 0.87



Kofleriaceae

non−ED vs. T1 adjusted p = 0.062
non−ED vs. T2 adjusted p = 0.071
T1 vs. T2 adjusted p = 0.87



Brucellaceae

non−ED vs. T1 adjusted p = 0.04
non−ED vs. T2 adjusted p = 0.073
T1 vs. T2 adjusted p = 0.73



Jonesiaceae

non−ED vs. T1 adjusted p = 0.41
non−ED vs. T2 adjusted p = 0.073
T1 vs. T2 adjusted p = 0.73



Thermoanaerobacterales Family III. Inc

non−ED vs. T1 adjusted p = 0.068
non−ED vs. T2 adjusted p = 0.073
T1 vs. T2 adjusted p = 0.98



Caldisphaeraceae

non−ED vs. T1 adjusted p = 0.076
non−ED vs. T2 adjusted p = 0.074
T1 vs. T2 adjusted p = 0.72



Pseudoalteromonadaceae

non−ED vs. T1 adjusted p = 0.048
non−ED vs. T2 adjusted p = 0.074
T1 vs. T2 adjusted p = 0.93



Candidatus Paracaedibacteraceae

non−ED vs. T1 adjusted p = 0.77
non−ED vs. T2 adjusted p = 0.077
T1 vs. T2 adjusted p = 0.72



Orbaceae

non−ED vs. T1 adjusted p = 0.07
non−ED vs. T2 adjusted p = 0.077
T1 vs. T2 adjusted p = 1



Cellulomonadaceae

non−ED vs. T1 adjusted p = 0.045
non−ED vs. T2 adjusted p = 0.079
T1 vs. T2 adjusted p = 0.81



Thermotogaceae

non−ED vs. T1 adjusted p = 0.21
non−ED vs. T2 adjusted p = 0.079
T1 vs. T2 adjusted p = 0.82



Salinisphaeraceae

non−ED vs. T1 adjusted p = 0.048
non−ED vs. T2 adjusted p = 0.08
T1 vs. T2 adjusted p = 0.72



Actinomycetaceae

non−ED vs. T1 adjusted p = 0.074
non−ED vs. T2 adjusted p = 0.082
T1 vs. T2 adjusted p = 0.87



Candidatus Nanopelagicaceae

non−ED vs. T1 adjusted p = 0.044
non−ED vs. T2 adjusted p = 0.082
T1 vs. T2 adjusted p = 0.87



Elusimicrobiaceae

non−ED vs. T1 adjusted p = 0.28
non−ED vs. T2 adjusted p = 0.082
T1 vs. T2 adjusted p = 0.77



Halanaerobiaceae

non−ED vs. T1 adjusted p = 0.093
non−ED vs. T2 adjusted p = 0.082
T1 vs. T2 adjusted p = 0.92



Moraxellaceae

non−ED vs. T1 adjusted p = 0.074
non−ED vs. T2 adjusted p = 0.082
T1 vs. T2 adjusted p = 0.97



Neisseriaceae

non−ED vs. T1 adjusted p = 0.068
non−ED vs. T2 adjusted p = 0.082
T1 vs. T2 adjusted p = 0.76



Parachlamydiaceae

non−ED vs. T1 adjusted p = 0.14
non−ED vs. T2 adjusted p = 0.082
T1 vs. T2 adjusted p = 0.8



Segniliparaceae

non−ED vs. T1 adjusted p = 0.044
non−ED vs. T2 adjusted p = 0.082
T1 vs. T2 adjusted p = 0.72



Staphylococcaceae

non−ED vs. T1 adjusted p = 0.32
non−ED vs. T2 adjusted p = 0.082
T1 vs. T2 adjusted p = 0.72



Thiotrichaceae

non−ED vs. T1 adjusted p = 0.055
non−ED vs. T2 adjusted p = 0.082
T1 vs. T2 adjusted p = 0.82



Unclassified Sphingomonadales Order

non−ED vs. T1 adjusted p = 0.047
non−ED vs. T2 adjusted p = 0.082
T1 vs. T2 adjusted p = 0.72



Ustilaginaceae

non−ED vs. T1 adjusted p = 0.045
non−ED vs. T2 adjusted p = 0.082
T1 vs. T2 adjusted p = 0.72



Woeseiaceae

non−ED vs. T1 adjusted p = 0.079
non−ED vs. T2 adjusted p = 0.082
T1 vs. T2 adjusted p = 0.91



Peptoniphilaceae

non−ED vs. T1 adjusted p = 0.11
non−ED vs. T2 adjusted p = 0.085
T1 vs. T2 adjusted p = 0.93



Comamonadaceae

non−ED vs. T1 adjusted p = 0.054
non−ED vs. T2 adjusted p = 0.085
T1 vs. T2 adjusted p = 0.72



Actinopolysporaceae

non−ED vs. T1 adjusted p = 0.064
non−ED vs. T2 adjusted p = 0.086
T1 vs. T2 adjusted p = 0.99



Aurantimonadaceae

non−ED vs. T1 adjusted p = 0.027
non−ED vs. T2 adjusted p = 0.086
T1 vs. T2 adjusted p = 0.72



Bradymonadaceae

non−ED vs. T1 adjusted p = 0.084
non−ED vs. T2 adjusted p = 0.086
T1 vs. T2 adjusted p = 0.92



Chelatococcaceae

non−ED vs. T1 adjusted p = 0.066
non−ED vs. T2 adjusted p = 0.086
T1 vs. T2 adjusted p = 0.72



Fervidobacteriaceae

non−ED vs. T1 adjusted p = 0.11
non−ED vs. T2 adjusted p = 0.086
T1 vs. T2 adjusted p = 0.84



Leptotrichiaceae

non−ED vs. T1 adjusted p = 0.14
non−ED vs. T2 adjusted p = 0.086
T1 vs. T2 adjusted p = 0.85



Methanobacteriaceae

non−ED vs. T1 adjusted p = 0.51
non−ED vs. T2 adjusted p = 0.086
T1 vs. T2 adjusted p = 0.72



Methanocorpusculaceae

non−ED vs. T1 adjusted p = 0.097
non−ED vs. T2 adjusted p = 0.086
T1 vs. T2 adjusted p = 0.98



Unclassified Blastocatellia Class

non−ED vs. T1 adjusted p = 0.21
non−ED vs. T2 adjusted p = 0.086
T1 vs. T2 adjusted p = 0.87



Unclassified Euryarchaeota Phylum

non−ED vs. T1 adjusted p = 0.13
non−ED vs. T2 adjusted p = 0.086
T1 vs. T2 adjusted p = 0.93



Unclassified Gammaproteobacteria Cla

non−ED vs. T1 adjusted p = 0.093
non−ED vs. T2 adjusted p = 0.086
T1 vs. T2 adjusted p = 0.73



Unclassified Sordariomycetes Class

non−ED vs. T1 adjusted p = 0.87
non−ED vs. T2 adjusted p = 0.086
T1 vs. T2 adjusted p = 0.72



Unikaryonidae

non−ED vs. T1 adjusted p = 0.64
non−ED vs. T2 adjusted p = 0.086
T1 vs. T2 adjusted p = 0.51



Unclassified Methanococcales Order

non−ED vs. T1 adjusted p = 0.019
non−ED vs. T2 adjusted p = 0.087
T1 vs. T2 adjusted p = 0.73



Bogoriellaceae

non−ED vs. T1 adjusted p = 0.034
non−ED vs. T2 adjusted p = 0.087
T1 vs. T2 adjusted p = 0.72



Opitutaceae

non−ED vs. T1 adjusted p = 0.066
non−ED vs. T2 adjusted p = 0.09
T1 vs. T2 adjusted p = 0.82



Phycisphaeraceae

non−ED vs. T1 adjusted p = 0.095
non−ED vs. T2 adjusted p = 0.09
T1 vs. T2 adjusted p = 0.97



Pichiaceae

non−ED vs. T1 adjusted p = 0.26
non−ED vs. T2 adjusted p = 0.09
T1 vs. T2 adjusted p = 0.72



Unclassified Epsilonproteobacteria Clas

non−ED vs. T1 adjusted p = 0.11
non−ED vs. T2 adjusted p = 0.09
T1 vs. T2 adjusted p = 0.93



Desulfohalobiaceae

non−ED vs. T1 adjusted p = 0.056
non−ED vs. T2 adjusted p = 0.09
T1 vs. T2 adjusted p = 0.82



Unclassified Cyanobacteria Phylum

non−ED vs. T1 adjusted p = 0.41
non−ED vs. T2 adjusted p = 0.094
T1 vs. T2 adjusted p = 0.72



Calotrichaceae

non−ED vs. T1 adjusted p = 0.054
non−ED vs. T2 adjusted p = 0.096
T1 vs. T2 adjusted p = 0.91



Shewanellaceae

non−ED vs. T1 adjusted p = 0.076
non−ED vs. T2 adjusted p = 0.096
T1 vs. T2 adjusted p = 0.87



Rickettsiaceae

non−ED vs. T1 adjusted p = 0.13
non−ED vs. T2 adjusted p = 0.098
T1 vs. T2 adjusted p = 0.82