
Differential correlation 1 of 10

Anaerostipes hadrus vs Lachnospira eligens

Differential p = 2.04e−04 | adjusted q = 2.15e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender and BMI separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.005.

Differential correlation 2 of 10

Bacteroides thetaiotaomicron vs Bacteroides uniformis

Differential p = 3.23e−04 | adjusted q = 2.15e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender and BMI separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.005.

Differential correlation 3 of 10

Agathobaculum butyriciproducens vs Collinsella aerofaciens

Differential p = 4.50e−04 | adjusted q = 2.15e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender and BMI separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.005.

Differential correlation 4 of 10

Bacteroides intestinalis vs Bifidobacterium longum

Differential p = 6.30e−04 | adjusted q = 2.15e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender and BMI separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.005.

Differential correlation 5 of 10

Fusicatenibacter saccharivorans vs Lachnospira eligens

Differential p = 6.36e−04 | adjusted q = 2.15e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender and BMI separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.005.

Differential correlation 6 of 10

Bacteroides intestinalis vs Prevotella copri

Differential p = 7.34e−04 | adjusted q = 2.15e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender and BMI separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.005.

Differential correlation 7 of 10

Bilophila wadsworthia vs Butyricimonas virosa

Differential p = 7.67e−04 | adjusted q = 2.15e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender and BMI separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.005.

Differential correlation 8 of 10

Alistipes shahii vs Dorea formicigenerans

Differential p = 8.54e−04 | adjusted q = 2.15e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender and BMI separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.005.

Differential correlation 9 of 10

[Ruminococcus] torques vs Alistipes shahii

Differential p = 1.22e−03 | adjusted q = 2.48e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender and BMI separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.005.

Differential correlation 10 of 10

Blautia wexlerae vs Lachnospira eligens

Differential p = 1.23e−03 | adjusted q = 2.48e−01



Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender and BMI separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.005.