
Differential correlation 1 of 10

Bacteroides thetaiotaomicron vs Bacteroides uniformis

Differential p = 3.18e−04 | adjusted q = 1.71e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender, BMI and age category separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.01.

Differential correlation 2 of 10

Alistipes putredinis vs Bilophila wadsworthia

Differential p = 3.49e−04 | adjusted q = 1.71e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender, BMI and age category separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.01.

Differential correlation 3 of 10

Flavonifractor plautii vs Parabacteroides distasonis

Differential p = 6.43e−04 | adjusted q = 2.10e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender, BMI and age category separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.01.

Differential correlation 4 of 10

Fusicatenibacter saccharivorans vs Phocaeicola vulgatus

Differential p = 8.99e−04 | adjusted q = 2.20e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender, BMI and age category separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.01.

Differential correlation 5 of 10

[Clostridium] innocuum vs Flavonifractor plautii

Differential p = 1.92e−03 | adjusted q = 3.75e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender, BMI and age category separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.01.

Differential correlation 6 of 10

Bacteroides uniformis vs Parabacteroides distasonis

Differential p = 2.44e−03 | adjusted q = 3.97e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender, BMI and age category separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.01.

Differential correlation 7 of 10

Eubacterium sp. CAG:38 vs Gemmiger formicilis

Differential p = 5.09e−03 | adjusted q = 6.74e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender, BMI and age category separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.01.

Differential correlation 8 of 10

Blautia wexlerae vs Enterocloster bolteae

Differential p = 5.51e−03 | adjusted q = 6.74e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender, BMI and age category separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.01.

Differential correlation 9 of 10

Bacteroides ovatus vs Bilophila wadsworthia

Differential p = 8.56e−03 | adjusted q = 8.45e−01




Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender, BMI and age category separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.01.

Differential correlation 10 of 10

Enterocloster bolteae vs Fusicatenibacter saccharivorans

Differential p = 8.91e−03 | adjusted q = 8.45e−01



Each point is one sample. Identical axis scales across groups within each page.
Inputs are robust CLR abundances residualized against gender, BMI and age category separately within each group.
Only complete residual pairs are plotted; abundance zeros remain missing.
Dashed lines: linear visual guides; annotations report saved Spearman correlations. Pairs ranked by differential q, then p; selected DISCO filter = 0.01.