



Furo[3,2-c]pyran-4-one




$p = 0.0072$
 $q = 0.36$

4,11,13,15–Tetrahydroidentin B




$p = 0.00085$
 $q = 0.27$

3-Aminosalicylic acid



$p = 0.0032$
 $q = 0.36$


Tyrosol glucuronide



$p = 0.01$
 $q = 0.4$


MG(22:4/0:0/0:0)

Normalized peak intensity




$p = 0.0073$
 $q = 0.36$

Thymine




$p = 0.0051$
 $q = 0.36$

1,24,25-trihydroxyvitamin D3




$p = 0.0076$
 $q = 0.36$

Dimethyl pimelimidate



$p = 0.0095$
 $q = 0.4$

5-Heneicosyl-1,3-benzenediol




$p = 0.0065$
 $q = 0.36$

A box plot showing HIV prevalence by gender. The y-axis represents HIV prevalence from 0 to 100. The x-axis has two categories: 'Non-HIV' and 'Pre-HIV'. For the 'Non-HIV' group, there are two box plots: a blue one for females (median ~55, IQR ~45-65) and a red one for males (median ~55, IQR ~45-65). For the 'Pre-HIV' group, there are two box plots: a blue one for females (median ~15, IQR ~10-20) and a red one for males (median ~15, IQR ~10-20). Individual data points are overlaid on the box plots.


$p = 0.0059$
 $q = 0.36$

N1,N10-Dicoumaroylspermidine




$p = 0.0071$
 $q = 0.36$

Osthenol




$p = 0.0043$
 $q = 0.36$

Docosapentaenoic acid (22n-3)




$p = 0.0043$
 $q = 0.36$

Flurandrenolide



$p = 0.0055$
 $q = 0.36$


Hypoxanthine



$p < 0.0001$
 $q = 0.056$


3-hydroxypristanic acid

Normalized peak intensity



$p = 0.0035$
 $q = 0.88$


Dibutyl phthalate



$p = 0.0097$
 $q = 0.88$

Frenolicin B

Normalized peak intensity



$p = 0.0071$
 $q = 0.88$