
Figure 1. STROBE patient flow: selection, development and temporal validation



Development group n is one third of the development cohort (training tertiles). Validation outcomes use the fixed cut-points.

Model performance and survival findings (temporal validation unless stated)

- Group separation: log-rank p = 2.0×10^{-52} in validation; p = 1.4×10^{-77} in development. Ordinal C-index 0.603.
- Discrimination: C-index 0.622 (95% CI 0.608–0.636) in validation vs 0.595 in development (apparent).
- Calibration: predicted 5-y survival 40.8% vs observed 46.6% (O/E 1.14); the model under-predicts survival.
- Evaluable at 5 years: 26% of validation patients (median follow-up: development 5.89 y, validation 1.47 y).
- Secondary outcome: median recurrence-free survival 2.03 years (95% CI 1.98–2.09).

Excluded counts are differences between consecutive steps.
Events = deaths within the 15-year analysis window. Development recorded deaths: 4,685; 10 of these fall beyond 15 years and are censored in the models.
TNM = tumour-node-metastasis; CRLM = colorectal liver metastases; M2a = stratified Cox model;
KM = Kaplan–Meier; O/E = observed/expected; IBS = integrated Brier score; RFS = recurrence-free survival.
Source: locked v8 analysis (results/final_v7) and Paper 1 model outputs (results/final_v9/paper1).
Every number is generated by make_figure1_strobe.py; see verification_report.md.