

Introduction to Discrete Math

Felipe P. Vista IV



Chonbuk National University

- 1 -

Global Frontier College

- Mathematical Thinking
 - Convincing Arguments, Find Example, Recursion, Logic, Invariants
- Probability & Combinatorics
 - Counting, Probability, Random Variables
- Graph Theory
 - Graphs (cycles, classes, parameters)
- Number Theory & Cryptography
 - Arithmetic in modular form
 - Intro to Cryptography

Mathematical Thinking – Invariants

INVARIANTS

- Invariants
- Coffee with milk
- More Coffee
- Debugging Problem

Invariants



vary = change

Invariants

- **Invariants** → properties that do not change/ remain constant

Invariants

- **Invariants** → properties that do not change/ remain constant
- Looking at the right property is important in general

Invariants

- **Invariants** → properties that do not change/ remain constant
- Looking at the right property is important in general
- It could be a number, or it could be something else

Invariants

- **Invariants** → properties that do not change/ remain constant
- Looking at the right property is important in general
- It could be a number, or it could be something else
- Double counting is a special case

- Invariants
- Coffee with milk
- More Coffee
- Debugging Problem

Coffee with milk

Problem

There are two cups, one with coffee and another with milk. We take a spoon of coffee and add it to the cup of milk. We then take a spoonful from the cup of milk and put it into the cup of coffee. Which is larger, the amount of milk in the cup of coffee or the amount of coffee in the cup of milk?

Coffee with milk

Problem

There are two cups, one with coffee and another with milk. We take a spoon of coffee and add it to the cup of milk. We then take a spoonful from the cup of milk and put it into the cup of coffee. Which is larger, the amount of milk in the cup of coffee or the amount of coffee in the cup of milk?

- It seems we need to do some serious calculation

Coffee with milk

Problem

There are two cups, one with coffee and another with milk. We take a spoon of coffee and add it to the cup of milk. We then take a spoonful from the cup of milk and put it into the cup of coffee. Which is larger, the amount of milk in the cup of coffee or the amount of coffee in the cup of milk?

- It seems we need to do some serious calculation
- We don't know size of cups not size of the spoon

Coffee with milk

Problem

There are two cups, one with coffee and another with milk. We take a spoon of coffee and add it to the cup of milk. We then take a spoonful from the cup of milk and put it into the cup of coffee. Which is larger, the amount of milk in the cup of coffee or the amount of coffee in the cup of milk?

- It seems we need to do some serious calculation
- We don't know size of cups not size of the spoon
 - Maybe the answer depends on these parameters?



Coffee with milk

Problem

There are two cups, one with coffee and another with milk. We take a spoon of coffee and add it to the cup of milk. We then take a spoonful from the cup of milk and put it into the cup of coffee. Which is larger, the amount of milk in the cup of coffee or the amount of coffee in the cup of milk?

- It seems we need to do some serious calculation
- We don't know size of cups not size of the spoon
 - Maybe the answer depends on these parameters?
- Well, it turns out we do not need to do any calculation 😊!




Coffee with milk

Before



Cup 1

After



Cup 2




Coffee with milk

Before



Cup 1

After



Cup 2



- The size of the drink in **cup 1** is invariant!


Coffee with milk

Before



Cup 1

After




Cup 2



- The size of the drink in **cup 1 is invariant!**
- So the amount of coffee missing in cup 1 is the same as the amount of milk added into cup 1

Coffee with milk



- The size of the drink in cup 1 is invariant!
- So the amount of coffee missing in cup 1 is the same as the amount of milk added into cup 1
- Conversely, amount of milk missing in cup 2 is the same as the amount of coffee added into cup 2

- Invariants
- Coffee with milk
- More Coffee
- Debugging Problem

More coffee

Problem

There are two equally-sized cups, one with coffee & another milk. Both cups are half-full. We want coffee w/ lots of milk on the 1st cup: $\frac{1}{3}$ coffee, $\frac{2}{3}$ milk. We can pour from one cup to another back & forth. Can we get our favorite coffee mix into our favorite cup? Any amount would do, right proportion matters.

More coffee

Problem

There are two equally-sized cups, one with coffee & another milk. Both cups are half-full. We want coffee w/ lots of milk on the 1st cup: $\frac{1}{3}$ coffee, $\frac{2}{3}$ milk. We can pour from one cup to another back & forth. Can we get our favorite coffee mix into our favorite cup? Any amount would do, right proportion matters.

- The previous problem doesn't help.



More coffee

Problem

There are two equally-sized cups, one with coffee & another milk. Both cups are half-full. We want coffee w/ lots of milk on the 1st cup: $\frac{1}{3}$ coffee, $\frac{2}{3}$ milk. We can pour from one cup to another back & forth. Can we get our favorite coffee mix into our favorite cup? Any amount would do, right proportion matters.

- The previous problem doesn't help.
 - It's good if second cup is mostly coffee



More coffee

Problem

There are two equally-sized cups, one with coffee & another milk. Both cups are half-full. We want coffee w/ lots of milk on the 1st cup: $\frac{1}{3}$ coffee, $\frac{2}{3}$ milk. We can pour from one cup to another back & forth. Can we get our favorite coffee mix into our favorite cup? Any amount would do, right proportion matters.

- The previous problem doesn't help.
 - It's good if second cup is mostly coffee
- Yet, invariants can help again



More coffee

Problem

There are two equally-sized cups, one with coffee & another milk. Both cups are half-full. We want coffee w/ lots of milk on the 1st cup: $\frac{1}{3}$ coffee, $\frac{2}{3}$ milk. We can pour from one cup to another back & forth. Can we get our favorite coffee mix into our favorite cup? Any amount would do, right proportion matters.

- The previous problem doesn't help.
 - It's good if second cup is mostly coffee
- Yet, invariants can help again
 - We just have to choose the right invariant



More coffee

Claim

The proportion of coffee in the 1st cup is always greater than the 2nd cup. That is, coffee in the 1st cup is stronger than 2nd cup.



More coffee

Claim

The proportion of coffee in the 1st cup is always greater than the 2nd cup. That is, coffee in the 1st cup is stronger than 2nd cup.

- Inequality bet proportions of coffee in cups is our invariant
 - Coffee in Cup 1 > Coffee in Cup 2

More coffee

Claim

The proportion of coffee in the 1st cup is always greater than the 2nd cup. That is, coffee in the 1st cup is stronger than 2nd cup.

- Inequality bet proportions of coffee in cups is **our invariant**
 - Coffee in Cup 1 > Coffee in Cup 2
- **Does not allow** us to have more milk than coffee in 1st cup

More coffee

Claim

The proportion of coffee in the 1st cup is always greater than the 2nd cup. That is, coffee in the 1st cup is stronger than 2nd cup.

- Inequality bet proportions of coffee in cups is **our invariant**
 - Coffee in Cup 1 $>$ Coffee in Cup 2
- **Does not allow** us to have more milk than coffee in 1st cup
 - Else we have more milk than coffee in both cups

More coffee

Claim

The proportion of coffee in the 1st cup is always greater than the 2nd cup. That is, coffee in the 1st cup is stronger than 2nd cup.

- Inequality bet proportions of coffee in cups is **our invariant**
 - Coffee in Cup 1 > Coffee in Cup 2
- **Does not allow** us to have more milk than coffee in 1st cup
 - Else we have more milk than coffee in both cups
- But, **total amount** of coffee and milk are the same



More coffee

Claim

The proportion of coffee in the 1st cup is always greater than the 2nd cup. That is, coffee in the 1st cup is stronger than 2nd cup.



More coffee

Claim

The proportion of coffee in the 1st cup is always greater than the 2nd cup. That is, coffee in the 1st cup is stronger than 2nd cup.

Why is this claim true?




More coffee

Claim

The proportion of coffee in the 1st cup is always greater than the 2nd cup. That is, coffee in the 1st cup is stronger than 2nd cup.

Why is this claim true?

Cup 1



Cup 2




More coffee

Claim

The proportion of coffee in the 1st cup is always greater than the 2nd cup. That is, coffee in the 1st cup is stronger than 2nd cup.

Why is this claim true?




More coffee

Claim

The proportion of coffee in the 1st cup is always greater than the 2nd cup. That is, coffee in the 1st cup is stronger than 2nd cup.

Why is this claim true?

Cup 1



Cup 2




More coffee

Claim

The proportion of coffee in the 1st cup is always greater than the 2nd cup. That is, coffee in the 1st cup is stronger than 2nd cup.

Why is this claim true?

Cup 1



Cup 2




We “milk down” drink
from Cup 1 into Cup 2

More coffee

Claim

The proportion of coffee in the 1st cup is always greater than the 2nd cup. That is, coffee in the 1st cup is stronger than 2nd cup.

Why is this claim true?




We “milk down” drink
from Cup 1 into Cup 2

More coffee

Claim

The proportion of coffee in the 1st cup is always greater than the 2nd cup. That is, coffee in the 1st cup is stronger than 2nd cup.

Why is this claim true?




We “**milk down**” drink
from Cup 1 into Cup 2

More coffee

Claim

The proportion of coffee in the 1st cup is always greater than the 2nd cup. That is, coffee in the 1st cup is stronger than 2nd cup.

Why is this claim true?



- Invariants
- Coffee with milk
- More Coffee
- Debugging Problem

Typical Debugging

Problem



1 → 3

Bob is debugging his code. There is only one bug when he starts. But three new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Typical Debugging

Problem

Bob is debugging his code. There is only one bug when he starts. But three new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Typical Debugging

Problem

Bob is debugging his code. There is only one bug when he starts. But three new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Fixed:

Pending:

Typical Debugging

Problem

Bob is debugging his code. There is only **one** bug when he starts. But three new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Fixed:

0					
1					

Pending:

Typical Debugging

Problem

Bob is debugging his code. There is only **one** bug when he starts. But **three** new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Fixed:	0	1				
Pending:	1	3				

Typical Debugging

Problem

Bob is debugging his code. There is only **one** bug when he starts. But **three** new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Fixed:	0	1	2			
Pending:	1	3	5			

Typical Debugging

Problem

Bob is debugging his code. There is only **one** bug when he starts. But **three** new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Fixed:	0	1	2	3		
Pending:	1	3	5	7		

Typical Debugging

Problem

Bob is debugging his code. There is only **one** bug when he starts. But **three** new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Fixed:	0	1	2	3	4	15
Pending:	1	3	5	7	9	?

Typical Debugging

Problem

Bob is debugging his code. There is only **one** bug when he starts. But **three** new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Fixed:	0	1	2	3	4	...
Pending:	1	3	5	7	9	...

Typical Debugging

Problem

Fixed:

0	1	2	3	4	...
1	3	5	7	9	...

Pending:



Typical Debugging

Problem

Fixed:	0	1	2	3	4	...
Pending:	1	3	5	7	9	...

- $\# \text{Pending} = \text{????}$

Typical Debugging

Problem

Fixed:	0	1	2	3	4	...
Pending:	1	3	5	7	9	...

- $\# \text{Pending} = 1 + 2 \times \# \text{Fixed}$
 - This is our invariant!

Typical Debugging

Problem

Fixed:	0	1	2	3	4	...
Pending:	1	3	5	7	9	...

- $\# \text{Pending} = 1 + 2 \times \# \text{Fixed}$
 - This is our invariant!
- Hence, if $\# \text{Fixed} = 15$

Typical Debugging

Problem

Fixed:	0	1	2	3	4	...
Pending:	1	3	5	7	9	...

- $\# \text{Pending} = 1 + 2 \times \# \text{Fixed}$
 - This is our invariant!
- Hence, if $\# \text{Fixed} = 15$
 - $\# \text{Pending} = 1 + 2 \times 15$

Typical Debugging

Problem

Fixed:	0	1	2	3	4	...
Pending:	1	3	5	7	9	...

- $\# \text{Pending} = 1 + 2 \times \# \text{Fixed}$
 - This is our invariant!
- Hence, if $\# \text{Fixed} = 15$
 - $\# \text{Pending} = 1 + 2 \times 15$
 - $\# \text{Pending} = 31$

Typical Debugging

Problem

Fixed:	0	1	2	3	4	...	15
Pending:	1	3	5	7	9	...	31

- $\#Pending = 1 + 2 \times \#Fixed$
 - This is our invariant!
- Hence, if $\#Fixed = 15$
 - $\#Pending = 1 + 2 \times 15$
 - $\#Pending = 31$

Mathematical Thinking – Invariants

TERMINATION

- Termination
- Football fans
- King Julien's Books

Termination

- We used invariants to show impossibility

Termination

- We used invariants to show **impossibility**
- For us, invariants are properties that **do not change**



Termination

- We used invariants to show **impossibility**
- For us, invariants are properties that **do not change**
- In a more wider sense, **invariant** → properties that change in **the right way**

Termination

- We used invariants to show impossibility
- For us, invariants are properties that do not change
- In a more wider sense, **invariant** → properties that change in the right way
- Another standard use of an invariant is showing the **termination** of a process

- Termination
- Football fans
- King Julien's Books

Football Fans

Problem

There are 2 football teams in a town. Each of the citizens support a team. If there are more fans of the other team among someone's friend other than his own, this person tends to switch to the other team. One such person switch teams every day. Is it possible that this switching process will go on forever? Assume the following: friendship is always mutual, population does not change, friendship does not change.

Football Fans

Problem

There are 2 football teams in a town. Each of the citizens support a team. If there are more fans of the other team among someone's friend other than his own, this person tends to switch to the other team. One such person switch teams every day. Is it possible that this switching process will go on forever? Assume the following: friendship is always mutual, population does not change, friendship does not change.

- It seems natural that the process will stop



Football Fans

Problem

There are 2 football teams in a town. Each of the citizens support a team. If there are more fans of the other team among someone's friend other than his own, this person tends to switch to the other team. One such person switch teams every day. Is it possible that this switching process will go on forever? Assume the following: friendship is always mutual, population does not change, friendship does not change.

- It seems natural that the process will stop
- How can we prove it though?
 - We need to look at the right value



Football Fans

Solution


Let us look at the number of opposite team's friendships, that is, the pairs of friends supporting opposite teams at the start of the 1st day.



Football Fans

Solution


Let us look at the number of opposite team's friendships, that is, the pairs of friends supporting opposite teams at the **start of the 1st day**.



Football Fans

Solution


Let us look at the number of opposite team's friendships, that is, the pairs of friends supporting opposite teams. Let's see what happens with this value at **the end of day 1**.



Football Fans

Solution


Let us look at the number of opposite team's friendships, that is, the pairs of friends supporting opposite teams. Let's see what happens with this value at the **start of 2nd day**.



Football Fans

Solution


Let us look at the number of opposite team's friendships, that is, the pairs of friends supporting opposite teams. Let's see what happens with this value at the **end of 2nd day**.



Football Fans

Solution

Let us look at the number of opposite team's friendships, that is, the pairs of friends supporting opposite teams. Let's see what happens with this value at the **end of 2nd day**.



- Termination
- Football fans
- King Julien's Books

King Julien's Books

Problem

King Julien has a shelf of his works consisting of 10 volumes, labeled 1, 2, ..., 10. These got jumbled & out of order over the years. King J asked Maurice to sort the collection but he can only take 2 books at once. The books are heavy hence he can switch only two per day. In how many days can Maurice guarantee that the volumes are sorted?



King Julien's Books

Problem

King Julien has a shelf of his works consisting of 10 volumes, labeled 1, 2, ..., 10. These got jumbled & out of order over the years. King J asked Maurice to sort the collection but he can only take 2 books at once. The books are heavy hence he can switch only two per day. In how many days can Maurice guarantee that the volumes are sorted?



King Julien's Books

Let's check

We can always place books in the right order in at most 9 days?



King Julien's Books

Let's check

We can always place books in the right order in at most 9 days?



- Day 1: place Vol 1 on it's place

King Julien's Books

Let's check

We can always place books in the right order in at most 9 days?



- Day 1: place Vol 1 on it's place

King Julien's Books

Let's check

We can always place books in the right order in at most 9 days?



- Day 1: place Vol 1 on it's place
- Day 2: place Vol 2 on it's place

King Julien's Books

Let's check

We can always place books in the right order in at most 9 days?




- Day 1: place Vol 1 on it's place
- Day 2: place Vol 2 on it's place

King Julien's Books

Let's check

We can always place books in the right order in at most 9 days?



- Day 1: place Vol 1 on it's place
- Day 2: place Vol 2 on it's place
- Day 3: place Vol 3 on it's place

King Julien's Books

Let's check

We can always place books in the right order in at most 9 days?



- Day 1: place Vol 1 on it's place
- Day 2: place Vol 2 on it's place
- Day 3: place Vol 3 on it's place
- And so on...

King Julien's Books

Let's check

We can always place books in the right order in at most 9 days?



- Day 1: place Vol 1 on it's place
- Day 2: place Vol 2 on it's place
- Day 3: place Vol 3 on it's place
- And so on...
- On Day 9, the first 9 volumes are on their proper places. The 10th volumes must also be in it's proper place since it's only one left.

King Julien's Books

Let's check

- Are 9 days the optimal?



King Julien's Books

Let's check

- Are 9 days the optimal?
- What could be the hardest permutation of the books?



King Julien's Books

Let's check

- Are 9 days the optimal?
- What could be the hardest permutation of the books?

Recall:

- Permutation – order of selection is a factor
- Combination – order of selection not a factor

Ex: Permutation/Combination pairs from the set {A, B, C, D, E}

Permutation: AB AC AD AE BA BC BD BE CA CB CD CE DA DB DC DE EA EB EC ED

Combination: AB AC AD AE BC BE CD CE DE



King Julien's Books

Let's check

- Are 9 days the optimal?
- What could be the hardest permutation of the books?
 - It seems that the hardest is when the books are in opposite order



Recall:

- Permutation – order of selection is a factor
- Combination – order of selection not a factor

Ex: Permutation/Combination pairs from the set {A, B, C, D, E}

Permutation: AB AC AD AE BA BC BD BE CA CB CD CE DA DB DC DE EA EB EC ED

Combination: AB AC AD AE BC BE CD CE DE

King Julien's Books

Let's check

- Are 9 days the optimal?
- What could be the hardest permutation of the books?
 - It seems that the hardest is when the books are in opposite order



King Julien's Books

Let's check

- Are 9 days the optimal?
- What could be the hardest permutation of the books?
 - It seems that the hardest is when the books are in opposite order



King Julien's Books

Let's check

- Are 9 days the optimal?
- What could be the hardest permutation of the books?
 - It seems that the hardest is when the books are in opposite order



King Julien's Books

Let's check

- Are 9 days the optimal?
- What could be the hardest permutation of the books?
 - It seems that the hardest is when the books are in opposite order



➤ All the books can be switched and sorted in just 5 days!

King Julien's Books

Let's check

- So what is the right number of days?



King Julien's Books

Let's check

- So what is the right number of days?
- And how to prove that it is the correct answer?



King Julien's Books

Let's check

- So what is the right number of days?
- And how to prove that it is the correct answer?
- To answer, we need to find some invariant that:

King Julien's Books

Let's check

- So what is the right number of days?
- And how to prove that it is the correct answer?
- To answer, we need to find some invariant that:
 - Do not change fast



King Julien's Books

Let's check

- So what is the right number of days?
- And how to prove that it is the correct answer?
- To answer, we need to find some invariant that:
 - Do not change fast
 - Should change substantially while ordering the book



King Julien's Books

Recall this:

Puzzle

- There is a sequence of 10 cells. The leftmost contains “1” while the rightmost has “30”. Is it possible to fill other cells with consecutive numbers in such a way that they differ by 3 at the most?

1									30
---	--	--	--	--	--	--	--	--	----

King Julien's Books

The **Invariant**:



King Julien's Books

The **Invariant**: The number of books staying to the right of their intended place



King Julien's Books

The **Invariant**: The number of books staying to the right of their intended place

- Small in the end: **equals 0**



King Julien's Books

The **Invariant**: The number of books staying to the right of their intended place

- Small in the end: equals 0
- Decreases slowly: by at most 1 each day

King Julien's Books

The **Invariant**: The number of books staying to the right of their intended place

- Small in the end: **equals 0**
- Decreases slowly: **by at most 1 each day**
- Large in the beginning? **Yes!**

King Julien's Books

The **Invariant**: The number of books staying to the right of their intended place

- Small in the end: **equals 0**
- Decreases slowly: **by at most 1 each day**
- Large in the beginning? **Yes!**



- The invariant is **9 in the beginning**, we need at **least 9 days**.

Thank you.