

# Introduction to Discrete Math

Felipe P. Vista IV



Chonbuk National University

- 1 -

Global Frontier College

- Mathematical Thinking
  - Convincing Arguments, Find Example, Recursion, Logic, Invariants
- Probability & Combinatorics
  - Counting, Probability, Random Variables
- Graph Theory
  - Graphs (cycles, classes, parameters)
- Number Theory & Cryptography
  - Arithmetic in modular form
  - Intro to Cryptography

Mathematical Thinking – Invariants

# **INVARIANTS**

- Invariants
- Coffee with milk
- More Coffee
- Debugging Problem

# Invariants



*vary = change*

# Invariants

- **Invariants** → properties that do not change/ remain constant

# Invariants

- **Invariants** → properties that do not change/ remain constant
- Looking at the right property is important in general

## Invariants

- **Invariants** → properties that do not change/ remain constant
- Looking at the right property is important in general
- It could be a number, or it could be something else



## Invariants

- **Invariants** → properties that do not change/ remain constant
- Looking at the right property is important in general
- It could be a number, or it could be something else
- Double counting is a special case

- Invariants
- Coffee with milk
- More Coffee
- Debugging Problem

## Coffee with milk

### Problem

There are two cups, one with coffee and another with milk. We take a spoon of coffee and add it to the cup of milk. We then take a spoonful from the cup of milk and put it into the cup of coffee. Which is larger, the amount of milk in the cup of coffee or the amount of coffee in the cup of milk?

## Coffee with milk

### Problem

There are two cups, one with coffee and another with milk. We take a spoon of coffee and add it to the cup of milk. We then take a spoonful from the cup of milk and put it into the cup of coffee. Which is larger, the amount of milk in the cup of coffee or the amount of coffee in the cup of milk?

- It seems we need to do some serious calculation

## Coffee with milk

### Problem

There are two cups, one with coffee and another with milk. We take a spoon of coffee and add it to the cup of milk. We then take a spoonful from the cup of milk and put it into the cup of coffee. Which is larger, the amount of milk in the cup of coffee or the amount of coffee in the cup of milk?

- It seems we need to do some serious calculation
- We don't know size of cups not size of the spoon

## Coffee with milk

### Problem

There are two cups, one with coffee and another with milk. We take a spoon of coffee and add it to the cup of milk. We then take a spoonful from the cup of milk and put it into the cup of coffee. Which is larger, the amount of milk in the cup of coffee or the amount of coffee in the cup of milk?

- It seems we need to do some serious calculation
- We don't know size of cups not size of the spoon
  - Maybe the answer depends on these parameters?



## Coffee with milk

### Problem

There are two cups, one with coffee and another with milk. We take a spoon of coffee and add it to the cup of milk. We then take a spoonful from the cup of milk and put it into the cup of coffee. Which is larger, the amount of milk in the cup of coffee or the amount of coffee in the cup of milk?

- It seems we need to do some serious calculation
- We don't know size of cups not size of the spoon
  - Maybe the answer depends on these parameters?
- Well, it turns out we do not need to do any calculation 😊!




## Coffee with milk

Before



Cup 1

After



Cup 2






## Coffee with milk

Before



Cup 1

After



Cup 2



- The size of the drink in **cup 1** is invariant!


## Coffee with milk

Before



Cup 1

After




Cup 2



- The size of the drink in **cup 1 is invariant!**
- So the amount of coffee missing in cup 1 is the same as the amount of milk added into cup 1

## Coffee with milk



- The size of the drink in cup 1 is invariant!
- So the amount of coffee missing in cup 1 is the same as the amount of milk added into cup 1
- Conversely, amount of milk missing in cup 2 is the same as the amount of coffee added into cup 2

- Invariants
- Coffee with milk
- More Coffee
- Debugging Problem

## More coffee

### Problem

There are two equally-sized cups, one with coffee & another milk. Both cups are half-full. We want coffee w/ lots of milk on the 1<sup>st</sup> cup:  $\frac{1}{3}$  coffee,  $\frac{2}{3}$  milk. We can pour from one cup to another back & forth. Can we get our favorite coffee mix into our favorite cup? Any amount would do, right proportion matters.

## More coffee

### Problem

There are two equally-sized cups, one with coffee & another milk. Both cups are half-full. We want coffee w/ lots of milk on the 1<sup>st</sup> cup:  $\frac{1}{3}$  coffee,  $\frac{2}{3}$  milk. We can pour from one cup to another back & forth. Can we get our favorite coffee mix into our favorite cup? Any amount would do, right proportion matters.

- The previous problem doesn't help.



## More coffee

### Problem

There are two equally-sized cups, one with coffee & another milk. Both cups are half-full. We want coffee w/ lots of milk on the 1<sup>st</sup> cup:  $\frac{1}{3}$  coffee,  $\frac{2}{3}$  milk. We can pour from one cup to another back & forth. Can we get our favorite coffee mix into our favorite cup? Any amount would do, right proportion matters.

- The previous problem doesn't help.
  - It's good if second cup is mostly coffee



## More coffee

### Problem

There are two equally-sized cups, one with coffee & another milk. Both cups are half-full. We want coffee w/ lots of milk on the 1<sup>st</sup> cup:  $\frac{1}{3}$  coffee,  $\frac{2}{3}$  milk. We can pour from one cup to another back & forth. Can we get our favorite coffee mix into our favorite cup? Any amount would do, right proportion matters.

- The previous problem doesn't help.
  - It's good if second cup is mostly coffee
- Yet, invariants can help again





## More coffee

### Problem

There are two equally-sized cups, one with coffee & another milk. Both cups are half-full. We want coffee w/ lots of milk on the 1<sup>st</sup> cup:  $\frac{1}{3}$  coffee,  $\frac{2}{3}$  milk. We can pour from one cup to another back & forth. Can we get our favorite coffee mix into our favorite cup? Any amount would do, right proportion matters.

- The previous problem doesn't help.
  - It's good if second cup is mostly coffee
- Yet, invariants can help again
  - We just have to choose the right invariant



## More coffee

### Claim

The proportion of coffee in the 1<sup>st</sup> cup is always greater than the 2<sup>nd</sup> cup. That is, coffee in the 1<sup>st</sup> cup is stronger than 2<sup>nd</sup> cup.



## More coffee

### Claim

The proportion of coffee in the 1<sup>st</sup> cup is always greater than the 2<sup>nd</sup> cup. That is, coffee in the 1<sup>st</sup> cup is stronger than 2<sup>nd</sup> cup.

- Inequality bet proportions of coffee in cups is our invariant
  - Coffee in Cup 1 > Coffee in Cup 2



## More coffee

### Claim

The proportion of coffee in the 1<sup>st</sup> cup is always greater than the 2<sup>nd</sup> cup. That is, coffee in the 1<sup>st</sup> cup is stronger than 2<sup>nd</sup> cup.

- Inequality bet proportions of coffee in cups is **our invariant**
  - Coffee in Cup 1  $>$  Coffee in Cup 2
- **Does not allow** us to have more milk than coffee in 1<sup>st</sup> cup

## More coffee

### Claim

The proportion of coffee in the 1<sup>st</sup> cup is always greater than the 2<sup>nd</sup> cup. That is, coffee in the 1<sup>st</sup> cup is stronger than 2<sup>nd</sup> cup.

- Inequality bet proportions of coffee in cups is **our invariant**
  - Coffee in Cup 1 > Coffee in Cup 2
- **Does not allow** us to have more milk than coffee in 1<sup>st</sup> cup
  - Else we have more milk than coffee in both cups



## More coffee

### Claim

The proportion of coffee in the 1<sup>st</sup> cup is always greater than the 2<sup>nd</sup> cup. That is, coffee in the 1<sup>st</sup> cup is stronger than 2<sup>nd</sup> cup.

- Inequality bet proportions of coffee in cups is **our invariant**
  - Coffee in Cup 1 > Coffee in Cup 2
- **Does not allow** us to have more milk than coffee in 1<sup>st</sup> cup
  - Else we have more milk than coffee in both cups
- But, **total amount** of coffee and milk are the same

## More coffee

### Claim

The proportion of coffee in the 1<sup>st</sup> cup is always greater than the 2<sup>nd</sup> cup. That is, coffee in the 1<sup>st</sup> cup is stronger than 2<sup>nd</sup> cup.



## More coffee

### Claim

The proportion of coffee in the 1<sup>st</sup> cup is always greater than the 2<sup>nd</sup> cup. That is, coffee in the 1<sup>st</sup> cup is stronger than 2<sup>nd</sup> cup.

Why is this claim true?




## More coffee

### Claim

The proportion of coffee in the 1<sup>st</sup> cup is always greater than the 2<sup>nd</sup> cup. That is, coffee in the 1<sup>st</sup> cup is stronger than 2<sup>nd</sup> cup.

Why is this claim true?

Cup 1



Cup 2




## More coffee

### Claim

The proportion of coffee in the 1<sup>st</sup> cup is always greater than the 2<sup>nd</sup> cup. That is, coffee in the 1<sup>st</sup> cup is stronger than 2<sup>nd</sup> cup.

Why is this claim true?




## More coffee

### Claim

The proportion of coffee in the 1<sup>st</sup> cup is always greater than the 2<sup>nd</sup> cup. That is, coffee in the 1<sup>st</sup> cup is stronger than 2<sup>nd</sup> cup.

Why is this claim true?

Cup 1



Cup 2




## More coffee

### Claim

The proportion of coffee in the 1<sup>st</sup> cup is always greater than the 2<sup>nd</sup> cup. That is, coffee in the 1<sup>st</sup> cup is stronger than 2<sup>nd</sup> cup.

Why is this claim true?

Cup 1



Cup 2




We “milk down” drink  
from Cup 1 into Cup 2

## More coffee

### Claim

The proportion of coffee in the 1<sup>st</sup> cup is always greater than the 2<sup>nd</sup> cup. That is, coffee in the 1<sup>st</sup> cup is stronger than 2<sup>nd</sup> cup.

Why is this claim true?




We “**milk down**” drink  
from Cup 1 into Cup 2

## More coffee

### Claim

The proportion of coffee in the 1<sup>st</sup> cup is always greater than the 2<sup>nd</sup> cup. That is, coffee in the 1<sup>st</sup> cup is stronger than 2<sup>nd</sup> cup.

Why is this claim true?




We “**milk down**” drink  
from Cup 1 into Cup 2

## More coffee

### Claim

The proportion of coffee in the 1<sup>st</sup> cup is always greater than the 2<sup>nd</sup> cup. That is, coffee in the 1<sup>st</sup> cup is stronger than 2<sup>nd</sup> cup.

Why is this claim true?



- Invariants
- Coffee with milk
- More Coffee
- Debugging Problem



## Typical Debugging

### Problem



1 → 3

Bob is debugging his code. There is only one bug when he starts. But three new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

## Typical Debugging

### Problem

Bob is debugging his code. There is only one bug when he starts. But three new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

# Typical Debugging

## Problem

Bob is debugging his code. There is only one bug when he starts. But three new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Fixed:

Pending:


# Typical Debugging

## Problem

Bob is debugging his code. There is only **one** bug when he starts. But three new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Fixed:	0					
Pending:	1					

# Typical Debugging

## Problem

Bob is debugging his code. There is only **one** bug when he starts. But **three** new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Fixed:	0	1				
Pending:	1	3				

# Typical Debugging

## Problem

Bob is debugging his code. There is only **one** bug when he starts. But **three** new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Fixed:	0	1	2			
Pending:	1	3	5			

# Typical Debugging

## Problem

Bob is debugging his code. There is only **one** bug when he starts. But **three** new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Fixed:	0	1	2	3		
Pending:	1	3	5	7		

# Typical Debugging

## Problem

Bob is debugging his code. There is only **one** bug when he starts. But **three** new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Fixed:	0	1	2	3	4	15
Pending:	1	3	5	7	9	?



# Typical Debugging

## Problem

Bob is debugging his code. There is only **one** bug when he starts. But **three** new bugs appear once he fixes a bug. Bob fixed 15 bugs after several hours. How many pending bugs (to be fixed) does he have at this point?

Let's see what is going on...

Fixed:	0	1	2	3	4	...
Pending:	1	3	5	7	9	...

# Typical Debugging

## Problem

Fixed:

0	1	2	3	4	...
1	3	5	7	9	...

Pending:



# Typical Debugging

## Problem

Fixed:	0	1	2	3	4	...
Pending:	1	3	5	7	9	...

- $\# \text{Pending} = \text{????}$

# Typical Debugging

## Problem

Fixed:	0	1	2	3	4	...
Pending:	1	3	5	7	9	...

- $\# \text{Pending} = 1 + 2 \times \# \text{Fixed}$ 
  - This is our invariant!

# Typical Debugging

## Problem

Fixed:	0	1	2	3	4	...
Pending:	1	3	5	7	9	...

- $\# \text{Pending} = 1 + 2 \times \# \text{Fixed}$ 
  - This is our invariant!
- Hence, if  $\# \text{Fixed} = 15$

# Typical Debugging

## Problem

Fixed:	0	1	2	3	4	...
Pending:	1	3	5	7	9	...

- $\# \text{Pending} = 1 + 2 \times \# \text{Fixed}$ 
  - This is our invariant!
- Hence, if  $\# \text{Fixed} = 15$ 
  - $\# \text{Pending} = 1 + 2 \times 15$

# Typical Debugging

## Problem

Fixed:	0	1	2	3	4	...
Pending:	1	3	5	7	9	...

- $\# \text{Pending} = 1 + 2 \times \# \text{Fixed}$ 
  - This is our invariant!
- Hence, if  $\# \text{Fixed} = 15$ 
  - $\# \text{Pending} = 1 + 2 \times 15$
  - $\# \text{Pending} = 31$

# Typical Debugging

## Problem

Fixed:	0	1	2	3	4	...	15
Pending:	1	3	5	7	9	...	31

- $\#Pending = 1 + 2 \times \#Fixed$ 
  - This is our invariant!
- Hence, if  $\#Fixed = 15$ 
  - $\#Pending = 1 + 2 \times 15$
  - $\#Pending = 31$



Mathematical Thinking – Invariants

# **TERMINATION**

- Termination
- Football fans
- King Julien's Books

# Termination

- We used invariants to show impossibility

## Termination

- We used invariants to show impossibility
- For us, invariants are properties that do not change



## Termination

- We used invariants to show **impossibility**
- For us, invariants are properties that **do not change**
- In a more wider sense, **invariant** → properties that change in **the right way**

## Termination

- We used invariants to show impossibility
- For us, invariants are properties that do not change
- In a more wider sense, **invariant** → properties that change in the right way
- Another standard use of an invariant is showing the **termination** of a process

- Termination
- Football fans
- King Julien's Books

## Football Fans

### Problem

There are 2 football teams in a town. Each of the citizens support a team. If there are more fans of the other team among someone's friend other than his own, this person tends to switch to the other team. One such person switch teams every day. Is it possible that this switching process will go on forever? Assume the following: friendship is always mutual, population does not change, friendship does not change.



## Football Fans

### Problem

There are 2 football teams in a town. Each of the citizens support a team. If there are more fans of the other team among someone's friend other than his own, this person tends to switch to the other team. One such person switch teams every day. Is it possible that this switching process will go on forever? Assume the following: friendship is always mutual, population does not change, friendship does not change.

- It seems natural that the process will stop

## Football Fans

### Problem

There are 2 football teams in a town. Each of the citizens support a team. If there are more fans of the other team among someone's friend other than his own, this person tends to switch to the other team. One such person switch teams every day. Is it possible that this switching process will go on forever? Assume the following: friendship is always mutual, population does not change, friendship does not change.

- It seems natural that the process will stop
- How can we prove it though?
  - We need to look at the right value



## Football Fans

### Solution


Let us look at the number of opposite team's friendships, that is, the pairs of friends supporting opposite teams at the start of the 1<sup>st</sup> day.



# Football Fans

## Solution


Let us look at the number of opposite team's friendships, that is, the pairs of friends supporting opposite teams at the **start of the 1<sup>st</sup> day**.



# Football Fans

## Solution


Let us look at the number of opposite team's friendships, that is, the pairs of friends supporting opposite teams. Let's see what happens with this value at **the end of day 1**.



# Football Fans

## Solution


Let us look at the number of opposite team's friendships, that is, the pairs of friends supporting opposite teams. Let's see what happens with this value at the **start of 2<sup>nd</sup> day**.



# Football Fans

## Solution


Let us look at the number of opposite team's friendships, that is, the pairs of friends supporting opposite teams. Let's see what happens with this value at the **end of 2<sup>nd</sup> day**.



# Football Fans

## Solution

Let us look at the number of opposite team's friendships, that is, the pairs of friends supporting opposite teams. Let's see what happens with this value at the **end of 2<sup>nd</sup> day**.





- Termination
- Football fans
- King Julien's Books

# King Julien's Books

## Problem

King Julien has a shelf of his works consisting of 10 volumes, labeled 1, 2, ..., 10. These got jumbled & out of order over the years. King J asked Maurice to sort the collection but he can only take 2 books at once. The books are heavy hence he can switch only two per day. In how many days can Maurice guarantee that the volumes are sorted?



# King Julien's Books

## Problem

King Julien has a shelf of his works consisting of 10 volumes, labeled 1, 2, ..., 10. These got jumbled & out of order over the years. King J asked Maurice to sort the collection but he can only take 2 books at once. The books are heavy hence he can switch only two per day. In how many days can Maurice guarantee that the volumes are sorted?



## King Julien's Books

Let's check

We can always place books in the right order in at most 9 days?



## King Julien's Books

Let's check

We can always place books in the right order in at most 9 days?



- Day 1: place Vol 1 on it's place

## King Julien's Books

Let's check

We can always place books in the right order in at most 9 days?



- Day 1: place Vol 1 on it's place

## King Julien's Books

Let's check

We can always place books in the right order in at most 9 days?



- Day 1: place Vol 1 on it's place
- Day 2: place Vol 2 on it's place

## King Julien's Books

Let's check

We can always place books in the right order in at most 9 days?




- Day 1: place Vol 1 on it's place
- Day 2: place Vol 2 on it's place



## King Julien's Books

### Let's check

We can always place books in the right order in at most 9 days?



- Day 1: place Vol 1 on it's place
- Day 2: place Vol 2 on it's place
- Day 3: place Vol 3 on it's place

## King Julien's Books

### Let's check

We can always place books in the right order in at most 9 days?



- Day 1: place Vol 1 on it's place
- Day 2: place Vol 2 on it's place
- Day 3: place Vol 3 on it's place
- And so on...

## King Julien's Books

### Let's check

We can always place books in the right order in at most 9 days?



- Day 1: place Vol 1 on it's place
- Day 2: place Vol 2 on it's place
- Day 3: place Vol 3 on it's place
- And so on...
- On Day 9, the first 9 volumes are on their proper places. The 10<sup>th</sup> volumes must also be in it's proper place since it's only one left.

## King Julien's Books

Let's check

- Are 9 days the optimal?



## King Julien's Books

### Let's check

- Are 9 days the optimal?
- What could be the hardest permutation of the books?



## King Julien's Books

### Let's check

- Are 9 days the optimal?
- What could be the hardest permutation of the books?

Recall:

- Permutation – order of selection is a factor
- Combination – order of selection not a factor

Ex: Permutation/Combination pairs from the set {A, B, C, D, E}

Permutation: AB AC AD AE BA BC BD BE CA CB CD CE DA DB DC DE EA EB EC ED

Combination: AB AC AD AE BC BE CD CE DE



## King Julien's Books

### Let's check

- Are 9 days the optimal?
- What could be the hardest permutation of the books?
  - It seems that the hardest is when the books are in opposite order



Recall:

- Permutation – order of selection is a factor
- Combination – order of selection not a factor

Ex: Permutation/Combination pairs from the set {A, B, C, D, E}

Permutation: AB AC AD AE BA BC BD BE CA CB CD CE DA DB DC DE EA EB EC ED

Combination: AB AC AD AE BC BE CD CE DE

## King Julien's Books

### Let's check

- Are 9 days the optimal?
- What could be the hardest permutation of the books?
  - It seems that the hardest is when the books are in opposite order





## King Julien's Books

### Let's check

- Are 9 days the optimal?
- What could be the hardest permutation of the books?
  - It seems that the hardest is when the books are in opposite order



## King Julien's Books

### Let's check

- Are 9 days the optimal?
- What could be the hardest permutation of the books?
  - It seems that the hardest is when the books are in opposite order



## King Julien's Books

### Let's check

- Are 9 days the optimal?
- What could be the hardest permutation of the books?
  - It seems that the hardest is when the books are in opposite order



➤ All the books can be switched and sorted in just 5 days!

## King Julien's Books

Let's check

- So what is the right number of days?



## King Julien's Books

### Let's check

- So what is the right number of days?
- And how to prove that it is the correct answer?

## King Julien's Books

### Let's check

- So what is the right number of days?
- And how to prove that it is the correct answer?
- To answer, we need to find some invariant that:



## King Julien's Books

### Let's check

- So what is the right number of days?
- And how to prove that it is the correct answer?
- To answer, we need to find some invariant that:
  - Do not change fast

## King Julien's Books

### Let's check

- So what is the right number of days?
- And how to prove that it is the correct answer?
- To answer, we need to find some invariant that:
  - Do not change fast
  - Should change substantially while ordering the book





## King Julien's Books

Recall this:

### *Puzzle*

- There is a sequence of 10 cells. The leftmost contains “1” while the rightmost has “30”. Is it possible to fill other cells with consecutive numbers in such a way that they differ by 3 at the most?

1									30
---	--	--	--	--	--	--	--	--	----

# King Julien's Books

The **Invariant**:



## King Julien's Books

The **Invariant**: The number of books staying to the right of their intended place



## King Julien's Books

The **Invariant**: The number of books staying to the right of their intended place

- Small in the end: **equals 0**

## King Julien's Books

The **Invariant**: The number of books staying to the right of their intended place

- Small in the end: **equals 0**
- Decreases slowly: **by at most 1 each day**

## King Julien's Books

The **Invariant**: The number of books staying to the right of their intended place

- Small in the end: **equals 0**
- Decreases slowly: **by at most 1 each day**
- Large in the beginning? **Yes!**

## King Julien's Books

The **Invariant**: The number of books staying to the right of their intended place

- Small in the end: **equals 0**
- Decreases slowly: **by at most 1 each day**
- Large in the beginning? **Yes!**



- The invariant is **9 in the beginning**, we need at **least 9 days**.

## Reminder

- Everybody, make sure that your name in ZOOM is in the following format:
  - Ex: 202054321 Juan Dela Cruz

Not changing your name to this format

\* you **will** be marked Absent \* → absent?

😊





Mathematical Thinking – Invariants

# **EVEN AND ODD NUMBERS**

- Even and Odd Numbers
- Piece on a Chessboard
- Summing up Digits
- Switching Signs
- Advanced Signs Switching

## Even and Odd Numbers

- Even numbers - integer numbers divisible by 2

## Even and Odd Numbers

- Even numbers - integer numbers divisible by 2
- Odd numbers - all others

## Even and Odd Numbers

- Even numbers - integer numbers divisible by 2
  - 2, 4, 6, 8, ... are even
- Odd numbers - all others
  - 1, 3, 5, 7, ... are even

## Even and Odd Numbers

- Even numbers - integer numbers divisible by 2
  - 2, 4, 6, 8, ... are even
- Odd numbers - all others
  - 1, 3, 5, 7, ... are even
- How about 0?

## Even and Odd Numbers

- Even numbers - integer numbers divisible by 2
  - 2, 4, 6, 8, ... are even
- Odd numbers - all others
  - 1, 3, 5, 7, ... are even
- How about 0?
  - Even! Recall section on “Connecting Points”

## Even and Odd Numbers

- Even numbers - integer numbers divisible by 2
  - 2, 4, 6, 8, ... are even
- Odd numbers - all others
  - 1, 3, 5, 7, ... are even
- How about 0?
  - Even! Recall section on “Connecting Points”

... -5 -4 -3 -2 -1 0 1 2 3 4 5 6 ...





## Even and Odd Numbers

- Even numbers - integer numbers divisible by 2
  - 2, 4, 6, 8, ... are even
- Odd numbers - all others
  - 1, 3, 5, 7, ... are even
- How about 0?
  - Even! Recall section on “Connecting Points”

... -5 -4 -3 -2 -1 0 1 2 3 4 5 6 ...



## Even and Odd Numbers

- The property of a number to be either even or odd is an important **variant**.



<https://i.pining.com/474x/0e/0f/54/0e0f54737e21ff81d07ce97ab90587ef.jpg>

- Even and Odd Numbers
- Piece on a Chessboard
- Summing up Digits
- Switching Signs
- Advanced Signs Switching

# Piece on a Chessboard


## Puzzle

- A piece on a chessboard can move to any cell adjacent by edge to the current one. Can it return to the original position after 17 moves? What about after 18 moves?

# Piece on a Chessboard

## Puzzle


- A piece on a chessboard can move to any cell adjacent by edge to the current one. Can it return to the original position after 17 moves? What about after 18 moves?



# Piece on a Chessboard

## Puzzle


- A piece on a chessboard can move to any cell adjacent by edge to the current one. Can it return to the original position after 17 moves? What about after 18 moves?



# Piece on a Chessboard

## Puzzle

- A piece on a chessboard can move to any cell adjacent by edge to the current one. Can it return to the original position after 17 moves? What about after 18 moves?




# Piece on a Chessboard

## Puzzle

- A piece on a chessboard can move to any cell adjacent by edge to the current one. Can it return to the original position after 17 moves? What about after 18 moves?

*17 moves, possible? §*






# Piece on a Chessboard

## Puzzle


- A piece on a chessboard can move to any cell adjacent by edge to the current one. Can it return to the original position after 17 moves? What about after 18 moves?



## Piece on a Chessboard

Let's check


- Start with the more simpler case of 18 moves



## Piece on a Chessboard

### Let's check

- Start with the more simpler case of 18 moves




- We can return after 2 steps
  - 1<sup>st</sup> step

## Piece on a Chessboard

### Let's check

- Start with the more simpler case of 18 moves




- We can return after 2 steps
  - 2<sup>nd</sup> step

## Piece on a Chessboard

### Let's check

- Start with the more simpler case of 18 moves




- We can return after 2 steps
  - Repeat 9 times

## Piece on a Chessboard

### Let's check

- Start with the more simpler case of 18 moves




- We can return after 2 steps
  - Repeat 9 times
- The piece would be back to its original place!

## Piece on a Chessboard

Let's check


- The same argument does not work for 17 steps



## Piece on a Chessboard

Let's check

- The same argument does not work for 17 steps




- Indeed, 17 is odd



## Piece on a Chessboard

### Let's check

- The same argument does not work for 17 steps




- Indeed, 17 is odd
- **Observation:** after even number of steps, a piece is on the lighter square

## Piece on a Chessboard

### Let's check

- The same argument does not work for 17 steps




- Indeed, 17 is odd
- **Observation:** after even number of steps, a piece is on the lighter square
- After odd number, it is on a dark square

## Piece on a Chessboard

### Let's check

- The same argument does not work for 17 steps



- Indeed, 17 is odd
- Observation:** after even number of steps, a piece is on the lighter square
- After odd number, it is on a dark square
- No way** to get back after odd number of steps

Mathematical Thinking – Invariants

# **SUMMING UP DIGITS**

- Even and Odd Numbers
- Piece on a Chessboard
- **Summing up Digits**
- Switching Signs
- Advanced Signs Switching

## Summing up Digits

### Problem I

- Is it possible to place signs in the form of the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$  to get as a result the sum 100? Can we get for 2?

## Summing up Digits

### Problem I

- Is it possible to place signs in the form of the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$  to get as a result the sum 100? Can we get for 2?

2-1.

$$\boxed{\pm 1} \quad \boxed{\pm 2} \quad \boxed{\pm 3} \quad \boxed{\pm 4} \quad \boxed{\pm 5} \quad \boxed{\pm 6} \quad \boxed{\pm 7} \quad \boxed{\pm 8} \quad \boxed{\pm 9} = \boxed{??}$$

100-?

# Summing up Digits

## Problem I

- Is it possible to place signs in the form of the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$  to get as a result the sum 100? Can we get for 2?
- Let's start with 100

0		1		2		3		4		5		6		7		8		9	=	
---	--	---	--	---	--	---	--	---	--	---	--	---	--	---	--	---	--	---	---	--

- We get largest sum if we place '+' everywhere





## Summing up Digits

### Problem I

- Is it possible to place signs in the form of the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$  to get as a result the sum 100? Can we get for 2?
  - Let's start with 100

0	+	1	+	2	+	3	+	4	+	5	+	6	+	7	+	8	+	9	=	45
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	----

45

- We get largest sum if we place '+' everywhere
- But we cannot get any value greater than 45, hence it is not possible have an expression that will give a sum of 100

## Summing up Digits

### Problem I

- Is it possible to place signs in the form of the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$  to get as a result the sum 100? **Can we get for 2?**
- Let's check for sum of 2

$$\boxed{\pm 1} \quad \boxed{\pm 2} \quad \boxed{\pm 3} \quad \boxed{\pm 4} \quad \boxed{\pm 5} \quad \boxed{\pm 6} \quad \boxed{\pm 7} \quad \boxed{\pm 8} \quad \boxed{\pm 9} = \boxed{??}$$

## Summing up Digits

### Problem I

- Is it possible to place signs in the form of the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$  to get as a result the sum 100? [Can we get for 2?](#)
- Let's check for sum of 2

$\pm 1$	$\pm 2$	$\pm 3$	$\pm 4$	$\pm 5$	$\pm 6$	$\pm 7$	$\pm 8$	$\pm 9$	$\square$
	/		/		/		/		

- Consider properties of the number sequence
  - Note that there are 5 odd and 4 even numbers

## Summing up Digits

### Problem I

- Is it possible to place signs in the form of the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$  to get as a result the sum 100? [Can we get for 2?](#)
- Let's check for sum of 2

$\pm 1$   $\pm 2$   $\pm 3$   $\pm 4$   $\pm 5$   $\pm 6$   $\pm 7$   $\pm 8$   $\pm 9$

- Consider properties of the number sequence
  - Note that there are 5 odd and 4 even numbers
  - $0 + 0 = E$ ,  $0 + E = O$ ,  $E + E = E$

## Summing up Digits

### Problem I

- Is it possible to place signs in the form of the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$  to get as a result the sum 100? **Can we get for 2?**
- Let's check for sum of 2

$\pm 1$   $\pm 2$   $\pm 3$   $\pm 4$   $\pm 5$   $\pm 6$   $\pm 7$   $\pm 8$   $\pm 9$

- Consider properties of the number sequence
  - Note that there are 5 odd and 4 even numbers
  - $O + O = E$ ,  $O + E = O$ ,  $E + E = E$
  - No matter how we put the signs, the sum will always be ODD



## Summing up Digits

### Problem I

- Is it possible to place signs in the form of the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$  to get as a result the sum 100? Can we get for 2?
- Let's check for sum of 2

$\pm 1$   $\pm 2$   $\pm 3$   $\pm 4$   $\pm 5$   $\pm 6$   $\pm 7$   $\pm 8$   $\pm 9$

- Consider properties of the number sequence
  - Note that there are 5 odd and 4 even numbers
  - $O + O = E$ ,  $O + E = O$ ,  $E + E = E$
  - No matter how we put the signs, the sum will always be ODD
- Hence the sum of 2 is impossible



# Summing up Digits

## Problem I

- Is it possible to place signs in the form of the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$  to get as a result the sum 100? Can we get for 2?
- Let's check for sum of 2

$\pm 1$   $\pm 2$   $\pm 3$   $\pm 4$   $\pm 5$   $\pm 6$   $\pm 7$   $\pm 8$   $\pm 9$

- Consider properties of the number
  - Note that there are 5 odd and 4 even numbers
  - $O + O = E$ ,  $O + E = O$ ,  $E + E = E$
  - No matter how we put the signs, the sum will always be ODD
- Hence the sum of 2 is impossible



## Summing up Digits

### Problem II

- Which sums is it possible to get by varying the signs in the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$ ?



## Summing up Digits

### Problem II

- Which sums is it possible to get by varying the signs in the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$ ?
  - We have seen two obstacles

## Summing up Digits

### Problem II

- Which sums is it possible to get by varying the signs in the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$ ?
  - We have seen two obstacles
    - Sum is ODD and sum is at **most 45** (symmetrically, at **least -45**)

## Summing up Digits

### Problem II

- Which sums is it possible to get by varying the signs in the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$ ?
  - We have seen two obstacles
    - Sum is **ODD** and sum is at **most 45** (symmetrically, at **least -45**)
    - Seems like this are the only restrictions

## Summing up Digits

### Problem II

- Which sums is it possible to get by varying the signs in the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$ ?
  - We have seen two obstacles
    - Sum is **ODD** and sum is at **most 45** (symmetrically, at **least -45**)
    - Seems like this are the only restrictions
  - How about the other sums?

## Summing up Digits

### Problem II

- Which sums is it possible to get by varying the signs in the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$ ?
  - We have seen two obstacles
    - Sum is **ODD** and sum is at **most 45** (symmetrically, at **least -45**)
    - Seems like this are the only restrictions
  - How about the other sums?
    - Turns out, all sums satisfying the found restrictions are possible!

## Summing up Digits

### Problem II

- Which sums is it possible to get by varying the signs in the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$ ? 45 + 50 = 0
- We have seen two obstacles 9/1 (+) all (-)
  - Sum is **ODD** and sum is at most 45 (symmetrically, at **least -45**)
  - Seems like this are the only restrictions
- How about the other sums?
  - Turns out, all sums satisfying the found restrictions **are possible!**
- You can check this out yourself.

## Summing up Digits

### Problem II

- Which sums is it possible to get by varying the signs in the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$ ? [-45, -43, ..., 43, 45]
- We have seen two obstacles
  - Sum is **ODD** and sum is at **most 45** (symmetrically, at **least -45**)
  - Seems like this are the only restrictions
- How about the other sums?
  - Turns out, all sums satisfying the found restrictions **are possible!**
- You can check this out yourself.
- Try this “greedy algorithm”, start from left to right and place the signs greedily through:

## Summing up Digits

### Problem II

- Which sums is it possible to get by varying the signs in the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$ ?
  - We have seen two obstacles
    - Sum is **ODD** and sum is at **most 45** (symmetrically, at **least -45**)
    - Seems like this are the only restrictions
  - How about the other sums?
    - Turns out, all sums satisfying the found restrictions **are possible!**
  - You can check this out yourself.
  - Try this “greedy algorithm”, start from left to right and place the signs greedily through:
    - If current sum is less than goal, **increase** the sum



## Summing up Digits

### Problem II

- Which sums is it possible to get by varying the signs in the expression  $\pm 1 \pm 2 \pm 3 \pm \dots \pm 9$ ?
  - We have seen two obstacles
    - Sum is **ODD** and sum is at **most 45** (symmetrically, at **least -45**)
    - Seems like this are the only restrictions
  - How about the other sums?
    - Turns out, all sums satisfying the found restrictions **are possible!**
  - You can check this out yourself.
  - Try this “greedy algorithm”, start from left to right and place the signs greedily through:
    - If current sum is less than goal, **increase** the sum
    - Else **decrease** it.

Mathematical Thinking – Invariants

# **SWITCHING SIGNS**

- Even and Odd Numbers
- Piece on a Chessboard
- Summing up Digits
- **Switching Signs**
- Advanced Signs Switching

## Switching Signs

### Puzzle


- Given a 4 x 4 table. The top-left corner has the ‘-’ sign and all other cells has “+” sign. For each step, we are allowed to switch all the signs in a column or row. Is it possible to switch all the signs to “+”?



# Switching Signs

## Puzzle

- Given a 4 x 4 table. The top-left corner has the ‘-’ sign and all other cells has “+” sign. For each step, we are allowed to switch all the signs in a column or row. Is it possible to switch all the signs to “+”?



-	+	(+)	+
(+)	(+)	(+)	(+)
+	+	(+)	+
+	+	(+)	+

## Switching Signs

Let's try (Case I)

+	+	+	+

- Let's take the case of switching a row

## Switching Signs

Let's try (Case I)

↓

( - )			
-	-	-	-

( )

- Let's take the case of switching a row
- If we try to do it, we will notice that the number of minus signs is always ODD
  - Take note that we started with a “-” sign in the upper-left cell

## Switching Signs

Let's try (Case II)

+	+	-	+

- Let's take another case



## Switching Signs

Let's try (Case II)

-	-	+	-

$(7) = 0 \text{ } 1 \text{ } 2$

-		(+)	
-	-	(+)	-
		(+)	
		(+)	

- Let's take another case
- If we try to do it, we will notice that the number of minus signs is still ODD

## Switching Signs

Let's try (Case III)

-	+	-	+

- Let's try another case

## Switching Signs

Let's try (Case III)

+	-	+	-

- Let's take another case
- If we try to do it, we will notice that the number of minus signs is still ODD
  - recall that we started with one “-” sign

## Switching Signs

Let's try

-	+	+	+
+	+	+	+
+	+	+	+
+	+	+	+

- After trying it, we will notice that the number of minus signs is always **ODD**



## Switching Signs

Let's try

-	+	+	+
+	+	+	+
+	+	+	+
+	+	+	+

$$= 14 = \text{even}$$


$$H(-) = 0217$$

- After trying it, we will notice that the number of minus signs is always **ODD**
- This is an invariant!

## Switching Signs

Let's try

-	+	+	+
+	+	+	+
+	+	+	+
+	+	+	+

- After trying it, we will notice that the number of minus signs is always **ODD** 
- This is an invariant!
- Hence, **we cannot switch all to “+”**, since the number of “-” signs would then be **zero**, which is even and violates this invariant

Mathematical Thinking – Invariants

# **ADVANCED SIGNS SWITCHING**

- Even and Odd Numbers
- Piece on a Chessboard
- Summing up Digits
- Switching Signs
- **Advanced Signs Switching**



# Advanced Signs Switching

## Puzzle

- Given a 4 x 4 table. All corner cells contain the ‘-’ sign and all other cells has “+” sign. For each step, we are allowed to switch all the signs in a column or row. Is it possible to switch all the signs to “+”?

-	+	+	-
+	+	+	+
+	+	+	+
-	+	+	-

# Advanced Signs Switching

Take note

-	+	+	-
+	+	+	+
+	+	+	+
-	+	+	-

- This looks more tricky. **The previous solution won't work.**

# Advanced Signs Switching

Take note

-	+	+	-
+	+	+	+
+	+	+	+
-	+	+	-

- This looks more tricky. **The previous solution won't work.**
- If you know about residues modulo 4 (we will learn later), you can use them but it also **does not help.**



## Advanced Signs Switching

Take note

-	+	+	-
+	+	+	+
+	+	+	+
-	+	+	-

- This looks more tricky. **The previous solution won't work.**
- If you know about residues modulo 4 (we will learn later), you can use them but it also **does not help**.
- There is a very simple solution

# Advanced Signs Switching

Take note

-	+	+	-
+	+	+	+
+	+	+	+
-	+	+	-

- This looks more tricky. **The previous solution won't work.** ✕
- If you know about residues modulo 4 (we will learn later), you can use them but it also **does not help.** ✕
- There is a very simple solution
  - Before proceeding, try to analyze more and you might find about it ☺



# Advanced Signs Switching


## Puzzle

- Given a 4 x 4 table. All corner cells contain the ‘-’ sign and all other cells has “+” sign. For each step, we are allowed to switch all the signs in a column or row. Is it possible to switch all the signs to “+”?

-	+	+	-
+	+	+	+
+	+	+	+
-	+	+	-

# Advanced Signs Switching

Let's try



-	+	+	-
+	+	+	+
+	+	+	+
-	+	+	-

- The idea is simple. Let's look at the small part of the problem

# Advanced Signs Switching

Let's try

-	+	+	-
+	+	+	+
+	+	+	+
-	+	+	-

- The idea is simple. Let's look at the small part of the problem
- If we switch a row or column in the large table, we switch a row or column in the smaller one (or do nothing)



# Advanced Signs Switching

Let's try

-	+	+	-
+	+	+	+
+	+	+	+
-	+	+	-

- To solve the big problem, we need to solve this small 2 x 2 problem

# Advanced Signs Switching

Let's try

-	+	+	-
+	+	+	+
+	+	+	+
-	+	+	-

- To solve the big problem, we need to solve this small 2 x 2 problem
- But this 2 x 2 problem cannot be solved using the previous argument!

## Advanced Signs Switching


Let's try

-	+	+	-
+	+	+	+
+	+	+	+
-	+	+	-

- We have a very strong obstacle – each 2 x 2 square should have even number of '-' signs for the puzzle to be solved

# Advanced Signs Switching

Let's try




A 4x4 grid of signs. The top-left 2x2 subgrid is highlighted with a red border. The signs in the grid are as follows:

-	+	+	-
+	+	+	+
+	+	+	+
-	+	+	-

- We have a very strong obstacle – each 2 x 2 square should have even number of '-' signs for the puzzle to be solved
- In fact, these are the only obstacles. That is, if all 2 x 2 squares has even number of '-' signs, then it is possible to switch all the signs into "+"

# Advanced Signs Switching


## Hint



- Assume that **our invariant** holds: each 2 x 2 square has even number of “-” signs

# Advanced Signs Switching

## Hint




- Assume that **our invariant** holds: each  $2 \times 2$  square has even number of “-” signs
- Switch the 1<sup>st</sup> row and 1<sup>st</sup> column to all “+”

# Advanced Signs Switching

## Hint

+	+	+	+
+			
+			
+			



- Assume that **our invariant** holds: each 2 x 2 square has even number of “-” signs
- Switch the 1<sup>st</sup> row and 1<sup>st</sup> column to all “+”

# Advanced Signs Switching

## Hint

+	+	+	+
+			
+			
+			

- Assume that **our invariant** holds: each  $2 \times 2$  square has even number of “-” signs
- Switch the 1<sup>st</sup> row and 1<sup>st</sup> column to all “+”
- Use invariant to show that the rest of the table is filled with “+”



# Advanced Signs Switching

## Hint

+	+	+	+
+	+		
+			
+			

- Assume that our invariant holds: each  $2 \times 2$  square has even number of “-” signs
- Switch the 1<sup>st</sup> row and 1<sup>st</sup> column to all “+”
- Use invariant to show that the rest of the table is filled with “+”
  - Zero is even

# Advanced Signs Switching

## Hint

+	+	+	+
+	+	+	+
+			
+			

- Assume that **our invariant** holds: each  $2 \times 2$  square has even number of “-” signs
- Switch the 1<sup>st</sup> row and 1<sup>st</sup> column to all “+”
- Use invariant to show that the rest of the table is filled with “+”
  - Zero is even

# Advanced Signs Switching

## Hint

+	+	+	+
+	+	+	+
+	+		
+	+		

- Assume that **our invariant** holds: each  $2 \times 2$  square has even number of “-” signs
- Switch the 1<sup>st</sup> row and 1<sup>st</sup> column to all “+”
- Use invariant to show that the rest of the table is filled with “+”
  - Zero is even

# Advanced Signs Switching

## Hint

+	+	+	+
+	+	+	+
+	+	+	+
+	+	+	+

- Assume that **our invariant** holds: each  $2 \times 2$  square has even number of “-” signs
- Switch the 1<sup>st</sup> row and 1<sup>st</sup> column to all “+”
- Use invariant to show that the rest of the table is filled with “+”
  - Zero is even



## Conclusion

- **Invariants** are important

## Conclusion

- Invariants are important
- Helps to prove impossibility

## Conclusion

- **Invariants** are important
- Helps to **prove impossibility**
- Also to establish **termination** and **running time bounds** (very important)

## Conclusion

- **Invariants** are important
- Helps to **prove impossibility**
- Also to establish **termination** and **running time bounds** (very important)
- Can take many forms



## Conclusion

- **Invariants** are important
- Helps to **prove impossibility**
- Also to establish **termination** and **running time bounds** (very important)
- Can take many forms
  - numbers, “even or odd”, equations, inequalities, etc...

## Conclusion

- **Invariants** are important
- Helps to **prove impossibility**
- Also to establish **termination** and **running time bounds** (very important)
- Can take many forms
  - numbers, “even or odd”, equations, inequalities, etc...
- **Double counting** is a special case

## Conclusion

- **Invariants** are important
- Helps to **prove impossibility**
- Also to establish **termination** and **running time bounds** (very important)
- Can take many forms
  - numbers, “even or odd”, equations, inequalities, etc...
- **Double counting** is a special case
- Yet, this is not all about invariants.

## Conclusion

- **Invariants** are important
- Helps to **prove impossibility**
- Also to establish **termination** and **running time bounds** (very important)
- Can take many forms
  - numbers, “even or odd”, equations, inequalities, etc...
- **Double counting** is a special case
- Yet, this is not all about invariants.
  - Still other forms (ex: residues); we will see later on

**Thank you.**