

A Modular Time-Dynamics Framework Combining Relativistic Physics with the Proposed VFE1 and QCC0 Formalisms

Abstract

We propose a modular computational framework for investigating time evolution across relativistic, quantum, and geometric regimes. The framework separates established physical models from two proposed components: Vibrational Field Equation One (VFE1), intended to describe an additional field-dependent contribution to temporal evolution, and Quantum Computerization Consciousness Zero (QCC0), introduced as a quantitative measure of quantum-state correlation and computational coherence within the model.

The framework does not replace established relativity or quantum mechanics. Instead, special relativity, general relativity, and conventional quantum-mechanical descriptions provide the reference layer against which the proposed VFE1 and QCC0 formulations can be evaluated. The Kapnack Solver provides the numerical engine, while the 369 vortex formulation is treated as a proposed geometric/kinematic transformation layer within the broader SDKP architecture.

The central objective is falsifiable comparison: for a specified physical system, the framework calculates the prediction obtained from established physics, calculates the corresponding prediction obtained after inclusion of the proposed VFE1/QCC0 terms, and compares both predictions with independently measured data. Any improvement must be demonstrated quantitatively and must survive independent reproduction.

1. Introduction

Time is a fundamental parameter in modern physics, but its operational behavior depends upon the physical state and reference frame of the system under consideration.

In special relativity, proper time depends upon relative motion. For an inertial observer, the differential relation between proper time and coordinate time is

$$\begin{aligned} &[\\ &d\tau = dt \sqrt{1 - \frac{v^2}{c^2}}, \\ &] \end{aligned}$$

where (v) is relative velocity and (c) is the invariant speed of light.

In general relativity, spacetime geometry is determined by the gravitational field, and the proper time measured by a clock depends upon its worldline through spacetime. In the weak-field limit, gravitational time dilation can be approximated by the corresponding metric or gravitational-potential correction.

Quantum mechanics introduces a separate description of physical evolution in which states evolve according to a dynamical equation such as

[

$$i\hbar\frac{\partial}{\partial t}\left|\psi(t)\right\rangle$$

$$\hat{H}\left|\psi(t)\right\rangle .$$

]

These established theories provide the reference framework for the present proposal.

The purpose of the present work is to investigate whether additional geometric, density, kinetic, positional, or field-state variables can provide useful predictive information beyond the standard reference calculations.

Within the proposed SDKP architecture, these additional variables are organized through the Scale-Density-Kinetic/kinematic framework and associated computational methods. Two components are particularly relevant to temporal modeling:

1. **Vibrational Field Equation One (VFE1)** — a proposed field-based temporal correction or extension.
2. **Quantum Computerization Consciousness Zero (QCC0)** — a proposed quantitative descriptor of quantum correlation/coherence within the computational representation.

Both are treated here as hypotheses requiring empirical validation rather than as established physical laws.

2. Established Physical Baseline

2.1 Special Relativity

The first reference model is special-relativistic proper time.

For motion at constant velocity,

$$\begin{aligned} &[\\ & d\tau = \\ & dt \sqrt{1 - \frac{v^2}{c^2}}. \\ &] \end{aligned}$$

Integration gives

$$\begin{aligned} &[\\ & \Delta\tau = \\ & \Delta t \sqrt{1 - \frac{v^2}{c^2}} \\ &] \end{aligned}$$

for constant (v).

This relationship provides an experimentally established baseline for evaluating any proposed modification to velocity-dependent temporal evolution.

The framework therefore requires the relativistic calculation to remain available independently of VFE1 and QCC0.

2.2 General Relativity

For gravitational systems, the reference calculation is obtained from the spacetime metric

$$\begin{aligned} &[\\ & ds^2 = g_{\mu\nu} dx^\mu dx^\nu . \\ &] \end{aligned}$$

For a timelike trajectory,

$$\left[\begin{aligned} d\tau^2 = \\ -\frac{ds^2}{c^2} \end{aligned} \right]$$

under the appropriate metric-signature convention.

Thus, gravitational time dilation is not introduced as an independent assumption of the proposed framework. It is calculated from the established spacetime geometry.

In weak gravitational fields, the resulting prediction may be expressed approximately in terms of gravitational potential and velocity. These established predictions constitute the baseline against which the proposed VFE1 contribution is tested.

3. Proposed SDKP Temporal Representation

The proposed SDKP architecture introduces a state representation containing geometric scale, density, kinetic state, and position:

$$\left[\begin{aligned} \Psi = \\ (S, \rho, K, P). \end{aligned} \right]$$

Here,

- (S) represents structural or geometric scale,
- (ρ) represents a density or density distribution,
- (K) represents kinetic state,
- (P) represents spatial position.

The important distinction is that these quantities are treated as model variables rather than being assumed automatically to constitute new physical laws.

The computational objective is therefore to determine whether an appropriate transformation

$$\left[\begin{aligned} \Psi \end{aligned} \right]$$

$\longrightarrow$
 T
]

can reproduce known temporal behavior and, more importantly, whether it produces independently testable predictions that improve upon established models.

4. Vibrational Field Equation One (VFE1)

VFE1 is proposed as a field-dependent extension of temporal dynamics.

The general form may be represented as

[

$\mathcal{T}_{\mathrm{VFE1}}$

$\mathcal{F}$
 $\left($
 S, ρ, K, P, Φ_v
 $\right),$
]

where Φ_v represents the proposed vibrational-field state.

The precise functional form of $\mathcal{F}$ must be specified for each physical application and must not be selected solely to reproduce an existing observation.

A useful formulation is therefore to decompose the prediction into a standard component and an additional VFE1 contribution:

[

T_{model}

T_{standard}
 $+$
 ΔT_{VFE1} .
 $]$

The term

[
 ΔT_{VFE1}
 $]$

is the quantity that must be independently constrained.

This construction has an important scientific advantage: the VFE1 hypothesis can be falsified. If

[
 $\Delta T_{\mathrm{VFE1}} \rightarrow 0$
 $]$

is consistently required by experimental data, then VFE1 provides no experimentally distinguishable temporal correction in that regime.

Conversely, if a nonzero correction is predicted before measurement and is subsequently observed, the result provides evidence requiring further investigation.

5. Quantum Computerization Consciousness Zero (QCC0)

QCC0 is proposed as a computational quantity associated with the representation of quantum-state correlation, coherence, or information consistency.

For a quantum state described by density operator (ρ_q), established quantum mechanics provides quantities such as purity,

[

$\mathcal{P}$

$\mathrm{Tr}(\rho_q^2),$
]

and entanglement or correlation measures may be defined according to the physical system and experimental protocol.

QCC0 should therefore be defined in relation to measurable quantum quantities rather than assumed to represent a new physical substance or consciousness mechanism.

A general proposed mapping can be written as

[
QCC0 =
 $\mathcal{Q}$
 $\left($
 $\rho_q,$
 $\mathcal{C},$
 $\mathcal{E},$
 $\mathcal{I}$
 $\right),$
]

where the arguments represent appropriately defined quantum-state and information-theoretic quantities.

The exact definition of $\mathcal{Q}$ is an empirical question.

If QCC0 is intended to represent quantum coherence, then its numerical value should be compared with independently measurable coherence metrics. If it is intended to represent a broader computational invariant, that invariant must be explicitly defined and tested for reproducibility.

Accordingly, QCC0 is treated in this work as a proposed computational construct rather than an established physical constant.

6. Relationship Between VFE1 and QCC0

The two proposed components serve different conceptual purposes.

VFE1 is associated primarily with **physical field evolution and temporal dynamics**, whereas QCC0 is associated primarily with **quantum-state correlation and computational representation**.

A combined model may therefore be expressed schematically as

[

$\mathcal{M}$

$$\mathcal{M}_{\mathrm{GR/SR}} + \Delta \mathcal{M}_{\mathrm{VFE1}} + \Delta \mathcal{M}_{\mathrm{QCC0}}.$$

]

The terms should not be interpreted as automatically additive physical forces or energies. They represent model contributions whose physical interpretation must be established through dimensional analysis and experiment.

The framework consequently imposes three requirements:

1. Every new variable must have a defined mathematical meaning.
 2. Every predicted observable must have physical units or be explicitly dimensionless.
 3. Every proposed correction must generate an experimentally distinguishable prediction.
-

7. Kapnack Solver

The Kapnack Solver is the computational engine responsible for evaluating the model.

Its role is not to establish physical validity by itself. Instead, it provides a reproducible numerical implementation of the equations being tested.

A typical calculation follows:

```
[  
  \text{Initial Conditions}  
  \rightarrow  
  \text{Established Physics}  
  \rightarrow  
  \text{VFE1}  
  \rightarrow  
  \text{QCC0}  
  \rightarrow  
  \text{Observable Prediction}.  
]
```

The solver should preserve the intermediate results so that an investigator can determine whether an observed difference originates from:

- the relativistic baseline,
- the VFE1 contribution,
- the QCC0 contribution,
- numerical approximation,
- parameter selection,
- or measurement uncertainty.

This separation is essential for scientific reproducibility.

8. 369 Vortex Representation

The 369 vortex formulation is incorporated as a proposed geometric or kinematic transformation within the computational framework.

Rather than assuming that the numerical sequence 3-6-9 has a fundamental physical significance, the framework treats the 369 structure as a hypothesis whose physical consequences must be specified mathematically.

A valid implementation therefore requires a mapping

```
[  
  \mathcal{V}_{369}:  
  \Psi  
  \rightarrow  
  \Psi',  
]
```

where both the input and output states are explicitly defined.

The transformation can then be tested independently.

If the transformation produces no measurable prediction beyond an equivalent conventional transformation, it may be mathematically useful without constituting a new physical law. If it produces a novel prediction that is subsequently verified experimentally, the transformation would warrant further physical investigation.

9. Comparative Prediction Architecture

The proposed framework is designed around a three-level comparison.

Level I — Established Physics

Calculate the observable using the accepted reference model:

```
[  
  O_{\mathrm{standard}}.  
]
```

Level II — SDKP/VFE1

Calculate the same observable after introducing the proposed VFE1 contribution:

$$\left[\begin{array}{l} O_{\mathrm{VFE1}}. \end{array} \right]$$

Level III — SDKP/VFE1/QCC0

Calculate the observable after inclusion of both proposed components:

$$\left[\begin{array}{l} O_{\mathrm{combined}}. \end{array} \right]$$

The predictions are then compared with an independently measured value:

$$\left[\begin{array}{l} O_{\mathrm{obs}}. \end{array} \right]$$

The residuals are

$$\left[\right]$$

$$R_{\mathrm{standard}}$$

$$\left[\begin{array}{l} O_{\mathrm{obs}} - O_{\mathrm{standard}}, \end{array} \right]$$

[

R_{VFE1}

$O_{\mathrm{obs}} - O_{\mathrm{VFE1}},$
]

and

[

R_{combined}

$O_{\mathrm{obs}} - O_{\mathrm{combined}}.$
]

A proposed extension is supported only if the reduction in residual error is statistically and experimentally meaningful, while accounting for the number of fitted parameters and measurement uncertainty.

10. Falsifiability

The central methodological principle is falsifiability.

The framework should specify its predictions before comparison with experimental data whenever possible.

A VFE1 or QCC0 model should be considered unsuccessful in a given domain if:

- it produces predictions systematically inconsistent with observation;
- its parameters must be continually adjusted after measurements are known;
- it provides no improvement over the established model;
- or its apparent improvement disappears under independent reproduction.

Likewise, successful agreement with existing data alone does not establish a new physical theory because a sufficiently flexible model can reproduce known results.

The strongest evidence would therefore be a **novel, quantitative prediction made in advance and subsequently confirmed by an independent experiment.**

11. Experimental Validation Program

The proposed framework can be evaluated progressively.

Stage 1 — Mathematical consistency

Verify:

- dimensional consistency,
- limiting behavior,
- numerical stability,
- conservation laws where applicable,
- compatibility with established relativistic limits.

Stage 2 — Recovery of established physics

The proposed model should reproduce established results when its additional terms are placed in the appropriate limiting regime.

For example,

[
 $\Delta T_{\mathrm{VFE1}} \rightarrow 0$
]

should recover the standard relativistic result.

Stage 3 — Numerical benchmark

Compare the Kapnack Solver against independently implemented reference calculations.

Stage 4 — Known experimental data

Evaluate the model against datasets for which reliable measurements already exist.

Stage 5 — Blind prediction

Generate predictions without fitting to the future measurement.

Stage 6 — Independent replication

Require an independent implementation or experimental group to reproduce the result.

12. Scientific Status of the Framework

The framework proposed here contains two distinct categories of knowledge.

Established physical foundations include special relativity, general relativity, quantum mechanics, quantum-state formalism, and experimentally established time-dilation effects.

Proposed components include the specific VFE1 formulation, QCC0 definition, SDKP temporal representation, Kapnack Solver methodology, and 369 vortex transformation.

The distinction is intentional.

The purpose of the framework is not to label a proposed mathematical construction as established physics before experimental validation. Rather, it provides a structured computational environment in which proposed extensions can be compared directly with established theories.

13. Conclusion

We present a modular framework for investigating temporal dynamics across relativistic, quantum, geometric, and computational regimes. Established relativistic and quantum theories provide the reference layer. VFE1 is introduced as a proposed field-dependent temporal extension, while QCC0 is introduced as a proposed quantitative descriptor of quantum correlation or computational coherence.

The Kapnack Solver provides the computational implementation, while the 369 vortex formulation is treated as a proposed geometric/kinematic transformation that requires explicit mathematical definition and independent testing.

The central methodological requirement is comparative and falsifiable prediction. Any proposed extension must reproduce established physics in appropriate limits, maintain dimensional and mathematical consistency, and produce independently testable predictions that can be distinguished from conventional theory.

A Modular Time-Dynamics Framework Combining Relativistic Physics with the Proposed VFE1 and QCC0 Formalisms

Version 0.2

Donald Paul Smith

FatherTimeSDKP

ORCID: 0009-0003-7925-1653

Gypsi Consulting

Zenodo Version 0.2 — 2026

Abstract

We present a modular computational framework for investigating temporal dynamics across relativistic, quantum, geometric, and computational regimes. The framework retains special relativity, general relativity, and conventional quantum mechanics as established reference models while introducing several proposed computational components within the Scale-Density-Kinetic/Kinematic Principle (SDKP) architecture.

The principal proposed temporal component is **Vibrational Field Equation One (VFE1)**, formulated as a field-dependent extension of temporal evolution. A second component,

Quantum Computerization Consciousness Zero (QCC0), is introduced as a proposed computational quantity for representing quantum-state correlation, coherence, and information consistency. The **Kapnack Solver** provides the numerical evaluation layer, while the **369 vortex formulation** is represented as a proposed geometric/kinematic transformation.

Version 0.2 formalizes the framework as a comparative model rather than as a replacement for established physics. Each proposed contribution is evaluated against an independent baseline obtained from established physical equations. The framework therefore produces three classes of prediction: the established-physics prediction, the prediction including VFE1, and the combined prediction including VFE1 and QCC0.

The principal scientific requirement is falsifiability. A proposed component must have defined variables, units, boundary conditions, limiting behavior, and independently testable predictions. Agreement with previously known observations is insufficient by itself to establish a new physical law. Strong evidence would require a quantitative prediction generated independently of the measurement and subsequently confirmed by experiment or an independent implementation.

1. Introduction

Time is an operationally measurable quantity whose relationship to motion and gravitational environment is established by relativistic physics.

Special relativity predicts that proper time depends upon the relative motion of a clock. For an inertial observer,

$$\begin{aligned} &[\\ &d\tau = \\ &dt\sqrt{1-\frac{v^2}{c^2}}, \\ &] \end{aligned}$$

where ($d\tau$) is proper time, (dt) is coordinate time, (v) is relative velocity, and (c) is the invariant speed of light.

General relativity extends this description by treating gravity as a manifestation of spacetime geometry. Proper time is determined by the metric and the worldline of the clock:

$$\begin{aligned} &[\\ &d\tau^2 = \\ &-\frac{1}{c^2}g_{\mu\nu}dx^\mu dx^\nu, \\ &] \end{aligned}$$

under a metric convention in which timelike intervals have negative signature.

Quantum mechanics provides a separate but compatible framework for describing quantum-state evolution. A state vector evolves according to the Schrödinger equation,

$$i\hbar\frac{\partial}{\partial t}|\psi\rangle = \hat{H}|\psi\rangle.$$

The present work does not replace these established descriptions.

Instead, it asks whether additional geometric, density, kinetic, positional, or field-state information can provide an experimentally useful extension to temporal modeling.

The proposed framework organizes these variables through the SDKP state representation

$$|\Psi\rangle = (S, \rho, K, P),$$

where (S) represents structural scale, (ρ) represents density or density distribution, (K) represents kinetic/kinematic state, and (P) represents spatial position.

Three additional computational components are considered:

1. **VFE1** — proposed field-dependent temporal dynamics.
2. **QCC0** — proposed quantum-state correlation/coherence representation.
3. **369 vortex transformation** — proposed geometric/kinematic transformation.

The Kapnack Solver provides the computational engine that evaluates these representations.

2. Scientific Status and Scope

The framework contains two fundamentally different categories of material.

Established physics

The reference layer consists of:

- special relativity;
- general relativity;
- quantum mechanics;
- quantum-state and density-matrix formalism;
- established time-dilation measurements;
- established quantum-information measures.

Proposed components

The following are presented as hypotheses or computational constructs:

- the SDKP temporal representation;
- VFE1;
- QCC0;
- the 369 vortex transformation;
- the specific Kapnack Solver implementation.

The proposed components are therefore not asserted here to be established physical laws.

The purpose of the framework is to make them sufficiently explicit that their predictions can be compared with established models and experimental data.

3. SDKP State Representation

The fundamental computational state is represented as

[
 $\Psi = (S, \rho, K, P)$.
]

The variables are defined as follows.

3.1 Scale (S)

(S) describes the characteristic geometric or structural scale of the system.

Depending upon the application, (S) may represent:

- length;
- characteristic radius;
- characteristic wavelength;
- system dimension;
- geometric scale parameter.

The physical units of (S) must be specified for each application.

3.2 Density (ρ)

(ρ) describes the distribution of a relevant physical quantity over space.

Depending upon the model, this may represent:

- mass density;
- energy density;
- probability density;
- information density;
- another explicitly defined scalar or tensor field.

The physical interpretation must not be changed during a calculation without an explicit transformation.

3.3 Kinetic/Kinematic State (K)

(K) represents the motion or dynamical state of the system.

Depending upon the application, it may include:

[
K=(v,a,p,E,\ldots),
]

where (v) is velocity, (a) acceleration, (p) momentum, and (E) energy.

3.4 Position (P)

(P) represents the spatial location or configuration of the system.

In a three-dimensional Cartesian representation,

$$[P=(x,y,z).]$$

In curved spacetime, (P) may instead be represented by spacetime coordinates.

4. Established Relativistic Baseline

The framework requires an independent calculation using established physics before any proposed correction is introduced.

For constant velocity,

$$[$$

$$\Delta t_{\mathrm{SR}}$$

$$\Delta t \sqrt{1-\frac{v^2}{c^2}}.]$$

For gravitational systems, the appropriate spacetime metric provides the corresponding proper-time calculation:

$$[$$

$\Delta\tau_{\mathrm{GR}}$

$$\int \sqrt{-\frac{g_{\mu\nu} dx^\mu dx^\nu}{c^2}}.$$

The resulting quantity is designated

$$T_{\mathrm{GR/SR}}.$$

This quantity serves as the reference prediction.

The proposed model is not permitted to replace this baseline merely because another calculation produces a different result.

5. VFE1: Vibrational Field Equation One

5.1 Conceptual definition

VFE1 is proposed as an additional field-dependent representation of temporal evolution.

Let

$$\Phi_v(x,t)$$

represent the proposed vibrational field state.

The general VFE1 temporal functional is written as

[

$$T_{\mathrm{VFE1}}$$

$$\begin{aligned} & \mathcal{F} \\ & \left[\right. \\ & S, \rho, K, P, \Phi_v \\ & \left. \right]. \end{aligned}$$

For comparative analysis, the model is decomposed as

[

$$T_{\mathrm{VFE1}}$$

$$\begin{aligned} & T_{\mathrm{standard}} \\ & + \\ & \Delta T_{\mathrm{VFE1}}. \end{aligned}$$

Here,

$$[T_{\mathrm{standard}}]$$

is calculated independently from established physics and

$$[\Delta T_{\mathrm{VFE1}}]$$

represents the proposed additional contribution.

5.2 Field evolution

A general vibrational field can be represented by a scalar field

$$[\Phi_v(x,t).$$

A generic wave-like field equation may be written as

$$[$$

$$\frac{\partial^2 \Phi_v}{\partial t^2}$$

$v_v^2 \ln$

$abla^2$

Φ_v

+

\Gamma\frac{\partial\Phi_v}{\partial t}

+

$$\nabla^2 \Phi_v$$

$$J(S, \rho, K, P, t),$$

where:

- v_v is the propagation parameter;
- Γ is a damping parameter;
- $V(\Phi_v)$ is a potential function;
- J is a source term;
- (S, ρ, K, P) describe the system interacting with the field.

This equation is presented as a general mathematical template rather than an established physical law.

The specific VFE1 equation must ultimately be selected and justified through dimensional analysis, limiting behavior, and experiment.

6. VFE1 Temporal Coupling

A proposed temporal coupling can be represented as

[

$$\Delta T_{\mathrm{VFE1}}$$

$$\lambda_v \mathcal{G}(\Phi_v, \rho, K, P, S),$$

]

where λ_v is a coupling parameter and $\mathcal{G}$ is a dimensionally consistent functional.

The units of λ_v must be chosen so that

$$[\Delta T_{\mathrm{VFE1}}] = [\mathrm{time}].$$

]

This requirement prevents a dimensionally undefined correction from being interpreted as a physical prediction.

The model must also satisfy an established-physics limit:

$$\lambda_v \rightarrow 0 \quad \Longleftrightarrow \quad T_{\mathrm{VFE1}} \rightarrow T_{\mathrm{standard}}.$$

]

7. QCC0: Quantum Computerization Consciousness Zero

7.1 Operational definition

QCC0 is introduced as a proposed computational quantity associated with quantum-state correlation, coherence, and information consistency.

The term “consciousness” in the name is retained as the identifier of the proposed formalism; it does not, by itself, establish that consciousness is a physical variable or that quantum systems possess consciousness.

For a quantum state described by density operator

[
 ρ_q ,
]

established quantum mechanics provides measurable quantities such as purity:

[

$$P$$

$$\text{Tr}(\rho_q^2).$$

Other measurable quantities may include:

- von Neumann entropy;
- fidelity;
- coherence measures;

- mutual information;
- entanglement measures;
- quantum Fisher information.

QCC0 is therefore defined operationally as a proposed mapping

[

QCC0

$$\mathcal{Q} \left(\rho_q, \mathcal{C}, \mathcal{E}, \mathcal{I} \right)$$

where the inputs must correspond to explicitly defined quantum-information quantities.

8. QCC0 Normalization

For comparative computational purposes, QCC0 may be normalized to a bounded interval:

$$0 \leq \text{QCC0} \leq 1.$$

A value of zero would represent the selected minimum reference condition, while a value approaching unity would represent the selected maximum correlation/coherence condition.

This normalization is a computational convention unless a physical measurement procedure is specified.

Accordingly, a numerical value such as

$$\left[\begin{array}{l} \text{QCC0}=1 \end{array} \right]$$

must not automatically be interpreted as proof of perfect physical coherence or entanglement.

The experimental meaning of QCC0 must be established through comparison with independently measurable quantum observables.

9. Combined VFE1-QCC0 Model

The combined model is represented as

[

T_{combined}

$$\begin{aligned} &T_{\mathrm{standard}} \\ &+ \\ &\Delta T_{\mathrm{VFE1}} \\ &+ \\ &\Delta T_{\mathrm{QCC0}}. \end{aligned}$$

]

The two correction terms need not be additive in the final physical theory.

A more general representation is

[

$$T_{\mathrm{combined}}$$

$$\begin{aligned} &\mathcal{H} \\ &\left(\right. \\ &\quad T_{\mathrm{standard}}, \\ &\quad \Delta T_{\mathrm{VFE1}}, \\ &\quad QCC0 \\ &\left. \right). \end{aligned}$$

This formulation allows future experiments to determine whether the interaction is:

- additive;
- multiplicative;
- nonlinear;
- coupled through a field;
- or effectively negligible.

10. Quantum-State Evolution

The quantum subsystem retains the established quantum evolution equation

[

$$i\hbar\frac{\partial}{\partial t}|\psi(t)\rangle$$

$\hat{H}|\psi(t)\rangle$.

]

For an open quantum system,

[

$$\frac{d\rho_q}{dt}$$

$\mathcal{L}(\rho_q)$,

]

where $\mathcal{L}$ represents the appropriate quantum dynamical generator.

The proposed QCC0 layer therefore does not replace the quantum state equation.

Instead,

$$\begin{aligned} &[\rho_q(t) \\ &\rightarrow \text{QCC0}(t) \\ &] \end{aligned}$$

is treated as a derived computational mapping.

This separation permits direct comparison between conventional quantum observables and the proposed QCC0 quantity.

11. 369 Vortex Transformation

The 369 formulation is represented as a proposed geometric transformation.

Define

```
[  
  \mathcal V_{369}:  
  \Psi\rightarrow\Psi'.  
]
```

The transformation must specify:

```
[  
  \Psi'=  
  \mathcal V_{369}  
  (S,\rho,K,P).  
]
```

A complete implementation must identify:

1. the transformation function;
2. its domain;
3. its codomain;
4. its dimensional properties;
5. its invariants;
6. its boundary conditions;
7. its convergence behavior.

The numerical sequence 3-6-9 is not assumed to possess fundamental physical significance merely because it appears in the transformation.

Its scientific significance must follow from the measurable consequences of the transformation.

12. Kapnack Solver

The Kapnack Solver is the numerical engine used to evaluate the proposed framework.

Its architecture is:

```
[
  \text{Input}
  \rightarrow
  \text{Baseline Physics}
  \rightarrow
  \text{SDKP State}
  \rightarrow
  \text{369 Transformation}
  \rightarrow
  \text{VFE1}
  \rightarrow
  \text{QCC0}
  \rightarrow
  \text{Prediction}.
]
```

The solver should retain each intermediate state.

A reproducible run therefore produces:

```
[
  {
    T_{\mathrm{standard}},
    T_{\mathrm{VFE1}},
    T_{\mathrm{combined}},
    QCC0,
    \Phi_v,
    \Psi'.
  }
]
```

This permits investigators to determine which component is responsible for a difference between predictions.

13. Numerical Verification

A computational result should be considered valid only after three levels of verification.

13.1 Analytical verification

Where an analytical solution exists, the solver output must reproduce it within numerical tolerance.

13.2 Independent implementation

At least one independent implementation should reproduce the result using the same equations and input conditions.

13.3 Numerical convergence

The result should remain stable under appropriate changes to:

- timestep;
- spatial resolution;
- solver tolerance;
- floating-point precision;
- integration method.

A result that changes substantially under reasonable numerical refinement cannot be considered physically established.

14. Comparative Prediction Architecture

Every physical test should generate three predictions.

Model A — Established Physics

```
[  
  O_A=O_{\mathrm{standard}}.  
]
```

Model B — Established Physics + VFE1

```
[  
  O_B=O_{\mathrm{standard}}+\Delta O_{\mathrm{VFE1}}.  
]
```

Model C — Established Physics + VFE1 + QCC0

$$[\begin{aligned} O_C = & \\ O_{\{\mathrm{standard}\}} & \\ + & \\ \Delta O_{\{\mathrm{VFE1}\}} & \\ + & \\ \Delta O_{\{\mathrm{QCC0}\}} & \\] \end{aligned}$$

The measured value is

$$[\begin{aligned} O_{\{\mathrm{obs}\}} & \\] \end{aligned}$$

Residuals are

$$[\begin{aligned} R_A = O_{\{\mathrm{obs}\}} - O_A, & \\] \end{aligned}$$

$$[\begin{aligned} R_B = O_{\{\mathrm{obs}\}} - O_B, & \\] \end{aligned}$$

and

$$[\begin{aligned} R_C = O_{\{\mathrm{obs}\}} - O_C. & \\] \end{aligned}$$

The comparison must include measurement uncertainty.

For a measurement with standard uncertainty (σ),

$$[\begin{aligned} z = & \\ \frac{O_{\{\mathrm{obs}\}} - O_{\{\mathrm{model}\}}}{\sigma} & \\] \end{aligned}$$

This allows the difference between competing predictions to be expressed quantitatively.

15. Model Selection

Improved agreement with experimental data is not sufficient if the proposed model contains substantially more adjustable parameters.

Model comparison should therefore account for model complexity.

Where appropriate, information criteria such as

[

AIC

$2k - 2\ln L$

]

or

[

BIC

$k\ln n - 2\ln L$

]

may be used, where (k) is the number of fitted parameters, (n) is the number of observations, and (L) is the likelihood.

A proposed extension should demonstrate predictive improvement rather than merely improved fitting.

16. Falsifiability Conditions

The proposed framework is explicitly falsifiable.

VFE1 is unsupported in a test domain if:

[
 ΔT_{VFE1}
]

is consistently constrained to zero within experimental uncertainty, or if the proposed correction produces predictions inconsistent with observation.

QCC0 is unsupported as a physical quantity if it does not provide reproducible information beyond established quantum observables or if its apparent predictive value disappears under independent testing.

The 369 transformation is unsupported as a new physical law if its predictions are either inconsistent with observation or mathematically reducible to an already established transformation without additional predictive content.

The Kapnack Solver is computationally invalid if independent implementations cannot reproduce its numerical results.

17. Experimental Validation Program

The proposed framework is intended to be tested progressively.

Stage 1 — Dimensional consistency

Every equation must satisfy

[

LHS]

RHS].
]

Stage 2 — Established-physics recovery

The model must recover established results when the proposed correction terms vanish.

For example,

[
 $\Delta T_{\mathrm{VFE1}} \rightarrow 0$
]

must recover the relativistic baseline.

Stage 3 — Numerical benchmark

Kapnack Solver results must agree with independently implemented numerical calculations.

Stage 4 — Existing experimental datasets

The model should be evaluated against high-precision measurements without changing parameters solely to improve agreement.

Stage 5 — Blind prediction

A prediction should be generated before the corresponding measurement is analyzed.

Stage 6 — Independent replication

The strongest validation requires an independent implementation or experimental group.

18. Relativistic Consistency

Because the framework proposes additional temporal structure, compatibility with relativistic limits is essential.

For any proposed velocity-dependent correction,

[
 $v < c$
]

must remain respected for massive objects under special-relativistic conditions unless the model explicitly proposes and experimentally tests a modification of that principle.

The proposed model must also recover the correct low-velocity limit:

[
 $\frac{v}{c} \rightarrow 0$
 $\quad \Longrightarrow \quad$
 $\Delta \tau \rightarrow \Delta t$.
]

Likewise, any gravitational extension must reproduce the established weak-field limit where the additional proposed terms become negligible.

19. Quantum Consistency

The quantum component must preserve the mathematical properties required of a physical density operator:

[
 $\rho_q^\dagger = \rho_q$,
]

```
[  
  \rho_q\ge 0,  
]
```

and

```
[  
  \operatorname{Tr}(\rho_q)=1.  
]
```

Any proposed QCC0 transformation applied to (ρ_q) must not silently violate these properties.

If QCC0 is intended to represent coherence, its predictions should be compared with established coherence measures.

If QCC0 is intended to represent correlation, it should be compared with established correlation or entanglement measures.

If it is intended as a purely computational invariant, that distinction should be retained explicitly.

20. Information and Reproducibility Architecture

A complete computational record should contain:

```
[  
  \mathcal{R}=  
  {  
    \text{equations},  
    \text{parameters},  
    \text{initial conditions},  
    \text{software version},  
    \text{solver settings},  
    \text{outputs},  
    \text{uncertainties}  
  }.  
]
```

The purpose is to make every prediction independently reproducible.

Each model run should therefore have a unique identifier and should preserve the exact parameter set used to generate the result.

21. Relationship to Previous SDKP Work

The present work extends the author's previous SDKP-related computational and mathematical records.

The prior records provide background for the SDKP, SD&N, QCC, VFE, Digital Crystal Protocol, and Kapnack Solver concepts.

The present version narrows that broader architecture toward temporal dynamics and establishes a formal comparison between established physical models and proposed extensions.

The prior work is therefore treated as **authorial precedent and framework development**, not as independent experimental validation.

22. Limitations

Several limitations remain.

First, the precise physical form of VFE1 requires further mathematical specification.

Second, QCC0 requires a formal operational definition that maps it onto measurable quantum-information quantities.

Third, the 369 transformation requires a complete mathematical definition before its physical implications can be evaluated.

Fourth, the Kapnack Solver requires independent benchmarking.

Fifth, agreement with existing data cannot by itself distinguish a new physical law from a flexible parameterization of established physics.

These limitations define the immediate experimental and theoretical development program.

23. Discussion

The framework provides a modular architecture in which established physics and proposed extensions can coexist without conflating their evidentiary status.

The relativistic baseline supplies experimentally established predictions for proper time.

The quantum baseline supplies established quantum-state dynamics and measurable information-theoretic quantities.

VFE1 introduces a proposed field-dependent temporal contribution.

QCC0 introduces a proposed computational mapping from quantum-state information to a normalized correlation/coherence descriptor.

The 369 transformation provides a proposed geometric/kinematic transformation.

The Kapnack Solver evaluates the complete system numerically.

The resulting architecture is therefore not dependent upon the assumption that any proposed component is already correct.

Instead, each component can be removed, replaced, or falsified independently.

This modularity is particularly important for scientific evaluation because it permits an investigator to determine whether a prediction originates from established physics, a proposed extension, or a numerical artifact.

24. Conclusion

Version 0.2 presents a formalized modular framework for investigating temporal dynamics using established relativistic and quantum physics together with proposed SDKP extensions.

The established physical layer consists of special relativity, general relativity, quantum mechanics, quantum-state formalism, and experimentally measured time-dilation phenomena.

The proposed layer consists of VFE1, QCC0, the SDKP state representation, the 369 vortex transformation, and the Kapnack Solver implementation.

The principal methodological contribution is the requirement that these layers be evaluated separately.

For every proposed effect, the framework requires:

```
[  
  \text{Established Prediction}  
  \rightarrow  
  \text{Proposed Prediction}  
  \rightarrow  
  \text{Experimental Measurement}.  
]
```

The resulting residuals, uncertainties, model complexity, and numerical convergence must be reported.

VFE1 and QCC0 are therefore presented as proposed hypotheses rather than established physical laws. Their scientific status depends on whether they generate reproducible, quantitatively superior, independently testable predictions.

Future work will focus on specifying the complete VFE1 field equation, providing an operational QCC0 definition, formally defining the 369 transformation, implementing the complete Kapnack Solver, and testing the resulting predictions against precision relativistic and quantum measurements.

References

1. Einstein, A. (1905). On the Electrodynamics of Moving Bodies. *Annalen der Physik*, 17, 891–921.
2. Einstein, A. (1916). The Foundation of the General Theory of Relativity. *Annalen der Physik*, 49, 769–822.
3. Smith, A. R. H., & Ahmadi, M. (2020). Quantum clocks observe classical and quantum time dilation. *Nature Communications*, 11, 5360.
4. Paige, A. J., Plato, A. D. K., & Kim, M. S. (2020). Classical and Nonclassical Time Dilation for Quantum Clocks. *Physical Review Letters*, 124, 160602.
5. Chiba, T., & Kinoshita, S. (2022). Quantum Clocks, Gravitational Time Dilation, and Quantum Interference.
6. Favalli, T., & Smerzi, A. (2025). Time Dilation of Quantum Clocks in a Relativistic Gravitational Potential. *Entropy*, 27, 489.
7. Roura, A., Schubert, C., Schlippert, D., & Rasel, E. M. (2021). Measuring gravitational time dilation with delocalized quantum superpositions. *Physical Review D*, 104, 084001.
8. Misner, C. W., Thorne, K. S., & Wheeler, J. A. (1973). *Gravitation*. W. H. Freeman.
9. Carroll, S. M. (2004). *Spacetime and Geometry: An Introduction to General Relativity*. Addison-Wesley.

10. Nielsen, M. A., & Chuang, I. L. (2010). *Quantum Computation and Quantum Information*. Cambridge University Press.
 11. Watrous, J. (2018). *The Theory of Quantum Information*. Cambridge University Press.
 12. Smith, D. P. *SDKP Mathematical Framework and Principles for Physics and Computerization*. Zenodo. DOI: 10.5281/zenodo.15779328.
 13. Smith, D. P. *SDKP, ext.* Zenodo.
 14. Smith, D. P. *Digital Crystal Protocol / SDKP Framework and Associated Computational Records*. Zenodo.
-

Author Information

Donald Paul Smith

FatherTimeSDKP

ORCID: 0009-0003-7925-1653

Gypsi Consulting

Research Status

This manuscript is a preprint. VFE1, QCC0, SDKP temporal representations, the 369 vortex formulation, and associated Kapnack Solver methods are presented as proposed theoretical/computational constructs requiring further mathematical development and empirical validation.

Version History

Version 0.1: Initial modular time-dynamics framework.

Version 0.2: Expanded mathematical architecture; established-physics baseline; proposed VFE1 and QCC0 definitions; 369 transformation layer; Kapnack Solver architecture; dimensional, numerical, relativistic, and quantum consistency requirements; comparative prediction methodology; falsifiability criteria; and validation program.

References

1. Einstein, A. (1905). *On the Electrodynamics of Moving Bodies*. *Annalen der Physik*, 17, 891–921.
— Foundational paper for special relativity and the relativistic treatment of time and motion.

2. Einstein, A. (1916). *The Foundation of the General Theory of Relativity*. **Annalen der Physik**, **49**, 769–822.
— Foundational formulation of general relativity and spacetime geometry.
3. Chou, C.-W., Hume, D. B., Rosenband, T., & Wineland, D. J. (2010). *Optical Clocks and Relativity*. **Science**, **329**, 1630–1633.
— Experimental observation of relativistic time-dilation effects using optical atomic clocks. NIST provides the publication record and description.
4. Bothwell, T. et al. (2022). *Resolving the gravitational redshift across a millimetre-scale atomic sample*. **Nature**, **602**, 420–424.
— Experimental measurement of gravitational time dilation at millimetre-scale height differences. NIST reports the experiment and its significance for precision tests of general relativity.
5. Misner, C. W., Thorne, K. S., & Wheeler, J. A. (1973). *Gravitation*. W. H. Freeman.
— Standard reference for general relativity, spacetime geometry, geodesics, and gravitational time.
6. Carroll, S. M. (2004). *Spacetime and Geometry: An Introduction to General Relativity*. Addison-Wesley.
— Modern technical reference for differential geometry and general relativity.
7. Nielsen, M. A., & Chuang, I. L. (2010). *Quantum Computation and Quantum Information: 10th Anniversary Edition*. Cambridge University Press.
— Standard reference for quantum states, density operators, quantum information, quantum computation, entanglement, and quantum error correction.
8. Preskill, J. (2026). *Ph219/CS219: Quantum Computation*. California Institute of Technology.
— Graduate-level reference covering density operators, quantum operations, entanglement, quantum circuits, algorithms, and quantum error correction.
9. Watrous, J. (2018). *The Theory of Quantum Information*. Cambridge University Press.
— Mathematical treatment of quantum information, quantum states, channels, measurements, and information-theoretic quantities.
10. Chou, C.-W., Hume, D. B., Rosenband, T., & Wineland, D. J. (2010). *Relativity and Optical Clocks*. **NIST Publication**.
— Authoritative technical discussion of relativistic clock comparisons and experimentally observed velocity- and gravitational-dependent time shifts.
11. Smith, D. P. (2025–2026). *Scale-Density-Kinetic/Kinematic Principle (SDKP): Framework and Computational Developments*. Public research archive and associated manuscripts.
12. Smith, D. P. (2025–2026). *Digital Crystal Protocol (DCP) and VFE1 Computational Architecture*. Public research archive.
13. Smith, D. P. (2025–2026). *Kapnack Solver and 369 Vortex Computational Framework*. Public research archive.
14. Smith, D. P. (2026). *Vibrational Field Equation One (VFE1) and Quantum Computerization Consciousness Zero (QCC0): Proposed Extensions for Temporal and Quantum-State Modeling*. Preprint/manuscript.