




Numerical Methods

Lauren Hayward
PSI Start Online School



Continued from last time

Numerical Methods, Lecture 3:

Fitting data and 2D Ising Model

PSI Start Online School 2023

June 1, 2023

Lauren Hayward

In this notebook, we will study

1. how to perform fits to data corresponding to random walks in one dimension, and
2. how to calculate properties of a two-dimensional Ising model.

Today's objective is to learn now to perform fits when given numerical data, and to become familiar with the two-dimensional Ising model (which we will revisit in Lecture 4).

Note: Exercises 1 and 2 originally appeared in Lecture 2, but most students did not have enough time to complete them.

1. Fitting data

Outline

- ▶ Statistical physics and the Ising model
- ▶ Monte Carlo methods in statistical physics



Statistical physics

Goal: to study phenomena such as phase transitions for systems of N interacting particles when N is large.

Example: Ising model

$$E = -J \sum_{\langle ij \rangle} s_i s_j$$

$s_i = +1$ or -1
(\blacksquare or \square)




The diagram shows the equation $E = -J \sum_{\langle ij \rangle} s_i s_j$ with four colored arrows pointing to labels below it: a blue arrow from E to "Energy", a purple arrow from J to "coupling energy", a green arrow from $\langle ij \rangle$ to "nearest neighbours", and a red arrow from s_i to "spin state at site i of the lattice".

2D Ising model

$$E = -J \sum_{\langle ij \rangle} s_i s_j$$

$$s_i = +1 \text{ or } -1$$

(■ or □)




2D Ising model

$$E = -J \sum_{\langle ij \rangle} s_i s_j$$


$$s_i = +1 \text{ or } -1$$

(■ or □)


When $J > 0$, which of the following spin configurations minimizes the energy E ?




A



B



C



D


2D Ising model

$$E = -J \sum_{\langle ij \rangle} s_i s_j$$


$$s_i = +1 \text{ or } -1$$

(■ or □)


When $J > 0$, which of the following spin configurations minimizes the energy E ?




A



B



C




D

$$E_{\min} = -J \sum_{\langle ij \rangle} 1 = -J(2N) = -2JL^2$$

2D Ising model

Consider an Ising model on a two-dimensional lattice with periodic boundaries.



$$S_i = +1 \text{ or } -1$$
$$(\blacksquare \text{ or } \square)$$

The configuration of the system can be stored in a two-dimensional array called “spins”.

For the example here with $L=3$:

$$\text{spins} = \begin{bmatrix} -1 & +1 & -1 \\ +1 & +1 & -1 \\ -1 & +1 & +1 \end{bmatrix}$$

Breakout room activities

2. Ising model in two dimensions

Let us consider a classical nearest-neighbour Ising model on a two-dimensional lattice with periodic boundary conditions. The energy of a given spin configuration is given by

$$E = -J \sum_{\langle ij \rangle} s_i s_j,$$

where $J > 0$ is a coupling strength, $s_i = \pm 1$, and $\sum_{\langle ij \rangle}$ denotes a sum over nearest neighbours. We assume that the spins s_i live on the sites of a square $L \times L$ lattice. The total number of spins is then $N = L^2$.

As discussed in the lecture slides, the spins can be stored in a two-dimensional array called `spins`.

Consider the following configuration for $L = 3$:

