

I dont know yet



Faustmann Christian

Faculty of Physics
Technical University of Vienna

This thesis is submitted for the degree of
Bachelor of Science

Abstract

This is where you write your abstract...

Table of contents

List of figures	vii
List of tables	ix
1 Theory	1
1.1 Rubidium	2
1.2 D2 line	3
1.3 Two-level atom	4
1.4 Laser absorbtion	4
1.5 Doppler shifts	4
1.6 Behavior of absorbtion coefficient	4
1.7 Non-linear differential equation	4
2 Experiment	5
2.1 Setup & Tools	5
2.2 Laser diameter measurement	5
2.3 Power / intensity measurement	5
2.4 Doppler-free measurement	5
3 Evaluation	7
3.1 Data processing	7
3.2 Temperature & saturation intensity	7
3.3 Comparison with theory	7
3.4 Compare Doppler-free measurement with theoretical values	7
References	9
Appendix A Theory	11

Appendix B	Experiment	13
Appendix C	Evaluation	15

List of figures

1.1	Rubidium Atom	2
1.2	$5^2S_{1/2} \rightarrow 6^2P_{3/2}$ transition of ^{85}Rb and ^{87}Rb with corresponding hyperfine structure	3
1.3	Doppler spectrum of D2 line	3

List of tables

1.1	Properties of rubidium isotopes	2
-----	---	---

Chapter 1

Theory

1.1 Rubidium




Fig. 1.1 Schematical representation of ⁸⁵Rb

Rubidium is a chemical element with symbol Rb and atomic number 37. It is a soft, silvery-white metallic element of the alkali metal group, with an atomic mass of 85.4678. Elemental rubidium is highly reactive, with properties similar to those of other alkali metals.

German chemists Robert Bunsen and Gustav Kirchhoff discovered rubidium in 1861 by the newly developed technique, flame spectroscopy. Because of the bright red lines in its emission spectrum, they chose a name derived from the Latin word rubidus, meaning “deep red”. [1]

Although rubidium is monoisotopic, rubidium in the Earth’s crust is composed of two isotopes: the stable ⁸⁵Rb and the radioactive ⁸⁷Rb. [2]

Rubidium		
Isotope	85	87
Atomic mass	84.911794	86.909187
in 10 ⁻²⁵ kg	1.40999	1.44316
Abundance	72.17%	27.83%
Spin I	5/2	3/2

Table 1.1 Properties of rubidium isotopes

1.2 D2 line




Fig. 1.2 $5^2S_{1/2} \rightarrow 6^2P_{3/2}$ transition of ^{85}Rb and ^{87}Rb with corresponding hyperfine structure

As we can see both isotopes have the same transition energy, but due to the different spin I (see table: 1.1) we get different energy levels for the groundstate [3]. This is the reason why we witness four doppler peaks in our spectrum (see figure:1.3)




Fig. 1.3 Doppler spectrum of D2 line

1.3 Two-level atom

1.4 Laser absorbtion

1.5 Doppler shifts

1.6 Behavior of absorbtion coefficient

1.7 Non-linear differential equation

Chapter 2

Experiment

2.1 Setup & Tools

2.2 Laser diameter measurement

2.3 Power / intensity measurement

2.4 Doppler-free measurement

Chapter 3

Evaluation

3.1 Data processing

3.2 Temperature & saturation intensity

3.3 Comparison with theory

3.4 Compare Doppler-free measurement with theoretical values

References

- [1] G. Kirchhoff and R. Bunsen. Chemische Analyse durch Spectralbeobachtungen. *Annalen der Physik*, 189:337–381, 1861.
- [2] G. Audi, O. Bersillon, J. Blachot, and A. H. Wapstra. The NUBASE evaluation of nuclear and decay properties. *Nuclear Physics A*, 729:3–128, December 2003.
- [3] J. Reader A. Kramida, Yu. Ralchenko and NIST ASD Team (2015). NIST atomic spectra database (ver. 5.3). *National Institute of Standards and Technology*, 2015.

Appendix A

Theory

Appendix B

Experiment

Appendix C

Evaluation

