Polisci 270 Final Cheat Sheet

Fayzaan Virk

2026-05-01

Contents

Data Wrangling . . . 2
- group_by() and summarize() . . . 2
- filter() and select() . . . 2
- mutate() and rename() . . . 3
- arrange() . . . 4
- Joining and pivot_longer() . . . 4

Average Treatment Effect . . . 6

Ggplot . . . 7
- Scatterplot . . . 7
- Bar plots . . . 8
- Histogram . . . 9
- Box and Whisker Plot . . . 10
- Colors . . . 11

Experimental Types . . . 12
- Research Designs . . . 12

Proportion Tables . . . 13

Z-Score . . . 13

Linear Models . . . 14
- Categorical and binary variables . . . 15
- Summary of a Linear Model . . . 15

Bootstrapping . . . 16
- Dplyr bootstrapping . . . 16
- Infer bootstrapping . . . 17

Confidence Intervals . . . 18
- Dplyr CIs . . . 18
- Infer CIs . . . 19

Hypothesis Testing . . . 20
Dplyr Testing . . . 20
Infer Testing . . . 21
Hypothesis test for a slope . . . 22
Scripts . . . 23
Interpret an ATE . . . 23
Interpret linear coefficients . . . 23
Interpret a CI . . . 23
Interpret a p-value . . . 24

Data Wrangling

Data wrangling is the process of taking a complicated data set and making it easier to work with. This can mean filtering or selecting only a certain number or rows or columns, adding new variables, or pivoting the table. In R, we use the group_by(), summarize(), pivot_wider(), mutate(), filter(), select(), pivot_longer(), and the three join() commands to wrangle our data.

I will use the gapminder data set to demonstrate some of these commands

group_by() and summarize()

First, I can use the group_by() and summarize() commands to find the mean GDP per capita for each continent.

When using mean(), median(), min(), max(), sd(), or sum() in summarize(), any NA values will lead to the whole result being NA. To avoid this, add "na.rm = T" in the summarize() call.

```
gapminder |>
  group_by(continent) |>
  summarize(mean_gdppc = mean(gdpPercap, na.rm = T))
```

```
## # A tibble: 5 x 2
##   continent mean_gdppc
##   <fct>          <dbl>
## 1 Africa         2194.
## 2 Americas       7136.
## 3 Asia           7902.
## 4 Europe        14469.
## 5 Oceania       18622.
```

filter() and select()

I can use the filter() command to only see certain rows of the data set.

```
gapminder |>
  filter(year == 2007)
```

```
## # A tibble: 142 x 6
##    country     continent  year lifeExp       pop gdpPercap
##    <fct>       <fct>     <int>   <dbl>     <int>     <dbl>
##  1 Afghanistan Asia       2007    43.8  31889923      975.
##  2 Albania     Europe     2007    76.4   3600523     5937.
##  3 Algeria     Africa     2007    72.3  33333216     6223.
##  4 Angola      Africa     2007    42.7  12420476     4797.
##  5 Argentina   Americas   2007    75.3  40301927    12779.
##  6 Australia   Oceania    2007    81.2  20434176    34435.
##  7 Austria     Europe     2007    79.8   8199783    36126.
##  8 Bahrain     Asia       2007    75.6    708573    29796.
##  9 Bangladesh  Asia       2007    64.1 150448339     1391.
## 10 Belgium     Europe     2007    79.4  10392226    33693.
## # i 132 more rows
```

The select() command works the same way as filter() but for choosing certain columns

```r
gapminder |>
  filter(year == 2007) |>
  select(country, gdpPercap)
```

```
## # A tibble: 142 x 2
##    country     gdpPercap
##    <fct>           <dbl>
##  1 Afghanistan      975.
##  2 Albania         5937.
##  3 Algeria         6223.
##  4 Angola          4797.
##  5 Argentina      12779.
##  6 Australia      34435.
##  7 Austria        36126.
##  8 Bahrain        29796.
##  9 Bangladesh      1391.
## 10 Belgium        33693.
## # i 132 more rows
```

mutate() and rename()

The mutate() command can create new columns using logicals or calculations and rename() can change existing columns' names.

```r
gapminder |>
  mutate(
    gdp = gdpPercap * pop,
    post_coldwar = if_else(year > 1991, 1, 0)
  ) |>
  rename("population" = pop)
```

```
## # A tibble: 1,704 x 8
##    country     continent  year lifeExp population gdpPercap     gdp post_coldwar
##    <fct>       <fct>     <int>   <dbl>      <int>     <dbl>   <dbl>        <dbl>
##  1 Afghanistan Asia       1952    28.8    8425333      779. 6.57e 9            0
```

```
##  2 Afghanistan Asia       1957    30.3    9240934      821. 7.59e 9            0
##  3 Afghanistan Asia       1962    32.0   10267083      853. 8.76e 9            0
##  4 Afghanistan Asia       1967    34.0   11537966      836. 9.65e 9            0
##  5 Afghanistan Asia       1972    36.1   13079460      740. 9.68e 9            0
##  6 Afghanistan Asia       1977    38.4   14880372      786. 1.17e10            0
##  7 Afghanistan Asia       1982    39.9   12881816      978. 1.26e10            0
##  8 Afghanistan Asia       1987    40.8   13867957      852. 1.18e10            0
##  9 Afghanistan Asia       1992    41.7   16317921      649. 1.06e10            1
## 10 Afghanistan Asia       1997    41.8   22227415      635. 1.41e10            1
## # i 1,694 more rows
```

arrange()

The arrange() function will reorder the entries in a column

```
gapminder |>
  filter(year == 2007) |>
  arrange(desc(gdpPercap))
```

```
## # A tibble: 142 x 6
##    country          continent  year lifeExp       pop gdpPercap
##    <fct>            <fct>     <int>   <dbl>     <int>     <dbl>
##  1 Norway           Europe     2007    80.2   4627926    49357.
##  2 Kuwait           Asia       2007    77.6   2505559    47307.
##  3 Singapore        Asia       2007    80.0   4553009    47143.
##  4 United States    Americas   2007    78.2 301139947    42952.
##  5 Ireland          Europe     2007    78.9   4109086    40676.
##  6 Hong Kong, China Asia       2007    82.2   6980412    39725.
##  7 Switzerland      Europe     2007    81.7   7554661    37506.
##  8 Netherlands      Europe     2007    79.8  16570613    36798.
##  9 Canada           Americas   2007    80.7  33390141    36319.
## 10 Iceland          Europe     2007    81.8    301931    36181.
## # i 132 more rows
```

Joining and pivot_longer()

pivot_longer() is the opposite of pivot_wider(). It takes data from columns and transforms them into rows. A good example of this is the mortality data set. It has a column for each year 1972 to 2020 for infant mortatility in that year. We could switch this to one row for each year per country.

```
mortality
```

```
## # A tibble: 217 x 52
##    country      country_code indicator `1972` `1973` `1974` `1975` `1976` `1977`
##    <chr>        <chr>        <chr>      <dbl>  <dbl>  <dbl>  <dbl>  <dbl>  <dbl>
##  1 Aruba        ABW          Mortalit~   NA     NA     NA     NA     NA     NA
##  2 Afghanistan  AFG          Mortalit~  291    285.   280.   274.   268    262.
##  3 Angola       AGO          Mortalit~   NA     NA     NA     NA     NA     NA
##  4 Albania      ALB          Mortalit~   NA     NA     NA     NA     NA     NA
##  5 Andorra      AND          Mortalit~   NA     NA     NA     NA     NA     NA
##  6 United Arab~ ARE          Mortalit~   80.1   72.6   65.7   59.4   53.6   48.3
##  7 Argentina    ARG          Mortalit~   69.7   68.2   66.1   63.3   59.8   55.7
```

```
##  8 Armenia      ARM          Mortalit~   NA     NA     NA     NA     87.1   83.6
##  9 American Sa~ ASM          Mortalit~   NA     NA     NA     NA     NA     NA
## 10 Antigua and~ ATG          Mortalit~   26.9   25.1   23.5   22.1   20.8   19.5
## # i 207 more rows
## # i 43 more variables: ‘1978‘ <dbl>, ‘1979‘ <dbl>, ‘1980‘ <dbl>, ‘1981‘ <dbl>,
## #   ‘1982‘ <dbl>, ‘1983‘ <dbl>, ‘1984‘ <dbl>, ‘1985‘ <dbl>, ‘1986‘ <dbl>,
## #   ‘1987‘ <dbl>, ‘1988‘ <dbl>, ‘1989‘ <dbl>, ‘1990‘ <dbl>, ‘1991‘ <dbl>,
## #   ‘1992‘ <dbl>, ‘1993‘ <dbl>, ‘1994‘ <dbl>, ‘1995‘ <dbl>, ‘1996‘ <dbl>,
## #   ‘1997‘ <dbl>, ‘1998‘ <dbl>, ‘1999‘ <dbl>, ‘2000‘ <dbl>, ‘2001‘ <dbl>,
## #   ‘2002‘ <dbl>, ‘2003‘ <dbl>, ‘2004‘ <dbl>, ‘2005‘ <dbl>, ‘2006‘ <dbl>, ...
```

```r
mortality_long <- mortality |>
  select(!indicator) |>
  pivot_longer(
    cols = `1972`:`2020`,
    names_to = "year",
    values_to = "child_mortality"
  ) |>
  drop_na(child_mortality)
mortality_long
```

```
## # A tibble: 9,157 x 4
##    country     country_code year  child_mortality
##    <chr>       <chr>        <chr>           <dbl>
##  1 Afghanistan AFG          1972             291
##  2 Afghanistan AFG          1973             285.
##  3 Afghanistan AFG          1974             280.
##  4 Afghanistan AFG          1975             274.
##  5 Afghanistan AFG          1976             268
##  6 Afghanistan AFG          1977             262.
##  7 Afghanistan AFG          1978             256
##  8 Afghanistan AFG          1979             250.
##  9 Afghanistan AFG          1980             244.
## 10 Afghanistan AFG          1981             237.
## # i 9,147 more rows
```

Joining comibnes two different data sets based on common variables, aka keys. There are three types of join functions: left_join() keeps all rows in the left data set and add columns from the right, right_join() does the same but for the right data set, and inner_join() only keeps rows which are present in both data sets. I'll demonstrate with gapminder and mortality.

```r
mortality_long <- mortality_long |> #First, we have to make sure the year column in
  # both data sets are integer so they can be joined
  mutate(
    year = as.integer(year)
  )
gapminder |>
  inner_join(mortality_long, by = "year") |>
  select(country.x, continent, year, lifeExp, pop, gdpPercap, child_mortality)
```

```
## Warning in inner_join(gapminder, mortality_long, by = "year"): Detected an unexpected many-to-many re
## i Row 5 of ‘x‘ matches multiple rows in ‘y‘.
```

```
## i Row 1 of `y` matches multiple rows in `x`.
## i If a many-to-many relationship is expected, set `relationship =
##   "many-to-many"` to silence this warning.
```

```
## # A tibble: 208,598 x 7
##    country.x   continent  year lifeExp      pop gdpPercap child_mortality
##    <fct>       <fct>     <int>   <dbl>    <int>     <dbl>           <dbl>
##  1 Afghanistan Asia       1972    36.1 13079460      740.           291
##  2 Afghanistan Asia       1972    36.1 13079460      740.            80.1
##  3 Afghanistan Asia       1972    36.1 13079460      740.            69.7
##  4 Afghanistan Asia       1972    36.1 13079460      740.            26.9
##  5 Afghanistan Asia       1972    36.1 13079460      740.            20.3
##  6 Afghanistan Asia       1972    36.1 13079460      740.            28.1
##  7 Afghanistan Asia       1972    36.1 13079460      740.           317.
##  8 Afghanistan Asia       1972    36.1 13079460      740.            22
##  9 Afghanistan Asia       1972    36.1 13079460      740.           251.
## 10 Afghanistan Asia       1972    36.1 13079460      740.           309.
## # i 208,588 more rows
```

See Code from Lecture 4, 5, 9, 12, Problem Set 2, 3, 4, 5, 6, 8

Average Treatment Effect

The average treatment effect is a way of calculation how much a treatment affects the outcome variable. The formula is the expected outcome variable if all subjects were treated minus the expected outcome if all subjects were not treated. To approximate this, we use the mean of the treatment group for the outcome variable minus the mean of the control group.

I'll use the trains experiments (Spanish-speakers in train stations in Boston).

```r
trains |>
  group_by(treatment) |>
  summarize(post_mean = mean(numberim.post)) |>
  pivot_wider(
    names_from = treatment,
    values_from = post_mean
  ) |>
  mutate(
    ate = `1` - `0`
  )
```

```
## # A tibble: 1 x 3
##     `0`   `1`   ate
##   <dbl> <dbl> <dbl>
## 1  2.73  3.12 0.383
```

Additionally, you can use select() and pull() to get just the numeric value

```r
trains |>
  group_by(treatment) |>
  summarize(post_mean = mean(numberim.post)) |>
  pivot_wider(
```

```r
    names_from = treatment,
    values_from = post_mean
  ) |>
  mutate(
    ate = `1` - `0`
  ) |>
  select(ate) |>
  pull()
```

```
## [1] 0.3832721
```

See Code from Lecture 7, Problem Set 3, 7

Ggplot

ggplot2 is a package that is part of tidyverse that is used to create many types of graphs and visual representations of data. In class, we learned scatterplots, box and whisker plots, barplots, and histograms. These are created using geom_point(), geom_box(), geom_bar() or geom_col(), and geom_histogram(). For a scatterplot, a line of best fit can be added using geom_smooth().

Scatterplot

```r
midwest |>
  ggplot(aes(x = percblack, y = percbelowpoverty)) +
  geom_point(alpha = 0.3) +
  geom_smooth(method = "lm", se = F) +
  labs(
    x = "Percentage of black population",
    y = "Percentage below the poverty line",
    title = "Relationship between black and impoverished population in Midwestern counties"
  )
```

```
## `geom_smooth()` using formula = 'y ~ x'
```



Bar plots

```
midwest |>
  group_by(state) |>
  summarize(population = sum(poptotal)) |>
  ggplot(aes(x = state, y = population)) +
  geom_col() +
  scale_y_continuous(labels = scales::comma) +
  labs(
    x = "State",
    y = "Total Population",
    title = "Population in each Midwestern State"
  )
```



Histogram

```
midwest |>
  ggplot(aes(x = percbelowpoverty)) +
  geom_histogram(binwidth = 2) +
  labs(
    x = "Percentage Below the Poverty Line",
    y = "Count",
    title = "Percentage Below the Poverty Line in Midwestern Counties"
  )
```



Box and Whisker Plot

```
midwest |>
  ggplot(aes(x = percbelowpoverty)) +
  geom_boxplot() +
  facet_wrap(~ state) +
  scale_y_discrete() +
  labs(
    x = "Percentage Below the Poverty Line",
    title = "Percentage Below the Poverty Line in Midwestern Counties by State"
  )
```



Colors

```
midwest |>
  ggplot(
    aes(
      x = percbelowpoverty,
      fill = state,
      color = state
    )) +
  geom_density(alpha = 0.3) +
  labs(
    x = "Percentage Below the Poverty Line",
    y = "Density",
    title = "Percentage Below the Poverty Line in Midwestern Counties by State"
  )
```



See Code from Lecture 3, Problem Set 1, 2, 3, 4, 5, 6, 7

Experimental Types

An observational study is where we simply observe how two variables interact in real life. We do not assign a treatment. This is very common in real life, since we often can't assign treatments due to ethical or logistical reasons. Because we cannot randomly assign treatments, it is much harder to establish causality from an observational study because of a greater chance of confounding variables. On the other hand, Randomized Controlled Trials (RCTs) do have randomly assigned treatment groups. RCTs tend to have a higher internal validity, meaning it is easier to conclude a causal relationship from them. Observational studies tend to have a higher external validity, which means the conclusion can be more easily generalized beyond this study.

Research Designs

There are three types of research designs. A cross-sectional design compares the control and treatment groups after the treatment has been administered. This assumes the treatment and control groups are on average identical. A before-and-after design compares the treatment group from before and after the treatment was administered. This assumes none of the confounders will vary with time. A difference-in-differences design used the before-and-after of the control group to infer what would've happened to the treatment group if no treatment was administered. This assumes that the trends that affect one group equally affect the other group. All three rely on assumptions that can't always be verified, and are thus more susceptible to confounding variables than a RCT. There are some ways to try and control for confounding variables using statistical procedures, one of which is subclassification. Subclassification is where treated and control groups are compared within the levels of a confounder.

See Code from Lecture 8, Problem Set 3

Proportion Tables

```
cces_2020 |>
  group_by(pres_vote) |>
  summarize(n = n()) |>
  mutate(proportion = n / sum(n))
```

```
## # A tibble: 7 x 3
##   pres_vote                        n proportion
##   <fct>                        <int>      <dbl>
## 1 Joe Biden (Democrat)         26188   0.508
## 2 Donald J. Trump (Republican) 17702   0.343
## 3 Other                         1458   0.0283
## 4 I did not vote in this race    100   0.00194
## 5 I did not vote                  13   0.000252
## 6 Not sure                       190   0.00369
## 7 <NA>                          5900   0.114
```

For multiple grouping variables, see below

```
cces_2020 |>
  filter(pres_vote %in% c("Joe Biden (Democrat)", "Donald J. Trump (Republican)")) |>
  mutate(pres_vote = if_else(pres_vote == "Joe Biden (Democrat)", "Biden", "Trump")) |>
  group_by(pid3, pres_vote) |>
  summarize(n = n()) |>
  mutate(prop = n/sum(n)) |>
  select(!n) |>
  pivot_wider(
    names_from = pid3,
    values_from = prop
  ) |>
  knitr::kable()
```

```
## `summarise()` has grouped output by 'pid3'. You can override using the
## `.groups` argument.
```

pres_vote	Democrat	Republican	Independent	Other	Not sure
Biden	0.9681295	0.0712146	0.5714162	0.4870775	0.5993092
Trump	0.0318705	0.9287854	0.4285838	0.5129225	0.4006908

See Code from Lecture 9, 10

Z-Score

The z-score represent how many standard deviations away from the mean a data point is. The formula is (x - mu)/sigma, where mu is variable's mean and sigma is the variable's standard deviation. Below I will find the z-score for Cook County, IL for the proportion of people 5 and older who got at least one dose of the COVID vaccine.

```r
dose5_stats <- covid_votes |>
  summarize(
    mean_dose5 = mean(one_dose_5plus_pct, na.rm=T),
    sd_dose5 = sd(one_dose_5plus_pct, na.rm=T)
  )

cook_dose5 <- covid_votes |>
  filter(county == "Cook County" & state == "IL") |>
  select(one_dose_5plus_pct) |>
  pull()

cook_z = (cook_dose5 - dose5_stats$mean_dose5) / dose5_stats$sd_dose5
cook_z
```

```
## [1] 1.695176
```

See Code from Lecture 11

Linear Models

A linear model is a great way to see if one variable has a positive, negative, or no effect on another variable. A linear model can be created using the lm() command. The parameters are lm(dependent variable(s) ~ independent, data set).

```r
lobf <- lm(percbelowpoverty ~ percblack, midwest)
coef(lobf)
```

```
## (Intercept)   percblack
##  11.9255333   0.2185736
```

For a county with a black percentage of zero, the percentage below the poverty line is predicted to be 11.93% For every percentage point increase in the black population, the percentage below the poverty line is predicted to increase by 0.22 percentage points.

```r
lobf_2 <- lm(percbelowpoverty ~ percblack + percasian + popdensity + poptotal, midwest)
coef(lobf_2)
```

```
##   (Intercept)     percblack     percasian    popdensity      poptotal
##  1.220449e+01  4.231014e-01 -3.700647e-01 -2.782850e-04  2.247291e-06
```

For a county with a black percentage of 0, and Asian percentage of 0, a population density of 0, and a total population of zero, the percentage below the poverty line is expected to be 12.2%. For every percentage point increase in the black population, the percentage below the poverty line is predicted to increase by 0.42 percentage points. For every percentage point increase in the Asian population, the percentage below the poverty line is predicted to decrease by 0.37 percentage points. For every increase by one in population density, the percentage below the poverty line is predicted to decrease by 0.0002 percentage points. For every increase by 1000 in the total population, the percentage below the poverty line is predicted to increase by 0.002 percentage points. Making a linear model with multiple variables can help us isolate confounding effect by comparing the different effects multiple response variables have on the outcome. In the example above, we see one potential confounder could be total population, and it becomes more difficult to draw a causal relationship between black population and percentage beneath the poverty line.

Categorical and binary variables

A linear model can be used to calculate the ATE if and only if the independent variable is a binary variable. In this case, the ATE would be the slope of the regression line. A linear model can also be used for categorical variables.

```
lobf_3 <- lm(nondiscrim_post ~ racename, trans)
coef(lobf_3)
```

```
##             (Intercept)           racenameAsian       racenameCaucasian
##              0.63492063              0.19841270              0.06244779
##        racenameHispanic racenameNative American
##              0.02936508             -0.63492063
```

This linear model means that the proportion of African Americans that didn't discriminate post-treatment was 63.5%. Every coefficient is how other races compare to that score. For example, the proportion of Asians that didn't discriminate post-treatment is 19.8 percentage points higher and the percentage of Native Americans that didn't discriminate post-treatment was 6.3 percentage points lower.

Summary of a Linear Model

```
summary(lobf)
```

```
##
## Call:
## lm(formula = percbelowpoverty ~ percblack, data = midwest)
##
## Residuals:
##    Min     1Q Median     3Q    Max
## -9.893 -3.367 -0.653  2.350 36.766
##
## Coefficients:
##             Estimate Std. Error t value Pr(>|t|)
## (Intercept) 11.92553    0.27151  43.923  < 2e-16 ***
## percblack    0.21857    0.04692   4.658 4.25e-06 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 5.032 on 435 degrees of freedom
## Multiple R-squared:  0.04751,    Adjusted R-squared:  0.04532
## F-statistic:  21.7 on 1 and 435 DF,  p-value: 4.25e-06
```

Calling summary() on a linear model gives us more information than just calling coef(). It gives us p-values for the intercept and the slope, which are in the Pr(>|t|) column. In this case, we can see both the intercept and the slope have nearly zero p-values. The Multiple R-squared is also useful. It shows how well our linear model fits the data set. For this R-squared of 0.047, 4.7% of the variation in percentage below the poverty line is explained by percentage of black population. This means our linear model is not a good fit for the data. A good R-Squared value would be greater than 0.8.

See Code from Lecture 14, 15, 16, 25, Problem Set 5

Bootstrapping

Bootstrapping is the process of creating a sampling distribution from a single sample. It is incredibly useful because in many cases, we don't have access to the full population data or multiple samples. It creates multiple samples of the same size as the one we have by repeatedly drawing with replacement from our sample. This creates a sampling distribution of the same shape (spread) as the sampling distribution would be, but centered at the sample statistic. We can then use the bootstrap for calculating confidence intervals or hypothesis tests because its shape is the same.

I will show both the dplyr and infer workflows for making a bootstrap to calculate the proportion of all Americans who want the government to do more about the environment using one sample from the American national election survey.

Dplyr bootstrapping

```
set.seed(53132)
dplyr_boots <- anes |>
  rep_slice_sample(reps = 1000, prop = 1, replace = T) |>
  group_by(replicate) |>
  summarize(mean_envir = mean(envir_doing_more))

dplyr_boots
```

```
## # A tibble: 1,000 x 2
##    replicate mean_envir
##        <int>      <dbl>
##  1         1      0.602
##  2         2      0.595
##  3         3      0.601
##  4         4      0.609
##  5         5      0.595
##  6         6      0.607
##  7         7      0.595
##  8         8      0.590
##  9         9      0.595
## 10        10      0.610
## # i 990 more rows
```

When being visualized, this bootstrap distribution should have an approximately normal distribution.

```
dplyr_boots |>
  ggplot(aes(x = mean_envir)) +
  geom_histogram(binwidth = 0.001) +
  labs(
    x = "Proportion who believes the government should be doing more",
    y = "Count"
  )
```



Infer bootstrapping

This can also be done using the infer package

```
set.seed(60004)
infer_boots <- anes |>
  specify(response = envir_doing_more) |>
  generate(reps = 1000, type = "bootstrap") |>
  calculate(stat = "mean")
infer_boots
```

```
## Response: envir_doing_more (numeric)
## # A tibble: 1,000 x 2
##    replicate  stat
##        <int> <dbl>
##  1         1 0.595
##  2         2 0.601
##  3         3 0.601
##  4         4 0.612
##  5         5 0.609
##  6         6 0.613
##  7         7 0.598
##  8         8 0.596
##  9         9 0.597
## 10        10 0.600
```

```
## # i 990 more rows
```

Infer can also be used to visualize the bootstrap distribution

```
visualize(infer_boots)
```



See Code from Lecture 17, 19, Problem Set 6, 7, 8

Confidence Intervals

A confidence interval gives a range of plausible values for a population parameter, given a sample statistic at a confidence level. It can be used to find a population mean or proportion. It is often used for ATEs to see if zero is in the interval or not.

Dplyr CIs

```
set.seed(414)
trans_ate_boots <- trans |>
  rep_slice_sample(prop = 1, replace = T, reps = 1000) |>
  group_by(replicate, treat_ind) |>
  summarize(post_mean = mean(nondiscrim_post)) |>
  pivot_wider(
    names_from = treat_ind,
```

```r
    values_from = post_mean
  ) |>
  mutate(
    ate = `1` - `0`
  ) |>
  select(replicate, ate)
```

```
## `summarise()` has grouped output by 'replicate'. You can override using the
## `.groups` argument.
```

```r
trans_ate_boots |>
  get_confidence_interval(type = "percentile", level = 0.95)
```

```
## # A tibble: 1 x 2
##   lower_ci upper_ci
##      <dbl>    <dbl>
## 1  -0.0399    0.111
```

For an ATE, a 95% confidence interval means that if the experiement were repeated many, many times, 95% of calculated intervals would contain the true population ATE. Since zero is contained in the interval, we do not have statistically significant evidence that having a conversation about trans people has an affect on peoples' opinions of trans people.

Infer CIs

The same can also be done with the infer package

```r
set.seed(414)
trans_boots <- trans |>
  mutate(
    treatment = if_else(treat_ind == 1, "Treated", "Control")
  ) |>
  specify(nondiscrim_post ~ treatment) |> # (response ~ explanatory)
  generate(reps = 1000, type = "bootstrap") |>
  calculate(stat = "diff in means", order = c("Treated", "Control"))

trans_ci <- trans_boots |>
  get_confidence_interval(level = 0.95, type = "percentile")
trans_ci
```

```
## # A tibble: 1 x 2
##   lower_ci upper_ci
##      <dbl>    <dbl>
## 1  -0.0399    0.111
```

See Code from Lecture 20, Problem Set 7

Hypothesis Testing

A hypothesis test provides statistically significant evidence to reject or fail to reject a claim about a population parameter. The null hypothesis is typically that no special relationship has been found. The alternative hypothesis is typically that the sample statistic is meaningful. Hypothesis tests do this by calculating a p-value, which states how rare an outcome as or more extreme than the sample statistic would be assuming the null hypothesis is true.

I will test if life expectancy in Africa is the same as in the Americas in 2007 using the gapminder data. The null hypothesis is that they will be the same.

Dplyr Testing

```
set.seed(414)
life_diff <- gapminder |>
  filter(year == 2007, continent %in% c("Africa", "Americas")) |>
  group_by(continent) |>
  summarize(mean_life = mean(lifeExp)) |>
  pivot_wider(
    names_from = continent,
    values_from = mean_life
  ) |>
  mutate(
    diff = Americas - Africa
  ) |>
  select(diff) |>
  pull()
life_diff
```

```
## [1] 18.80208
```

```
dplyr_null <- gapminder |>
  filter(year == 2007, continent %in% c("Africa", "Americas")) |>
  rep_slice_sample(prop = 1, replace = FALSE, reps = 1000) |>
  group_by(replicate) |>
  mutate(
    continent_shuffled = sample(continent)
  ) |>
  group_by(replicate, continent_shuffled) |>
  summarize(mean_life = mean(lifeExp)) |>
  pivot_wider(
    names_from = continent_shuffled,
    values_from = mean_life
  ) |>
  mutate(
    diff = Americas - Africa
  )
```

```
## `summarise()` has grouped output by 'replicate'. You can override using the
## `.groups` argument.
```

```r
dplyr_null |>
  ungroup() |>
  summarize(p_value = mean(abs(diff) >= abs(life_diff)))
```

```
## # A tibble: 1 x 1
##   p_value
##     <dbl>
## 1       0
```

Infer Testing

Here's how to do it using the infer package

```r
set.seed(414)
infer_diff <- gapminder |>
  filter(year == 2007, continent %in% c("Africa", "Americas")) |>
  specify(lifeExp ~ continent) |>
  calculate(stat = "diff in means", order = c("Americas", "Africa"))
```

```
## Dropping unused factor levels c("Asia", "Europe", "Oceania") from the supplied
## explanatory variable 'continent'.
```

```r
infer_null <- gapminder |>
  filter(year == 2007, continent %in% c("Africa", "Americas")) |>
  specify(lifeExp ~ continent) |>
  hypothesize(null = "independence") |>
  generate(reps = 1000, type = "permute") |>
  calculate(stat = "diff in means", order = c("Americas", "Africa"))
```

```
## Dropping unused factor levels c("Asia", "Europe", "Oceania") from the supplied
## explanatory variable 'continent'.
```

```r
infer_null |>
  get_p_value(obs_stat = infer_diff, direction = "both")
```

```
## Warning: Please be cautious in reporting a p-value of 0. This result is an approximation
## based on the number of `reps` chosen in the `generate()` step.
## i See `get_p_value()` (`?infer::get_p_value()`) for more information.
```

```
## # A tibble: 1 x 1
##   p_value
##     <dbl>
## 1       0
```

```r
infer_null |>
  visualize() +
  shade_p_value(obs_stat = infer_diff, direction = "both")
```



In both cases, the p-value is approximately zero. This means that assuming the null hypothesis is true, there is a near zero percent chance of obtaining a sample difference in means as or more extreme as 18.8. Therefore, we can reject the null hypothesis at the alpha = 0.05 level.

Hypothesis test for a slope

Now we will see if GDP per capita has an affect on life expectancy in 2007

```
set.seed(414)
gap_2007 <- gapminder |>
  filter(year == 2007)

lobf_4 <- lm(lifeExp ~ gdpPercap, gap_2007)
coef(lobf_4)
```

```
##  (Intercept)    gdpPercap
## 5.956565e+01 6.371341e-04
```

```
# Or could use infer to find observed slope

infer_slope <- gap_2007 |>
  specify(lifeExp ~ gdpPercap) |>
  calculate(stat = "slope")

slope_null <- gap_2007 |>
```

```r
  specify(lifeExp ~ gdpPercap) |>
  hypothesize(null = "independence") |>
  generate(reps = 1000, type = "permute") |>
  calculate(stat = "slope")

slope_null |>
  get_p_value(obs_stat = infer_slope, direction = "both")
```

```
## Warning: Please be cautious in reporting a p-value of 0. This result is an approximation
## based on the number of ‘reps‘ chosen in the ‘generate()‘ step.
## i See ‘get_p_value()‘ (‘?infer::get_p_value()‘) for more information.
```

```
## # A tibble: 1 x 1
##   p_value
##     <dbl>
## 1       0
```

Since the p-value is approximately zero, we reject the null hypothesis at the alpha = 0.05 level. We have statistically significant evidence that GDP per capita has a non-zero effect on life expectancy.

See Code from Lecture 21, 22, 25, Problem Set 8

Scripts

Interpret an ATE

The average treatment effect is [ATE]. This means that [treatment group] had a [higher/lower] [outcome variable] by [absolute value of ATE with units], on average.

Example: The average treatment effect is 0.097. This means that registered Democrats had a higher COVID vaccination rate by 9.7 percentage points, on average.

Interpret linear coefficients

When [all independent variables] are equal to zero, the value of [outcome variable] is predicted to be [intercept value with units].

Example: When a country's GDP per capita is equal to zero, it's life expectancy is predicted to be 11.3 years.

For every [one unit] increase of [independent variable], the [outcome variable] is predicted to [increase/decrease] by [absolute value of slope with units], holding all other variables constant.

Example: For every percentage point increase of COVID vaccination, the Democrat vote share of a county is predicted to increase by 0.89 percentage points, holding all other variables constant.

Interpret a CI

We are [confidence level]% confident that the true population [parameter] of [variable/relationship] is between [lower bound] and [upper bound with units].

FOR AN ATE CONFIDENCE INTERVAL, ADD: Since zero [is/is not] contained in the interval, we [do not have/have] statistically significant evidence that [treatment] has an effect on [outcome variable] at the alpha = [1 - confidence level] level.

Example: We are 95% confident that the true population ATE of having a conversation about trans people on non-discrimination attitudes is between -0.04 and 0.11. Since zero is contained in the interval, we do not have statistically significant evidence that having a conversation about trans people has an effect on nondiscrimination attitudes at the alpha = 0.05 level.

OPTIONAL: If we were to repeat this sampling process many, many times and construct a confidence interval each time, [confidence level]% of those intervals would contain the true population [parameter].

Interpret a p-value

The p-value is [p-value]. This means that, assuming the null hypothesis of [state null hypothesis] is true, there is a [p-value * 100] chance of observing a sample [statistic] as or more extreme than [observed statistic value]. Since the p-value [is less than/is greater than] alpha = [significance level], we [reject/fail to reject] the null hypothesis. We [have/do not have] statistically significant evidence that [alternative hypothesis].

Example: The p-value is 0.038. This means that, assuming the null hypothesis of no relationship between percentage of black population and percentage below the poverty line is true, there is a 3.8% chance of observing a sample slope as or more extreme than 0.006. Since the p-value is less than alpha = 0.05, we reject the null hypothesis. We have statistically significant evidence that there is a relationship between relationship between percentage of black population and percentage below the poverty line.