

# Some Notes about the things I encountered

Simon Pohmann

July 11, 2022

## Notation

If  $E$  is an Elliptic Curve defined over a finite field of characteristic  $p$ , we write  $E^{(p)}$  for the curve defined by the equations of  $E$  after replacing all coefficients by their  $p$ -th power. Similarly, for an isogeny  $\phi : E \rightarrow E'$ , write  $\phi^{(p)} : E^{(p)} \rightarrow E'^{(p)}$  for the isogeny defined by the polynomials of  $\phi$  after replacing all coefficients by their  $p$ -th power. Furthermore, for a point  $P = (x : y : z) \in \mathbb{P}^2$  write  $P^{(p)} := (x^p : y^p : z^p)$ . Finally, for a set of points or endomorphisms  $S$  write  $S^{(p)} := \{s^{(p)} \mid s \in S\}$ . Note that

$$\begin{aligned} \cdot^{(p)} : \mathbf{Ell} &\rightarrow \mathbf{Ell}, & E &\mapsto E^{(p)} \\ \mathrm{Hom}_{\mathbf{Ell}}(E, E') &\ni \phi &\rightarrow \phi^{(p)} \end{aligned}$$

is a covariant endofunctor on the category  $\mathbf{Ell}$  of Elliptic Curves defined over  $\bar{\mathbb{F}}_p$  and their isogenies.

Sometimes, we abuse terminology and speak of Elliptic Curves when we mean isomorphism classes of Elliptic Curves.

Many examples will be over  $\mathbb{F}_{101^2}$ . Let  $p = 101$  and  $q = p^2$ . We usually use the generator  $\alpha \in \mathbb{F}_q$  with minimal polynomial  $x^2 + 97x + 2$ .

## 1 Example - The cases I, II and III

For the analysis of these cases, we consider an ordinary Elliptic Curve  $E/\mathbb{F}_{p^2}$ , a prime  $l$  and the curve  $E_0$ , defined as the curve on the crater of the  $l$ -isogeny volcano that is the root of the subtree containing  $E$ . In other words,  $E_0$  is the curve on the crater that is closest to  $E$ , and there is a path of ascending  $l$ -isogenies  $E \rightarrow \dots \rightarrow E_0$ . Finally, if  $E$  is already on the crater, then clearly  $E = E_0$ .

### 1.1 Case I

We say that we are in *Case I*, if  $E_0 = E_0^{(p)}$ , i.e.  $E_0$  is defined over  $\mathbb{F}_p$ .

Finding examples of case I is trivial - just take a curve  $E$  with  $j(E) \in \mathbb{F}_p$ . Then clearly  $E^{(p)} = E$  and so also  $E_0^{(p)} = E_0$  (since  $\cdot^{(p)}$  maps the path  $E \rightarrow E_0$  to  $E = E^{(p)} \rightarrow E_0^{(p)}$ ).



are more or less two possibilities for the  $\cdot^{(p)}$  conjugate path<sup>1</sup>.

- It could be that the conjugate of  $E_i \rightarrow E_{i+1}$  is the dual of  $E_{n-i-1} \rightarrow E_{n-i}$ , hence we just go the path  $E \rightarrow \dots \rightarrow E^{(p)}$  backwards.
- It could be that the conjugate of  $E_i \rightarrow E_{i+1}$  is  $E_{n+i} \rightarrow E_{n+i+1}$ , where

$$E_0, \dots, E_n, E_{n+1}, \dots, E_{n+m} = E_0$$

is the cycle along the whole crater.

However, as we will see, the first case is impossible.

Note that we have

**Proposition 1.1.** *Let  $[\mathfrak{b}] \in \text{Cl}(\mathcal{O})$  where  $\mathcal{O} = \text{End}(E)$  for an ordinary Elliptic Curve  $E/\mathbb{F}_{p^2}$  such that  $[\mathfrak{b}].E = E^{(p)}$ . Then  $[\mathfrak{b}]^2 = [(1)]$ .*

*Proof.* I think there is some mistake with my definition of the class group action, see also the next paragraph. With the current (probably wrong) definition, the following works. Otherwise, I suppose that anyway we have  $[\mathfrak{b}] = [(p, \pi)]$  and then the claim follows by Lemma 2.5.

We recall the definition of the class group action in the case  $[\mathfrak{b}].E^{(p)}$ . For an ideal  $\mathfrak{b}' \leq \text{End}(E^{(p)})$ , have by definition

$$[\mathfrak{b}'].E^{(p)} = E^{(p)}/E^{(p)}[\mathfrak{b}'] = E^{(p)} / \bigcap_{\beta \in \mathfrak{b}'} \ker(\beta)$$

However,  $\mathfrak{b}$  is an ideal in  $\text{End}(E)$ , which is only isomorphic to  $\text{End}(E^{(p)})$ . Since  $\text{End}^0(E)$  is a quadratic imaginary number field, it has one nontrivial field automorphism, and thus the isomorphism  $\text{End}(E) \cong \text{End}(E^{(p)})$  is not unique. But there is a unique canonical isomorphism, i.e. an isomorphism that is induced by an (equivalently any) isogeny  $\phi : E \rightarrow E^{(p)}$  as

$$\Phi_* : \text{End}(E) \rightarrow \text{End}(E^{(p)}), \quad \alpha \mapsto \frac{1}{\deg(\phi)} \phi \circ \alpha \circ \hat{\phi}$$

This is the isomorphism we use, i.e. we say

$$E^{(p)}[\mathfrak{b}] = E^{(p)}[\Phi_*(\mathfrak{b})] \quad \text{and} \quad [\mathfrak{b}].E^{(p)} = [\Phi_*(\mathfrak{b})].E^{(p)} = E^{(p)}/E^{(p)}[\mathfrak{b}]$$

Now let  $\phi : E \rightarrow E/E[\mathfrak{b}] = E^{(p)}$  be a separable isogeny with kernel  $E[\mathfrak{b}]$  (by choosing the representative  $\mathfrak{b}$  of  $[\mathfrak{b}] \in \text{Cl}(\mathcal{O})$  correspondingly, we can assume that). We have

$$\ker(\phi^{(p)}) = E[\mathfrak{b}]^{(p)} = \bigcap_{\beta \in \mathfrak{b}} \ker(\beta)^{(p)} = \bigcap_{\beta \in \mathfrak{b}} \ker(\beta^{(p)}) = \bigcap_{\beta \in \mathfrak{b}^{(p)}} \ker(\beta)$$

---

<sup>1</sup>Remember that  $\cdot^{(p)}$  is functorial, hence we can also apply to isogenies  $E_i \rightarrow E_{i+1}$




Figure 2: A 3-isogeny vulcano over  $\mathbb{F}_{101^2} = \mathbb{F}_{101}[\alpha]$  that satisfies case III (in the plot have  $z = \alpha$ ).

Now note that the Frobenius isogeny  $\pi : E \rightarrow E^{(p)}$ ,  $P \mapsto P^{(p)}$  induces the canonical isomorphism  $\text{End}(E) \rightarrow \text{End}(E^{(p)})$  and so the image of  $\mathfrak{b}$  under that isomorphism is  $\mathfrak{b}' = \mathfrak{b}^{(p)} \leq \text{End}(E^{(p)})$ . Thus

$$\bigcap_{\beta \in \mathfrak{b}^{(p)}} \ker(\beta) = \bigcap_{\beta \in \mathfrak{b}'} \ker(\beta) = E^{(p)}[\mathfrak{b}'] = E^{(p)}[\mathfrak{b}]$$

So by the uniqueness of the image curve for an isogeny with fixed kernel yields that  $E = \text{im}(\phi^{(p)}) = [\mathfrak{b}].E^{(p)}$ . Thus  $[\mathfrak{b}]^2.E = [\mathfrak{b}].E^{(p)} = E$  and since the class group action is free, we see that  $[\mathfrak{b}]^2 = [(1)]$ .  $\square$

From this we get the

**Corollary 1.2.** *Assume that  $E = E_0 \rightarrow E_1 \rightarrow \dots \rightarrow E_n = E$  is the cycle once around the crater (and  $j(E) \in \mathbb{F}_{p^2} \setminus \mathbb{F}_p$ ). If  $E^{(p)} = E_i$  then  $n$  is even and  $i = n/2$ , i.e.  $E^{(p)}$  is*

on the other side of the crater<sup>2</sup>.

*Proof.* If  $l$  does not split in  $\mathcal{O}_K$ , then the crater has at most two elements and this is trivial. So assume  $(l) = \mathfrak{l}_1 \mathfrak{l}_2$ . It is known that then the action of  $[\mathfrak{l}_1]$  resp.  $[\mathfrak{l}_2]$  corresponds to walking around the crater in one direction resp. the other. So wlog  $[\mathfrak{l}_1].E_i = E_{i+1}$ .

Now assume that  $E^{(p)} = E_i$ , so  $[\mathfrak{b}].E = E_i = [\mathfrak{l}_1]^i.E$ . Since the action is free, it follows that  $[\mathfrak{b}] = [\mathfrak{l}_1]^i$ . By the previous theorem, we have now  $[\mathfrak{l}_1]^{2i} = [\mathfrak{b}]^2 = [(1)]$  and so  $[\mathfrak{l}_1]^{2i}.E = E_{2i} = E$ . Thus  $i = n/2$  and the claim follows.  $\square$

In particular, the path between  $E$  and  $E^{(p)}$  is likely to have length  $\omega(\log(p))$ , since the crater is usually large. This is displayed e.g. Figure 1.

### 1.3 Case III

We say we are in *Case III*, if  $E_0$  and  $E_0^{(p)}$  are in different  $l$ -isogeny volcanos, i.e. there is no  $l$ -isogeny path between them.

We give the example displayed in Figure 3. Consider  $E$  with  $j(E) = 64\alpha + 5$ . Then  $j(E^{(p)}) = j(E)^p = 37\alpha + 59$ . However, we have that  $E$  lies on the crater, together with curve of  $j$ -invariants

$$88\alpha + 70, 54\alpha + 52, 95\alpha + 11$$

Hence there is no 3-isogeny path from  $E$  to  $E^{(p)}$ . Note that  $[\mathcal{O}_K : \mathbb{Z}[\pi]] = 2^2 \cdot 3^2$  but  $[\mathcal{O}_K : \text{End}(E)] = 2^2$ , which shows that  $E$  lies on the crater.

Now we want to have a closer look onto the class group action in this case. Have  $d(\text{End}(E)) = -320$ , so  $K = \mathbb{Q}(\sqrt{-5})$  and  $d(\mathcal{O}_K) = -5$ . Hence, we have  $\text{End}(E) \cong \mathbb{Z}[4\sqrt{-5}]$  and  $\mathcal{O}_K \cong \mathbb{Z}[\sqrt{-5}]$ .

Sage tells us that  $h(\mathcal{O}_K) = 2$  and  $h(\text{End}(E)) = 8$ . With this, we can already see that

$$64\alpha + 5, 88\alpha + 70, 54\alpha + 52, 95\alpha + 11$$

and

$$(64\alpha + 5)^p, (88\alpha + 70)^p, (54\alpha + 52)^p, (95\alpha + 11)^p$$

is the set of  $j$ -invariants of all Elliptic Curves with endomorphism ring  $\cong \text{End}(E)$ . On this set,  $\text{Cl}(\mathbb{Z}[4\sqrt{-5}])$  then acts freely and transitively. Now it would be of course interesting to find out how  $\text{Cl}(\mathbb{Z}[4\sqrt{-5}])$  really looks like.

## 2 Properties of the endomorphism ring vs the cases

**Proposition 2.1.** *Let  $E$  be an ordinary Elliptic Curve defined over a finite field of characteristic  $p$ .*

- $\text{End}(E)$  has an element of norm  $p$  iff  $j(E) \in \mathbb{F}_p$ .
- $\text{End}(E)$  has a nontrivial element (i.e.  $\neq \epsilon p$  for a unit  $\epsilon$ ) of norm  $p^2$  iff  $j(E) \in \mathbb{F}_{p^2}$ .

<sup>2</sup>Note that this does not hold if  $E, E^{(p)}$  are not in the same crater, see Figure 2.




Figure 3: A 3-isogeny volcano over  $\mathbb{F}_{101^2} = \mathbb{F}_{101}[\alpha]$  that satisfies case III (in the plot have  $z = \alpha$ ).

*Proof.* The directions  $\Leftarrow$  is clear, as the norm of the  $q$ -th power Frobenius endomorphism is  $q$ .

For the direction  $\Rightarrow$ , assume there is an element  $\alpha \in \text{End}(E)$  with  $N(\alpha) = p$ . If  $\alpha$  is inseparable (as isogeny), then we have that it factors through the  $p$ -th power Frobenius endomorphism  $\pi$ , and thus  $\alpha = \lambda \circ \pi$  for an isomorphism  $\lambda : E^{(p)} \rightarrow E$ . Thus  $j(E^{(p)}) = j(E)$ .

On the other hand, if  $\alpha$  is separable, it must have kernel of size  $p$ , so  $\ker(\alpha) = E[p]$  since  $\#E[p] = p$  ( $E$  is ordinary). Thus  $\ker(\alpha) \subseteq \ker([p])$  and we see that  $[p]$  factors through  $\alpha$  as  $[p] = \psi \circ \alpha$ . Now have that  $\deg(\psi) = p = p^2/\deg(\alpha)$  and clearly  $\psi$  is inseparable. The claim follows as above.

For the second point, assume  $\alpha \in \text{End}(E)$  has norm  $N(\alpha) = p^2$  and  $\alpha \neq \pm p$ . If  $\alpha$  is purely inseparable, we are done. If  $\alpha$  is separable, its kernel must be  $E[p^2]$  and so it factors through  $[p^2]$ . Since  $[p^2]$  has inseparability degree  $p^2$ , we see that  $[p^2] = \pi^2 \circ \alpha$  where  $\pi$  is the  $p$ -th power Frobenius morphism. Since  $\alpha$  is an endomorphism of  $E$ , find  $\pi^2 : E \rightarrow E$ , thus  $j(E) \in \mathbb{F}_{p^2}$ .

Finally, if  $\alpha$  has inseparability degree  $p$ , then its kernel must be  $E[p]$  and so  $\alpha = \beta \circ \pi$  where  $\beta : E^{(p)} \rightarrow E$  is separable with kernel  $E^{(p)}[p]$ . However, by the uniqueness of the separable isogeny with kernel  $E^{(p)}[p]$ , we know that (up to isomorphism) also  $[p]$  is  $\beta \circ \pi$ . This now implies that  $\alpha = \epsilon p$  for some unit  $\epsilon$ .  $\square$

**Corollary 2.2.** *Let  $D < 0$ . Then the curves  $E$  with  $\text{End}(E) = \mathbb{Z}[\sqrt{D}]$  have  $j(E) \in \mathbb{F}_{p^2} \setminus \mathbb{F}_p$  if and only if*

$$a^2 - b^2 D = p$$

*has no solution  $a, b \in \mathbb{Z}$  and*

$$a^2 - b^2 D = p^2$$

*has a nontrivial solution  $a, b \in \mathbb{Z}$ .*

Let  $E/\mathbb{F}_{p^2}$  be an ordinary Elliptic Curve and write  $\mathcal{O} := \text{End}(E)$ ,  $\mathcal{K} := \text{End}(E) \otimes \mathbb{Q}$ . Let  $\pi$  be the  $q = p^2$ -th power Frobenius endomorphism and let  $t$  be its trace. Assume  $p \neq 2$ .

**Lemma 2.3.** *Have  $(p) = (p, \pi)(p, \pi - t)$  in  $\mathbb{Z}[\pi]$ ,  $\mathcal{O}$  resp.  $\mathcal{O}_{\mathcal{K}}$ .*

*Proof.* Have

$$(p, \pi)(p, \pi - t) = (p^2, p\pi, p\pi - pt, \pi^2 - t\pi) = (p^2, pt, p\pi, -p^2) = (up^2 + vtp, \dots) = (p)$$

where  $up + vt = 1$  (note that  $t \perp p$  since  $E$  is ordinary).  $\square$

**Lemma 2.4.**  *$(p, \pi)$  is principal (in  $\mathcal{O}$ ) if and only if  $E/\mathbb{F}_p$ .*

*Proof.* If  $(p, \pi)$  is principal, then its generator is an element of norm  $p$ , so  $E/\mathbb{F}_p$ . On the other hand, if  $E/\mathbb{F}_p$ , then the  $p$ -th power Frobenius endomorphism  $\pi_p$  satisfies  $p = \pi_p(t_p - \pi_p)$ ,  $\pi = \pi_p^2$  and  $\pi_p = u(\pi + p) + v\pi_p p$ , where  $t_p$  is its trace and  $ut + vp = 1$ .  $\square$

There must be some problem in my definition of the class group action, as it can happen that  $[(p, \pi)]$  is not  $[(1)]$ , but  $E[(p, \pi)]$  is clearly trivial, so<sup>3</sup>  $(p, \pi).E = E/E[(p, \pi)] = E$ . However, this contradicts the freeness of the class group action.

**Lemma 2.5.** *Assume  $j(E) \neq 0, 1728$  (and  $E$  is defined over  $\mathbb{F}_{p^2}$ ).  $[(p, \pi)]$  has order  $\leq 2$  in  $\text{Cl}(\mathcal{O})$  resp.  $\text{Cl}(\mathcal{O}_K)$ .*

*Proof.* Since  $E/\mathbb{F}_{p^2}$ , we know that there is a nontrivial element  $\alpha$  of norm  $p^2$ . Now have in  $\mathcal{O}_K$  that  $(\alpha)|(p)^2$  and with  $p = (p, \pi)(p, \pi - t)$  have thus  $(\alpha) = (p)$  or  $(\alpha) = (p, \pi)^2$  or  $(\alpha) = (p, \pi - t)^2$ . However, by assumption we only have units  $\pm 1$  in  $\mathcal{O}_K$  resp.  $\mathcal{O}$ , so the first case is impossible, as it implies  $\alpha = \pm p$ .

Note that  $[(p, \pi)] = [(p, \pi - t)^{-1}]$ , so wlog assume  $(\alpha) = (p, \pi)^2$ . It follows that  $(p, \pi)^2$  is principal, so  $[(p, \pi)]^2 = [(1)]$ .  $\square$

### 3 Ideals in $\mathcal{O}$ resp. $\mathcal{O}_K$

In this paragraph, I tried to improve my understanding of ideals, invertible ideals and the class group in a nonmaximal order in a quadratic imaginary number field. In particular, I was interested in which properties carry over from  $\mathcal{O}_K$  and which do not hold anymore.

Consider an ordinary Elliptic Curve  $E/\mathbb{F}_q$ ,  $\mathcal{O} := \text{End}(E)$ ,  $K := \mathcal{O} \otimes \mathbb{Q}$  and  $\mathcal{O}_K$  the ring of integers in  $K$ . Assume that  $j(E) \in \mathbb{F}_q$  is not contained in any proper subfield of  $\mathbb{F}_q$  and let  $\pi$  be the  $q$ -th power Frobenius endomorphism. Let  $t$  be its trace.

**Proposition 3.1.**  $p \nmid d(\mathcal{O})$

*Proof.* Have that

$$d(\mathbb{Z}[\pi]) = t^2 - 4q \perp p$$

since  $t \perp p$  as  $E$  is ordinary. The claim follows since  $d(\mathcal{O}) \mid d(\mathcal{O})[\mathcal{O} : \mathbb{Z}[\pi]]^2 = d(\mathbb{Z}[\pi])$ .  $\square$

**Proposition 3.2.** *Let  $\mathfrak{a} \leq \mathcal{O}$ . Then  $\mathfrak{a} \cap \mathbb{Z} = (a)$  with  $a \mid [\mathcal{O} : \mathfrak{a}] \mid a^2$ . Note that if  $\mathfrak{a} = \mathfrak{p}$  is prime, then trivially  $a$  must be prime.*

*Proof.* Clearly  $[\mathcal{O} : \mathfrak{a}] \in \mathfrak{a}$  as  $1 \in \mathcal{O}/\mathfrak{a}$  has order dividing  $\#(\mathcal{O}/\mathfrak{a}) = [\mathcal{O} : \mathfrak{a}]$ , so  $a \mid [\mathcal{O} : \mathfrak{a}]$ . On the other hand, have  $[\mathcal{O} : \mathfrak{a}] \mid [\mathcal{O} : a\mathcal{O}] = a^2$ .  $\square$

**Lemma 3.3.** *Let  $\mathfrak{p} \leq \mathcal{O}_K$  be a prime with  $\mathfrak{N}(\mathfrak{p}) \perp [\mathcal{O}_K : \mathcal{O}]$ . Then  $\mathfrak{p}$  has a set of generators in  $\mathcal{O}$ .*

*Proof.* Suppose  $\mathfrak{p}$  is a prime over  $p$ , and let  $\mathcal{O} = \mathbb{Z}[\phi]$ . We use the decomposition law in Dedekind ring extensions. Since  $\mathfrak{N}(\mathfrak{p}) \perp [\mathcal{O}_K : \mathcal{O}]$  are coprime, we can apply it with generator  $\phi$  of  $\mathcal{O}$ .

If  $\text{MiPo}(\phi) = f(X)g(X) \pmod{p}$  splits, then have

$$p\mathcal{O}_K = (p, f(\phi))(p, g(\phi))$$

---

<sup>3</sup>We know that  $(p, \pi)$  is invertible, as  $\frac{1}{p}(p, \pi)(p, \pi - t) = (1)$ .



and so the prime ideals over  $p$  are  $(p, f(\phi))$  and  $(p, g(\phi))$ . If  $\text{MiPo}(\phi) \bmod p$  is irreducible, then have that  $p\mathcal{O}_K$  is prime and thus the only prime ideal over  $p$ . Hence, all prime ideals over  $p$  (including  $\mathfrak{p}$ ) have a set of generators in  $\mathcal{O}$ .  $\square$

**Corollary 3.4.** *Let  $\mathfrak{a} \leq \mathcal{O}_K$  be an ideal with  $\mathfrak{N}(\mathfrak{a}) \perp [\mathcal{O}_K : \mathcal{O}]$ . Then  $\mathfrak{a}$  has a set of generators in  $\mathcal{O}$ .*

**Proposition 3.5.** *Let  $\mathfrak{p} \leq \mathcal{O}$  be a prime ideal with  $\mathfrak{N}(\mathfrak{p}) \perp [\mathcal{O}_K : \mathcal{O}]$  and  $\mathfrak{p}' = \mathfrak{p}\mathcal{O}_K$ . Then  $\mathcal{O}_{\mathfrak{p}} = (\mathcal{O}_K)_{\mathfrak{p}'}$ .*

*Proof.* We have  $\mathcal{O}_K = \mathbb{Z}[\alpha]$  and  $\mathcal{O} = \mathbb{Z}[f\alpha]$  where  $f = [\mathcal{O}_K : \mathcal{O}]$ . Thus  $f \notin \mathfrak{p}$  and so  $f \in \mathcal{O}_{\mathfrak{p}}^*$ . Therefore  $\mathcal{O}_K \subseteq \mathcal{O}_{\mathfrak{p}}$  and thus  $(\mathcal{O}_K)_{\mathfrak{p}'} \subseteq \mathcal{O}_{\mathfrak{p}}$ .  $\square$

**Proposition 3.6.** *Let  $\mathfrak{I}(\mathcal{O})$  resp.  $\mathfrak{I}(\mathcal{O}_K)$  denote the set of invertible ideals of norm  $\perp [\mathcal{O}_K : \mathcal{O}]$ . Then*

$$\mathfrak{I}(\mathcal{O}) \rightarrow \mathfrak{I}(\mathcal{O}_K), \quad \mathfrak{a} \mapsto \mathfrak{a}\mathcal{O}_K$$

*is a monoid isomorphism with inverse*

$$\mathfrak{I}(\mathcal{O}_K) \rightarrow \mathfrak{I}(\mathcal{O}), \quad \mathfrak{a} \mapsto \mathfrak{a} \cap \mathcal{O}$$

*Proof.* Clearly, this is a well-defined monoid homomorphism. Hence, we have to show that it is bijective.

By Corollary 3.4, we know that any  $\mathfrak{a} \leq \mathcal{O}_K$  with  $\mathfrak{N}(\mathfrak{a}) \perp [\mathcal{O}_K : \mathcal{O}]$  has generators in  $\mathcal{O}$ , thus  $(\mathfrak{a} \cap \mathcal{O})\mathcal{O}_K = \mathfrak{a}$ . This shows that  $\mathfrak{a} \cap \mathcal{O}$  is a preimage of  $\mathfrak{a}$ , and so the map is surjective.

Assume now  $\mathfrak{a}, \mathfrak{b} \leq \mathcal{O}$  with  $\mathfrak{a}\mathcal{O}_K = \mathfrak{b}\mathcal{O}_K$  and  $\mathfrak{N}(\mathfrak{a}), \mathfrak{N}(\mathfrak{b}) \perp [\mathcal{O}_K : \mathcal{O}]$ . We show that  $\mathfrak{a}_{\mathfrak{p}} = \mathfrak{b}_{\mathfrak{p}}$  for all primes  $\mathfrak{p} \leq \mathcal{O}$ . Note that if  $\mathfrak{N}(\mathfrak{p}) \not\perp [\mathcal{O}_K : \mathcal{O}]$ , this holds trivially, as  $\mathfrak{a}_{\mathfrak{p}} = \mathcal{O}_{\mathfrak{p}} = \mathfrak{b}_{\mathfrak{p}}$ . Otherwise, note that

$$\mathfrak{a}_{\mathfrak{p}} = \mathfrak{a}_{\mathfrak{p}}\mathcal{O}_{\mathfrak{p}} = \mathfrak{a}(\mathcal{O}_K)_{\mathfrak{p}} = \mathfrak{a}\mathcal{O}_K(\mathcal{O}_K)_{\mathfrak{p}} = \mathfrak{b}\mathcal{O}_K(\mathcal{O}_K)_{\mathfrak{p}} = \mathfrak{b}_{\mathfrak{p}}(\mathcal{O}_K)_{\mathfrak{p}} = \mathfrak{b}_{\mathfrak{p}}$$

as  $\mathcal{O}_{\mathfrak{p}} = (\mathcal{O}_K)_{\mathfrak{p}}$ . This shows that  $\mathfrak{a}_{\mathfrak{p}} = \mathfrak{b}_{\mathfrak{p}}$  at all primes, so  $\mathfrak{a} = \mathfrak{b}$  and our map is injective. Furthermore, since  $(\mathfrak{a} \cap \mathcal{O})\mathcal{O}_K = \mathfrak{a}$ , we see that it has the inverse

$$\mathfrak{I}(\mathcal{O}_K) \rightarrow \mathfrak{I}(\mathcal{O}), \quad \mathfrak{a} \mapsto \mathfrak{a} \cap \mathcal{O}$$

which must then be well-defined.  $\square$

**Proposition 3.7.** *Let  $\mathfrak{a} \leq R$  be a radical ideal in a commutative unital ring  $R$ . If  $\alpha \in \mathfrak{p}$  for all primes  $\mathfrak{p} \supseteq \mathfrak{a}$  then  $\alpha \in \mathfrak{a}$ .*

*Proof.* We have that  $\mathfrak{a}_{\alpha} \neq (1)$  otherwise  $\alpha^n \in \mathfrak{a}$ , so  $\alpha \in \mathfrak{a}$ . Thus  $\mathfrak{a}_{\alpha} \subseteq \mathfrak{m}$  for a maximal ideal  $\mathfrak{m} \leq R_{\alpha}$ . A preimage under  $R \rightarrow R_{\alpha}$  is now a prime  $\mathfrak{p}$  with  $\mathfrak{a} \subseteq \mathfrak{p}$  and  $\alpha \notin \mathfrak{p}$ .  $\square$

**Corollary 3.8.** *If  $q \perp d(\mathcal{O})$  is an integer and  $q \mid \alpha$  in  $\mathcal{O}_K$ , then also  $q \mid \alpha$  in  $\mathcal{O}$ .*

*Proof.* It suffices to prove this for primes  $q$ . Since  $q \nmid d(\mathcal{O})$ , we know that  $(q)$  is unramified, hence radical. Now observe that  $\mathcal{O}_{\mathfrak{p}} = (\mathcal{O}_K)_{\mathfrak{p}}$  for primes  $\mathfrak{p}$  over  $q$  and so  $\alpha \in \mathfrak{p}$  for all primes  $\mathfrak{p}$  over  $q$ . The previous proposition now shows that  $\alpha \in (q)$ .  $\square$

## 4 Norm equations

**Lemma 4.1.** *Let  $D < 0$  and  $\mathcal{O}$  be the imaginary quadratic order of discriminant  $D$ . Then  $1, \alpha$  with  $\alpha = \frac{D+\sqrt{D}}{2}$  is a  $\mathbb{Z}$ -basis of  $\mathcal{O}$  and*

$$N(a + b\alpha) = \left(a + \frac{D}{2}b\right)^2 - \frac{D}{4}b^2$$

*Proof.*

$$\begin{aligned} N(\alpha) &= a^2 + ab\frac{D+\sqrt{D}}{2} + ab\frac{D-\sqrt{D}}{2} + \frac{D^2-D}{4}b^2 = a^2 + Dab + \frac{D^2-D}{4}b^2 \\ &= \left(a + \frac{D}{2}b\right)^2 - \frac{D}{4}b^2 \end{aligned}$$

□

**Corollary 4.2.** *Let  $l$  be a prime and  $D < 0$ . Let  $\mathcal{O}$  be the quadratic imaginary order of discriminant  $D$ . If there exists a nontrivial element  $\alpha \in \mathcal{O}$  of norm  $l^e$  (i.e.  $\alpha \notin \mathbb{Z}$ ), then*

$$e \geq \log_l(-D) - \log_l(4)$$

**Corollary 4.3.** *Let  $E/\bar{\mathbb{F}}_p$  be an ordinary Elliptic Curve such that  $\text{End}(E)$  has discriminant  $D$ . Suppose that  $j(E_0) \notin \mathbb{F}_p$  and  $l$  is not ramified in  $\text{End}(E) \otimes \mathbb{Q}$ . Then the shortest  $l$ -isogeny path between  $E$  and  $E^{(p)}$  has length at least*

$$\frac{1}{2} \log_l(-D) - \log_l(2)$$

*Proof.* First, define  $E_0$  to be the corresponding curve on the crater of the  $l$ -isogeny volcano. Let  $v \in \mathfrak{N}$  be maximal with  $l^v \mid [\text{End}(E_0) : \text{End}(E)]$ , so  $\text{End}(E_0)$  has discriminant  $D/l^v$ . Now we know that  $E_0^{(p)}$  is at the opposite side of the crater, and the size of the crater is the order of  $[l_1]$  in  $\text{Cl}(\text{End}(E_0))$  where  $l = l_1 l_2$ . If  $e$  is this order, then have that  $l^e = (\alpha)$  is principal. Note that  $\alpha \notin \mathbb{Z}$ , otherwise we would have  $\alpha = \pm l^{e/2}$ , but since  $(\alpha) = l_1^e$ , we know that  $l_2 \nmid (\alpha)$  (here we use that  $l_1 \neq l_2$ , i.e.  $l$  is unramified).

So

$$e \geq \log_l(-D/l^v) - \log_l(4)$$

Thus, the distance of  $E_0$  and  $E_0^{(p)}$  is at least

$$\frac{1}{2}e \geq \frac{1}{2} \log_l(-D/l^v) - \frac{1}{2} \log_l(4) = \frac{1}{2} \log_l(-D) - \frac{1}{2}v - \log_l(2)$$

However, the shortest path from  $E$  to  $E_0$  has length  $v$ , and similarly for the shortest path from  $E^{(p)}$  to  $E_0^{(p)}$ . Thus we find that the length of the shortest path from  $E$  to  $E^{(p)}$  is at least

$$\frac{1}{2} \log_l(-D) - \frac{1}{2}v - \log_l(2) + 2v \geq \frac{1}{2} \log_l(-D) - \log_l(2)$$

□

### More sophisticated bounds

First of all, note that the supersingular  $l$ -isogeny graph is  $(l+1)$ -regular and Ramajuan. Hence, the statistical  $\ell_1$ -distance of the random walk distribution after  $t$  steps from the uniform distribution is

$$\leq \lambda^t \sqrt{n} = \left( \frac{d}{2\sqrt{d}-1} \right)^t \sqrt{n} \approx \left( \frac{\sqrt{l}}{2} \right)^t \sqrt{n}$$

We look for the minimal number of steps  $t$  such that every node has nonzero probability (i.e. there exists a length  $\leq t$  path), this is then

$$\log_{\sqrt{l}/2}(n/\sqrt{n}) = \frac{1}{2} \log_{\sqrt{l}/2}(n) = \frac{\ln(n)}{2 \ln(\sqrt{l}/2)} = \frac{\ln(n)}{\ln(l) - 2 \ln(2)} \approx \log_l(n)$$

Since  $n \approx p/12$ , assume for now that we consider paths of length  $\log_l(p)$ .

Hence, we ask how many ordinary vulcanos have  $l$ -isogeny crater  $\leq 2 \log_l(n)$ . This is then equivalent to

$$x^2 + Dy^2 = \underbrace{l^{2 \log_l(p)}}_{=:m} = \Theta(p^2) \text{ is solvable with } x, y \in \mathbb{Z}$$

Now consider a fixed  $0 \leq y \leq \sqrt{p}$ . We have

$$\begin{aligned} & |\{p \leq D \leq p^2 \mid x^2 + Dy^2 = m \text{ solvable with } x \in \mathbb{Z}\}| \\ &= |\{p \leq D \leq p^2 \mid m - Dy^2 \text{ square}\}| \\ &\leq |\{p \leq D \leq p^2 \mid \left( \frac{m - Dy^2}{q_i} \right) = 1 \ \forall i\}| \end{aligned}$$

where  $q_i$  are primes  $\perp y$ . Now note that if  $q := \prod q_i \leq p^2 - p$  and we assume  $D$  to be uniformly distributed on  $\mathbb{Z}/q\mathbb{Z}$ , then also  $Dy^2$  is uniformly distributed on  $\mathbb{Z}/q\mathbb{Z}$  and so is  $m - Dy^2$ . Hence the events  $(m - Dy^2/q_i) = 1$  are independent with probability  $1/2$ .

To find out how many such primes we can choose, note that the product of the first  $n$  primes is  $\approx \prod_{i \leq n} n \ln(n) \approx \exp(n \ln(n) + n \ln \ln(n)) \approx n^n$ . To invert the function  $x \mapsto x^x$ , we use the Lambert W function, and find that the inverse is  $x \mapsto \exp(W(\ln(x)))$ . There is the known bound

$$\ln(x) - \ln \ln(x) < W(x)$$

and so we see that the product of the first

$$\exp(\ln \ln(p^2) - \ln \ln \ln(p^2)) \approx 2 \ln(p) / \ln \ln(p)$$

primes is  $\leq p^2$ . Hence we have the bound on the probability

$$\begin{aligned} \Pr \left[ \left( \frac{m - Dy^2}{q_i} \right) = 1 \ \forall i \right] &\approx 2^{-\exp(\ln \ln(p^2) - \ln \ln \ln(p^2))} \\ &= \exp(-2 \ln(2) \ln(p) / \ln \ln(p)) = p^{-\frac{2 \ln(2)}{\ln \ln(p)}} \end{aligned}$$

where the  $\ln(\sqrt{p})$  factor comes from excluding primes not coprime to  $y$ . Now summing over all  $y \leq \sqrt{p}$  yields

$$\Pr[x^2 + Dy^2 = m \text{ solvable}] \leq \sqrt{p} p^{-\frac{2 \ln(2)}{\ln \ln(p)}} \sqrt{p} = p^{1/2 - \frac{2 \ln(2)}{\ln \ln(p)}}$$

This results in a bound on the number of  $D$  of

$$p^2 p^{1/2 - \frac{2 \ln(2)}{\ln \ln(p)}} = p^{5/2 - \frac{2 \ln(2)}{\ln \ln(p)}}$$

which is of course totally useless.

## 5 Example - j-invariant $61\alpha + 16$

Let  $E$  be an Elliptic Curve defined over  $\mathbb{F}_{101^2}$  with j-invariant  $61\alpha + 16$ . Then the  $q$ -th power Frobenius  $\pi$  has minimal polynomial

$$X^2 - 190X + 10201$$

Furthermore, we find that  $\text{End}(E) = \mathcal{O}_{\mathcal{K}}$  for  $\mathcal{K} = \mathbb{Q}(\sqrt{-6})$ . So

$$\pi = \frac{190 + 28\sqrt{-6}}{2} = 95 + 14\sqrt{-6}$$

or

$$\sqrt{-6} = \frac{\pi - 95}{14}$$

Note that  $\pi - 95$  has norm  $2^3 \cdot 3 \cdot 7^2$ . The class group of  $\mathcal{O}$  has order 2, and a generator is e.g. the coset of  $(2, \sqrt{-6})$ . Hence, to find  $E[(2, \sqrt{-6})]$  we need to find

$$\ker\left(\frac{\pi - 95}{14}\right) \cap E[2] = 14 \ker(\pi - 95) \cap E[2] = 14(\ker(\pi - 95) \cap E[28])$$

Now choose a  $\mathbb{Z}/4\mathbb{Z}$ -basis  $P_1, P_2$  of  $E[4]$  and a  $\mathbb{Z}/7\mathbb{Z}$ -basis  $Q_1, Q_2$  of  $E[7]$ . Have that w.r.t. these basis,  $\pi$  is given by the matrices

$$\begin{pmatrix} 3 & 2 \\ 0 & 3 \end{pmatrix} \quad \text{resp.} \quad \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix}$$

Since  $95 \equiv 3 \pmod{4}$  and  $95 \equiv 4 \pmod{7}$ , we see that  $\ker(\pi - 95) \cap E[28]$  projects to

$$\langle P_1, 2P_2 \rangle \subseteq E[4] \quad \text{and} \quad E[7] \subseteq E[7]$$

and thus is  $E[14] + \langle P_1 \rangle$ . This implies that  $E[(2, \sqrt{-6})] = \langle 2P_1 \rangle$ . Note that we picked

$$\begin{aligned} P_1 &= (59 + 7\alpha, 48 + 75\alpha + (73 + 3\alpha)t), \\ P_2 &= (7 + 17\alpha + 100t, 71 + 72\alpha + (31 + 88\alpha)t) \end{aligned}$$

$j(E)$	$h(\text{End}(E))$	$[\mathcal{O}_K : \text{End}(E)]$	$d(\text{End}(E))$	$[\text{End}(E) : \mathbb{Z}[\pi]]$
$\alpha$	36	3	-36315	1
$4\alpha + 99$	64	1	-40020	1
$61\alpha + 16$	2	1	-24	28
$48\alpha + 73$	64	?	-37440	?
$12\alpha + 79$	12	?	-2548	?
$91\alpha + 34$	24	?	-16468	?
$95\alpha + 20$	64	?	-40548	?
$97\alpha + 12$	48	?	-35475	?
$97\alpha + 8$	48	?	-35620	?
$93\alpha + 8$	24	?	-23643	?
$77\alpha + 16$	16	?	-2340	?
$21\alpha + 48$	30	?	-35179	?
$31\alpha + 59$	48	?	-29355	?
$82\alpha + 39$	24	?	-18603	?
$64\alpha + 38$	36	?	-40075	?
$92\alpha + 74$	32	?	-30195	?
$38\alpha + 18$	16	?	-2340	?
$69\alpha + 25$	40	?	-31588	?
$99\alpha + 64$	32	?	-30195	?
$56\alpha + 4$	32	?	-30195	?
$26\alpha + 90$	12	?	-2548	?
$93\alpha + 49$	48	?	-36708	?
$17\alpha + 16$	32	?	-13908	?
$84\alpha + 67$	4	?	-180	?
$100\alpha + 34$	56	?	-40788	?
$30\alpha + 2$	16	?	-2244	?
$21\alpha + 41$	2	?	-52	?
$24\alpha + 59$	24	?	-26643	?
$67\alpha + 94$	64	?	-37204	?
$88\alpha + 99$	2	?	-88	?
$47\alpha + 26$	48	?	-24420	?
$12\alpha + 7$	16	?	-2520	?
$55\alpha + 77$	24	?	-17395	?
$95\alpha + 92$	8	?	-987	?
$68\alpha + 12$	12	?	-756	?
$82\alpha + 66$	28	?	-4532	?
$91\alpha + 38$	16	?	-6948	?
$99\alpha + 20$	24	?	-18603	?
$52\alpha + 77$	80	?	-40404	?

Table 1: Table of class numbers of  $\text{End}(E)$  for Elliptic Curves  $E/\mathbb{F}_{101^2} = \mathbb{F}_{101}[\alpha]$ . Note that the  $j$ -values are not uniformly chosen, in particular,  $j$ -values that lead to a conductor  $[\mathcal{O}_K : \mathbb{Z}[\pi]]$  with “big” prime power divisors have been ignored, as the current implementation of computing the endomorphism ring would take ages for them.




Figure 4: The 2-isogeny vulcano containing  $61\alpha+16$ . Note that 2 is ramified in  $\text{End}(E) \otimes \mathbb{Q}$ , thus the crater only has size 2.

before, where  $t$  has minimal polynomial  $(24+51\alpha)+(94+84\alpha)T+T^2$ . Hence,  $\overline{(2, \sqrt{-6})}.E$  is the (isomorphism class of the) image of the 2-isogeny  $\phi : E \rightarrow E/\langle 2P_1 \rangle$ , which is

$$j(E/\langle 2P_1 \rangle) = 40\alpha + 58 = (61\alpha + 16)^{101}$$

The 2-isogeny vulcano containing  $61\alpha + 16$  is shown in

To find the whole kernel, pick a  $\mathbb{Z}/8\mathbb{Z}$ -basis  $P_1, P_2$  of  $E[8]$ , a  $\mathbb{Z}/3\mathbb{Z}$ -basis  $Q_1, Q_2$  of  $E[3]$  and a  $\mathbb{Z}/49\mathbb{Z}$ -basis  $R_1, R_2$  of  $E[49]$ . Find then that modulo 8, 3 resp. 49,  $\pi$  is given by the matrix

$$\begin{pmatrix} 3 & 6 \\ 4 & 3 \end{pmatrix} \quad \text{resp.} \quad \begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix} \quad \text{resp.} \quad \begin{pmatrix} 32 & 14 \\ 0 & 11 \end{pmatrix}$$

## 6 Example - The ordinary endomorphism ring

The information in this section is all known material - I just wanted to understand properly how one can compute the endomorphism ring, and what problems occur.

Consider the finite field

$$\mathbb{F}_q = \mathbb{F}_{37^2} = \mathbb{F}_{37} + \alpha\mathbb{F}_{37}$$

where  $\alpha^2 + 33\alpha + 2 = 0$ . Further, consider the Elliptic Curve  $E/\mathbb{F}_q$  with  $j$ -invariant  $3\alpha$ , given by

$$E : y^2 = x^3 + (15\alpha + 17)x + (5\alpha + 3)$$

Then we find that the  $q$ -th power Frobenius endomorphism  $\pi$  satisfies the minimal equation

$$\pi^2 + 47\pi + 1369$$

and in particular, its trace is  $-47$ . Hence, the number field  $\mathcal{K} := \mathcal{O} \otimes \mathbb{Q}$  where  $\mathcal{O} = \text{End}(E)$  contains  $\sqrt{47^2 - 4 \cdot 1369} = \sqrt{-3^3 \cdot 11^2}$ . We observe that  $\mathcal{K} = \mathbb{Q}(\sqrt{-3})$  and has discriminant  $-3$ . Furthermore the ring of integers is  $\mathcal{O}_{\mathcal{K}} = \mathbb{Z}[\frac{1}{2}(1 + \sqrt{-3})]$ .

Knowing the number field, we want to find the endomorphism ring. First, observe that the Frobenius order  $\mathbb{Z}[\pi]$  has conductor 33. Now consider the endomorphism

$$\phi := 2\pi + 47$$

The advantage is that we can evaluate  $\phi$  on points of  $E$ , but evaluating  $\pi + 47/2$  is not so easy. Clearly  $[\mathbb{Z}[\pi] : \mathbb{Z}[\phi]] = 2$  and so  $\mathbb{Z}[\phi]$  has conductor 66.

### Torsion points

In order to find whether  $\phi/n \in \mathcal{O}$ , we factor  $66 = 2 \cdot 3 \cdot 11$  and compute the corresponding torsion groups. This turns out to be quite difficult.

Assume  $\mathbb{F}_{37^{12}} = \mathbb{F}_{37}[\beta]$  with

$$\text{MiPo}_{\mathbb{F}_{37}}(\beta) = x^{12} + 4x^7 + 31x^6 + 10x^5 + 23x^4 + 18x^2 + 33x + 2$$

Then  $E[2]$  is generated by

$$\begin{aligned} P_1 &= (11\beta^{11} + 19\beta^{10} + \beta^9 + 27\beta^8 + 8\beta^7 + 16\beta^6 + 17\beta^5 + 32\beta^4 + 12\beta^3 + 14\beta^2 + 24\beta + 32 : 0 : 1) \\ Q_1 &= (15\beta^{11} + 7\beta^{10} + 33\beta^9 + 11\beta^8 + 6\beta^7 + 12\beta^6 + 26\beta^5 + 7\beta^4 + 33\beta^3 + 25\beta^2 + 8\beta + 19 : 0 : 1) \end{aligned}$$

Further  $E[3]$  is generated by

$$\begin{aligned} P_2 &= (19\beta^{11} + 34\beta^{10} + 3\beta^9 + 29\beta^8 + 7\beta^7 + 3\beta^6 + 18\beta^5 + 21\beta^4 + 23\beta^3 + 30\beta^2 + 23\beta + 25 \\ &\quad : 6\beta^{11} + 25\beta^{10} + 4\beta^9 + 13\beta^8 + 10\beta^7 + 23\beta^6 + 20\beta^5 + 30\beta^4 + 24\beta^3 + 6\beta^2 + 17\beta + 5 : 1) \\ Q_2 &= (31\beta^{11} + 24\beta^{10} + 35\beta^9 + 32\beta^8 + 2\beta^7 + 10\beta^6 + 23\beta^5 + 35\beta^4 + 22\beta^3 + 13\beta^2 + 12\beta + 12 \\ &\quad : 18\beta^{11} + 2\beta^{10} + 32\beta^9 + 26\beta^8 + 17\beta^7 + 5\beta^6 + 19\beta^5 + 31\beta^4 + 31\beta^3 + \beta^2 + 22\beta + 1 : 1) \end{aligned}$$

For  $E[11]$  we must even go to the extension degree 110. So assume  $\mathbb{F}_{37^{220}} = \mathbb{F}_{37}[\gamma]$ . Then  $E[11]$  is generated by  $P_3$  and  $Q_3$ . For the values of  $\text{MiPo}_{\mathbb{F}_{37}}(\gamma)$  and  $P_3, Q_3$  see Section 7.

Now we can compute  $\phi(P_1), \phi(Q_1), \phi(P_2), \phi(Q_2), \phi(P_3), \phi(Q_3)$  and see that none of them is zero. Since  $\deg(\phi) = [\mathcal{O} : \mathbb{Z}[\phi]] \mid [\mathcal{O}_{\mathcal{K}} : \mathbb{Z}[\phi]] = 2 \cdot 3 \cdot 11$ , we see that the kernel of  $\phi$  is trivial. Thus no  $\phi/n$  is contained in  $\mathcal{O}$ . Therefore we see that

$$\mathcal{O} \cap \mathbb{Z}[\sqrt{D}] = \mathbb{Z}[\phi]$$

The inclusion  $\supseteq$  is clear, and for the other direction, note that  $\mathcal{O} \cap \mathbb{Z}[\sqrt{D}] = \mathbb{Z} + t\sqrt{D}\mathbb{Z}$  and  $\mathbb{Z}[\phi] = \mathbb{Z} + s\sqrt{D}\mathbb{Z}$ . Since  $\mathbb{Z}[\phi] \subseteq \mathcal{O} \cap \mathbb{Z}[\phi]$  find thus  $t \mid s$ . Now observe that by choice of  $\phi$ , have  $\phi^2 \in \mathbb{Z}$  and so  $\phi = s\sqrt{D}$ . However,  $\phi/\frac{s}{t} = t\sqrt{D} \in \mathcal{O}$ . By the above, it follows that  $\frac{s}{t} = 1$ , i.e.  $s = t$ .

### The index $[\mathcal{O} : \mathbb{Z}[\phi]]$

From the consideration of the torsion points, we see that  $\mathcal{O} \cap \mathbb{Z}[\sqrt{D}] = \mathbb{Z}[\phi]$ . However, since  $[\mathcal{O}_{\mathcal{K}} : \mathbb{Z}[\sqrt{D}]] \leq 2$ , we deduce that  $[\mathcal{O} : \mathbb{Z}[\phi]] \leq 2$  and so

$$\mathcal{O} = \mathbb{Z}[\pi]$$

### My implementation

I have implemented the above algorithm, with computing the torsion groups via division polynomials in Rust, as the SAGE functions for that are extremely slow (apparently, SAGE is not fast when working with polynomials over  $\mathbb{F}_{p^2}$ , I assume because elements of  $\mathbb{F}_{p^2}$  are considered objects and require dynamic memory allocation etc.). The advantage is that this can be extended to compute kernels of all endomorphisms, and so be used to compute the class group action.



## 7 $P_3$ and $Q_3$

The minimal polynomial of  $\gamma$  is

$$\begin{aligned}
& x^{220} + 31x^{219} + 13x^{218} + 21x^{217} + 23x^{216} + 9x^{215} \\
& + 2x^{214} + 35x^{212} + 10x^{211} + 29x^{210} + 25x^{209} + 20x^{208} \\
& + 17x^{207} + 30x^{206} + 5x^{205} + 15x^{204} + 11x^{203} + 10x^{202} \\
& + 11x^{201} + 32x^{200} + 5x^{199} + 28x^{198} + 7x^{197} + 13x^{196} \\
& + 10x^{195} + 32x^{194} + 17x^{193} + 19x^{192} + 36x^{191} \\
& + 17x^{190} + 31x^{189} + 14x^{188} + 6x^{187} + 30x^{186} + 8x^{185} \\
& + 22x^{184} + 2x^{183} + 9x^{182} + 11x^{181} + 6x^{180} + 23x^{179} \\
& + 14x^{178} + 36x^{177} + 16x^{176} + 34x^{175} + 14x^{174} \\
& + 33x^{173} + 14x^{172} + 7x^{171} + 36x^{170} + 18x^{169} + 27x^{168} \\
& + 5x^{167} + 31x^{166} + 6x^{165} + 15x^{164} + 14x^{163} + 17x^{162} \\
& + 7x^{161} + 16x^{160} + 6x^{159} + 29x^{158} + 11x^{157} + 8x^{156} \\
& + 15x^{155} + 20x^{154} + 17x^{153} + 7x^{152} + 8x^{151} + 6x^{150} \\
& + 12x^{149} + 36x^{148} + 7x^{147} + 3x^{146} + 25x^{145} + 13x^{144} \\
& + 6x^{143} + 17x^{142} + 22x^{141} + 9x^{140} + 18x^{139} + 36x^{138} \\
& + x^{137} + 6x^{136} + 36x^{135} + 33x^{134} + 32x^{133} + 35x^{132} \\
& + 33x^{131} + 7x^{130} + 3x^{129} + 7x^{128} + 20x^{127} + 31x^{126} \\
& + 26x^{125} + 6x^{124} + 9x^{123} + 10x^{122} + 25x^{121} + 33x^{120} \\
& + 33x^{119} + 30x^{118} + 34x^{117} + 22x^{116} + 8x^{115} + 10x^{114} \\
& + 36x^{113} + 26x^{112} + 8x^{111} + 33x^{110} + 30x^{109} + 11x^{108} \\
& + 14x^{107} + 22x^{106} + 26x^{105} + 11x^{104} + 35x^{103} \\
& + 34x^{102} + 33x^{101} + 27x^{100} + 14x^{99} + 31x^{98} + 24x^{97} \\
& + x^{96} + 6x^{95} + 36x^{93} + 32x^{92} + 18x^{91} + 36x^{90} + 3x^{89} \\
& + 22x^{88} + 36x^{87} + 6x^{86} + 20x^{85} + 25x^{84} + 8x^{82} \\
& + 34x^{81} + 7x^{80} + 25x^{79} + 21x^{78} + 17x^{77} + 29x^{76} \\
& + 5x^{75} + 19x^{74} + 19x^{73} + 8x^{72} + 8x^{71} + 26x^{70} \\
& + 7x^{69} + 27x^{68} + 10x^{67} + 31x^{66} + 4x^{65} + 29x^{64} \\
& + 36x^{62} + 3x^{61} + 27x^{60} + 13x^{59} + 23x^{58} + 33x^{57} \\
& + 14x^{56} + 19x^{55} + 12x^{54} + 20x^{53} + 32x^{52} + 18x^{51} \\
& + 20x^{49} + 20x^{48} + x^{47} + 17x^{46} + 16x^{45} + 4x^{44} \\
& + 12x^{43} + 7x^{42} + 34x^{41} + 9x^{40} + 16x^{39} + 10x^{38} \\
& + 25x^{37} + 10x^{36} + 10x^{35} + 28x^{34} + 33x^{33} + 22x^{32} \\
& + 24x^{31} + 33x^{30} + 6x^{29} + 8x^{28} + 8x^{27} + 16x^{26} \\
& + 31x^{25} + 7x^{24} + 26x^{23} + 36x^{22} + 29x^{21} + 36x^{20} \\
& + 7x^{19} + x^{18} + 26x^{17} + 18x^{16} + 23x^{15} + 10x^{14} \\
& + 4x^{13} + x^{12} + 24x^{11} + 25x^{10} + 34x^9 + 33x^8 \\
& + 33x^7 + 8x^6 + 12x^5 + x^4 + 15x^3 + 27x^2 + 9x + 2
\end{aligned}$$

$P_3$  is given by

$$\begin{aligned}
& (23z^{220} + 5z^{219} + 26z^{218} + 27z^{217} + 27z^{216} \\
& + 26z^{215} + 12z^{214} + 11z^{213} + 10z^{212}
\end{aligned}$$

$$\begin{aligned}
& + 29*z^{220}_{211} + 9*z^{220}_{210} + 16*z^{220}_{209} + 24*z^{220}_{208} \\
& + 18*z^{220}_{207} + 11*z^{220}_{206} + 11*z^{220}_{205} + 6*z^{220}_{204} \\
& + 24*z^{220}_{203} + 3*z^{220}_{202} + 34*z^{220}_{201} + 18*z^{220}_{200} \\
& + 17*z^{220}_{199} + 9*z^{220}_{198} + 26*z^{220}_{197} + 2*z^{220}_{196} \\
& + 31*z^{220}_{195} + 7*z^{220}_{194} + 15*z^{220}_{193} + 11*z^{220}_{192} \\
& + 15*z^{220}_{191} + 28*z^{220}_{190} + 13*z^{220}_{189} + 6*z^{220}_{188} \\
& + 7*z^{220}_{187} + 28*z^{220}_{186} + 9*z^{220}_{185} + 9*z^{220}_{184} \\
& + 7*z^{220}_{183} + 27*z^{220}_{182} + 36*z^{220}_{181} + 35*z^{220}_{180} \\
& + 30*z^{220}_{179} + 32*z^{220}_{178} + 16*z^{220}_{177} + 15*z^{220}_{176} \\
& + 16*z^{220}_{175} + 9*z^{220}_{174} + 21*z^{220}_{173} + 6*z^{220}_{172} \\
& + 15*z^{220}_{171} + 3*z^{220}_{170} + 25*z^{220}_{169} + 23*z^{220}_{168} \\
& + z^{220}_{167} + 8*z^{220}_{166} + 34*z^{220}_{165} + 14*z^{220}_{164} \\
& + 12*z^{220}_{163} + 20*z^{220}_{162} + 4*z^{220}_{161} + 9*z^{220}_{160} \\
& + z^{220}_{159} + 25*z^{220}_{158} + 16*z^{220}_{157} + z^{220}_{156} \\
& + 21*z^{220}_{155} + 10*z^{220}_{154} + 7*z^{220}_{153} + 13*z^{220}_{152} \\
& + 32*z^{220}_{151} + 31*z^{220}_{150} + 17*z^{220}_{148} + 24*z^{220}_{147} \\
& + 26*z^{220}_{146} + 28*z^{220}_{145} + 27*z^{220}_{144} + 4*z^{220}_{143} \\
& + 5*z^{220}_{142} + 14*z^{220}_{141} + 26*z^{220}_{140} + 10*z^{220}_{139} \\
& + 14*z^{220}_{138} + 19*z^{220}_{137} + 20*z^{220}_{136} + 18*z^{220}_{135} \\
& + 16*z^{220}_{134} + 11*z^{220}_{133} + 23*z^{220}_{132} + 35*z^{220}_{131} \\
& + 22*z^{220}_{130} + 31*z^{220}_{129} + 34*z^{220}_{128} + 17*z^{220}_{127} \\
& + z^{220}_{126} + 15*z^{220}_{125} + 2*z^{220}_{124} + 22*z^{220}_{123} \\
& + 27*z^{220}_{122} + 6*z^{220}_{121} + 10*z^{220}_{120} + 7*z^{220}_{119} \\
& + 4*z^{220}_{118} + 26*z^{220}_{117} + z^{220}_{116} + 32*z^{220}_{115} \\
& + 29*z^{220}_{114} + 32*z^{220}_{113} + 18*z^{220}_{112} + 3*z^{220}_{111} \\
& + 28*z^{220}_{110} + 20*z^{220}_{109} + 17*z^{220}_{108} + 17*z^{220}_{107} \\
& + 32*z^{220}_{106} + 32*z^{220}_{105} + 26*z^{220}_{104} + 24*z^{220}_{103} \\
& + 17*z^{220}_{102} + 8*z^{220}_{101} + 3*z^{220}_{100} + 2*z^{220}_{99} \\
& + 16*z^{220}_{98} + 29*z^{220}_{97} + 19*z^{220}_{96} + 27*z^{220}_{95} \\
& + 4*z^{220}_{94} + 29*z^{220}_{93} + 24*z^{220}_{92} + 19*z^{220}_{91} \\
& + 2*z^{220}_{90} + 2*z^{220}_{89} + 32*z^{220}_{88} + 23*z^{220}_{87} \\
& + 32*z^{220}_{86} + 15*z^{220}_{85} + 24*z^{220}_{84} + 36*z^{220}_{83} \\
& + 29*z^{220}_{82} + 18*z^{220}_{81} + 2*z^{220}_{80} + z^{220}_{79} \\
& + 33*z^{220}_{78} + 34*z^{220}_{77} + 4*z^{220}_{76} + 11*z^{220}_{75} \\
& + 21*z^{220}_{74} + 15*z^{220}_{73} + 10*z^{220}_{72} + 24*z^{220}_{71} \\
& + 22*z^{220}_{70} + 22*z^{220}_{69} + 31*z^{220}_{68} + 32*z^{220}_{67} \\
& + 28*z^{220}_{66} + z^{220}_{65} + 17*z^{220}_{64} + 13*z^{220}_{63} \\
& + 32*z^{220}_{62} + 20*z^{220}_{61} + 32*z^{220}_{60} + 21*z^{220}_{59} \\
& + 34*z^{220}_{58} + 11*z^{220}_{57} + 29*z^{220}_{56} + 12*z^{220}_{55} \\
& + 22*z^{220}_{54} + 11*z^{220}_{53} + 36*z^{220}_{52} + 35*z^{220}_{51} \\
& + 19*z^{220}_{50} + 35*z^{220}_{49} + 8*z^{220}_{48} + 16*z^{220}_{47} \\
& + 16*z^{220}_{46} + 27*z^{220}_{45} + 32*z^{220}_{44} + 12*z^{220}_{43} \\
& + 15*z^{220}_{42} + 6*z^{220}_{41} + 36*z^{220}_{40} + 27*z^{220}_{39}
\end{aligned}$$

$$\begin{aligned}
& + 17*z^{220}_{220}^{38} + 20*z^{220}_{220}^{37} + 33*z^{220}_{220}^{36} + 34*z^{220}_{220}^{35} \\
& + 34*z^{220}_{220}^{34} + 3*z^{220}_{220}^{33} + 12*z^{220}_{220}^{32} + 12*z^{220}_{220}^{31} \\
& + 12*z^{220}_{220}^{30} + 5*z^{220}_{220}^{29} + 10*z^{220}_{220}^{28} + 13*z^{220}_{220}^{27} \\
& + 36*z^{220}_{220}^{26} + 16*z^{220}_{220}^{25} + 16*z^{220}_{220}^{24} + 15*z^{220}_{220}^{23} \\
& + 36*z^{220}_{220}^{22} + 18*z^{220}_{220}^{21} + 13*z^{220}_{220}^{20} + 26*z^{220}_{220}^{19} \\
& + 25*z^{220}_{220}^{18} + 21*z^{220}_{220}^{17} + 35*z^{220}_{220}^{16} + 3*z^{220}_{220}^{14} \\
& + 31*z^{220}_{220}^{13} + 8*z^{220}_{220}^{12} + 7*z^{220}_{220}^{11} + 10*z^{220}_{220}^{10} \\
& + 10*z^{220}_{220}^9 + 6*z^{220}_{220}^8 + 5*z^{220}_{220}^7 + 33*z^{220}_{220}^6 \\
& + 6*z^{220}_{220}^5 + 4*z^{220}_{220}^4 + 31*z^{220}_{220}^3 + 27*z^{220}_{220}^2 + 27*z^{220}_{220} + 14
\end{aligned}$$

$$\begin{aligned}
& : 8*z^{220}_{220}^{219} + 17*z^{220}_{220}^{218} + 27*z^{220}_{220}^{217} + 14*z^{220}_{220}^{216} \\
& + 6*z^{220}_{220}^{215} + 19*z^{220}_{220}^{214} + 18*z^{220}_{220}^{213} + 6*z^{220}_{220}^{212} \\
& + 30*z^{220}_{220}^{211} + 24*z^{220}_{220}^{210} + 33*z^{220}_{220}^{209} + 19*z^{220}_{220}^{208} \\
& + 27*z^{220}_{220}^{207} + 16*z^{220}_{220}^{206} + 24*z^{220}_{220}^{205} + 3*z^{220}_{220}^{204} \\
& + 4*z^{220}_{220}^{203} + 25*z^{220}_{220}^{202} + 29*z^{220}_{220}^{201} + 31*z^{220}_{220}^{200} \\
& + 23*z^{220}_{220}^{199} + 7*z^{220}_{220}^{198} + 28*z^{220}_{220}^{197} + 4*z^{220}_{220}^{196} \\
& + 26*z^{220}_{220}^{195} + 36*z^{220}_{220}^{194} + 18*z^{220}_{220}^{193} + 24*z^{220}_{220}^{192} \\
& + 29*z^{220}_{220}^{191} + 25*z^{220}_{220}^{190} + 23*z^{220}_{220}^{189} + 14*z^{220}_{220}^{188} \\
& + 33*z^{220}_{220}^{187} + 19*z^{220}_{220}^{186} + 14*z^{220}_{220}^{184} + 21*z^{220}_{220}^{183} \\
& + 10*z^{220}_{220}^{182} + 13*z^{220}_{220}^{181} + 21*z^{220}_{220}^{180} + 24*z^{220}_{220}^{179} \\
& + 33*z^{220}_{220}^{178} + 19*z^{220}_{220}^{177} + 7*z^{220}_{220}^{176} + 36*z^{220}_{220}^{175} \\
& + 30*z^{220}_{220}^{174} + 34*z^{220}_{220}^{173} + 27*z^{220}_{220}^{172} + 3*z^{220}_{220}^{171} \\
& + 34*z^{220}_{220}^{170} + 5*z^{220}_{220}^{169} + 36*z^{220}_{220}^{168} + 19*z^{220}_{220}^{167} \\
& + 27*z^{220}_{220}^{166} + 14*z^{220}_{220}^{165} + 10*z^{220}_{220}^{164} + 2*z^{220}_{220}^{163} \\
& + 31*z^{220}_{220}^{162} + 22*z^{220}_{220}^{161} + 7*z^{220}_{220}^{160} + 14*z^{220}_{220}^{159} \\
& + 5*z^{220}_{220}^{158} + 3*z^{220}_{220}^{157} + 22*z^{220}_{220}^{156} + 32*z^{220}_{220}^{155} \\
& + 21*z^{220}_{220}^{154} + 17*z^{220}_{220}^{153} + 34*z^{220}_{220}^{152} + 9*z^{220}_{220}^{151} \\
& + 33*z^{220}_{220}^{150} + 32*z^{220}_{220}^{149} + 24*z^{220}_{220}^{148} + 16*z^{220}_{220}^{147} \\
& + 19*z^{220}_{220}^{146} + 6*z^{220}_{220}^{145} + 26*z^{220}_{220}^{144} + 24*z^{220}_{220}^{143} \\
& + 34*z^{220}_{220}^{141} + 25*z^{220}_{220}^{140} + 17*z^{220}_{220}^{139} + 25*z^{220}_{220}^{138} \\
& + 19*z^{220}_{220}^{137} + 36*z^{220}_{220}^{136} + 7*z^{220}_{220}^{134} + 32*z^{220}_{220}^{133} \\
& + 24*z^{220}_{220}^{132} + 6*z^{220}_{220}^{131} + 12*z^{220}_{220}^{130} + 30*z^{220}_{220}^{129} \\
& + 35*z^{220}_{220}^{128} + 13*z^{220}_{220}^{127} + 29*z^{220}_{220}^{126} + 2*z^{220}_{220}^{125} \\
& + 24*z^{220}_{220}^{124} + 36*z^{220}_{220}^{123} + 34*z^{220}_{220}^{122} + 2*z^{220}_{220}^{121} \\
& + 33*z^{220}_{220}^{120} + 10*z^{220}_{220}^{119} + 33*z^{220}_{220}^{118} + 2*z^{220}_{220}^{117} \\
& + 17*z^{220}_{220}^{116} + 33*z^{220}_{220}^{115} + 14*z^{220}_{220}^{114} + 22*z^{220}_{220}^{113} \\
& + 27*z^{220}_{220}^{112} + 20*z^{220}_{220}^{111} + 23*z^{220}_{220}^{110} + 34*z^{220}_{220}^{109} \\
& + 6*z^{220}_{220}^{108} + 33*z^{220}_{220}^{107} + 14*z^{220}_{220}^{106} + 28*z^{220}_{220}^{105} \\
& + 29*z^{220}_{220}^{104} + 36*z^{220}_{220}^{103} + 22*z^{220}_{220}^{102} + 35*z^{220}_{220}^{101} \\
& + 8*z^{220}_{220}^{100} + 10*z^{220}_{220}^{99} + 10*z^{220}_{220}^{98} + 16*z^{220}_{220}^{97} \\
& + 19*z^{220}_{220}^{96} + 17*z^{220}_{220}^{95} + 21*z^{220}_{220}^{94} + 13*z^{220}_{220}^{93} \\
& + 24*z^{220}_{220}^{92} + 36*z^{220}_{220}^{91} + 25*z^{220}_{220}^{90} + 25*z^{220}_{220}^{89} \\
& + 22*z^{220}_{220}^{88} + 27*z^{220}_{220}^{87} + 28*z^{220}_{220}^{86} + 11*z^{220}_{220}^{85}
\end{aligned}$$

$$\begin{aligned}
& + 3*z^{220^84} + 14*z^{220^82} + 31*z^{220^81} + 7*z^{220^80} \\
& + 33*z^{220^79} + 33*z^{220^78} + 2*z^{220^77} + 15*z^{220^76} \\
& + 17*z^{220^75} + 32*z^{220^74} + 4*z^{220^73} + 18*z^{220^72} \\
& + 10*z^{220^71} + 34*z^{220^70} + 9*z^{220^69} + 3*z^{220^68} \\
& + 20*z^{220^67} + 33*z^{220^66} + 23*z^{220^65} + 5*z^{220^64} \\
& + 20*z^{220^63} + 36*z^{220^62} + 29*z^{220^61} + 2*z^{220^60} \\
& + 25*z^{220^59} + 14*z^{220^58} + 16*z^{220^57} + 31*z^{220^56} \\
& + 22*z^{220^55} + 31*z^{220^54} + 33*z^{220^53} + 19*z^{220^52} \\
& + 22*z^{220^51} + 23*z^{220^50} + 36*z^{220^49} + 11*z^{220^48} \\
& + 15*z^{220^47} + 15*z^{220^46} + 35*z^{220^45} + 7*z^{220^44} \\
& + 27*z^{220^43} + 28*z^{220^42} + 15*z^{220^41} + 31*z^{220^40} \\
& + 12*z^{220^39} + 19*z^{220^38} + 21*z^{220^37} + 18*z^{220^36} \\
& + 3*z^{220^35} + 36*z^{220^33} + z^{220^32} + 35*z^{220^31} \\
& + 21*z^{220^30} + 2*z^{220^29} + 13*z^{220^28} + 19*z^{220^27} \\
& + 6*z^{220^26} + 22*z^{220^24} + 26*z^{220^23} + 9*z^{220^22} \\
& + 7*z^{220^21} + 31*z^{220^20} + 31*z^{220^19} + 9*z^{220^18} \\
& + 23*z^{220^17} + 23*z^{220^16} + 6*z^{220^15} + 27*z^{220^14} \\
& + 36*z^{220^13} + 4*z^{220^12} + 26*z^{220^11} + 30*z^{220^10} \\
& + 9*z^{220^9} + 8*z^{220^8} + 15*z^{220^7} + 26*z^{220^6} \\
& + 17*z^{220^5} + 29*z^{220^4} + 24*z^{220^3} + 8*z^{220^2} \\
& + 29*z^{220} : 1)
\end{aligned}$$

$Q_3$  is given by

$$\begin{aligned}
& (35*z^{220^219} + 22*z^{220^218} + 36*z^{220^216} + 24*z^{220^215} \\
& + 19*z^{220^214} + 32*z^{220^213} + 13*z^{220^212} + 19*z^{220^211} \\
& + 3*z^{220^210} + 36*z^{220^209} + 29*z^{220^208} + 35*z^{220^206} \\
& + 31*z^{220^205} + 32*z^{220^204} + 23*z^{220^203} + 21*z^{220^202} \\
& + 10*z^{220^201} + 32*z^{220^200} + 32*z^{220^199} + 21*z^{220^198} \\
& + 16*z^{220^197} + 23*z^{220^196} + 32*z^{220^195} + 12*z^{220^194} \\
& + 9*z^{220^193} + 35*z^{220^192} + 8*z^{220^191} + 19*z^{220^190} \\
& + 33*z^{220^189} + 13*z^{220^188} + 11*z^{220^187} + 35*z^{220^186} \\
& + 25*z^{220^185} + 28*z^{220^184} + 5*z^{220^183} + 7*z^{220^182} \\
& + 24*z^{220^181} + 35*z^{220^180} + 33*z^{220^179} + 18*z^{220^178} \\
& + 5*z^{220^177} + 31*z^{220^176} + 18*z^{220^175} + 30*z^{220^174} \\
& + 27*z^{220^173} + 3*z^{220^172} + 8*z^{220^171} + 24*z^{220^170} \\
& + 14*z^{220^169} + 2*z^{220^168} + 16*z^{220^167} + 14*z^{220^166} \\
& + 18*z^{220^165} + 22*z^{220^164} + 32*z^{220^163} + 28*z^{220^162} \\
& + 7*z^{220^161} + 19*z^{220^160} + 3*z^{220^159} + 14*z^{220^158} \\
& + 27*z^{220^157} + 35*z^{220^156} + 8*z^{220^155} + 25*z^{220^154} \\
& + 11*z^{220^153} + 19*z^{220^152} + 21*z^{220^151} + 10*z^{220^150} \\
& + 2*z^{220^149} + 4*z^{220^148} + 4*z^{220^147} + 31*z^{220^146} \\
& + 26*z^{220^145} + 17*z^{220^143} + 14*z^{220^142} + 12*z^{220^141} \\
& + 17*z^{220^140} + 22*z^{220^139} + 30*z^{220^138} + 30*z^{220^137} \\
& + 15*z^{220^136} + 16*z^{220^135} + 25*z^{220^134} + 8*z^{220^133}
\end{aligned}$$

$$\begin{aligned}
& + 28*z^{220}_{132} + 5*z^{220}_{131} + 14*z^{220}_{130} + 26*z^{220}_{129} \\
& + 13*z^{220}_{128} + 10*z^{220}_{127} + 13*z^{220}_{126} + 10*z^{220}_{125} \\
& + 17*z^{220}_{124} + 33*z^{220}_{123} + 9*z^{220}_{122} + 9*z^{220}_{121} \\
& + 10*z^{220}_{120} + 12*z^{220}_{119} + 4*z^{220}_{118} + 6*z^{220}_{117} \\
& + 33*z^{220}_{116} + 21*z^{220}_{115} + 14*z^{220}_{114} + 33*z^{220}_{113} \\
& + 11*z^{220}_{112} + 4*z^{220}_{111} + 3*z^{220}_{110} + 3*z^{220}_{109} \\
& + 3*z^{220}_{108} + 3*z^{220}_{107} + 27*z^{220}_{106} + 8*z^{220}_{105} \\
& + 25*z^{220}_{104} + 10*z^{220}_{103} + 24*z^{220}_{102} + 2*z^{220}_{101} \\
& + 12*z^{220}_{100} + 35*z^{220}_{99} + 30*z^{220}_{98} + 14*z^{220}_{97} \\
& + 8*z^{220}_{96} + 16*z^{220}_{95} + 24*z^{220}_{94} + 23*z^{220}_{93} \\
& + 34*z^{220}_{91} + 3*z^{220}_{90} + 13*z^{220}_{89} + 10*z^{220}_{88} \\
& + 20*z^{220}_{87} + 14*z^{220}_{86} + 9*z^{220}_{85} + 36*z^{220}_{84} \\
& + 33*z^{220}_{83} + 12*z^{220}_{82} + 20*z^{220}_{81} + 5*z^{220}_{80} \\
& + 27*z^{220}_{79} + 27*z^{220}_{78} + 9*z^{220}_{77} + 23*z^{220}_{76} \\
& + 4*z^{220}_{75} + 26*z^{220}_{74} + 8*z^{220}_{73} + 11*z^{220}_{72} \\
& + 25*z^{220}_{71} + 35*z^{220}_{70} + 19*z^{220}_{69} + 36*z^{220}_{68} \\
& + 35*z^{220}_{67} + 24*z^{220}_{66} + 8*z^{220}_{65} + 32*z^{220}_{64} \\
& + 10*z^{220}_{63} + 3*z^{220}_{62} + 18*z^{220}_{61} + 35*z^{220}_{60} \\
& + 17*z^{220}_{59} + 30*z^{220}_{58} + 2*z^{220}_{57} + 25*z^{220}_{56} \\
& + 7*z^{220}_{55} + 20*z^{220}_{54} + 27*z^{220}_{53} + z^{220}_{52} \\
& + 10*z^{220}_{51} + 2*z^{220}_{50} + 18*z^{220}_{49} + 30*z^{220}_{48} \\
& + 32*z^{220}_{47} + 20*z^{220}_{46} + 4*z^{220}_{45} + 16*z^{220}_{43} \\
& + 16*z^{220}_{42} + 11*z^{220}_{41} + 8*z^{220}_{40} + 12*z^{220}_{39} \\
& + 15*z^{220}_{38} + 25*z^{220}_{37} + 33*z^{220}_{36} + 4*z^{220}_{35} \\
& + 11*z^{220}_{34} + 6*z^{220}_{33} + 7*z^{220}_{32} + 32*z^{220}_{31} \\
& + 19*z^{220}_{30} + 19*z^{220}_{29} + 16*z^{220}_{28} + 10*z^{220}_{27} \\
& + 7*z^{220}_{26} + 10*z^{220}_{25} + 33*z^{220}_{24} + 25*z^{220}_{23} \\
& + 21*z^{220}_{22} + 35*z^{220}_{21} + 15*z^{220}_{20} + z^{220}_{19} \\
& + 19*z^{220}_{18} + 16*z^{220}_{17} + 10*z^{220}_{16} + 18*z^{220}_{15} \\
& + 17*z^{220}_{14} + 2*z^{220}_{13} + 35*z^{220}_{12} + 30*z^{220}_{11} \\
& + 17*z^{220}_{10} + 30*z^{220}_9 + 26*z^{220}_8 + 9*z^{220}_7 \\
& + 34*z^{220}_6 + 4*z^{220}_5 + 12*z^{220}_4 + 16*z^{220}_3 \\
& + 27*z^{220}_2 + 12*z^{220} + 36 \\
\\
& : 21*z^{220}_{219} + 24*z^{220}_{218} \\
& + 33*z^{220}_{217} + 31*z^{220}_{216} + 29*z^{220}_{215} + 16*z^{220}_{214} \\
& + 26*z^{220}_{213} + 7*z^{220}_{212} + 15*z^{220}_{211} + 9*z^{220}_{210} \\
& + 19*z^{220}_{209} + 18*z^{220}_{208} + 16*z^{220}_{207} + 23*z^{220}_{206} \\
& + 27*z^{220}_{205} + 16*z^{220}_{204} + 5*z^{220}_{203} + 10*z^{220}_{202} \\
& + 2*z^{220}_{201} + 19*z^{220}_{200} + 19*z^{220}_{199} + 8*z^{220}_{198} \\
& + 30*z^{220}_{197} + 9*z^{220}_{196} + 27*z^{220}_{195} + 7*z^{220}_{194} \\
& + 20*z^{220}_{193} + 8*z^{220}_{192} + 29*z^{220}_{191} + 10*z^{220}_{190} \\
& + 32*z^{220}_{189} + 9*z^{220}_{188} + 4*z^{220}_{187} + 31*z^{220}_{186}
\end{aligned}$$

$$\begin{aligned}
& + 8*z^{220}_{185} + 4*z^{220}_{184} + 8*z^{220}_{183} + 11*z^{220}_{182} \\
& + 13*z^{220}_{181} + 5*z^{220}_{180} + 29*z^{220}_{179} + 13*z^{220}_{178} \\
& + 20*z^{220}_{177} + 9*z^{220}_{176} + 3*z^{220}_{175} + 32*z^{220}_{174} \\
& + 3*z^{220}_{173} + 25*z^{220}_{172} + 33*z^{220}_{171} + 36*z^{220}_{170} \\
& + 11*z^{220}_{169} + 22*z^{220}_{168} + 18*z^{220}_{167} + 7*z^{220}_{166} \\
& + 4*z^{220}_{165} + 9*z^{220}_{164} + 33*z^{220}_{163} + 33*z^{220}_{162} \\
& + 18*z^{220}_{161} + 3*z^{220}_{160} + 35*z^{220}_{159} + 31*z^{220}_{158} \\
& + 20*z^{220}_{157} + 28*z^{220}_{155} + 33*z^{220}_{154} + 30*z^{220}_{153} \\
& + 28*z^{220}_{152} + 18*z^{220}_{151} + z^{220}_{150} + 34*z^{220}_{149} \\
& + 16*z^{220}_{148} + 23*z^{220}_{147} + 30*z^{220}_{146} + 3*z^{220}_{144} \\
& + 28*z^{220}_{143} + 8*z^{220}_{142} + 35*z^{220}_{140} + 11*z^{220}_{139} \\
& + 16*z^{220}_{138} + 20*z^{220}_{137} + 31*z^{220}_{136} + 11*z^{220}_{135} \\
& + 24*z^{220}_{134} + 29*z^{220}_{133} + 29*z^{220}_{132} + 8*z^{220}_{131} \\
& + 25*z^{220}_{130} + 11*z^{220}_{129} + 35*z^{220}_{128} + 36*z^{220}_{127} \\
& + 33*z^{220}_{126} + 18*z^{220}_{125} + 8*z^{220}_{124} + 9*z^{220}_{123} \\
& + 31*z^{220}_{122} + 29*z^{220}_{121} + 7*z^{220}_{120} + 4*z^{220}_{119} \\
& + 3*z^{220}_{118} + 13*z^{220}_{117} + 35*z^{220}_{116} + 17*z^{220}_{115} \\
& + 6*z^{220}_{114} + 3*z^{220}_{113} + 13*z^{220}_{112} + 5*z^{220}_{111} \\
& + 31*z^{220}_{110} + 32*z^{220}_{109} + 17*z^{220}_{108} + 28*z^{220}_{107} \\
& + 21*z^{220}_{106} + 14*z^{220}_{105} + 25*z^{220}_{104} + 17*z^{220}_{103} \\
& + 33*z^{220}_{102} + 19*z^{220}_{101} + 4*z^{220}_{100} + 2*z^{220}_{99} \\
& + 7*z^{220}_{98} + 34*z^{220}_{97} + 15*z^{220}_{96} + 7*z^{220}_{95} \\
& + 34*z^{220}_{94} + 22*z^{220}_{93} + 22*z^{220}_{92} + 11*z^{220}_{91} \\
& + 33*z^{220}_{90} + 32*z^{220}_{89} + 19*z^{220}_{88} + 21*z^{220}_{87} \\
& + 23*z^{220}_{86} + 34*z^{220}_{85} + 35*z^{220}_{84} + 23*z^{220}_{83} \\
& + 27*z^{220}_{82} + 25*z^{220}_{81} + 26*z^{220}_{80} + 2*z^{220}_{79} \\
& + 33*z^{220}_{78} + 32*z^{220}_{77} + 8*z^{220}_{76} + 32*z^{220}_{75} \\
& + 15*z^{220}_{74} + 17*z^{220}_{73} + 31*z^{220}_{72} + 7*z^{220}_{71} \\
& + 8*z^{220}_{70} + 8*z^{220}_{69} + 22*z^{220}_{68} + 7*z^{220}_{67} \\
& + 14*z^{220}_{66} + 15*z^{220}_{65} + 26*z^{220}_{64} + 26*z^{220}_{63} \\
& + 35*z^{220}_{62} + 19*z^{220}_{61} + 18*z^{220}_{60} + 22*z^{220}_{59} \\
& + 25*z^{220}_{57} + 4*z^{220}_{56} + 5*z^{220}_{55} + 4*z^{220}_{54} \\
& + 20*z^{220}_{53} + 32*z^{220}_{52} + 17*z^{220}_{51} + 14*z^{220}_{50} \\
& + 31*z^{220}_{49} + 9*z^{220}_{48} + 30*z^{220}_{47} + 20*z^{220}_{46} \\
& + 7*z^{220}_{45} + 16*z^{220}_{43} + 23*z^{220}_{42} + 12*z^{220}_{41} \\
& + 21*z^{220}_{40} + 14*z^{220}_{39} + 8*z^{220}_{38} + 14*z^{220}_{37} \\
& + 35*z^{220}_{36} + 14*z^{220}_{35} + 22*z^{220}_{34} + 8*z^{220}_{33} \\
& + z^{220}_{32} + 24*z^{220}_{31} + 21*z^{220}_{30} + 33*z^{220}_{29} \\
& + 21*z^{220}_{28} + 22*z^{220}_{26} + 33*z^{220}_{25} + 13*z^{220}_{24} \\
& + 13*z^{220}_{23} + 5*z^{220}_{22} + 35*z^{220}_{21} + 3*z^{220}_{20} \\
& + 31*z^{220}_{19} + 13*z^{220}_{18} + 33*z^{220}_{17} + 30*z^{220}_{16} \\
& + 16*z^{220}_{15} + 30*z^{220}_{14} + 16*z^{220}_{13} + 11*z^{220}_{12} \\
& + 35*z^{220}_{11} + 22*z^{220}_{10} + 11*z^{220}_9 + 8*z^{220}_8
\end{aligned}$$

$$\begin{aligned}
& + z^{220^7} + 25z^{220^6} + 8z^{220^5} + 27z^{220^4} + z^{220^3} \\
& + 29z^{220^2} + 34z^{220} + 29 : 1)
\end{aligned}$$