

Table 1. Population characteristics

	Controls n = 544	Cases n = 136	P value
Female	432 (79.4%)	112 (20.6%)	1
Age (years), mean (SD)	82.3 (8.6)	82.3 (8.7)	0.935
51-65 years	28 (5.1%)	7 (5.1%)	
66-81 years	176 (32.4%)	44 (32.4%)	
82-95 years	340 (62.5%)	85 (62.5%)	
Weight (kg), mean (SD)	67.5 (13)	65 (11.6)	0.163
Height (cm), mean (SD)	155.5 (8)	152.4 (7.8)	0.035
BMI (kg/m ²), mean (SD)	27.8 (4.9)	27.5 (4.9)	0.035
Referred smoking habit			0
Non-smoker/not stated	522 (96%)	121 (89%)	
Active smoker	16 (2.9%)	6 (4.4%)	
Former smoker	6 (1.1%)	9 (6.6%)	
Referred alcoholic habit			0.042
Nondrinker/not stated	482 (88.6%)	128 (94.1%)	
Light drinker	59 (10.8%)	6 (4.4%)	
Moderate drinker	3 (0.6%)	2 (1.5%)	
Rheumatoid arthritis	5 (0.9%)	1 (0.7%)	0.827
Previous fracture	2 (0.4%)	58 (42.6%)	0
Diabetes mellitus	123 (22.6%)	46 (33.8%)	0.013
Diabetes mellitus type 1	4 (0.7%)	0 (0%)	0.01
Diabetes mellitus type 2	119 (21.9%)	46 (33.8%)	0.01
Osteoporosis	13 (2.4%)	19 (14%)	0
Neoplasia	65 (11.9%)	13 (9.6%)	0.409
Hiperthyroidism	3 (0.6%)	4 (2.9%)	0.111
Malnutrition	1 (0.2%)	3 (2.2%)	0.116
Malabsorption	1 (0.2%)	0 (0%)	0.318
Chronic liver disease	7 (1.3%)	5 (3.7%)	0.159
# active prescriptions			
Patients with >= act. pres.	229 (42.1%)	104 (76.5%)	0