

Table 1: Summary descriptives table by groups of ‘Grup’

	[ALL] N=408	Control N=272	Case N=136	p.overall
Age (years)	84.0 [78.0;88.0]	84.0 [78.0;88.0]	84.0 [78.0;88.0]	0.871
Sex. F: Female, M: Male:				1.000
F	300 (79.2%)	200 (79.1%)	100 (79.4%)	
M	79 (20.8%)	53 (20.9%)	26 (20.6%)	
Hospital discharge department:				<0.001
CGURG	31 (7.60%)	31 (11.4%)	0 (0.00%)	
GOURG	2 (0.49%)	2 (0.74%)	0 (0.00%)	
MDURG	225 (55.1%)	225 (82.7%)	0 (0.00%)	
MED	2 (0.49%)	0 (0.00%)	2 (1.47%)	
PSURG	1 (0.25%)	1 (0.37%)	0 (0.00%)	
TOURG	13 (3.19%)	13 (4.78%)	0 (0.00%)	
TRA	134 (32.8%)	0 (0.00%)	134 (98.5%)	
Weight (kg)	67.5 [57.1;74.5]	68.5 [59.0;76.7]	65.0 [57.0;71.0]	0.089
Height (cm)	153 [148;160]	154 [148;160]	152 [147;159]	0.344
BMI (kg/m ²)	28.1 [25.4;31.7]	28.7 [25.5;32.1]	27.4 [24.7;29.3]	0.136
ALCOHOL:				0.197
No	379 (92.9%)	251 (92.3%)	128 (94.1%)	
Light	26 (6.37%)	20 (7.35%)	6 (4.41%)	
Moderate	3 (0.74%)	1 (0.37%)	2 (1.47%)	
Smoker:				0.756
No	362 (88.7%)	241 (88.6%)	121 (89.0%)	
Active	15 (3.68%)	9 (3.31%)	6 (4.41%)	
Former	31 (7.60%)	22 (8.09%)	9 (6.62%)	
Hypogonadism: No	408 (100%)	272 (100%)	136 (100%)	.
Rheumatoid arthritis:				1.000
No	404 (99.0%)	269 (98.9%)	135 (99.3%)	
Yes	4 (0.98%)	3 (1.10%)	1 (0.74%)	
Previous fracture:				<0.001
No	296 (72.5%)	218 (80.1%)	78 (57.4%)	
Yes	112 (27.5%)	54 (19.9%)	58 (42.6%)	
Number of previous fractures	0.00 [0.00;1.00]	0.00 [0.00;0.00]	0.00 [0.00;1.00]	<0.001
Family history of fractures: No	408 (100%)	272 (100%)	136 (100%)	.
Diabetes:				0.473
No	281 (68.9%)	191 (70.2%)	90 (66.2%)	
Yes	127 (31.1%)	81 (29.8%)	46 (33.8%)	
tipus_diabetis:				0.433
No	281 (68.9%)	191 (70.2%)	90 (66.2%)	
Type 1	2 (0.49%)	2 (0.74%)	0 (0.00%)	
Type 2	125 (30.6%)	79 (29.0%)	46 (33.8%)	
Densitometry performed:				0.201
No	390 (95.6%)	263 (96.7%)	127 (93.4%)	
Yes	18 (4.41%)	9 (3.31%)	9 (6.62%)	
Osteoporosis:				0.006
No	374 (91.7%)	257 (94.5%)	117 (86.0%)	
Yes	34 (8.33%)	15 (5.51%)	19 (14.0%)	
Neoplasia:				0.311
No	358 (87.7%)	235 (86.4%)	123 (90.4%)	
Yes	50 (12.3%)	37 (13.6%)	13 (9.56%)	

Table 1: Summary descriptives table by groups of ‘Grup’ (*continued*)

	[ALL]	Control	Case	p.overall
Hyperthyroidism:				0.449
No	400 (98.0%)	268 (98.5%)	132 (97.1%)	
Yes	8 (1.96%)	4 (1.47%)	4 (2.94%)	
Malnutrition:				0.690
No	401 (98.3%)	268 (98.5%)	133 (97.8%)	
Yes	7 (1.72%)	4 (1.47%)	3 (2.21%)	
Malabsorption:				1.000
No	407 (99.8%)	271 (99.6%)	136 (100%)	
Yes	1 (0.25%)	1 (0.37%)	0 (0.00%)	
Chronic liver disease:				0.517
No	397 (97.3%)	266 (97.8%)	131 (96.3%)	
Yes	11 (2.70%)	6 (2.21%)	5 (3.68%)	
Other fracture risk factors:				0.751
No	334 (81.9%)	221 (81.2%)	113 (83.1%)	
Yes	74 (18.1%)	51 (18.8%)	23 (16.9%)	
Number of drugs ≥ 4 :				0.967
No	94 (23.0%)	62 (22.8%)	32 (23.5%)	
Yes	314 (77.0%)	210 (77.2%)	104 (76.5%)	
No. of previous fractures (categorised):				<0.001
0	296 (72.5%)	218 (80.1%)	78 (57.4%)	
1	95 (23.3%)	44 (16.2%)	51 (37.5%)	
≥ 2	17 (4.17%)	10 (3.68%)	7 (5.15%)	
No. of previous fractures (continuous)	0.00 [0.00;1.00]	0.00 [0.00;0.00]	0.00 [0.00;1.00]	<0.001