

UNIVERSITY OF STRATHCLYDE
DEPARTMENT OF MATHEMATICS AND STATISTICS

MM201 Linear Algebra and Differential Equations

1 Exercise Solutions: Chapter 1

1. (a) Not defined. (b) Defined: 4×2 (c) Not defined. (d) Not defined.
 (e) Defined: 5×5 (f) Defined: 5×2 (g) Not defined. (h) Defined: 5×2 .

2. (a)

$$4E - 2D = \begin{bmatrix} 24 & 4 & 12 \\ -4 & 4 & 8 \\ 16 & 4 & 12 \end{bmatrix} - \begin{bmatrix} 2 & 10 & 4 \\ -2 & 0 & 2 \\ 6 & 4 & 8 \end{bmatrix} = \begin{bmatrix} 22 & -6 & 8 \\ -2 & 4 & 6 \\ 10 & 0 & 4 \end{bmatrix}$$

- (b) $2B - C$ is not defined.

- (c)

$$AB = \begin{bmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 4 & -1 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} 12 & -3 \\ -4 & 5 \\ 4 & 1 \end{bmatrix}$$

- (d) BA is not defined.

- (e)

$$(3E)D = 3(ED) = 3 \begin{bmatrix} 14 & 36 & 25 \\ 4 & -1 & 7 \\ 12 & 26 & 21 \end{bmatrix} = \begin{bmatrix} 42 & 108 & 75 \\ 12 & -3 & 21 \\ 36 & 78 & 63 \end{bmatrix}$$

- (f)

$$(AB)C = \begin{bmatrix} 12 & -3 \\ -4 & 5 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix} = \begin{bmatrix} 3 & 45 & 9 \\ 11 & -11 & 17 \\ 7 & 17 & 13 \end{bmatrix}$$

- (g)

$$A(BC) = (AB)C \quad (\text{same as (f)})$$

- (h)

$$DA = \begin{bmatrix} 0 & 12 \\ -2 & 1 \\ 11 & 8 \end{bmatrix} \Rightarrow (DA)^T = \begin{bmatrix} 0 & -2 & 11 \\ 12 & 1 & 8 \end{bmatrix}$$

(i)

$$A^T D^T = (DA)^T \quad (\text{same as (h)})$$

(j)

$$4E^T - D = 4 \begin{bmatrix} 6 & -1 & 4 \\ 1 & 1 & 1 \\ 3 & 2 & 3 \end{bmatrix} - \begin{bmatrix} 1 & 5 & 2 \\ -1 & 0 & 1 \\ 3 & 2 & 4 \end{bmatrix} = \begin{bmatrix} 23 & -9 & 14 \\ 5 & 4 & 3 \\ 9 & 6 & 8 \end{bmatrix}$$

$$\Rightarrow \text{tr}(4E^T - D) = 23 + 4 + 8 = 35$$

(k)

$$CC^T = \begin{bmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 4 & 1 \\ 2 & 5 \end{bmatrix} = \begin{bmatrix} 21 & 17 \\ 17 & 35 \end{bmatrix} \Rightarrow \text{tr}(CC^T) = 21 + 35 = 56$$

3. Equations are

$$\begin{array}{rcl} a & - & b & = & 8 \\ & & b & + & c & = & 1 \\ & & & & c & + & 3d & = & 7 \\ 2a & & & & & - & 4d & = & 6 \end{array}$$

so augmented matrix is

$$\left[\begin{array}{cccc|c} 1 & -1 & 0 & 0 & 8 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 3 & 7 \\ 2 & 0 & 0 & -4 & 6 \end{array} \right] \begin{array}{l} \\ \\ r'_4 = r_4 - 2r_1 \end{array} \rightarrow \left[\begin{array}{cccc|c} 1 & -1 & 0 & 0 & 8 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 3 & 7 \\ 0 & 2 & 0 & -4 & -10 \end{array} \right] \begin{array}{l} \\ \\ r'_4 = r_4 - 2r_2 \end{array}$$

$$\rightarrow \left[\begin{array}{cccc|c} 1 & -1 & 0 & 0 & 8 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 3 & 7 \\ 0 & 0 & -2 & -4 & -12 \end{array} \right] \begin{array}{l} \\ \\ r'_4 = r_4 + 2r_3 \end{array} \rightarrow \left[\begin{array}{cccc|c} 1 & -1 & 0 & 0 & 8 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 3 & 7 \\ 0 & 0 & 0 & 2 & 2 \end{array} \right]$$

Back substitute:

$$d = 1, \quad c + 3d = 7 \Rightarrow c = 4, \quad b + c = 1 \Rightarrow b = -3, \quad a - b = 8 \Rightarrow a = 5.$$

4.

$$A^{-1} = \frac{1}{6-5} \begin{bmatrix} 2 & -1 \\ -5 & 3 \end{bmatrix} = \begin{bmatrix} 2 & -1 \\ -5 & 3 \end{bmatrix}$$

$$B^{-1} = \frac{1}{8+12} \begin{bmatrix} 4 & 3 \\ -4 & 2 \end{bmatrix} = \frac{1}{20} \begin{bmatrix} 4 & 3 \\ -4 & 2 \end{bmatrix}$$

$$C^{-1} = \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{3} \end{bmatrix}$$

$$AB = \begin{bmatrix} 10 & -5 \\ 18 & -7 \end{bmatrix} \Rightarrow (AB)^{-1} = \frac{1}{-70 + 90} \begin{bmatrix} -7 & 5 \\ -18 & 10 \end{bmatrix} = \frac{1}{20} \begin{bmatrix} -7 & 5 \\ -18 & 10 \end{bmatrix}$$

$$B^{-1}A^{-1} = \frac{1}{20} \begin{bmatrix} 4 & 3 \\ -4 & 2 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ -5 & 3 \end{bmatrix} = \frac{1}{20} \begin{bmatrix} -7 & 5 \\ -18 & 10 \end{bmatrix}$$

$$ABC = (AB)C = \begin{bmatrix} 10 & -5 \\ 18 & -7 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 20 & -15 \\ 36 & -21 \end{bmatrix} \Rightarrow (ABC)^{-1} = \frac{1}{120} \begin{bmatrix} -21 & 15 \\ -36 & 20 \end{bmatrix}$$

$$C^{-1}B^{-1}A^{-1} = C^{-1}(B^{-1}A^{-1}) = \frac{1}{20} \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{3} \end{bmatrix} \begin{bmatrix} -7 & 5 \\ -18 & 10 \end{bmatrix} = \frac{1}{120} \begin{bmatrix} -21 & 15 \\ -36 & 20 \end{bmatrix}$$

5.

$$(7D)^{-1} = \begin{bmatrix} -3 & 7 \\ 1 & -2 \end{bmatrix} \Rightarrow 7D = \begin{bmatrix} -3 & 7 \\ 1 & -2 \end{bmatrix}^{-1} = \frac{1}{6-7} \begin{bmatrix} -2 & -7 \\ -1 & -3 \end{bmatrix} = \begin{bmatrix} 2 & 7 \\ 1 & 3 \end{bmatrix}$$

so

$$D = \frac{1}{7} \begin{bmatrix} 2 & 7 \\ 1 & 3 \end{bmatrix}.$$

$$6. \quad \text{tr}(0_{n \times n}) = 0, \quad \text{tr}(I_n) = n$$

$$(a) \quad \text{tr}(A + B) = \sum_{i=1}^n (a_{ii} + b_{ii}) = \sum_{i=1}^n a_{ii} + \sum_{i=1}^n b_{ii} = \text{tr}(A) + \text{tr}(B).$$

(b) Write $C = AB$ and $D = BA$. Then $C = [c_{ij}]_{n \times n}$ with $c_{ij} = \sum_{k=1}^n a_{ik}b_{kj}$ and $D = [d_{ij}]_{n \times n}$ with $d_{ij} = \sum_{k=1}^n b_{ik}a_{kj}$.

$$\text{tr}(AB) = \text{tr}(C) = \sum_{i=1}^n c_{ii} = \sum_{i=1}^n \sum_{k=1}^n a_{ik}b_{ki} = \sum_{k=1}^n \sum_{i=1}^n b_{ki}a_{ik} = \sum_{k=1}^n d_{kk} = \text{tr}(D) = \text{tr}(BA).$$

$$(c) \quad \text{tr}(\lambda A) = \sum_{i=1}^n (\lambda a_{ii}) = \lambda \sum_{i=1}^n a_{ii} = \lambda \text{tr}(A).$$

7. (a) Both. (b) Neither (leading 1 in r_2 to the right of leading 1 in r_3).
 (c) Both. (d) Echelon (NOT reduced echelon: nonzero entry above leading 1 in c_2).
 (e) Neither (leading 1 in r_2 in same column as leading 1 in r_1). (f) Both.

8. (a)

$$\begin{array}{lcl}
\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1 & 1 \\ 0 & 1 & 0 & 3 & 1 & 2 \\ 0 & 0 & 1 & 5 & 1 & 3 \end{array} & \begin{array}{l} r'_2 = r_2 - 3r_1 \\ r'_3 = r_3 - 5r_1 \end{array} & \rightarrow \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 & -2 & -1 \\ 0 & 0 & 1 & 0 & -4 & -2 \end{array} \begin{array}{l} r'_1 = r_1 - r_2 \\ r'_3 = r_3 - 2r_2 \end{array} \\
\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1 & 1 \\ -3 & 1 & 0 & 0 & -2 & -1 \\ 1 & -2 & 1 & 0 & 0 & 0 \end{array} & \begin{array}{l} r'_2 = -\frac{1}{2}r_2 \\ r'_3 = -\frac{1}{2}r_2 \end{array} & \rightarrow \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1 & 1 \\ \frac{3}{2} & -\frac{1}{2} & 0 & 0 & 1 & \frac{1}{2} \\ 1 & -2 & 1 & 0 & 0 & 0 \end{array} \\
& & \rightarrow \begin{array}{ccc|ccc} -\frac{1}{2} & \frac{1}{2} & 0 & 1 & 0 & \frac{1}{2} \\ \frac{3}{2} & -\frac{1}{2} & 0 & 0 & 1 & \frac{1}{2} \\ 1 & -2 & 1 & 0 & 0 & 0 \end{array}
\end{array}$$

Reduced echelon matrix with

$$P = \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} & 0 \\ \frac{3}{2} & -\frac{1}{2} & 0 \\ 1 & -2 & 1 \end{bmatrix}.$$

(b)

$$\begin{array}{lcl}
\begin{array}{ccc|cccc} 1 & 0 & 0 & 1 & 3 & 4 & 3 \\ 0 & 1 & 0 & 1 & 2 & 2 & 2 \\ 0 & 0 & 1 & -1 & 6 & 4 & 1 \end{array} & \begin{array}{l} r'_2 = r_2 - r_1 \\ r'_3 = r_3 + r_1 \end{array} & \rightarrow \begin{array}{ccc|cccc} 1 & 0 & 0 & 1 & 3 & 4 & 3 \\ -1 & 1 & 0 & 0 & -1 & -2 & -1 \\ 1 & 0 & 1 & 0 & 9 & 8 & 4 \end{array} \begin{array}{l} r'_1 = r_1 - 4r_3 \\ r'_2 = r_2 - 2r_3 \end{array} \\
\begin{array}{ccc|cccc} 1 & 0 & 0 & 1 & 3 & 4 & 3 \\ -1 & 1 & 0 & 0 & -1 & -2 & -1 \\ -8 & 9 & 1 & 0 & 0 & -10 & -5 \end{array} & \begin{array}{l} r'_2 = -r_2 \\ r'_3 = -\frac{1}{10}r_3 \end{array} & \rightarrow \begin{array}{ccc|cccc} 1 & 0 & 0 & 1 & 3 & 4 & 3 \\ 1 & -1 & 0 & 0 & 1 & 2 & 1 \\ \frac{4}{5} & -\frac{9}{10} & -\frac{1}{10} & 0 & 0 & 1 & \frac{1}{2} \end{array} \\
\begin{array}{ccc|cccc} -\frac{11}{5} & \frac{18}{5} & \frac{2}{5} & 1 & 3 & 0 & 1 \\ -\frac{3}{5} & \frac{4}{5} & \frac{1}{5} & 0 & 1 & 0 & 0 \\ \frac{4}{5} & -\frac{9}{10} & -\frac{1}{10} & 0 & 0 & 1 & \frac{1}{2} \end{array} & \begin{array}{l} r'_1 = r_1 - 3r_2 \\ r'_3 = r_3 - 3r_2 \end{array} & \rightarrow \begin{array}{ccc|cccc} -\frac{2}{5} & \frac{6}{5} & -\frac{1}{5} & 1 & 0 & 0 & 1 \\ -\frac{3}{5} & \frac{4}{5} & \frac{1}{5} & 0 & 1 & 0 & 0 \\ \frac{4}{5} & -\frac{9}{10} & -\frac{1}{10} & 0 & 0 & 1 & \frac{1}{2} \end{array}
\end{array}$$

Reduced echelon form with

$$P = \frac{1}{10} \begin{bmatrix} -4 & 12 & -2 \\ -6 & 8 & 2 \\ 8 & -9 & -1 \end{bmatrix}.$$

(c)

$$\begin{array}{lcl}
\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 & 1 & 2 & 4 \\ 0 & 0 & 1 & 0 & 1 & 3 & 9 \\ 0 & 0 & 0 & 1 & 1 & 4 & 16 \end{array} & \begin{array}{l} r'_2 = r_2 - r_1 \\ r'_3 = r_3 - r_1 \\ r'_4 = r_4 - r_1 \end{array} & \rightarrow \begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 1 & 1 & 1 \\ -1 & 1 & 0 & 0 & 0 & 1 & 3 \\ -1 & 0 & 1 & 0 & 0 & 2 & 8 \\ -1 & 0 & 0 & 1 & 0 & 3 & 15 \end{array} \begin{array}{l} r'_1 = r_1 - r_2 \\ r'_3 = r_3 - 2r_2 \\ r'_4 = r_4 - 3r_2 \end{array}
\end{array}$$

$$\begin{array}{ccc}
\begin{array}{cccc|cccc}
1 & 0 & 0 & 0 & 1 & 1 & 1 \\
-1 & 1 & 0 & 0 & 0 & 1 & 3 \\
1 & -2 & 1 & 0 & 0 & 0 & 2 \\
2 & -3 & 0 & 1 & 0 & 0 & 6
\end{array} & \xrightarrow{r'_4 = r_4 - 3r_3} & \begin{array}{cccc|cccc}
1 & 0 & 0 & 0 & 1 & 1 & 1 \\
-1 & 1 & 0 & 0 & 0 & 1 & 3 \\
1 & -2 & 1 & 0 & 0 & 0 & 2 \\
-1 & 3 & -3 & 1 & 0 & 0 & 0
\end{array}
\end{array} \quad r'_3 = \frac{1}{2}r_3$$

$$\begin{array}{ccc}
\begin{array}{cccc|cccc}
1 & 0 & 0 & 0 & 1 & 1 & 1 \\
-1 & 1 & 0 & 0 & 0 & 1 & 3 \\
\frac{1}{2} & -1 & \frac{1}{2} & 0 & 0 & 0 & 1 \\
-1 & 3 & -3 & 1 & 0 & 0 & 0
\end{array} & \xrightarrow{r'_1 = r_1 - r_3, r'_2 = r_2 - 3r_1} & \begin{array}{cccc|cccc}
\frac{1}{2} & 1 & -\frac{1}{2} & 0 & 1 & 1 & 0 \\
-\frac{5}{2} & 4 & -\frac{3}{2} & 0 & 0 & 1 & 0 \\
\frac{1}{2} & -1 & \frac{1}{2} & 0 & 0 & 0 & 1 \\
-1 & 3 & -3 & 1 & 0 & 0 & 0
\end{array}
\end{array} \quad r'_1 = r_1 - r_2$$

$$\begin{array}{ccc}
\begin{array}{cccc|cccc}
3 & -3 & 1 & 0 & 1 & 0 & 0 \\
-\frac{5}{2} & 4 & -\frac{3}{2} & 0 & 0 & 1 & 0 \\
\frac{1}{2} & -1 & \frac{1}{2} & 0 & 0 & 0 & 1 \\
-1 & 3 & -3 & 1 & 0 & 0 & 0
\end{array} & \xrightarrow{} & \begin{array}{cccc|cccc}
3 & -3 & 1 & 0 & 1 & 0 & 0 \\
-\frac{5}{2} & 4 & -\frac{3}{2} & 0 & 0 & 1 & 0 \\
\frac{1}{2} & -1 & \frac{1}{2} & 0 & 0 & 0 & 1 \\
-1 & 3 & -3 & 1 & 0 & 0 & 0
\end{array}
\end{array}$$

Reduced echelon matrix with

$$P = \frac{1}{2} \begin{bmatrix} 6 & -6 & 2 & 0 \\ -5 & 8 & -3 & 0 \\ 1 & -2 & 1 & 0 \\ -2 & 6 & -6 & 2 \end{bmatrix}.$$

(d)

$$\begin{array}{ccc}
\begin{array}{cccc|cccc}
1 & 0 & 0 & 0 & 1 & 2 & 1 & 2 \\
0 & 1 & 0 & 0 & 2 & 4 & 3 & 4 \\
0 & 0 & 1 & 0 & 1 & 3 & 2 & 3 \\
0 & 0 & 0 & 1 & 0 & 3 & 1 & 3
\end{array} & \xrightarrow{r'_2 = r_2 - 2r_1, r'_3 = r_3 - r_1} & \begin{array}{cccc|cccc}
1 & 0 & 0 & 0 & 1 & 2 & 1 & 2 \\
-2 & 1 & 0 & 0 & 0 & 0 & 1 & 0 \\
-1 & 0 & 1 & 0 & 0 & 1 & 1 & 1 \\
0 & 0 & 0 & 1 & 0 & 3 & 1 & 3
\end{array}
\end{array} \quad r'_2 = r_2, r'_3 = r_3$$

$$\begin{array}{ccc}
\begin{array}{cccc|cccc}
1 & 0 & 0 & 0 & 1 & 2 & 1 & 2 \\
-1 & 0 & 1 & 0 & 0 & 1 & 1 & 1 \\
-2 & 1 & 0 & 0 & 0 & 0 & 1 & 0 \\
0 & 0 & 0 & 1 & 0 & 3 & 1 & 3
\end{array} & \xrightarrow{r'_4 = r_4 - 3r_2} & \begin{array}{cccc|cccc}
1 & 0 & 0 & 0 & 1 & 2 & 1 & 2 \\
-1 & 0 & 1 & 0 & 0 & 1 & 1 & 1 \\
-2 & 1 & 0 & 0 & 0 & 0 & 1 & 0 \\
3 & 0 & -3 & 1 & 0 & 0 & -2 & 0
\end{array}
\end{array} \quad r'_4 = r_4 + 2r_3$$

$$\begin{array}{ccc}
\begin{array}{cccc|cccc}
1 & 0 & 0 & 0 & 1 & 2 & 1 & 2 \\
-1 & 0 & 1 & 0 & 0 & 1 & 1 & 1 \\
-2 & 1 & 0 & 0 & 0 & 0 & 1 & 0 \\
-1 & 2 & -3 & 1 & 0 & 0 & 0 & 0
\end{array} & \xrightarrow{r'_1 = r_1 - r_3, r'_2 = r_2 - r_3} & \begin{array}{cccc|cccc}
3 & -1 & 0 & 0 & 1 & 2 & 0 & 2 \\
1 & -1 & 1 & 0 & 0 & 1 & 0 & 1 \\
-2 & 1 & 0 & 0 & 0 & 0 & 1 & 0 \\
-1 & 2 & -3 & 1 & 0 & 0 & 0 & 0
\end{array}
\end{array} \quad r'_1 = r_1 - 2r_2$$

$$\begin{array}{ccc}
\begin{array}{cccc|cccc}
1 & 1 & -2 & 0 & 1 & 0 & 0 & 0 \\
1 & -1 & 1 & 0 & 0 & 1 & 0 & 1 \\
-2 & 1 & 0 & 0 & 0 & 0 & 1 & 0 \\
-1 & 2 & -3 & 1 & 0 & 0 & 0 & 0
\end{array} & \xrightarrow{} & \begin{array}{cccc|cccc}
1 & 1 & -2 & 0 & 1 & 0 & 0 & 0 \\
1 & -1 & 1 & 0 & 0 & 1 & 0 & 1 \\
-2 & 1 & 0 & 0 & 0 & 0 & 1 & 0 \\
-1 & 2 & -3 & 1 & 0 & 0 & 0 & 0
\end{array}
\end{array}$$

Reduced echelon matrix with

$$P = \begin{bmatrix} 1 & 1 & -2 & 0 \\ 1 & -1 & 1 & 0 \\ -2 & 1 & 0 & 0 \\ -1 & 2 & -3 & 1 \end{bmatrix}.$$

(e)

$$\begin{array}{l} \begin{array}{ccc|ccc} 1 & 0 & 0 & 2 & 2 & 3 & r'_1 = r_2 \\ 0 & 1 & 0 & 1 & -1 & 0 & r'_2 = r_1 \\ 0 & 0 & 1 & -1 & 2 & 1 & \end{array} \rightarrow \begin{array}{ccc|ccc} 1 & 0 & 0 & 2 & 2 & 3 & r'_2 = r_2 - 2r_1 \\ 0 & 1 & 0 & -1 & 2 & 1 & r'_3 = r_3 + r_1 \\ 0 & 0 & 1 & 1 & -4 & 3 & \end{array} \\ \\ \begin{array}{ccc|ccc} 0 & 1 & 0 & 1 & -1 & 0 \\ \rightarrow 1 & -2 & 0 & 0 & 4 & 3 & r'_2 = \frac{1}{4}r_2 \\ 0 & 1 & 1 & 0 & 1 & 1 & \end{array} \rightarrow \begin{array}{ccc|ccc} 0 & 1 & 0 & 1 & -1 & 0 \\ 1 & -2 & 0 & 0 & 4 & 3 & r'_2 = \frac{1}{4}r_2 \\ 0 & 1 & 1 & 0 & 1 & 1 & \end{array} \\ \\ \begin{array}{ccc|ccc} 0 & 1 & 0 & 1 & -1 & 0 \\ \rightarrow \frac{1}{4} & -\frac{1}{2} & 0 & 0 & 1 & \frac{3}{4} \\ 0 & 1 & 1 & 0 & 1 & 1 & r'_3 = r_3 - r_2 \end{array} \rightarrow \begin{array}{ccc|ccc} 0 & 1 & 0 & 1 & -1 & 0 \\ \frac{1}{4} & -\frac{1}{2} & 0 & 0 & 1 & \frac{3}{4} & r'_2 = r_2 - 3r_3 \\ -\frac{1}{4} & \frac{3}{2} & 1 & 0 & 0 & \frac{1}{4} & \end{array} \\ \\ \begin{array}{ccc|ccc} 0 & 1 & 0 & 1 & -1 & 0 & r'_1 = r_1 + r_2 \\ \rightarrow 1 & -5 & -3 & 0 & 1 & 0 \\ -\frac{1}{4} & \frac{3}{2} & 1 & 0 & 0 & \frac{1}{4} & \end{array} \rightarrow \begin{array}{ccc|ccc} 1 & -4 & -3 & 1 & 0 & 0 \\ 1 & -5 & -3 & 0 & 1 & 0 \\ -\frac{1}{4} & \frac{3}{2} & 1 & 0 & 0 & \frac{1}{4} & r'_3 = 4r_3 \end{array} \\ \\ \begin{array}{ccc|ccc} 1 & -4 & -3 & 1 & 0 & 0 \\ \rightarrow 1 & -5 & -3 & 0 & 1 & 0 \\ -1 & 6 & 4 & 0 & 0 & 1 \end{array} \end{array}$$

Reduced echelon matrix with

$$P = \begin{bmatrix} 1 & -4 & -3 \\ 1 & -5 & -3 \\ -1 & 6 & 4 \end{bmatrix}. \quad (\text{Here } P \text{ is the inverse of the given matrix. Why?})$$

9. (a)

$$x_3 = 5; \quad x_2 + 2x_3 = 2 \Rightarrow x_2 = -8; \quad x_1 - 3x_2 + 4x_3 = 7 \Rightarrow x_1 = -37$$

so

$$\mathbf{x} = \begin{bmatrix} -37 \\ -8 \\ 5 \end{bmatrix}.$$

(b) Set $x_4 = \lambda$. Then

$$x_3 + x_4 = 2 \Rightarrow x_3 = 2 - \lambda; \quad x_2 + 4x_3 - 9x_4 = 3 \Rightarrow x_2 = -5 + 13\lambda;$$

$$x_1 + 8x_3 - 5x_4 = 6 \Rightarrow x_1 = -10 + 13\lambda$$

so

$$\mathbf{x} = \begin{bmatrix} -10 \\ -5 \\ 2 \\ 0 \end{bmatrix} + \lambda \begin{bmatrix} 13 \\ 13 \\ -1 \\ 1 \end{bmatrix}.$$

(c) Set $x_5 = \lambda$. Then

$$x_4 + 3x_5 = 9 \Rightarrow x_4 = 9 - 3\lambda; \quad x_3 + x_4 + 6x_5 = 5 \Rightarrow x_3 = -4 - 3\lambda.$$

Now set $x_2 = \mu$ to get

$$x_1 + 7x_2 - 2x_3 - 8x_5 = -3 \Rightarrow x_1 = -11 - 7\mu + 2\lambda,$$

so

$$\mathbf{x} = \begin{bmatrix} -11 \\ 0 \\ -4 \\ 9 \\ 0 \end{bmatrix} + \lambda \begin{bmatrix} 2 \\ 0 \\ -3 \\ -3 \\ 1 \end{bmatrix} + \mu \begin{bmatrix} -7 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}.$$

(d) Inconsistent: no solution.

10. (a)

$$\begin{array}{cccc|cccc} 2 & 1 & 3 & | & 0 & r'_1 = r_3 & 1 & -3 & -2 & | & 0 \\ 3 & -2 & 1 & | & 0 & & \rightarrow & 3 & -2 & 1 & | & 0 & r'_2 = r_2 - 3r_1 \\ 1 & -3 & -2 & | & 0 & r'_3 = r_1 & 2 & 1 & 3 & | & 0 & r'_3 = r_3 - 2r_1 \\ & & & & & & & & & & & \\ & 1 & -3 & -2 & | & 0 & & 1 & -3 & -2 & | & 0 \\ \rightarrow & 0 & 7 & 7 & | & 0 & & \rightarrow & 0 & 7 & 7 & | & 0 \\ & 0 & 7 & 7 & | & 0 & r'_3 = r_3 - r_2 & 0 & 0 & 0 & | & 0 \end{array}$$

Let $x_3 = \lambda$. Then

$$x_2 = -\lambda; \quad x_1 - 3x_2 - 2x_3 = 0 \Rightarrow x_1 = -\lambda$$

so solution is

$$\mathbf{x} = \lambda \begin{bmatrix} -1 \\ -1 \\ 1 \end{bmatrix}.$$

(b)

$$\begin{array}{cccc|cccc} 2 & 3 & 4 & | & 1 & r'_1 = \frac{1}{2}r_1 & 1 & \frac{3}{2} & 2 & | & \frac{1}{2} \\ 5 & 6 & 7 & | & 2 & & \rightarrow & 5 & 6 & 7 & | & 2 & r'_2 = r_2 - 5r_1 \\ 8 & 9 & 10 & | & 4 & & & 8 & 9 & 10 & | & 4 & r'_3 = r_3 - 8r_1 \\ & & & & & & & & & & & \\ & 1 & \frac{3}{2} & 2 & | & \frac{1}{2} & & 1 & \frac{3}{2} & 2 & | & \frac{1}{2} \\ \rightarrow & 0 & -\frac{3}{2} & -3 & | & -\frac{1}{2} & & \rightarrow & 0 & -\frac{3}{2} & -3 & | & -\frac{1}{2} \\ & 0 & -3 & -6 & | & 0 & r'_3 = r_3 - 2r_2 & 0 & 0 & 0 & | & 1 \end{array}$$

Equations are inconsistent, and have no solution.

(c)

$$\begin{array}{ccc}
\begin{array}{cccc|c} 1 & -3 & 5 & -7 & -2 \\ -2 & 4 & -6 & 8 & 2 \\ 1 & 1 & 1 & 1 & 2 \\ 1 & 5 & 2 & 5 & 7 \end{array} & \begin{array}{l} r'_2 = r_2 + 2r_1 \\ r'_3 = r_3 - r_1 \\ r'_4 = r_4 - r_1 \end{array} & \rightarrow \begin{array}{cccc|c} 1 & -3 & 5 & -7 & -2 \\ 0 & -2 & 4 & -6 & -2 \\ 0 & 4 & -4 & 8 & 4 \\ 0 & 8 & -3 & 12 & 9 \end{array} \begin{array}{l} r'_2 = -\frac{1}{2}r_2 \\ \\ \\ \end{array} \\
\\
\begin{array}{cccc|c} 1 & -3 & 5 & -7 & -2 \\ 0 & 1 & -2 & 3 & 1 \\ 0 & 4 & -4 & 8 & 4 \\ 0 & 8 & -3 & 12 & 9 \end{array} & \begin{array}{l} r'_3 = r_3 - 4r_2 \\ r'_4 = r_4 - 8r_2 \end{array} & \rightarrow \begin{array}{cccc|c} 1 & -3 & 5 & -7 & -2 \\ 0 & 1 & -2 & 3 & 1 \\ 0 & 0 & 4 & -4 & 0 \\ 0 & 0 & 13 & -12 & 1 \end{array} \begin{array}{l} r'_3 = \frac{1}{4}r_3 \\ \\ \\ \end{array} \\
\\
\begin{array}{cccc|c} 1 & -3 & 5 & -7 & -2 \\ 0 & 1 & -2 & 3 & 1 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 13 & -12 & 1 \end{array} & \begin{array}{l} r'_4 = r_4 - 13r_3 \end{array} & \rightarrow \begin{array}{cccc|c} 1 & -3 & 5 & -7 & -2 \\ 0 & 1 & -2 & 3 & 1 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 & 1 \end{array}
\end{array}$$

so

$$x_4 = 1; \quad x_3 - x_4 = 0 \Rightarrow x_3 = 1; \quad x_2 - 2x_3 + 3x_4 = 1 \Rightarrow x_2 = 0;$$

$$x_1 - 3x_2 + 5x_3 - 7x_4 = -2 \Rightarrow x_1 = 0$$

and solution is

$$\mathbf{x} = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}.$$

(d)

$$\begin{array}{ccc}
\begin{array}{cccc|c} 3 & 1 & 2 & 4 & 3 \\ 5 & 2 & 3 & 6 & 5 \\ 4 & 1 & 3 & 6 & 4 \\ 5 & 1 & 4 & 8 & 5 \end{array} & \begin{array}{l} r'_1 = \frac{1}{3}r_1 \\ \\ \\ \end{array} & \rightarrow \begin{array}{cccc|c} 1 & \frac{1}{3} & \frac{2}{3} & \frac{4}{3} & 1 \\ 5 & 2 & 3 & 6 & 5 \\ 4 & 1 & 3 & 6 & 4 \\ 5 & 1 & 4 & 8 & 5 \end{array} \begin{array}{l} r'_2 = r_2 - 5r_1 \\ r'_3 = r_3 - 4r_1 \\ r'_4 = r_4 - 5r_1 \end{array} \\
\\
\begin{array}{cccc|c} 1 & \frac{1}{3} & \frac{2}{3} & \frac{4}{3} & 1 \\ 0 & \frac{1}{3} & -\frac{1}{3} & -\frac{2}{3} & 0 \\ 0 & -\frac{1}{3} & \frac{1}{3} & \frac{2}{3} & 0 \\ 0 & -\frac{2}{3} & \frac{2}{3} & \frac{4}{3} & 0 \end{array} & \begin{array}{l} r'_3 = r_3 + r_2 \\ r'_4 = r_4 + 2r_2 \end{array} & \rightarrow \begin{array}{cccc|c} 1 & \frac{1}{3} & \frac{2}{3} & \frac{4}{3} & 1 \\ 0 & \frac{1}{3} & -\frac{1}{3} & -\frac{2}{3} & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \begin{array}{l} r'_2 = 3r_2 \\ \\ \\ \end{array} \\
\\
\begin{array}{cccc|c} 1 & \frac{1}{3} & \frac{2}{3} & \frac{4}{3} & 1 \\ 0 & 1 & -1 & -2 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array}
\end{array}$$

Let $x_3 = \mu$, $x_4 = \lambda$ to get

$$x_2 - x_3 - 2x_4 = 0 \Rightarrow x_2 = \mu + 2\lambda; \quad x_1 + \frac{1}{3}x_2 + \frac{2}{3}x_3 + \frac{4}{3}x_4 = 1 \Rightarrow x_1 = 1 - \mu - 2\lambda$$

so the solution is

$$\mathbf{x} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + \mu \begin{bmatrix} -1 \\ 1 \\ 1 \\ 0 \end{bmatrix} + \lambda \begin{bmatrix} -2 \\ 2 \\ 0 \\ 1 \end{bmatrix}.$$

(e)

$$\begin{array}{ccc|c} 2 & 1 & 3 & 5 & | & 6 \\ 3 & 2 & 4 & 6 & | & 8 \\ -1 & 3 & 2 & 7 & | & -3 \end{array} \begin{array}{l} r'_1 = r_3 \\ r'_3 = r_1 \end{array} \rightarrow \begin{array}{ccc|c} -1 & 3 & 2 & 7 & | & -3 \\ 3 & 2 & 4 & 6 & | & 8 \\ 2 & 1 & 3 & 5 & | & 6 \end{array} \begin{array}{l} r'_2 = r_2 + 3r_1r'_3 = r_3 + 2r_1 \end{array}$$

$$\begin{array}{ccc|c} -1 & 3 & 2 & 7 & | & -3 \\ 0 & 11 & 10 & 27 & | & -1 \\ 0 & 7 & 7 & 19 & | & 0 \end{array} \begin{array}{l} r'_2 = \frac{1}{11}r_2 \end{array} \rightarrow \begin{array}{ccc|c} -1 & 3 & 2 & 7 & | & -3 \\ 0 & 1 & \frac{10}{11} & \frac{27}{11} & | & -\frac{1}{11} \\ 0 & 7 & 7 & 19 & | & 0 \end{array} \begin{array}{l} r'_3 = r_3 - 7r_2 \end{array}$$

$$\begin{array}{ccc|c} -1 & 3 & 2 & 7 & | & -3 \\ 0 & 1 & \frac{10}{11} & \frac{27}{11} & | & -\frac{1}{11} \\ 0 & 0 & \frac{7}{11} & \frac{20}{11} & | & \frac{7}{11} \end{array}$$

Let $x_4 = \lambda$ so

$$\frac{7}{11}x_3 + \frac{20}{11}x_4 = \frac{7}{11} \Rightarrow x_3 = 1 - \frac{20}{7}\lambda; \quad x_2 + \frac{10}{11}x_3 + \frac{27}{11}x_4 = -\frac{1}{11} \Rightarrow x_2 = -1 + \frac{1}{7}\lambda;$$

$$-x_1 + 3x_2 + 2x_3 + 7x_4 = -3 \Rightarrow x_1 = 2 + \frac{12}{7}\lambda$$

so solution is

$$\mathbf{x} = \begin{bmatrix} 2 \\ -1 \\ 1 \\ 0 \end{bmatrix} + \frac{\lambda}{7} \begin{bmatrix} 12 \\ 1 \\ -20 \\ 7 \end{bmatrix}.$$

11. We have $A(\mathbf{x}_1 + \alpha\mathbf{x}_2) = A\mathbf{x}_1 + A(\alpha\mathbf{x}_2) = A\mathbf{x}_1 + \alpha A\mathbf{x}_2 = \mathbf{0} + \mathbf{0} = \mathbf{0}$ so $\mathbf{x}_1 + \alpha\mathbf{x}_2$ is also a solution of the system.

12. One possible system is $x + y = 0$, $z - y = 3$.

13. We assume $|A| \neq 0$ and consider two separate cases. Let us first suppose that $a \neq 0$. Then

$$\begin{array}{ccc|c} a & b \\ c & d \end{array} \begin{array}{l} r'_1 = \frac{1}{a}r_1 \\ r'_2 = r_2 - cr_1 \end{array} \rightarrow \begin{array}{cc|c} 1 & \frac{b}{a} \\ c & d \end{array} \rightarrow \begin{array}{cc|c} 1 & \frac{b}{a} \\ 0 & d - \frac{bc}{a} \end{array} \begin{array}{l} r'_2 = \frac{a}{ad-bc}r_2 \end{array} \\ \rightarrow \begin{array}{cc|c} 1 & \frac{b}{a} \\ 0 & 1 \end{array} \begin{array}{l} r'_1 = r_1 - \frac{b}{a}r_2 \end{array} \rightarrow \begin{array}{cc|c} 1 & 0 \\ 0 & 1 \end{array}.$$

Secondly, if $a = 0$, then $ad - bc = -bc \neq 0 \Rightarrow b \neq 0$ and $c \neq 0$, so we have

$$\begin{array}{cc|c} 0 & b \\ c & d \end{array} \begin{array}{l} r'_1 = r_2 \\ r'_2 = r_1 \end{array} \rightarrow \begin{array}{cc|c} c & d \\ 0 & b \end{array} \begin{array}{l} r'_1 = \frac{1}{c}r_1 \\ r'_2 = \frac{1}{b}r_2 \end{array} \rightarrow \begin{array}{cc|c} 1 & \frac{d}{c} \\ 0 & 1 \end{array} \begin{array}{l} r'_1 = r_1 - \frac{d}{c}r_2 \end{array} \rightarrow \begin{array}{cc|c} 1 & 0 \\ 0 & 1 \end{array}.$$

14. Terms in $\det(A)$ are of the form $\text{sgn}(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} a_{3\sigma(3)} a_{4\sigma(4)}$, where σ is a permutation of $\{1, 2, 3, 4\}$.

$a_{13} a_{24} a_{41} a_{32} = a_{13} a_{24} a_{32} a_{41}$ corresponds to $\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 3 & 4 & 2 & 1 \end{pmatrix}$, which has $\text{sign} = (-1)^5 = -1$.

Hence $-a_{13} a_{24} a_{41} a_{32}$ occurs in $\det(A)$.

$a_{31} a_{42} a_{14} a_{33}$ contains no term $a_{2\sigma(2)}$ and hence it does **not** occur in $\det(A)$.

$a_{12} a_{22} a_{31} a_{44}$ does not correspond to $a_{1\sigma(1)} a_{2\sigma(2)} a_{3\sigma(3)} a_{4\sigma(4)}$ since a permutation is one-to-one: $\sigma(1)$ cannot have same value as $\sigma(2)$. Hence this term is **not** in $\det(A)$.

15. (a) $I_n = [\delta_{ij}]_{n \times n}$, where δ_{ij} is the Kronecker delta. Hence $\det(I_n) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) \delta_{1\sigma(1)} \delta_{2\sigma(2)} \dots \delta_{n\sigma(n)}$.

The only non-zero term occurs when $\sigma(1) = 1, \sigma(2) = 2, \dots, \sigma(n) = n$, and this is the identity permutation, with $\text{sign} = (-1)^0 = 1$.

Hence $\det(I_n) = \delta_{11} \delta_{22} \dots \delta_{nn} = 1$.

$0_{n \times n} = [\theta_{ij}]_{n \times n}$ where $\theta_{ij} = 0 \quad \forall i, j = 1, 2, \dots, n$.

$$\begin{aligned} \Rightarrow \det(0_{n \times n}) &= \sum_{\sigma \in S_n} \text{sgn}(\sigma) \theta_{1\sigma(1)} \theta_{2\sigma(2)} \dots \theta_{n\sigma(n)} \\ &= \sum_{\sigma \in S_n} \text{sgn}(\sigma) 0 \cdot 0 \dots 0 = 0. \end{aligned}$$

- (b) $A = [a_{ij}]_{n \times n}$. Let $a_{lj} = 0$ for $j = 1, 2, \dots, n$ —row l of A is zero.

$$\begin{aligned} \det(A) &= \sum_{\sigma \in S_n} \text{sgn}(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \dots a_{l\sigma(l)} \dots a_{n\sigma(n)} \\ &= 0 \quad \text{since } a_{l\sigma(l)} = 0. \end{aligned}$$

Proof for a **zero column** is similar.

- (c) *Prove by induction.* Result holds for $n = 2$ since

$$\det \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = a_{11} a_{22} - a_{12} a_{21} = \det \begin{bmatrix} a_{11} & a_{21} \\ a_{12} & a_{22} \end{bmatrix}.$$

Assume result to be true for any $n \times n$ matrix.

Consider $A = [a_{ij}]_{(n+1) \times (n+1)}$

$$\begin{aligned} \det(A) &= \sum_{j=1}^{n+1} a_{1j} A_{1j} \quad (\text{expand by row 1}) \\ &= \sum_{j=1}^{n+1} a_{1j} (-1)^{1+j} \det(M_{1j}) \quad (\det(M_{1j}) \text{ is minor of } a_{1j}) \end{aligned}$$

$$\begin{aligned}
&= \sum_{j=1}^{n+1} a_{1j} (-1)^{1+j} \det(M_{j1}^T) \quad (\text{by assumption}) \\
&\quad \{\text{Here } M_{j1}^T \text{ is submatrix of } A^T \text{ obtained by deleting} \\
&\quad \text{element in row } j \text{ and column 1 from } A^T. \\
&= \sum_{j=1}^{n+1} a_{j1}^T A_{j1}^T, \\
&\quad \text{where } a_{j1}^T = a_{1j} = \text{element in row } j \text{ and column 1 of } A^T, \\
&\quad \text{and } A_{j1}^T = \text{cofactor of } a_{j1}^T \text{ in } A^T. \\
&= \det(A^T). \text{ Result follows by induction.}
\end{aligned}$$

(d) *Prove by induction.* Result holds for $n = 2$.

If $A = [a_{ij}]_{2 \times 2}$, then $\det(A) = a_{11} a_{22} - a_{12} a_{21}$. Now interchange rows 1 and 2 to obtain $E_{12} A = \begin{bmatrix} a_{21} & a_{22} \\ a_{11} & a_{12} \end{bmatrix}$ which has determinant $a_{21} a_{12} - a_{11} a_{22} = -\det(A)$.

Assume result is true for any $n \times n$ matrix and consider $A = [a_{ij}]_{(n+1) \times (n+1)}$. Interchanging rows r and s produces $B = E_{rs} A$.

Let M_{ij} denote the $n \times n$ matrix obtained by deleting row i and column j of A and let M_{ij}^B be the analogous $n \times n$ matrix obtained by deleting row i and column j of matrix B . If $i \neq r, s$ then M_{ij}^B is derived from M_{ij} by interchanging rows in M_{ij} corresponding to rows r and s of A .

Now

$$\begin{aligned}
\det(B) &= \sum_{j=1}^{n+1} b_{ij} B_{ij} = \sum_{j=1}^{n+1} a_{ij} B_{ij} \quad (i \neq r, s) \\
&= - \sum_j a_{ij} A_{ij}
\end{aligned}$$

(from assumption, noting that $B_{ij} = (-1)^{i+j} \det(M_{ij}^B) = -(-1)^{i+j} \det(M_{ij}) = -A_{ij}$).

Hence $\det(B) = -\det(A)$. Result follows by induction.

(e) If σ is not the identity permutation, then for at least one $i, i < \sigma(i)$, in which case $a_{i\sigma(i)} = 0$ since A is lower triangular. Hence the only non-zero term in $\det(A)$ is that corresponding to identity perm.

Hence $\det(A) = a_{11} a_{22} a_{33} \dots a_{nn}$.

$$\begin{aligned}
16. \quad \det(C) &= \sum_{\sigma \in S_n} (\operatorname{sgn} \sigma) c_{1\sigma(1)} c_{2\sigma(2)} \cdots c_{n\sigma(n)} \\
&= \sum_{\sigma} (\operatorname{sgn} \sigma) (a_{1\sigma(1)} + b_{1\sigma(1)}) c_{2\sigma(2)} c_{3\sigma(3)} \cdots c_{n\sigma(n)} \\
&= \sum_{\sigma} (\operatorname{sgn} \sigma) a_{1\sigma(1)} c_{2\sigma(2)} \cdots c_{n\sigma(n)} + \sum_{\sigma} (\operatorname{sgn} \sigma) b_{1\sigma(1)} c_{2\sigma(2)} \cdots c_{n\sigma(n)} \\
&= \sum_{\sigma \in S_n} (\operatorname{sgn} \sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{n\sigma(n)} + \sum_{\sigma \in S_n} (\operatorname{sgn} \sigma) b_{1\sigma(1)} b_{2\sigma(2)} \cdots b_{n\sigma(n)} \\
&= \det(A) + \det(B).
\end{aligned}$$

$$\begin{aligned}
17. \quad (a) \quad & \begin{vmatrix} 2 & 1 & 2 & -1 \\ 3 & -1 & 0 & 2 \\ 4 & 2 & 3 & 1 \\ -2 & 1 & 0 & 2 \end{vmatrix} = 2 \begin{vmatrix} -1 & 0 & 2 \\ 2 & 3 & 1 \\ 1 & 0 & 2 \end{vmatrix} - \begin{vmatrix} 3 & 0 & 2 \\ 4 & 3 & 1 \\ -2 & 0 & 2 \end{vmatrix} \\
& \quad + 2 \begin{vmatrix} 3 & -1 & 2 \\ 4 & 2 & 1 \\ -2 & 1 & 2 \end{vmatrix} + \begin{vmatrix} 3 & -1 & 0 \\ 4 & 2 & 3 \\ -2 & 1 & 0 \end{vmatrix} \\
&= 2[-1(6) + 2(-3)] - [3(6) + 2(6)] + 2[3(3) + 10 + 2(8)] + [3(-3) + 6] \\
&= -24 - 30 + 70 - 3 = 13.
\end{aligned}$$

$$\begin{aligned}
(b) \quad \det &= 2 \begin{vmatrix} 3 & -1 & 2 \\ 4 & 2 & 1 \\ -2 & 1 & 2 \end{vmatrix} + 3 \begin{vmatrix} 2 & 1 & -1 \\ 3 & -1 & 2 \\ -2 & 1 & 2 \end{vmatrix} = 70 + 3(-8 - 10 - 1) \\
&= 70 - 57 = 13 \\
&\quad \text{(see above!)}
\end{aligned}$$

$$\begin{aligned}
(c) \quad & \begin{vmatrix} 2 & 1 & 2 & -1 \\ 3 & -1 & 0 & 2 \\ 4 & 2 & 3 & 1 \\ -2 & 1 & 0 & 2 \end{vmatrix} = \begin{vmatrix} 2 & 1 & 2 & -1 \\ 5 & 0 & 2 & 1 \\ 0 & 0 & -1 & 3 \\ -4 & 0 & -2 & 3 \end{vmatrix} = - \begin{vmatrix} 5 & 2 & 1 \\ 0 & -1 & 3 \\ -4 & -2 & 3 \end{vmatrix} \\
& \quad \nearrow \qquad \qquad \qquad \nearrow \\
& \quad \left\{ \begin{array}{l} r_2 := r_2 + r_1 \\ r_3 := r_3 - 2r_1 \\ r_4 := r_4 - r_1 \end{array} \right\} \qquad c_3 := c_3 + 3c_2
\end{aligned}$$

$$= - \begin{vmatrix} 5 & 2 & 7 \\ 0 & -1 & 0 \\ -4 & -2 & -3 \end{vmatrix} = + \begin{vmatrix} 5 & 7 \\ -4 & -3 \end{vmatrix} = -15 + 28 = 13$$

$$\begin{aligned}
18. \quad (a) \quad & \begin{vmatrix} 17 & 25 & 13 \\ 9 & 11 & 8 \\ 19 & -58 & 57 \end{vmatrix} = \begin{vmatrix} 17 & 25 & -38 \\ 9 & 11 & -19 \\ 19 & -58 & 0 \end{vmatrix} = -19 \begin{vmatrix} 17 & 25 & 2 \\ 9 & 11 & 1 \\ 19 & -58 & 0 \end{vmatrix} \\
& \quad c_3 := c_3 - 3c_1 \qquad \qquad \qquad r_1 := r_1 - 2r_2
\end{aligned}$$

$$= -19 \begin{vmatrix} -1 & 3 & 0 \\ 9 & 11 & 1 \\ 19 & -58 & 0 \end{vmatrix} = +19 \begin{vmatrix} -1 & 3 \\ 19 & -58 \end{vmatrix} = 19.$$

$$(b) \begin{vmatrix} 299 & 199 & 99 \\ 597 & 397 & 197 \\ 999 & 599 & 299 \end{vmatrix} = \begin{vmatrix} 100 & 100 & 99 \\ 200 & 200 & 197 \\ 400 & 300 & 299 \end{vmatrix} = \begin{vmatrix} 0 & 100 & 99 \\ 0 & 200 & 197 \\ 100 & 300 & 299 \end{vmatrix}$$

$$\{c_1 := c_1 - c_2; c_2 := c_2 - c_3\} \{c_1 := c_1 - c_2\}$$

$$= 10^4 \begin{vmatrix} 1 & 99 \\ 2 & 197 \end{vmatrix} = 10^4(197 - 198) = -10^4$$

$$(c) \begin{vmatrix} 11 & 3 & -2 & 6 \\ 20 & 0 & 36 & 17 \\ 38 & 9 & 3 & 22 \\ 30 & 6 & 11 & 19 \end{vmatrix} = \begin{vmatrix} 11 & 3 & -2 & 6 \\ 20 & 0 & 36 & 17 \\ 5 & 0 & 9 & 4 \\ 8 & 0 & 15 & 7 \end{vmatrix}$$

$$\{r_3 := r_3 - 3r_1; r_4 := r_4 - 2r_1\}$$

$$= -3 \begin{vmatrix} 20 & 36 & 17 \\ 5 & 9 & 4 \\ 8 & 15 & 7 \end{vmatrix} = -3 \begin{vmatrix} 0 & 0 & 1 \\ 5 & 9 & 4 \\ 8 & 15 & 7 \end{vmatrix} = -3(75 - 72)$$

$$= -9$$

$$r_1 := r_1 - 4r_2$$

$$19. (a) \begin{vmatrix} 1 & 1 & 1 \\ x & y & z \\ x^2 & y^2 & z^2 \end{vmatrix} = \begin{vmatrix} 1 & 0 & 0 \\ x & y-x & z-x \\ x^2 & y^2-x^2 & z^2-x^2 \end{vmatrix} = (y-x)(z-x) \begin{vmatrix} 1 & 1 \\ y+x & z+x \end{vmatrix}$$

$$= (y-x)(z-x)(z-y)$$

$$\{= (x-y)(y-z)(z-x)\}$$

$$(b) \begin{vmatrix} 1 & 1 & 1 \\ x^2 & y^2 & z^2 \\ x^3 & y^3 & z^3 \end{vmatrix} = \begin{vmatrix} 1 & 0 & 0 \\ x^2 & y^2-x^2 & z^2-x^2 \\ x^3 & y^3-x^3 & z^3-x^3 \end{vmatrix} = (y-x)(z-x) \begin{vmatrix} y+x & z+x \\ y^2+xy+x^2 & z^2+xz+x^2 \end{vmatrix}$$

$$r_2 := r_2 - xr_1$$

$$= (y-x)(z-x) \begin{vmatrix} y+x & z+x \\ y^2 & z^2 \end{vmatrix} = (y-x)(z-x) \begin{vmatrix} y-z & z+x \\ y^2-z^2 & z^2 \end{vmatrix}$$

$$c_1 := c_1 - c_2$$

$$= (y-x)(z-x)(y-z) \begin{vmatrix} 1 & z+x \\ y+z & z^2 \end{vmatrix} = (x-y)(y-z)(z-x)(xy + yz + zx)$$

$$(c) \begin{vmatrix} 1 & 1 & 1 \\ x+y & x+z & y+z \\ xy & xz & yz \end{vmatrix} = \begin{vmatrix} 1 & 0 & 0 \\ x+y & z-y & z-x \\ xy & x(z-y) & y(z-x) \end{vmatrix}$$

$$= (z - y)(z - x) \begin{vmatrix} 1 & 1 \\ x & y \end{vmatrix} = (z - y)(z - x)(y - x) = (x - y)(y - z)(z - x)$$

$$(d) \begin{vmatrix} x & a & a & a \\ a & x & a & a \\ a & a & x & a \\ a & a & a & x \end{vmatrix} = \begin{vmatrix} x+3a & a & a & a \\ x+3a & x & a & a \\ x+3a & a & x & a \\ x+3a & a & a & x \end{vmatrix} = (x+3a) \begin{vmatrix} 1 & a & a & a \\ 0 & (x-a) & 0 & 0 \\ 0 & 0 & (x-a) & 0 \\ 0 & 0 & 0 & (x-a) \end{vmatrix}$$

$$\{c_1 := c_1 + c_2 + c_3 + c_4\} \{r_2 := r_2 - r_1; r_3 := r_3 - r_1; r_4 := r_4 - r_1\}$$

$$= (x+3a)(x-a)^3.$$

$$(e) \begin{vmatrix} 1 & 1 & 1 & 1 \\ w & x & y & z \\ w^2 & x^2 & y^2 & z^2 \\ w^3 & x^3 & y^3 & z^3 \end{vmatrix} = \begin{vmatrix} 1 & 0 & 0 & 0 \\ w & x-w & y-w & z-w \\ w^2 & x^2-w^2 & y^2-w^2 & z^2-w^2 \\ w^3 & x^3-w^3 & y^3-w^3 & z^3-w^3 \end{vmatrix}$$

$$= (x-w)(y-w)(z-w) \begin{vmatrix} 1 & 1 & 1 \\ x+w & y+w & z+w \\ x^2+xw+w^2 & y^2+yw+w^2 & z^2+zw+w^2 \end{vmatrix}$$

$$= (x-w)(y-w)(z-w) \begin{vmatrix} 1 & 0 & 0 \\ x+w & y-x & z-x \\ x^2+xw+w^2 & (y-x)(x+y+w) & (z-x)(z+x+w) \end{vmatrix}$$

$$= (x-w)(y-w)(z-w)(y-x)(z-x) \begin{vmatrix} 1 & 1 \\ x+y+w & z+x+w \end{vmatrix}$$

$$= (x-w)(y-w)(z-w)(y-x)(z-x)(z-y)$$

$$20. \begin{vmatrix} 1+x_1 & x_2 & \cdots & x_n \\ x_1 & 1+x_2 & \cdots & x_n \\ \vdots & \vdots & \ddots & \vdots \\ x_1 & x_2 & \cdots & 1+x_n \end{vmatrix} = \begin{vmatrix} 1 + \sum_{i=1}^n x_i & x_2 & \cdots & x_n \\ 1 + \sum_{i=1}^n x_i & 1+x_2 & \cdots & x_n \\ \vdots & \vdots & \ddots & \vdots \\ 1 + \sum_{i=1}^n x_i & x_2 & \cdots & 1+x_n \end{vmatrix}$$

$$= \begin{vmatrix} 1 + \sum_{i=1}^n x_i & x_2 & x_3 & \cdots & x_n \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{vmatrix} = \left(1 + \sum_{i=1}^n x_i\right) \det(I_{n-1})$$

$$= 1 + \sum_{i=1}^n x_i$$

$$21. \quad (a) \quad |A_1| = \begin{vmatrix} b_1 & a_{12} & a_{13} \\ b_2 & a_{22} & a_{23} \\ b_3 & a_{32} & a_{33} \end{vmatrix}, \text{ where } \mathbf{b} \in \mathbb{R}^3.$$

If $A\mathbf{x} = \mathbf{b}$, then

$$\begin{aligned} |A_1| &= \begin{vmatrix} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 & a_{12} & a_{13} \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 & a_{22} & a_{23} \\ a_{31}x_1 + a_{32}x_2 + a_{33}x_3 & a_{32} & a_{33} \end{vmatrix} \\ &\quad c_1 := c_1 - x_2c_2 - x_3c_3 \\ &= \begin{vmatrix} a_{11}x_1 & a_{12} & a_{13} \\ a_{21}x_1 & a_{22} & a_{23} \\ a_{31}x_1 & a_{32} & a_{33} \end{vmatrix} = x_1 |A| \end{aligned}$$

Thus, $x_1 = |A_1|/|A|$ since $|A| \neq 0$.

Similar results for x_2 and x_3 .

Hence, if $A\mathbf{x} = \mathbf{b}$ then $x_1 = |A_1|/|A|$, $x_2 = |A_2|/|A|$, $x_3 = |A_3|/|A|$.

(b) Here,

$$A = \begin{bmatrix} 1 & 1 & -3 \\ -1 & 0 & 2 \\ -3 & 5 & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{b} = \begin{bmatrix} -6 \\ 5 \\ 7 \end{bmatrix}$$

$$|A| = (-10) - (6) - 3(-5) = -1$$

$$|A_1| = \begin{vmatrix} -6 & 1 & -3 \\ 5 & 0 & 2 \\ 7 & 5 & 0 \end{vmatrix} = -1; \quad |A_2| = \begin{vmatrix} 1 & -6 & -3 \\ -1 & 5 & 2 \\ -3 & 7 & 0 \end{vmatrix} = -2$$

$$|A_3| = \begin{vmatrix} 1 & 1 & -6 \\ -1 & 0 & 5 \\ -3 & 5 & 7 \end{vmatrix} = -3 \quad \text{Hence } x_1 = 1, \quad x_2 = 2, \quad x_3 = 3.$$