EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH (CERN)



CERN-EP-20XX-YYY LHCb-PAPER-20XX-YYY May 8, 2016

Template for writing LHCb papers

The LHCb collaboration[†]

Abstract

Guidelines for the preparation of LHCb documents are given. This is a "living" document that should reflect our current practice. It is expected that these guidelines are implemented for papers before they go into the first collaboration wide review. Please contact the Editorial Board chair if you have suggestions for modifications. This is the title page for journal publications (PAPER). For a CONF note or ANA note, switch to the appropriate template by uncommenting the corresponding line in the file main.tex.

Submitted to JHEP / Phys. Rev. D / Phys. Rev. Lett. / Phys. Lett. B / Eur. Phys. J. C / Nucl. Phys. B

© CERN on behalf of the LHCb collaboration, licence CC-BY-4.0.

[†]Authors are listed at the end of this paper.

1 Introduction

This is the template for typesetting LHCb notes and journal papers. It should be used for any document in LHCb [1] that is to be publicly available. The format should be used for uploading to preprint servers and only afterwards should specific typesetting required for journals or conference proceedings be applied. The main Latex file contains several options as described in the Latex comment lines.

It is expected that these guidelines are implemented for papers already before they go into the first collaboration wide review.

This template also contains the guidelines for how publications and conference reports should be written. The symbols defined in lhcb-symbols-def.tex are compatible with LHCb guidelines.

The front page should be adjusted according to what is written. Default versions are available for papers, conference reports and analysis notes. Just comment out what you require in the main.tex file.

This directory contains a file called Makefile. Typing make will apply all Latex and Bibtex commands in the correct order to produce a pdf file of the document. The default Latex compliler is pdflatex, which requires figures to be in pdf format. To change to plain Latex, edit line 9 of Makefile. Typing make clean will remove all temporary files generated by (pdf)latex.

There is also a PRL template, which is called main-prl.tex. You need to have REVTEX 4.1 installed [2] to compile this. Typing make prl produces a PRL-style PDF file. Note that this version is not meant for LHCb-wide circulation, nor for submission to the arXiv. It is just available to have a look-and-feel of the final PRL version. Typing make count will count the words in the main body.

2 General principles

The main goal is for a paper to be clear. It should be as brief as possible, without sacrificing clarity. For all public documents, special consideration should be given to the fact that the reader will be less familiar with LHCb than the author.

Here follow a list of general principles that should be adhered to:

- 1. Choices that are made concerning layout and typography should be consistently applied throughout the document.
- 2. Standard English should be used (British rather than American) for LHCb notes and preprints. Examples: colour, flavour, centre, metre, modelled and aluminium. Words ending on -ise or -isation (polarise, hadronisation) can be written with -ize or -ization ending. The punctuation normally follows the closing quote mark of quoted text, rather than being included before the closing quote. Footnotes come after punctuation. Papers to be submitted to an American journal can be written in American English instead. Under no circumstance should the two be mixed.

- 3. Use of jargon should be avoided where possible. "Systematics" are "systematic uncertainties", "L0" is "hardware trigger", "penguin" diagrams are best introduced with an expression like "electroweak loop (penguin) diagrams".
- 4. Avoid using quantities that are internal jargon and/or are impossible to reproduce without the full simulation: instead of 'It is required that $\chi^2_{\rm vtx} < 3$ ', say 'A good quality vertex is required'; instead of 'It is required that $\chi^2_{\rm IP} > 16$ ', say 'The track is inconsistent with originating from a PV'; instead of 'A DLL greater than 20 is required' say 'Tracks are required to be identified as kaons'.
- 5. Latex should be used for typesetting. Line numbering should be switched on for drafts that are circulated for comments.
- 6. The abstract should be concise, and not include citations or numbered equations, and should give the key results from the paper.
- 7. Apart from descriptions of the detector, the trigger and the simulation, the text should not be cut-and-pasted from other sources that have previously been published.
- 8. References should usually be made only to publicly accessible documents. References to LHCb conference reports and public notes should be avoided in journal publications, instead including the relevant material in the paper itself.
- 9. The use of tenses should be consistent. It is recommended to mainly stay in the present tense, for the abstract, the description of the analysis, *etc.*; the past tense is then used where necessary, for example when describing the data taking conditions.
- 10. It is recommended to use the passive rather than active voice: "the mass is measured", rather than "we measure the mass". Limited use of the active voice is acceptable, in situations where re-writing in the passive form would be cumbersome, such as for the acknowledgements. Some leeway is permitted to accommodate different author's styles, but "we" should not appear excessively in the abstract or the first lines of introduction or conclusion.
- 11. A sentence should not start with a variable, a particle or an acronym. A title or caption should not start with an article.
- 12. Incorrect punctuation around conjunctive adverbs and the use of dangling modifiers are the two most common mistakes of English grammar in LHCb draft papers. If in doubt, read the wikipedia articles on conjunctive adverb and dangling modifier.

3 Layout

1. Unnecessary blank space should be avoided, between paragraphs or around figures and tables.

- 2. Figure and table captions should be concise and use a somewhat smaller typeface than the main text, to help distinguish them. This is achieved by inserting \small at the beginning of the caption. (NB with the latest version of the file premable.tex this is automatic) Figure captions go below the figure, table captions go above the table.
- 3. Captions and footnotes should be punctuated correctly, like normal text. The use of too many footnotes should be avoided: typically they are used for giving commercial details of companies, or standard items like coordinate system definition or the implicit inclusion of charge-conjugate processes.^{1,2}
- 4. Tables should be formatted in a simple fashion, without excessive use of horizontal and vertical lines. See Table 1 for an example.
 - 5. Figures and tables should normally be placed so that they appear on the same page as their first reference, but at the top or bottom of the page; if this is not possible, they should come as soon as possible afterwards. They must all be referred to from the text.
- 6. If one or more equations are referenced, all equations should be numbered using parentheses as shown in Eq. 1,

$$V_{us}V_{ub}^* + V_{cs}V_{cb}^* + V_{ts}V_{tb}^* = 0. (1)$$

7. Displayed results like

84

85

86

87

90

92

$$\mathcal{B}(B_s^0 \to \mu^+ \mu^-) < 1.5 \times 10^{-8} \text{ at } 95\% \text{ CL}$$

- should in general not be numbered.
 - 8. Numbered equations should be avoided in captions and footnotes.
- 93. Displayed equations are part of the normal grammar of the text. This means that
 the equation should end in full stop or comma if required when reading aloud. The
 line after the equation should only be indented if it starts a new paragraph.
- 10. Sub-sectioning should not be excessive: sections with more than three levels of index (1.1.1) should be avoided.
- 11. Acronyms should be defined the first time they are used, e.g. "Monte Carlo (MC) events containing a doubly Cabibbo-suppressed (DCS) decay have been generated."

 The abbreviated words should not be capitalised if it is not naturally written with

¹If placed at the end of a sentence, the footnote symbol normally follows the punctuation; if placed in the middle of an equation, take care to avoid any possible confusion with an index.

²The standard footnote reads: "The inclusion of charge-conjugate processes is implied throughout." This may need to be modified, for example with "except in the discussion of asymmetries."

Table 1: Background-to-signal ratio estimated in a $\pm 50 \,\text{MeV}/c^2$ mass window for the prompt and long-lived backgrounds, and the minimum bias rate.

Channel	$B_{\rm pr}/S$	$B_{\rm LL}/S$	MB rate
$B_s^0 \to J/\psi \phi$	1.6 ± 0.6	0.51 ± 0.08	$\sim 0.3~\mathrm{Hz}$
$B^0 \! o J \! / \! \psi K^{*0}$	5.2 ± 0.3	1.53 ± 0.08	$\sim 8.1~\mathrm{Hz}$
$B^+ \rightarrow J/\psi K^{*+}$	1.6 ± 0.2	0.29 ± 0.06	$\sim 1.4~\mathrm{Hz}$

capitals, e.g. quantum chromodynamics (QCD), impact parameter (IP), boosted decision tree (BDT). Avoid acronyms if they are used three times or less. A sentence should never start with an acronym and its better to avoid it as the last word of a sentence as well.

4 Typography

101

102

103

104

105

109

119

120

121

122

123

124

125

The use of the Latex typesetting symbols defined in the file lhcb-symbols-def.tex and detailed in the appendices of this document is strongly encouraged as it will make it much easier to follow the recommendation set out below.

- 1. LHCb is typeset with a normal (roman) lowercase b.
- 2. Titles are in bold face, and usually only the first word is capitalised.
- 3. Mathematical symbols and particle names should also be typeset in bold when appearing in titles.
- 4. Units are in roman type, except for constants such as c or h that are italic: GeV, GeV/c^2 . The unit should be separated from the value with a thin space ("\,"), and they should not be broken over two lines. Correct spacing is automatic when using predefined units inside math mode: \$3.0\gev\$ \rightarrow 3.0 GeV. Spacing goes wrong when using predefined units outside math mode AND forcing extra space: 3.0\,\gev\$ \rightarrow 3.0 GeV or worse: 3.0\\gev\$ \rightarrow 3.0 GeV.
 - 5. If factors of c are kept, they should be used both for masses and momenta, e.g. $p = 5.2 \,\text{GeV}/c$ (or $\,\text{GeV}c^{-1}$), $m = 3.1 \,\text{GeV}/c^2$ (or $\,\text{GeV}c^{-2}$). If they are dropped this should be done consistently throughout, and a note should be added at the first instance to indicate that units are taken with c = 1.
 - 6. The % sign should not be separated from the number that precedes it: 5%, not 5 %. A thin space is also acceptable: 5%, but should be applied consistently throughout the paper.

- 7. Ranges should be formatted consistently. The recommendend form is to use a dash with no spacing around it: 7–8 GeV, obtained as 7––8\gev.
- 8. Italic is preferred for particle names (although roman is acceptable, if applied 128 consistently throughout). Particle Data Group conventions should generally be 129 followed: B^0 (no need for a "d" subscript), $B_s^0 \to J/\psi \phi$, \overline{B}_s^0 , (note the long bar, 130 obtained with \overline, in contrast to the discouraged short \bar{B} resulting in 131 \bar{B}), $K_{\rm s}^0$ (note the uppercase roman type "S"). This is most easily achieved by using 132 the predefined symbols described in Appendix C. Unless there is a good reason not 133 to, the charge of a particle should be specified if there is any possible ambiguity 134 $(m(K^+K^-))$ instead of m(KK), which could refer to neutral kaons). 135
- 9. Decay chains can be written in several ways, depending on the complexity and the number of times it occurs. Unless there is a good reason not to, usage of a particular type should be consistent within the paper. Examples are: $D_s^+ \to \phi \pi^+$, with $\phi \to K^+K^-$; $D_s^+ \to \phi \pi^+$ ($\phi \to K^+K^-$); $D_s^+ \to \phi (\to K^+K^-) \pi^+$; or $D_s^+ \to [K^+K^-]_{\phi} \pi^+$.
- 10. Variables are usually italic: V is a voltage (variable), while 1 V is a volt (unit). Also in combined expressions: Q-value, z-scale, R-parity etc.
- 11. Subscripts and superscripts are roman type when they refer to a word (such as T for transverse) and italic when they refer to a variable (such as t for time): $p_{\rm T}$, Δm_s , $t_{\rm rec}$.
- 12. Standard function names are in roman type: e.g. cos, sin and exp.
- 13. Figure, Section, Equation, Chapter and Reference should be abbreviated as Fig.,
 Sect. (or alternatively Sec.), Eq., Chap. and Ref. respectively, when they refer to a
 particular (numbered) item, except when they start a sentence. Table and Appendix
 are not abbreviated. The plural form of abbreviation keeps the point after the s,
 e.g. Figs. 1 and 2. Equations may be referred to either with ("Eq. (1)") or without
 ("Eq. 1") parentheses, but it should be consistent within the paper.
- 14. Common abbreviations derived from Latin such as "for example" (e.g.), "in other words" (i.e.), "and so forth" (etc.), "and others" (et al.), "versus" (vs.) can be used, with the typography shown, but not excessively; other more esoteric abbreviations should be avoided.
- 15. Units, material and particle names are usually lower case if spelled out, but often capitalised if abbreviated: amps (A), gauss (G), lead (Pb), silicon (Si), kaon (K), but proton (p).
- 16. Counting numbers are usually written in words if they start a sentence or if they have a value of ten or below in descriptive text (*i.e.* not including figure numbers such as "Fig. 4", or values followed by a unit such as "4 cm"). The word 'unity' can be useful to express the special meaning of the number one in expressions such as:

 "The BDT output takes values between zero and unity".

- 17. Numbers larger than 9999 have a comma (or a small space, but not both) between the multiples of thousand: e.g. 10,000 or 12,345,678. The decimal point is indicated with a point rather than a comma: e.g. 3.141.
- 18. We apply the rounding rules of the PDG [3]. The basic rule states that if the three 167 highest order digits of the uncertainty lie between 100 and 354, we round to two 168 significant digits. If they lie between 355 and 949, we round to one significant digit. 169 Finally, if they lie between 950 and 999, we round up and keep two significant digits. 170 In all cases, the central value is given with a precision that matches that of the 171 uncertainty. So, for example, the result 0.827 ± 0.119 should be written as 0.83 ± 0.12 , 172 0.827 ± 0.367 should turn into 0.8 ± 0.4 , and 14.674 ± 0.964 becomes 14.7 ± 1.0 . When 173 writing numbers with uncertainty components from different sources, i.e. statistical 174 and systematic uncertainties, the rule applies to the uncertainty with the best 175 precision, so $0.827 \pm 0.367 \, (\text{stat}) \pm 0.179 \, (\text{syst})$ goes to $0.83 \pm 0.37 \, (\text{stat}) \pm 0.18 \, (\text{syst})$ 176 and $8.943 \pm 0.123 \, (\text{stat}) \pm 0.995 \, (\text{syst})$ goes to $8.94 \pm 0.12 \, (\text{stat}) \pm 1.00 \, (\text{syst})$. 177
- 19. When rounding numbers, it should be avoided to pad with zeroes at the end. So 51237 ± 4561 should be rounded as $(5.12 \pm 0.46) \times 10^4$ and not 51200 ± 4600 .
- when rounding numbers in a table, some variation of the rounding rules above may be required to achieve uniformity.

182

183

184

185

186

187

188

189

190

191

192

193

- 21. Hyphenation should be used where necessary to avoid ambiguity, but not excessively. For example: "big-toothed fish" (to indicate that big refers to the teeth, not to the fish), but "big white fish". A compound modifier often requires hyphenation (CP-violating observables, b-hadron decays, final-state radiation, second-order polynomial), even if the same combination in an adjective-noun combination does not (direct CP violation, heavy b hadrons, charmless final state). Adverb-adjective combinations are not hyphenated if the adverb ends with 'ly': oppositely charged pions, kinematically similar decay. Cross-section, cross-check, and two-dimensional are hyphenated. Semileptonic, pseudorapidity, pseudoexperiment, multivariate, multidimensional, reweighted, preselection, nonresonant, nonzero, nonparametric, nonrelativistic, misreconstructed and misidentified are single words and should not be hyphenated.
- 194 22. Minus signs should be in a proper font (-1), not just hyphens (-1); this applies to figure labels as well as the body of the text. In Latex, use math mode (between \$\$'s) or make a dash ("--"). In ROOT, use #font[122]{-} to get a normal-sized minus sign.
- 23. Inverted commas (around a title, for example) should be a matching set of left- and right-handed pairs: "Title". The use of these should be avoided where possible.
- 24. Single symbols are preferred for variables in equations, $e.g. \mathcal{B}$ rather than BF for a branching fraction.

- 25. Parentheses are not usually required around a value and its uncertainty, before the unit, unless there is possible ambiguity: so $\Delta m_s = 20 \pm 2 \,\mathrm{ps^{-1}}$ does not need parentheses, whereas $f_d = (40 \pm 4)\%$ or $x = (1.7 \pm 0.3) \times 10^{-6}$ does. The unit does not need to be repeated in expressions like $1.2 < E < 2.4 \,\mathrm{GeV}$.
- 26. The same number of decimal places should be given for all values in any one expression (e.g. $5.20 < m_B < 5.34 \,\text{GeV}/c^2$).
- 27. Apostrophes are best avoided for abbreviations: if the abbreviated term is capitalised or otherwise easily identified then the plural can simply add an s, otherwise it is best to rephrase: e.g. HPDs, π^0 s, pions, rather than HPD's, π^0 's, π s.
- 28. Particle labels, decay descriptors and mathematical functions are not nouns, and need often to be followed by a noun. Thus "background from $B^0 \to \pi^+\pi^-$ decays" instead of "background from $B^0 \to \pi^+\pi^-$ ", and "the width of the Gaussian function" instead of "the width of the Gaussian".
- 29. In equations with multidimensional integrations or differentiations, the differential terms should be separated by a thin space. Thus $\int f(x,y)dx\,dy$ instead $\int f(x,y)dxdy$ and $\frac{d^2\Gamma}{dx\,dQ^2}$ instead of $\frac{d^2\Gamma}{dxdQ^2}$. The d's are allowed in either roman or italic font, but should be consistent throughout the paper.

5 Detector and simulation

219

220

221

222

223

225

226

227

228

229

230

The paragraph below can be used for the detector description. Modifications may be required in specific papers to fit within page limits, to enhance particular detector elements or to introduce acronyms used later in the text. For journals where strict word counts are applied (for example, PRL), and space is at a premium, it may be sufficient to write, as a minimum: "The LHCb detector is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, described in detail in Refs. [1,4]". A slightly longer version could specify the most relevant sub-detectors, e.g "The LHCb detector [1,4] is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector elements that are particularly relevant to this analysis are: a silicon-strip vertex detector surrounding the pp interaction region that allows c- and b-hadrons to be identified from their characteristically long flight distance; a tracking system that provides a measurement of momentum, p, of charged particles; and two ring-imaging Cherenkov detectors that are able to discriminate between different species of charged hadrons."

In the following paragraph, references to the individual detector performance papers are marked with a * and should only be included if the analysis relies on numbers or methods described in the specific papers. Otherwise, a reference to the overall detector performance paper \cite{LHCb-DP-2014-002} will suffice. Note also that the text

defines the acronyms for primary vertex, PV, and impact parameter, IP. Remove either of those in case it is not used later on.

The LHCb detector [1, 4] is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector includes a high-precision tracking system consisting of a siliconstrip vertex detector surrounding the pp interaction region [5]*, a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes [6]* placed downstream of the magnet. The tracking system provides a measurement of momentum, p, of charged particles with a relative uncertainty that varies from 0.5% at low momentum to 1.0% at $200 \,\mathrm{GeV}/c$. The minimum distance of a track to a primary vertex (PV), the impact parameter (IP), is measured with a resolution of $(15 + 29/p_T) \mu m$, where p_T is the component of the momentum transverse to the beam, in GeV/c. Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov detectors [7]*. Photons, electrons and hadrons are identified by a calorimeter system consisting of scintillatingpad and preshower detectors, an electromagnetic calorimeter and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers [8]*. The online event selection is performed by a trigger [9]*, which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction.

A more detailed description of the 'full event reconstruction' could be:

• The trigger [9]* consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, in which all charged particles with $p_{\rm T} > 500\,(300)$ MeV are reconstructed for 2011 (2012) data. For triggers that require neutral particles, energy deposits in the electromagnetic calorimeter are analysed to reconstruct π^0 and γ candidates.

The trigger description has to be specific for the analysis in question. In general, you should not attempt to describe the full trigger system. Below are a few variations that inspiration can be taken from. First from a hadronic analysis, and second from an analysis with muons in the final state. A detailed description of the trigger conditions for Run 1 is available in Ref. [10].

• At the hardware trigger stage, events are required to have a muon with high $p_{\rm T}$ or a hadron, photon or electron with high transverse energy in the calorimeters. For hadrons, the transverse energy threshold is 3.5 GeV. The software trigger requires a two-, three- or four-track secondary vertex with a significant displacement from the primary pp interaction vertices. At least one charged particle must have a transverse momentum $p_{\rm T} > 1.7 \, {\rm GeV}/c$ and be inconsistent with originating from a PV. A multivariate algorithm [11] is used for the identification of secondary vertices consistent with the decay of a b hadron.

• Candidate events are first required to pass the hardware trigger, which selects muons with a transverse momentum $p_{\rm T} > 1.48\,{\rm GeV}/c$ in the 7 TeV data or $p_{\rm T} > 1.76\,{\rm GeV}/c$ in the 8 TeV data. In the subsequent software trigger, at least one of the final-state particles is required to have both $p_{\rm T} > 0.8\,{\rm GeV}/c$ and impact parameter larger than 100 μ m with respect to all of the primary pp interaction vertices (PVs) in the event. Finally, the tracks of two or more of the final-state particles are required to form a vertex that is significantly displaced from the PVs.

An example to describe the use of both TOS and TIS events:

• In the offline selection, trigger signals are associated with reconstructed particles. Selection requirements can therefore be made on the trigger selection itself and on whether the decision was due to the signal candidate, other particles produced in the pp collision, or a combination of both.

A good example of a description of long and downstream K_s^0 is given in Ref. [12]:

• Decays of $K_s^0 \to \pi^+\pi^-$ are reconstructed in two different categories: the first involving K_s^0 mesons that decay early enough for the daughter pions to be reconstructed in the vertex detector; and the second containing K_s^0 that decay later such that track segments of the pions cannot be formed in the vertex detector. These categories are referred to as *long* and *downstream*, respectively. The long category has better mass, momentum and vertex resolution than the downstream category.

The description of our software stack for simulation is often causing trouble. The following paragraph can act as inspiration but with variations according to the level of detail required and if mentioning of *e.g.* Photos is required.

• In the simulation, pp collisions are generated using PYTHIA [13] (In case only PYTHIA 6 is used, remove *Sjostrand:2007gs from this citation; if only PYTHIA 8 is used, then reverse the order of the papers in the citation.) with a specific LHCb configuration [14]. Decays of hadronic particles are described by EVTGEN [15], in which final-state radiation is generated using PHOTOS [16]. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [17] as described in Ref. [18].

Many analyses depend on boosted decision trees. It is inappropriate to use TMVA as the reference as that is merely an implementation of the BDT algorithm. Rather it is suggested to write

In this paper we use a boosted decision tree (BDT) [19, 20] to separate signal from background.

When describing the integrated luminosity of the data set, do not use expressions like " $1.0 \,\mathrm{fb^{-1}}$ of data", but *e.g.* "data corresponding to an integrated luminosity of $1.0 \,\mathrm{fb^{-1}}$ ", or "data obtained from $3 \,\mathrm{fb^{-1}}$ of integrated luminosity".

For analyses where the periodical reversal of the magnetic field is crucial, e.g. in measurements of direct CP violation, the following description can be used as an example




Figure 1: Example plots for (a) data and (b) simulation using the LHCb style from the URANIA package RootTools/LHCbStyle. The signal data is shown as points with the signal component as yellow (light shaded), background 1 as green (medium shaded) and background 2 as blue (dark shaded).

phrase: "The polarity of the dipole magnet is reversed periodically throughout datataking. The configuration with the magnetic field vertically upwards, MagUp (downwards, MagDown), bends positively (negatively) charged particles in the horizontal plane towards the centre of the LHC." Only use the MagUp, MagDown symbols if they are used extensively in tables or figures.

6 Figures

A standard LHCb style file for use in production of figures in ROOT is in the URANIA package RootTools/LHCbStyle or directly in SVN at svn+ssh://svn.cern.ch/reps/lhcb/Urania/trunk/RootTools/LHCbStyle. It is not mandatory to use this style, but it makes it easier to follow the recommendations below. Figure 1 shows an example of how to include an eps or pdf figure with the

Figure 1 shows an example of how to include an eps or pdf figure with the \includegraphics command (eps figures will not work with pdflatex). Note that if the graphics sits in figs/myfig.pdf, you can just write \includegraphics{myfig} as the figs subdirectory is searched automatically and the extension .pdf (.eps) is automatically added for pdflatex (latex).

- 1. Figures should be legible at the size they will appear in the publication, with suitable line width. Their axes should be labelled, and have suitable units (e.g. avoid a mass plot with labels in MeV/c^2 if the region of interest covers a few GeV/c^2 and all the numbers then run together). Spurious background shading and boxes around text should be avoided.
- 2. For the y-axis, "Entries" or "Candidates" is approriate in case no background subtraction has been applied. Otherwise "Yield" or "Decays" may be more appropriate. If the unit on the y-axis corresponds to the yield per bin, indicate so, for example "Entries / ($5 \text{ MeV}/c^2$)" or "Entries per $5 \text{ MeV}/c^2$ ".

- 341 3. Fit curves should not obscure the data points, and data points are best (re)drawn over the fit curves.
- 4. Colour may be used in figures, but the distinction between differently coloured areas or lines should be clear also when the document is printed in black and white, for example through differently dashed lines. The LHCb style mentioned above implements a colour scheme that works well but individual adjustments might be required.
- 5. Using different hatching styles helps to disinguished filled areas, also in black and white prints. Hatching styles 3001-3025 should be avoided since they behave unpredictably under zooming and scaling. Good styles for "falling hatched" and "rising hatched" are 3345 and 3354.
- 6. Figures with more than one part should have the parts labelled (a), (b) etc., with a corresponding description in the caption; alternatively they should be clearly referred to by their position, e.g. Fig. 1 (left). In the caption, the labels (a), (b) etc. should precede their description. When referencing specific sub-figures, use "see Fig. 1(a)" or "see Figs. 2(b)-(e)".
 - 7. All figures containing LHCb data should have LHCb written on them. For preliminary results, that should be replaced by "LHCb preliminary". Figures that only have simulated data should display "LHCb simulation". Figures that do not depend on LHCb-specific software (e.g. only on PYTHIA) should not have any label.

7 References

357

358

359

360

361

367

368

369

References should be made using BibTEX [21]. A special style LHCb.bst has been created to achieve a uniform style. Independent of the journal the paper is submitted to, the preprint should be created using this style. Where arXiv numbers exist, these should be added even for published articles. In the PDF file, hyperlinks will be created to both the arXiv and the published version.

- 1. Citations are marked using square brackets, and the corresponding references should be typeset using BibTEX and the official LHCb BibTEX style. An example is in Ref. [13].
- 2. For references with four or less authors all of the authors' names are listed [22], otherwise the first author is given, followed by *et al.*. The LHCb BibTEX style will take care of this.
- 373 3. The order of references should be sequential when reading the document. This is automatic when using BibT_EX.

4. The titles of papers should in general be included. To remove them, change $\scalebox{setboolean{articletitles}{false}}$ to true at the top of this template. Note that the titles in LHCb-PAPER. bib are in plain LaTex, in order to correspond to the actual title on the arXiv record. Some differences in style can thus be noticed with respect to the main text, for example particle names that use capital Greek letters are not slanted in the reference titles (Λ vs Λ)

- 5. Whenever possible, use references from the supplied files main.bib, LHCb-PAPER.bib, LHCb-CONF.bib, and LHCB-DP.bib. These are kept up-to-date by the EB. If you see a mistake, do not edit these files, but let the EB know. This way, for every update of the paper, you save yourself the work of updating the references. Instead, you can just copy or check in the latest versions of the .bib files from the repository.
- 6. For those references not provided by the EB, the best is to copy the BibTEX entry directly from Inspire. Often these need to be edited to get the correct title, author names and formatting. For authors with multiple initials, add a space between them (change R.G.C. to R. G. C.), otherwise only the first initial will be taken. Also, make sure to eliminate unnecessary capitalisation. Apart from that, the title should be respected as much as possible (e.g. do not change particle names to PDG convention nor introduce/remove factors of c). Check that both the arXiv and the journal index are clickable and point to the right article.
- 7. The mciteplus [23] package is used to enable multiple references to show up as a single item in the reference list. As an example \cite{Mohapatra:1979ia,*Pascoli:2007qh} where the * indicates that the reference should be merged with the previous one. The result of this can be seen in Ref. [24]. Be aware that the mciteplus package should be included as the very last item before the \begin{document} to work correctly.
- 8. It should be avoided to make references to public notes and conference reports in public documents. Exceptions can be discussed on a case-by-case basis with the review committee for the analysis. In internal reports they are of course welcome and can be referenced as seen in Ref. [25] using the lhcbreport category. For conference reports, omit the author field completely in the BibTFX record.
 - 9. To get the typesetting and hyperlinks correct for LHCb reports, the category lhcbreport should be used in the BibTEX file. See Refs. [26] for some examples. It can be used for LHCb documents in the series CONF, PAPER, PROC, THESIS, LHCC, TDR and internal LHCb reports. Papers sent for publication, but not published yet, should be referred with their arXiv number, so the PAPER category should only be used in the rare case of a forward reference to a paper.
- 10. Proceedings can be used for references to items such as the LHCb simulation [18], where we do not yet have a published paper.
- There is a set of standard references to be used in LHCb that are listed in Appendix A.

₁₄ 8 Inclusion of supplementary material

Three types of supplementary material should be distinguised:

- A regular appendix: lengthy equations or long tables are sometimes better put in an appendix in order not to interrupt the main flow of a paper. Appendices will appear in the final paper, on arXiv and on the cds record and should be considered integral part of a paper, and are thus to be reviewed like the rest of the paper. An example of an LHCb paper with an appendix is Ref. [27].
- Supplementary material for cds: plots or tables that would make the paper exceed the page limit or are not appropriate to include in the paper itself, but are desireable to be shown in public should be added to the paper drafts in an appendix, and removed from the paper before submitting to arXiv or the journal. See Appendix D for further instructions. Examples are: comparison plots of the new result with older results, plots that illustrate cross-checks. An example of an LHCb paper with supplementary material for cds is Ref. [28]. Supplementary material for cds cannot be referenced in the paper. Supplementary material should be included in the draft paper to be reviewed by the collaboration.
- Supplementary material for the paper. This is usually called "supplemental material", which distinguishes it from supplementary material for cds only. Most journals allow to submit files along with the paper that will not be part of the text of the article, but will be stored on the journal server. Examples are plain text files with numerical data corresponding to the plots in the paper. The supplemental material should be cited in the paper by including a reference which should say "See supplemental material at [link] for [give brief description of material]." The journal will insert a specific link for [link]. The arXiv version will usually include the supplemental material as part of the paper and so should not contain the words "at [link]". Supplemental material should be included in the draft paper to be reviewed by the collaboration. An example of an LHCb paper with supplemental material is Ref. [29]

441 Acknowledgements

The text below are the acknowledgements as approved by the collaboration board. Extend-ing the acknowledgements to include individuals from outside the collaboration who have contributed to the analysis should be approved by the EB. The extra acknowledgements are normally placed before the standard acknowledgements, unless it matches better with the text of the standard acknowledgements to put them elsewhere. They should be included in the draft for the first circulation. Except in exceptional circumstances, to be approved by the EB chair, authors of the paper should not be named in extended acknowledgements. We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies:

CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 (France); 452 BMBF, DFG and MPG (Germany); INFN (Italy); FOM and NWO (The Netherlands); 453 MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FANO (Russia); MinECo 454 (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF 455 (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 456 (France), KIT and DESY (Germany), INFN (Italy), SURF (The Netherlands), PIC (Spain), 457 GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-458 HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to 459 the communities behind the multiple open source software packages on which we depend. 460 Individual groups or members have received support from AvH Foundation (Germany), 461 EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union), Conseil Général 462 de Haute-Savoie, Labex ENIGMASS and OCEVU, Région Auvergne (France), RFBR and 463 Yandex LLC (Russia), GVA, XuntaGal and GENCAT (Spain), Herchel Smith Fund, The 464 Royal Society, Royal Commission for the Exhibition of 1851 and the Leverhulme Trust 465 (United Kingdom).

467 Appendices

A Standard References

Below is a list of common references, as well as a list of all LHCb publications. As they are already in prepared bib files, they can be used as simply as \cite{Alves:2008zz} to get the LHCb detector paper. The references are defined in the files main.bib, LHCb-PAPER.bib, LHCb-CONF.bib, LHCb-DP.bib LHCb-TDR.bib files, with obvious contents. Each of these have their LHCb-ZZZ-20XX-0YY number as their cite code. If you believe there is a problem with the formatting or content of one of the entries, then get in contact with the Editorial Board rather than just editing it in your local file, since you are likely to need the latest version just before submiting the article.

Description	cite code	Reference
LHCb detector	Alves:2008zz	[1]
LHCb simulation	LHCb-PROC-2011-006	[18]
PDG 2014	PDG2014	[3]
HFAG	HFAG	[30]
Рутніа	Sjostrand:2006za, *Sjostrand:2007gs	[13]
LHCb Pythia tuning	LHCb-PROC-2010-056	[14]
Geant4	Allison:2006ve, *Agostinelli:2002hh	[17]
EVTGEN	Lange: 2001uf	[15]
Photos	Golonka:2005pn	[16]
DIRAC	Tsaregorodtsev:2010zz, *BelleDIRACAmazon	[31]
Crystal Ball function ³	Skwarnicki:1986xj	[32]
Wilks' theorem	Wilks:1938dza	[33]
BDT	Breiman	[19]
BDT training	AdaBoost	[20]
HLT2 topo	BBDT	[11]
DecayTreeFitter	Hulsbergen:2005pu	[34]
sPlot	Pivk:2004ty	[35]
Punzi's optimization	Punzi:2003bu	[36]
f_s/f_d	fsfd	[37]

477

³A valid alternative for most papers where the normalisation is not critical is to use the expression "Gaussian function with a low-mass power-law tail" or "Gaussian function with power-law tails". In that case, no citation is needed

	LHCb-DP number	Title
	LHCb-DP-2014-002 [4]	LHCb detector performance
	LHCb-DP-2014-001 [5]	Performance of the LHCb Vertex Locator
	LHCb-DP-2013-004 [38]	Performance of the LHCb calorimeters
	LHCb-DP-2013-003 [6]	Performance of the LHCb Outer Tracker
	LHCb-DP-2013-002 [39]	Measurement of the track reconstruction efficiency at LHCb
	LHCb-DP-2013-001 [40]	Performance of the muon identification at LHCb
478	LHCb-DP-2012-005 [41]	Radiation damage in the LHCb Vertex Locator
	LHCb-DP-2012-004 [9]	The LHCb trigger and its performance in 2011
	LHCb-DP-2012-003 [7]	Performance of the LHCb RICH detector at the LHC
	LHCb-DP-2012-002 [8]	Performance of the LHCb muon system
	LHCb-DP-2012-001 [42]	Radiation hardness of the LHCb Outer Tracker
	LHCb-DP-2011-002 [43]	Simulation of machine induced background
	LHCb-DP-2011-001 [44]	Performance of the LHCb muon system with cosmic rays
	LHCb-DP-2010-001 [45]	First spatial alignment of the LHCb VELO

LHCb-TDR number	Title
LHCb-TDR-016 [46]	Trigger and online upgrade
LHCb-TDR-015 $[47]$	Tracker upgrade
LHCb-TDR-014 $[48]$	PID upgrade
LHCb-TDR-013 $[49]$	VELO upgrade
LHCb-TDR-012 $[50]$	Framework TDR for the upgrade
LHCb-TDR-011 $[51]$	Computing
LHCb-TDR-010 $[52]$	Trigger
LHCb-TDR-009 $[53]$	Reoptimized detector
LHCb-TDR-008 $[54]$	Inner Tracker
LHCb-TDR-007 $[55]$	Online, DAQ, ECS
LHCb-TDR-006 $[56]$	Outer Tracker
LHCb-TDR-005 $[57]$	VELO
LHCb-TDR-004 $[58]$	Muon system
LHCb-TDR-003 $[59]$	RICH
LHCb-TDR-002 $[60]$	Calorimeters
LHCb-TDR-001 [61]	Magnet

Table 3: LHCb-PAPERs (which have their identifier as their cite code). Note that LHCb-PAPER- 2011-039 does not exist.

LHCb-PAPER-2016-014 [62]	LHCb-PAPER-2016-013 [63]
LHCb-PAPER-2016-012 [64]	LHCb-PAPER-2016-011 [65]
LHCb-PAPER-2016-010 [66]	LHCb-PAPER-2016-009 [67]
LHCb-PAPER-2016-008 [68]	LHCb-PAPER-2016-007 [69]
LHCb-PAPER-2016-006 [70]	LHCb-PAPER-2016-005 [71]
LHCb-PAPER-2016-004 [72]	LHCb-PAPER-2016-003 [73]

– continued from previous page.

- continued from previous page.							
LHCb-PAPER-2016-002	[74]	LHCb-PAPER-2016-001	[75]				
LHCb-PAPER-2015-060	[76]	LHCb-PAPER-2015-059	[77]				
LHCb-PAPER-2015-058	[78]	LHCb-PAPER-2015-057	[79]				
LHCb-PAPER-2015-056	[80]	LHCb-PAPER-2015-055	[81]				
LHCb-PAPER-2015-054	[82]	LHCb-PAPER-2015-053	[83]				
LHCb-PAPER-2015-052	[84]	LHCb-PAPER-2015-051	[85]				
LHCb-PAPER-2015-050	[86]	LHCb-PAPER-2015-049	[87]				
LHCb-PAPER-2015-048	[88]	LHCb-PAPER-2015-047	[89]				
LHCb-PAPER-2015-046	[90]	LHCb-PAPER-2015-045	[91]				
LHCb-PAPER-2015-044	[92]	LHCb-PAPER-2015-043	[93]				
LHCb-PAPER-2015-042	[94]	LHCb-PAPER-2015-041	[95]				
LHCb-PAPER-2015-040	[96]	LHCb-PAPER-2015-039	[97]				
LHCb-PAPER-2015-038	[98]	LHCb-PAPER-2015-037	[99]				
LHCb-PAPER-2015-036	[100]	LHCb-PAPER-2015-035	[101]				
LHCb-PAPER-2015-034	[102]	LHCb-PAPER-2015-033	[103]				
LHCb-PAPER-2015-032	[104]	LHCb-PAPER-2015-031	[105]				
LHCb-PAPER-2015-030	[106]	LHCb-PAPER-2015-029	[29]				
LHCb-PAPER-2015-028	[107]	LHCb-PAPER-2015-027	[108]				
LHCb-PAPER-2015-026	[109]	LHCb-PAPER-2015-025	[110]				
LHCb-PAPER-2015-024	[111]	LHCb-PAPER-2015-023	[112]				
LHCb-PAPER-2015-022	[113]	LHCb-PAPER-2015-021	[114]				
LHCb-PAPER-2015-020	[115]	LHCb-PAPER-2015-019	[116]				
LHCb-PAPER-2015-018	[117]	LHCb-PAPER-2015-017	[118]				
LHCb-PAPER-2015-016	[119]	LHCb-PAPER-2015-015	[120]				
LHCb-PAPER-2015-014	[121]	LHCb-PAPER-2015-013	[122]				
LHCb-PAPER-2015-012	[123]	LHCb-PAPER-2015-011	[124]				
LHCb-PAPER-2015-010	[125]	LHCb-PAPER-2015-009	[126]				
LHCb-PAPER-2015-008	[127]	LHCb-PAPER-2015-007	[128]				
LHCb-PAPER-2015-006	[129]	LHCb-PAPER-2015-005	[130]				
LHCb-PAPER-2015-004		LHCb-PAPER-2015-003					
LHCb-PAPER-2015-002	[133]	LHCb-PAPER-2015-001	[134]				
LHCb-PAPER-2014-070	[135]	LHCb-PAPER-2014-069	[136]				
LHCb-PAPER-2014-068	[137]	LHCb-PAPER-2014-067	[138]				
LHCb-PAPER-2014-066	[139]	LHCb-PAPER-2014-065	[140]				
LHCb-PAPER-2014-064	[141]	LHCb-PAPER-2014-063	[142]				
LHCb-PAPER-2014-062	[143]	LHCb-PAPER-2014-061	[144]				
LHCb-PAPER-2014-060		LHCb-PAPER-2014-059	: :				
LHCb-PAPER-2014-058	[147]	LHCb-PAPER-2014-057	[148]				
LHCb-PAPER-2014-056		LHCb-PAPER-2014-055					
LHCb-PAPER-2014-054	: :	LHCb-PAPER-2014-053	: :				
LHCb-PAPER-2014-052	: :	LHCb-PAPER-2014-051	1 1				
LHCb-PAPER-2014-050	[155]	LHCb-PAPER-2014-049	[156]				

– continued from previous page.

```
LHCb-PAPER-2014-048 [157]
                            LHCb-PAPER-2014-047 [158]
LHCb-PAPER-2014-046 [159]
                            LHCb-PAPER-2014-045 [160]
LHCb-PAPER-2014-044 [161]
                            LHCb-PAPER-2014-043 [162]
LHCb-PAPER-2014-042 [163]
                            LHCb-PAPER-2014-041 [164]
LHCb-PAPER-2014-040 |165|
                            LHCb-PAPER-2014-039 |166|
LHCb-PAPER-2014-038 [167]
                            LHCb-PAPER-2014-037 [168]
LHCb-PAPER-2014-036 [169]
                            LHCb-PAPER-2014-035 [170]
LHCb-PAPER-2014-034 |171|
                            LHCb-PAPER-2014-033 |172|
LHCb-PAPER-2014-032 [173]
                            LHCb-PAPER-2014-031 [174]
LHCb-PAPER-2014-030 [175]
                            LHCb-PAPER-2014-029 [176]
LHCb-PAPER-2014-028 |177
                            LHCb-PAPER-2014-027 | 178
LHCb-PAPER-2014-026 [179]
                            LHCb-PAPER-2014-025 [180]
LHCb-PAPER-2014-024 [181]
                            LHCb-PAPER-2014-023 [182]
LHCb-PAPER-2014-022 [183]
                            LHCb-PAPER-2014-021 [184]
LHCb-PAPER-2014-020 [185]
                            LHCb-PAPER-2014-019 [186]
LHCb-PAPER-2014-018 [187]
                            LHCb-PAPER-2014-017 [188]
LHCb-PAPER-2014-016 [189]
                            LHCb-PAPER-2014-015 [190]
LHCb-PAPER-2014-014 [191]
                            LHCb-PAPER-2014-013 [192]
LHCb-PAPER-2014-012 |193|
                            LHCb-PAPER-2014-011 | 194
LHCb-PAPER-2014-010 |195|
                            LHCb-PAPER-2014-009 | 196
LHCb-PAPER-2014-008 [197]
                            LHCb-PAPER-2014-007 [198]
LHCb-PAPER-2014-006 [12]
                            LHCb-PAPER-2014-005 [199]
LHCb-PAPER-2014-004 [200]
                            LHCb-PAPER-2014-003 |201
LHCb-PAPER-2014-002 [202]
                            LHCb-PAPER-2014-001 [203]
LHCb-PAPER-2013-070 [27]
                            LHCb-PAPER-2013-069 [204]
LHCb-PAPER-2013-068 [205]
                            LHCb-PAPER-2013-067 [206]
LHCb-PAPER-2013-066 [207]
                            LHCb-PAPER-2013-065 [208]
LHCb-PAPER-2013-064 [209]
                            LHCb-PAPER-2013-063 [210]
LHCb-PAPER-2013-062 |211|
                            LHCb-PAPER-2013-061 |212
LHCb-PAPER-2013-060 [213]
                            LHCb-PAPER-2013-059 [214]
LHCb-PAPER-2013-058 [215]
                            LHCb-PAPER-2013-057 [216]
LHCb-PAPER-2013-056 |217|
                            LHCb-PAPER-2013-055 |218
LHCb-PAPER-2013-054 [219]
                            LHCb-PAPER-2013-053 [220]
LHCb-PAPER-2013-052 [221]
                            LHCb-PAPER-2013-051 [222]
LHCb-PAPER-2013-050 [223]
                            LHCb-PAPER-2013-049 [224]
LHCb-PAPER-2013-048 |225|
                            LHCb-PAPER-2013-047 | 226
LHCb-PAPER-2013-046 [227]
                            LHCb-PAPER-2013-045 [228]
LHCb-PAPER-2013-044 [229]
                            LHCb-PAPER-2013-043 [230]
LHCb-PAPER-2013-042 |231|
                            LHCb-PAPER-2013-041 |232
LHCb-PAPER-2013-040 [233]
                            LHCb-PAPER-2013-039 [234]
LHCb-PAPER-2013-038 [235]
                            LHCb-PAPER-2013-037 [236]
LHCb-PAPER-2013-036 [237]
                            LHCb-PAPER-2013-035 [28]
```

- continued from previous page.

```
LHCb-PAPER-2013-034 [238]
                            LHCb-PAPER-2013-033 [239]
LHCb-PAPER-2013-032 [240]
                            LHCb-PAPER-2013-031 [241]
LHCb-PAPER-2013-030 [242]
                            LHCb-PAPER-2013-029 [243]
LHCb-PAPER-2013-028 [244]
                            LHCb-PAPER-2013-027 |245
LHCb-PAPER-2013-026 |246|
                            LHCb-PAPER-2013-025 |247
LHCb-PAPER-2013-024 [248]
                            LHCb-PAPER-2013-023 [249]
LHCb-PAPER-2013-022 [250]
                            LHCb-PAPER-2013-021 [251]
LHCb-PAPER-2013-020 |252|
                            LHCb-PAPER-2013-019 |253|
LHCb-PAPER-2013-018 [254]
                            LHCb-PAPER-2013-017 [255]
LHCb-PAPER-2013-016 [256]
                            LHCb-PAPER-2013-015 [257]
LHCb-PAPER-2013-014 [258]
                            LHCb-PAPER-2013-013 | 259
LHCb-PAPER-2013-012 [260]
                            LHCb-PAPER-2013-011 [261]
LHCb-PAPER-2013-010 [262]
                            LHCb-PAPER-2013-009 [263]
LHCb-PAPER-2013-008 [264]
                            LHCb-PAPER-2013-007 [265]
LHCb-PAPER-2013-006 [266]
                            LHCb-PAPER-2013-005 [267]
LHCb-PAPER-2013-004 [268]
                            LHCb-PAPER-2013-003 [269]
                            LHCb-PAPER-2013-001 [271
LHCb-PAPER-2013-002 |270|
LHCb-PAPER-2012-057 [272]
LHCb-PAPER-2012-056 [273]
                            LHCb-PAPER-2012-055 [274]
LHCb-PAPER-2012-054 |275|
                            LHCb-PAPER-2012-053 |276
LHCb-PAPER-2012-052 [277]
                            LHCb-PAPER-2012-051 [278]
LHCb-PAPER-2012-050 [279]
                            LHCb-PAPER-2012-049 [280]
LHCb-PAPER-2012-048 [281]
                            LHCb-PAPER-2012-047 [282]
LHCb-PAPER-2012-046 [283]
                            LHCb-PAPER-2012-045 [284]
LHCb-PAPER-2012-044 [285]
                            LHCb-PAPER-2012-043 [286]
LHCb-PAPER-2012-042 [287]
                            LHCb-PAPER-2012-041 [288]
LHCb-PAPER-2012-040 [289]
                            LHCb-PAPER-2012-039 [290]
LHCb-PAPER-2012-038 [291]
                            LHCb-PAPER-2012-037 [292]
LHCb-PAPER-2012-036 [293]
                            LHCb-PAPER-2012-035 |294
LHCb-PAPER-2012-034 [295]
                            LHCb-PAPER-2012-033 [296]
LHCb-PAPER-2012-032 [297]
                            LHCb-PAPER-2012-031 [298]
LHCb-PAPER-2012-030 [299]
                            LHCb-PAPER-2012-029 [300]
LHCb-PAPER-2012-028 [301]
                            LHCb-PAPER-2012-027 [302]
LHCb-PAPER-2012-026 [303]
                            LHCb-PAPER-2012-025 [304]
LHCb-PAPER-2012-024 [305]
                            LHCb-PAPER-2012-023 [306]
LHCb-PAPER-2012-022 |307|
                            LHCb-PAPER-2012-021 |308|
LHCb-PAPER-2012-020 [309]
                            LHCb-PAPER-2012-019 [310]
LHCb-PAPER-2012-018 [311]
                            LHCb-PAPER-2012-017 [312]
LHCb-PAPER-2012-016 |313|
                            LHCb-PAPER-2012-015 |314|
LHCb-PAPER-2012-014 [315]
                            LHCb-PAPER-2012-013 [316]
LHCb-PAPER-2012-012 [317]
                            LHCb-PAPER-2012-011 [318]
LHCb-PAPER-2012-010 [319]
                            LHCb-PAPER-2012-009 [320]
```

- continued from previous page.

- commue	за поп	n previous page.
LHCb-PAPER-2012-008	[321]	LHCb-PAPER-2012-007 [322]
LHCb-PAPER-2012-006	[323]	LHCb-PAPER-2012-005 [324]
LHCb-PAPER-2012-004	[325]	LHCb-PAPER-2012-003 [326]
LHCb-PAPER-2012-002	[327]	LHCb-PAPER-2012-001 [328]
LHCb-PAPER-2011-045	[329]	LHCb-PAPER-2011-044 [330]
LHCb-PAPER-2011-043	[331]	LHCb-PAPER-2011-042 [332]
LHCb-PAPER-2011-041	[333]	LHCb-PAPER-2011-040 [334]
LHCb-PAPER-2011-038	[335]	LHCb-PAPER-2011-037 [336]
LHCb-PAPER-2011-036	[337]	LHCb-PAPER-2011-035 [338]
LHCb-PAPER-2011-034	[339]	LHCb-PAPER-2011-033 [340]
LHCb-PAPER-2011-032	[341]	LHCb-PAPER-2011-031 [342]
LHCb-PAPER-2011-031	·	LHCb-PAPER-2011-029 [344]
LHCb-PAPER-2011-028	[345]	LHCb-PAPER-2011-027 [346]
LHCb-PAPER-2011-026	[347]	LHCb-PAPER-2011-025 [348]
LHCb-PAPER-2011-024	F	LHCb-PAPER-2011-023 [350]
LHCb-PAPER-2011-023	[351]	LHCb-PAPER-2011-021 [352]
LHCb-PAPER-2011-020	[353]	LHCb-PAPER-2011-019 [354]
LHCb-PAPER-2011-018		LHCb-PAPER-2011-017 [356]
LHCb-PAPER-2011-016	F 1	LHCb-PAPER-2011-015 [358]
LHCb-PAPER-2011-014	[359]	LHCb-PAPER-2011-013 [360]
LHCb-PAPER-2011-012		LHCb-PAPER-2011-011 [362]
LHCb-PAPER-2011-010	[363]	LHCb-PAPER-2011-009 [364]
LHCb-PAPER-2011-008		LHCb-PAPER-2011-007 [366]
LHCb-PAPER-2011-006		LHCb-PAPER-2011-005 [368]
LHCb-PAPER-2011-004		LHCb-PAPER-2011-003 [370]
LHCb-PAPER-2011-002	[371]	LHCb-PAPER-2011-001 [372]
LHCb-PAPER-2010-002	[373]	LHCb-PAPER-2010-001 [374]
LHCb-PAPER-2010-002	[373]	LHCb-PAPER-2010-001 [374]

480

Table 4: LHCb-CONFs (which have their identifier as their cite code). Note that LHCb-CONF-2011-032 does not exist.

LHCb-CONF-2016-004 [375]	LHCb-CONF-2016-003 [376]
LHCb-CONF-2016-002 [377]	LHCb-CONF-2016-001 [378]
LHCb-CONF-2015-005 [379]	
LHCb-CONF-2015-004 [380]	LHCb-CONF-2015-003 [381]
LHCb-CONF-2015-002 [382]	LHCb-CONF-2015-001 [383]
LHCb-CONF-2014-004 [384] ⁴	LHCb-CONF-2014-003 [385]
LHCb-CONF-2014-002 [386]	LHCb-CONF-2014-001 [387]

⁴If you cite the gamma combination, always also cite the latest gamma paper as \cite{LHCb-PAPER-2013-020,*LHCb-CONF-2014-004} (unless you cite LHCb-PAPER-2013-020 separately too).

– continued from previous page.

– continu	ed from	previous page.
LHCb-CONF-2013-013	[388]	
LHCb-CONF-2013-012	[389]	LHCb-CONF-2013-011 [390]
LHCb-CONF-2013-010	[391]	LHCb-CONF-2013-009 [392]
LHCb-CONF-2013-008	[393]	LHCb-CONF-2013-007 [394]
LHCb-CONF-2013-006	Ē	LHCb-CONF-2013-005 [396]
LHCb-CONF-2013-004	[397]	LHCb-CONF-2013-003 [398]
LHCb-CONF-2013-002		LHCb-CONF-2013-001 [400]
LHCb-CONF-2012-034	[401]	LHCb-CONF-2012-033 [402]
LHCb-CONF-2012-032	[403]	LHCb-CONF-2012-031 [404]
LHCb-CONF-2012-030	[405]	LHCb-CONF-2012-029 [406]
LHCb-CONF-2012-028	[407]	LHCb-CONF-2012-027 [408]
LHCb-CONF-2012-026	[409]	LHCb-CONF-2012-025 [410]
LHCb-CONF-2012-024	[411]	LHCb-CONF-2012-023 [412]
LHCb-CONF-2012-022		LHCb-CONF-2012-021 [414]
LHCb-CONF-2012-020	[415]	LHCb-CONF-2012-019 [416]
LHCb-CONF-2012-018	[417]	LHCb-CONF-2012-017 [418]
LHCb-CONF-2012-016	[419]	LHCb-CONF-2012-015 [420]
LHCb-CONF-2012-014	[421]	LHCb-CONF-2012-013 [422]
LHCb-CONF-2012-012	[423]	LHCb-CONF-2012-011 [424]
LHCb-CONF-2012-010	[425]	LHCb-CONF-2012-009 [426]
LHCb-CONF-2012-008	[427]	LHCb-CONF-2012-007 [428]
LHCb-CONF-2012-006	[429]	LHCb-CONF-2012-005 [430]
LHCb-CONF-2012-004	[431]	${\tt LHCb-CONF-2012-003} \ [432]$
LHCb-CONF-2012-002	[433]	LHCb-CONF-2012-001 [434]
LHCb-CONF-2011-062	$[\overline{435}]$	LHCb-CONF-2011-061 [436]
LHCb-CONF-2011-060	[437]	LHCb-CONF-2011-059 [438]
LHCb-CONF-2011-058	[439]	LHCb-CONF-2011-057 [440]
LHCb-CONF-2011-056	[441]	LHCb-CONF-2011-055 [442]
LHCb-CONF-2011-054	[443]	LHCb-CONF-2011-053 [444]
LHCb-CONF-2011-052		LHCb-CONF-2011-051 [446]
LHCb-CONF-2011-050	[447]	LHCb-CONF-2011-049 [448]
LHCb-CONF-2011-048	[449]	LHCb-CONF-2011-047 [450]
LHCb-CONF-2011-046	[451]	LHCb-CONF-2011-045 [452]
LHCb-CONF-2011-044	[453]	LHCb-CONF-2011-043 [454]
LHCb-CONF-2011-042	[455]	LHCb-CONF-2011-041 [456]
LHCb-CONF-2011-040	[457]	LHCb-CONF-2011-039 [458]
LHCb-CONF-2011-038	[459]	LHCb-CONF-2011-037 [460]
LHCb-CONF-2011-036	[461]	LHCb-CONF-2011-035 [462]
LHCb-CONF-2011-034	: :	LHCb-CONF-2011-033 [464]
LHCb-CONF-2011-031	[465]	
LHCb-CONF-2011-030	Ē ī	LHCb-CONF-2011-029 [467]
LHCb-CONF-2011-028	[468]	LHCb-CONF-2011-027 [469]

- continued from previous page.

```
LHCb-CONF-2011-026
                           LHCb-CONF-2011-025 [471]
                    |470|
LHCb-CONF-2011-024
                    |472|
                           LHCb-CONF-2011-023 [473]
LHCb-CONF-2011-023 [474]
                           LHCb-CONF-2011-021 [475]
LHCb-CONF-2011-020 [476]
                           LHCb-CONF-2011-019 [477]
LHCb-CONF-2011-018
                    |478|
                           LHCb-CONF-2011-017 |479|
LHCb-CONF-2011-016
                    [480]
                           LHCb-CONF-2011-015 [481]
LHCb-CONF-2011-014 [482
                           LHCb-CONF-2011-013 [483]
LHCb-CONF-2011-012 |484
                           LHCb-CONF-2011-011 |485|
LHCb-CONF-2011-010
                    [486]
                           LHCb-CONF-2011-009 [487]
LHCb-CONF-2011-008
                    488
                           LHCb-CONF-2011-007 [489]
LHCb-CONF-2011-006
                    |490|
                           LHCb-CONF-2011-005 [491]
LHCb-CONF-2011-004
                           LHCb-CONF-2011-003 [25]
                    |492|
LHCb-CONF-2011-002 [493
                           LHCb-CONF-2011-001 [494]
LHCb-CONF-2010-014
                           LHCb-CONF-2010-013 [496]
                    |495|
LHCb-CONF-2010-012
                           LHCb-CONF-2010-011 [498]
                    [497]
                           LHCb-CONF-2010-009 [500]
LHCb-CONF-2010-010
                    [499]
LHCb-CONF-2010-008 [501
```

Some LHCb papers quoted together will look like [366–370]. The combination of CMS and LHCb results on $B_{(s)}^0 \to \mu^+ \mu^-$ should be cited like [389].

⁴⁸⁴ B Standard symbols

481

485

486

487

488

489

490

492

493

494

495

496

497

498

499

As explained in Sect. 4 this appendix contains standard typesetting of symbols, particle names, units etc. in LHCb documents.

In the file lhcb-symbols-def.tex, which is included, a large number of symbols is defined. While they can lead to quicker typing, the main reason is to ensure a uniform notation within a document and between different LHCb documents. If a symbol like \CP to typeset CP violation is available for a unit, particle name, process or whatever, it should be used. If you do not agree with the notation you should ask to get the definition in lhcb-symbols-def.tex changed rather than just ignoring it.

All the main particles have been given symbols. The B mesons are thus named B^+ , B^0 , B_s^0 , and B_c^+ . There is no need to go into math mode to use particle names, thus saving the typing of many \$ signs. By default particle names are typeset in italic type to agree with the PDG preference. To get roman particle names you can just change \setboolean{uprightparticles}{false} to true at the top of this template.

There is a large number of units typeset that ensures the correct use of fonts, capitals and spacing. As an example we have $m_{B_s^0} = 5366.3 \pm 0.6 \,\mathrm{MeV}/c^2$. Note that $\,\mu\mathrm{m}$ is typeset with an upright $\,\mu$, even if the particle names have slanted greek letters.

A set of useful symbols are defined for working groups. More of these symbols can be included later. As an example in the Rare Decay group we have several different analyses looking for a measurement of $\mathcal{C}_7^{'(\mathrm{eff})}$ and $\mathcal{O}_7^{'}$.

C List of all symbols

505 C.1 Experiments

	$\backslash exttt{lhcb}$	LHCb	\setminus atlas	ATLAS	$\backslash \mathtt{cms}$	CMS
\	\alice	ALICE	\babar	BaBar	\belle	Belle
	\cleo	CLEO	\cdf	CDF	\dzero	D0
506	$ar{ ext{aleph}}$	ALEPH	\delphi	DELPHI	\opal	OPAL
\	ackslash 1three	L3	\sld	SLD	\cern	CERN
	\lhc	LHC	\lep	LEP	tevatron	Tevatron

C.1.1 LHCb sub-detectors and sub-systems

	\velo	VELO	$\$ rich	RICH	$\backslash { t richone}$	RICH1
\	richtwo	RICH2	ttracker	TT	\intr	IT
	\st	ST	\ot	OT	\spd	SPD
508 \	presh	PS	\ecal	ECAL	\hcal	HCAL
	$\backslash \texttt{MagUp}$	MagUp	$\backslash \texttt{MagDown}$	MagDown	\ode	ODE
	\daq	DAQ	\tfc	TFC	\ecs	ECS
١	\lone	L0	$ackslash exttt{hlt}$	HLT	$ackslash ext{hltone}$	HLT1
	\hlttwo	HLT2	•			

509 C.2 Particles

510 C.2.1 Leptons

	ackslashelectron	e	\en	e^{-}	\ep	e^+
	\epm	e^{\pm}	$\backslash \mathtt{epem}$	e^+e^-	$\backslash \mathtt{muon}$	μ
	$\backslash \mathtt{mup}$	μ^+	$\mbox{\tt mun}$	μ^-	$\backslash mumu$	$\mu^+\mu^-$
	ackslashtauon	au	\setminus taup	$ au^+$	$\backslash \mathtt{taum}$	$ au^-$
511	\tautau	$ au^+ au^-$	\setminus lepton	ℓ	$\backslash \mathtt{ellm}$	ℓ^-
	\ellp	ℓ^+	\ellell	$\ell^+\ell^-$	\neu	ν
	\neub	$\overline{ u}$	\neue	$ u_e$	$\new $	$\overline{ u}_e$
	\setminus neum	$ u_{\mu}$	$\new neumb$	$\overline{ u}_{\mu}$	\nextit{neut}	$ u_{ au}$
	$ackslash ext{neutb}$	$\overline{ u}_{ au}$	$\ne $	$ u_{\ell}$	$\ne $	$\overline{ u}_\ell$

$_{512}$ C.2.2 Gauge bosons and scalars

	\g	γ	$\backslash H$	H^0	$\backslash \mathtt{Hp}$	H^+
513	$\backslash \mathtt{Hm}$	H^{-}	$\backslash \mathtt{Hpm}$	H^\pm	$\backslash \mathtt{W}$	W
	$\backslash \mathtt{Wp}$	W^+	\Wm	W^-	$\backslash \mathtt{Wpm}$	W^{\pm}
	$\backslash Z$	Z				

514 C.2.3 Quarks

,	$ackslash ext{quark}$	q	\quarkbar	\overline{q}	$\backslash qqbar$	$q\overline{q}$
,	$ackslash ext{uquark}$	u	\uquarkbar	\overline{u}	\uubar	$u\overline{u}$
,	$\backslash \mathtt{dquark}$	d	\dquarkbar	\overline{d}	\ddbar	$d\overline{d}$
515	$ackslash ext{squark}$	S	\setminus squarkbar	\overline{S}	\setminus ssbar	$s\overline{s}$
,	ackslash cquark	c	\cquarkbar	\overline{c}	\ccbar	$c\overline{c}$
,	\bquark	b	\bquarkbar	\overline{b}	\bbbar	$b\overline{b}$
,	$ackslash ag{tquark}$	t	$ackslash ag{tquarkbar}$	\overline{t}	$ackslash ag{ttbar}$	$t\overline{t}$

516 C.2.4 Light mesons

	$\$ hadron	h	\pion	π	\piz	π^0
	\pizs	$\pi^0\mathrm{s}$	\pip	π^+	\pim	π^-
	\pipm	π^\pm	\pimp	π^{\mp}	$\$ rhomeson	ρ
	$\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ $	$ ho^0$	$\backslash { t rhop}$	$ ho^+$	rhom	$ ho^-$
	\rdown	$ ho^\pm$	$\backslash \texttt{rhomp}$	$ ho^{\mp}$	\kaon	K
	\Kb	\overline{K}	\backslash KorKbar	(\overline{K})	\Kz	K^0
517	\Kzb	\overline{K}^0	\Kp	K^+	\Km	K^{-}
	Kpm	K^{\pm}	\Kmp	K^{\mp}	\KS	$K_{ m \scriptscriptstyle S}^0$
	\KL	$K_{ m \scriptscriptstyle L}^0$	\setminus Kstarz	K^{*0}	\Kstarzb	\overline{K}^{*0}
	Kstar	K^*	Kstarb	\overline{K}^*	\Kstarp	K^{*+}
,	Kstarm	K^{*-}	Kstarpm	$K^{*\pm}$	Kstarmp	$K^{*\mp}$
	\etaz	η	\etapr	η'	\phiz	ϕ
	\omegaz	ω				

Heavy mesons 518 **C.2.5**

523 \Lbbar

 \X ib

 $\backslash \texttt{Xic}$

 $\backslash \texttt{Xibbar}$

 $\backslash \texttt{Xicbar}$

 $\backslash \mathtt{Omegac}$

 $\backslash \mathtt{Omegabbar}$

519	\D \Dz \Dm \Dstar \Dstarzb \Dstarpm \Dsp \Dsmp \Dssm \B \BorBbar \Bu \Bm \Bd \Bdb \Bdb	$D \\ D^{0} \\ D^{-} \\ D^{*} \\ \overline{D}^{*0} \\ D^{*\pm} \\ D^{*\pm} \\ D^{+}_{s} \\ D^{*-}_{s} \\ B \\ (\overline{B}) \\ B^{+} \\ B^{-} \\ B^{0} \\ \overline{B}^{0} \\ B^{-}_{c}$	\Db \Dzb \Dpm \Dstarb \Dstarp \Dstarmp \Dsm \Dss \Dsspm \Bbar \Bz \Bub \Bpm \Bs \Bc \Bc	$ar{D}$ $ar{D}^0$ D^\pm $ar{D}^*$ D^{*+} D^{*+} D^s^{++} D^s^{*+} B^* B^0 $B^ B^\pm$ B^0 B^c B^c B^c	\DorDbar \Dp \Dmp \Dstarz \Dstarm \Ds \Dspm \Dssp \Dssmp \Bb \Bzb \Bp \Bmp \Bsb \Bcp	$\stackrel{(\overline{D})}{D}^{+}$ D^{+} D^{*0} D^{*-} D^{+}_{s} D^{*}_{s} D^{*+}_{s} \overline{B}^{0} B^{+} B^{0}_{s} B^{+}_{c}
520	C.2.6 On	ia				
521	\jpsi \etac \chictwo \ThreeS \chic	J/ψ η_c χ_{c2} $\Upsilon(3S)$ χ_c	\psitwos \chiczero \OneS \FourS	$\psi(2S)$ χ_{c0} $\Upsilon(1S)$ $\Upsilon(4S)$	\chicone χ_{c1} \TwoS $\Upsilon($	3770) 1 2S) 5S)
522	C.2.7 Ba	ryons				
	\proton \antineutr \Xires \Lbar \Lambdares \Omegares	$rac{arnothing}{ar{arLambda}}$	\antiproto \Deltares \Xiresbar \LorLbar \Sigmares \Omegaresb	$\frac{\Delta}{\overline{\Xi}}$ $\overline{\Lambda}$ Σ	\neutron \Deltaresbar \Lz \Lambdares \Sigmaresbar \Lb	$ \frac{n}{\Delta} $ $ \Lambda $ $ \frac{\Lambda}{\Sigma} $ $ \frac{\Lambda_b^0}{\delta} $

\Lc

\Xibz

 $\backslash \mathtt{Xicz}$

\Xibbarz

\Xicbarz

\Omegacbar

 \Lcbar

 $\backslash Xibm$

 $\backslash \texttt{Xicp}$

\Xibbarp

 $\backslash \texttt{Xicbarm}$

 $\backslash \mathtt{Omegab}$

⁵²⁴ C.3 Physics symbols

525 C.3.1 Decays

527 C.3.2 Lifetimes

529 C.3.3 Masses

=	$\backslash \mathtt{mBd}$	m_{B^0}	$\backslash \mathtt{mBp}$	m_{B^+}	$\backslash \mathtt{mBs}$	$m_{B_s^0}$
530	$\backslash \mathtt{mBc}$	$m_{B_c^+}$	\mbox{mLb}	$m_{A_b^0}$		

531 C.3.4 EW theory, groups

	\grpsuthree	SU(3)	$\backslash \mathtt{grpsutw}$	SU(2)	\grpuone	U(1)
,	\setminus ssqtw	$\sin^2\!\theta_{ m W}$	ackslash csqtw	$\cos^2 \theta_{ m W}$	$\backslash \mathtt{stw}$	$\sin \theta_{ m W}$
	\ctw	$\cos heta_{ m W}$	\setminus ssqtwef	$\sin^2\! heta_{ m W}^{ m eff}$	$\backslash \mathtt{csqtwef}$	$\cos^2 \theta_{ m W}^{ m eff}$
532	\stwef	$\sin heta_{ m W}^{ m eff}$	ackslash ctwef	$\cos heta_{ m W}^{ m eff}$	\gv	$g_{ m \scriptscriptstyle V}$
\	\ga	$g_{ m A}$	$\backslash \mathtt{order}$	\mathcal{O}	$\backslash \mathtt{ordalph}$	$\mathcal{O}(\alpha)$
	\ordalsq	$\mathcal{O}(lpha^2)$	$\backslash \mathtt{ordalcb}$	$\mathcal{O}(lpha^3)$		

533 C.3.5 QCD parameters

,	\setminus as	α_s	$ackslash ext{MSb}$	$\backslash ext{lqcd}$	$\Lambda_{ m QCD}$
534	\qsq	q^2			

535 C.3.6 CKM, CP violation

	ackslasheps	ε	\epsK	$arepsilon_K$	$\backslash \mathtt{epsB}$	ε_B
	$\backslash \mathtt{epsp}$	$arepsilon_K'$	\CP	CP	$\backslash \mathtt{CPT}$	CPT
	$\$ rhobar	$\overline{ ho}$	\etabar	$\overline{\eta}$	$\setminus Vud$	V_{ud}
	$\backslash Vcd$	V_{cd}	\Vtd	V_{td}	$\setminus \mathtt{Vus}$	V_{us}
536	\Vcs	V_{cs}	\Vts	V_{ts}	$\setminus Vub$	V_{ub}
	$\backslash exttt{Vcb}$	V_{cb}	\Vtb	V_{tb}	$\setminus \mathtt{Vuds}$	V_{ud}^*
	$\backslash \mathtt{Vcds}$	V_{cd}^*	\Vtds	V_{td}^*	$ackslash exttt{Vuss}$	V_{us}^*
	$ackslash extsf{Vcss}$	V_{cs}^*	\Vtss	V_{ts}^*	$\setminus extsf{Vubs}$	V_{ub}^*
	$ackslash extsf{Vcbs}$	V_{cb}^{st}	\Vtbs	V_{tb}^*		

537 C.3.7 Oscillations

	\dm	Δm	$\backslash \mathtt{dms}$	Δm_s	\dmd	Δm_d
	\DG	$\Delta\Gamma$	\DGs	$\Delta\Gamma_s$	\DGd	$\Delta\Gamma_d$
	\Gs	Γ_s	\Gd	Γ_d	\ \ MBq	M_{B_a}
	DGq	$\Delta\Gamma_q$	\Gq	Γ_q	$\backslash dmq$	Δm_q
	\GL	$\Gamma_{ m L}$	\GH	$\Gamma_{ m H}$	DGsGs	$\Delta\Gamma_s/\Gamma_s$
538	Delm	Δm	ACP	\mathcal{A}^{CP}	Adir	$\mathcal{A}^{ ext{dir}}$
	\Amix	$\mathcal{A}^{ ext{mix}}$	\ADelta	\mathcal{A}^{Δ}	\ \phid	ϕ_d
	sinphid	$\sin \phi_d$	\phis	ϕ_s	\betas	β_s
,	\sbetas	$\sigma(\beta_s)$	\stbetas	$\sigma(2\beta_s)$	stphis	$\sigma(\phi_s)$
	sinphis	$\sin \phi_s$	•			, ,

539 C.3.8 Tagging

,	\edet	$arepsilon_{ m det}$	\erec	$\varepsilon_{ m rec/det}$	\esel	$\varepsilon_{ m sel/rec}$
,	\etrg	$\varepsilon_{ m trg/sel}$	\etot	$arepsilon_{ ext{tot}}$	ackslashmistag	ω
540	$\backslash \mathtt{wcomb}$	ω^{comb}	\etag	$arepsilon_{ ext{tag}}$	ackslashetagcomb	$\varepsilon_{ m tag}^{ m comb}$
,	\effeff	$arepsilon_{ ext{eff}}$	\effeffcomb	$arepsilon_{ ext{eff}}^{ ext{comb}}$	\efftag	$\varepsilon_{\rm tag}(1-2\omega)^2$
,	\effD	$\varepsilon_{\mathrm{tag}}D^2$	\etagprompt	$arepsilon_{ ext{tag}}^{ ext{Pr}}$	\etagLL	$arepsilon_{ m tag}^{ m LL}$

541 C.3.9 Key decay channels

	$\backslash \texttt{BdToKstmm}$	$B^0 \rightarrow K^{*0} \mu^+ \mu^-$	$^-ackslash BdbToKstmm$	$\overline B{}^0 \to \overline K^{*0} \mu^+ \mu^-$	−\BsToJPsiPhi	$B_s^0 \to J/\psi \phi$
	\BdToJPsiKst	$B^0 \rightarrow J/\psi K^{*0}$	$\backslash \texttt{BdbToJPsiKst}$	$\overline B{}^0\! o J\!/\!\psi\overline K^{*0}$	ackslash BsPhiGam	$B_s^0 \to \phi \gamma$
542	$\backslash \texttt{BdKstGam}$	$B^0\! o K^{*0}\gamma$	$\backslash \mathtt{BTohh}$	$B \rightarrow h^+ h^{\prime -}$	$\backslash \texttt{BdTopipi}$	$B^0 \rightarrow \pi^+\pi^-$
	\BdToKpi	$B^0 \rightarrow K^+\pi^-$	\BsToKK	$B_a^0 \rightarrow K^+K^-$	\BsTopiK	$B_a^0 \rightarrow \pi^+ K^-$

543 C.3.10 Rare decays

545 C.3.11 Wilson coefficients and operators

547 C.3.12 Charm

```
549 C.3.13 QM
```

 $\texttt{b} \quad \texttt{bra[1] } \\ \texttt{b} \quad \texttt{a} \quad \texttt{a} \quad \texttt{b} \quad \texttt{b} \quad \texttt{b} \quad \texttt{b} \quad \texttt{braket[2] } \\ \texttt{b} \quad \texttt{a} \\ \texttt{b} \quad \texttt{a} \\ \texttt{b} \quad \texttt{b} \quad \texttt{b} \quad \texttt{b} \\ \texttt{b} \\ \texttt{b} \quad \texttt{b} \\ \texttt{b} \\$

551 C.4 Units

 $_{552}$ \unit[1] \unit{kg} kg

553 C.4.1 Energy and momentum

	\tev	TeV	\gev	GeV	\mev	MeV
,	\kev	keV	\ev	eV	\gevc	GeV/c
554	$\backslash \mathtt{mevc}$	MeV/c	\gevcc	GeV/c^2	\gevgevccc	GeV^2/c^4
	$\backslash \mathtt{mevcc}$	MeV/c^2				

555 C.4.2 Distance and area

	$\backslash \mathtt{km}$	km	$\backslash m$	m	$\backslash \mathtt{ma}$	m^2
\	$\backslash \mathtt{cm}$	cm	$\backslash \mathtt{cma}$	cm^2	$\backslash mm$	mm
	$\backslash \mathtt{mma}$	mm^2	$\backslash \mathtt{mum}$	μm	$\backslash \mathtt{muma}$	$\mu\mathrm{m}^2$
	$\backslash \mathtt{nm}$	nm	$\backslash \mathtt{fm}$	fm	\barn	b
556	ackslashmbarn	mb	\mub	μb	\nb	nb
\	\setminus invnb	nb^{-1}	\pb	pb '	\invpb	pb^{-1}
	fb	fb	\invfb	fb^{-1}	\ab	ab
	\invab	ab^{-1}	•			

557 C.4.3 Time

	\sec	S	$\backslash \mathtt{ms}$	ms	$\backslash \mathtt{mus}$	μs
	$\backslash \mathtt{ns}$	ns	\ps	ps	\fs	fs
558	mhz	MHz	\khz	kHz	\hz	Hz
	\invps	ps^{-1}	invns	ns^{-1}	\yr	yr
	\hr	hr				

559 C.4.4 Temperature

 $_{\text{560}}$ \degc $^{\circ}\mathrm{C}$ \degk K

61 C.4.5 Material lengths, radiation

	\Xrad	X_0	\NIL	λ_{int}	$\backslash exttt{mip}$	MIP
562	\setminus neutroneq	n_{eq}	\setminus neqcmcm	$n_{\rm eq}/{ m cm}^2$	\kRad	kRad
	\MRad	MRad	\ci	Ci	\mci	mCi

563 C.4.6 Uncertainties

565 C.4.7 Maths

	\order	\mathcal{O}	ackslashchisq	χ^2	ackslashchisqndf	χ^2/ndf
	$\backslash \mathtt{chisqip}$	$\chi^2_{ m IP}$	ackslashchisqvs	$\chi^2_{ m VS}$	$\backslash \mathtt{chisqvtx}$	$\chi^2_{ m vtx}$
566 \	$\backslash \mathtt{chisqvtxndf}$	$\chi^2_{ m vtx}/{ m ndf}$	\deriv	d	$\gray gsim$	\gtrsim
	$\$ lsim	\lesssim	$\mathbb{1} \operatorname{mean}[1]$	$\langle x \rangle$	$\abs[1] \abs[x]$	x
	\Real	$\mathcal{R}e$	$\backslash { t Imag}$	$\mathcal{I}m$	\PDF	PDF
	\sPlot	sPlot				

567 C.5 Kinematics

568 C.5.1 Energy, Momenta

	ackslash Ebeam	$E_{\scriptscriptstyle m BEAM}$	\sqs	\sqrt{S}	$\backslash \mathtt{ptot}$	p
569	\pt	$p_{ m T}$	\et	$E_{ m T}$	$\backslash \mathtt{mt}$	$M_{ m T}$
	\dpp	$\Delta p/p$	$\backslash \mathtt{msq}$	m^2	$\backslash dedx$	dE/dx

570 C.5.2 PID

\dllkpi
$$\mathrm{DLL}_{K\pi}$$
 \dllppi $\mathrm{DLL}_{p\pi}$ \dllepi $\mathrm{DLL}_{e\pi}$

572 C.5.3 Geometry

$$\$$
 \degrees $^{\circ}$ \krad krad \mrad mrad 573 \rad rad

574 C.5.4 Accelerator

$$_{75}$$
 \betastar eta^* \lum ${\cal L}$ \intlum[1] \intlum{2 fb $^{-1}$ } $\int {\cal L} = 2 ext{ fb}^{-1}$

576 C.6 Software

577 C.6.1 Programs

	\bcvegpy	BCVEGPY	ackslashboole	BOOLE	\brunel	Brunel
	\davinci	DAVINCI	\dirac	DIRAC	\evtgen	EVTGEN
	\fewz	FEWZ	\fluka	FLUKA	$\backslash \mathtt{ganga}$	Ganga
578	\gaudi	GAUDI	\gauss	Gauss	$\backslash { t geant}$	Geant4
	$\backslash \texttt{hepmc}$	НЕРМС	ackslashherwig	HERWIG	$\backslash \mathtt{moore}$	Moore
	\setminus neurobayes	NeuroBayes	$ackslash exttt{photos}$	Photos	\setminus powheg	Powheg
,	$\protect\pro$	Рутніа	\resbos	ResBos	$\backslash { t roofit}$	RooFit
	\root	Root	\spice	SPICE	\setminus urania	Urania

579 C.6.2 Languages

		C++	$\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ $	Ruby	$\setminus \texttt{fortran}$	FORTRAN
580	\svn	SVN				

581 C.6.3 Data processing

	ackslash kbytes	kbytes	ackslash kbsps	kbits/s	$ackslash ext{kbits}$	kbits
	ackslash kbsps	kbits/s	$ackslash {\tt mbsps}$	Mbytes/s	$ackslash \mathtt{mbytes}$	Mbytes
582	$\backslash \mathtt{mbps}$	Mbyte/s	$ackslash {\tt mbsps}$	Mbytes/s	ackslash gbsps	Gbytes/s
	ackslash gbytes	Gbytes	ackslash gbsps	Gbytes/s	$ackslash ext{tbytes}$	Tbytes
	$ackslash ag{tbpy}$	Tbytes/yr	$\backslash exttt{dst}$	DST		

583 C.7 Detector related

584 C.7.1 Detector technologies

'	$\setminus \mathtt{nonn}$	n^+ -on- n	\setminus ponn	p^+ -on- n	\setminus nonp	n^+ -on- p
585 _\	\cvd	CVD	$\backslash \mathtt{mwpc}$	MWPC	$\backslash \mathtt{gem}$	GEM

⁵⁸⁶ C.7.2 Detector components, electronics

,	ackslashtell1	$\mathrm{TELL1}$	\ukl1	UKL1	\beetle	Beetle
,	ackslashotis	OTIS	\croc	CROC	\carioca	CARIOCA
,	$ackslash ext{dialog}$	DIALOG	\setminus sync	SYNC	\cardiac	CARDIAC
,	\gol	GOL	\vcsel	VCSEL	\ttc	TTC
,	ackslash ttcrx	TTCrx	\hpd	HPD	\pmt	PMT
587	\specs	SPECS	\elmb	ELMB	\fpga	FPGA
,	\plc	PLC	$\backslash { t rasnik}$	RASNIK	\elmb	ELMB
,	\setminus can	CAN	\lvds	LVDS	\ntc	NTC
,	\adc	ADC	$\backslash ext{led}$	LED	\ccd	CCD
,	\hv	HV	\lv	LV	\pvss	PVSS
,	$\backslash \mathtt{cmos}$	CMOS	\fifo	FIFO	\ccpc	CCPC

⁵⁸ C.7.3 Chemical symbols

590 C.8 Special Text

592 D Supplementary material for LHCb-PAPER-20XX-593 YYY

This appendix contains supplementary material that will posted on the public cds record but will not appear in the paper.

594


595

597

598

599

Please leave the above sentence in your draft for first and second circulation and replace what follows by your actual supplementary material. For more information about other types of supplementary material, see Section 8. Plots and tables that follow should be well described, either with captions or with additional explanatory text.



 $\varphi_s^{J/\psi\varphi}$ (rad) Figure 2: Comparison of our result to those from other experiments. Note that the style of this figure differs slightly from that of Figure 1

\sim References

- [1] LHCb collaboration, A. A. Alves Jr. et al., The LHCb detector at the LHC, JINST 3 (2008) S08005.
- [2] American Physical Society, APS REVTEX package, https://journals.aps.org/revtex.
- [3] Particle Data Group, K. A. Olive et al., Review of particle physics, Chin. Phys. C38 (2014) 090001, and 2015 update.
- [4] LHCb collaboration, R. Aaij et al., LHCb detector performance, Int. J. Mod. Phys.
 A30 (2015) 1530022, arXiv:1412.6352.
- [5] R. Aaij *et al.*, *Performance of the LHCb Vertex Locator*, JINST **9** (2014) P09007, arXiv:1405.7808.
- [6] R. Arink et al., Performance of the LHCb Outer Tracker, JINST 9 (2014) P01002, arXiv:1311.3893.
- [7] M. Adinolfi et al., Performance of the LHCb RICH detector at the LHC, Eur. Phys. J. C73 (2013) 2431, arXiv:1211.6759.
- [8] A. A. Alves Jr. et al., Performance of the LHCb muon system, JINST 8 (2013) P02022, arXiv:1211.1346.
- [9] R. Aaij et al., The LHCb trigger and its performance in 2011, JINST 8 (2013) P04022, arXiv:1211.3055.
- 618 [10] A. Puig, The LHCb trigger in 2011 and 2012, LHCb-PUB-2014-046.
- [11] V. V. Gligorov and M. Williams, Efficient, reliable and fast high-level triggering using a bonsai boosted decision tree, JINST 8 (2013) P02013, arXiv:1210.6861.
- [12] LHCb collaboration, R. Aaij et al., Differential branching fractions and isospin asymmetries of $B \to K^{(*)} \mu^+ \mu^-$ decays, JHEP **06** (2014) 133, arXiv:1403.8044.
- [13] T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026, arXiv:hep-ph/0603175; T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, Comput. Phys. Commun. **178** (2008) 852, arXiv:0710.3820.
- [14] I. Belyaev et al., Handling of the generation of primary events in Gauss, the LHCb simulation framework, J. Phys. Conf. Ser. **331** (2011) 032047.
- [15] D. J. Lange, The EvtGen particle decay simulation package, Nucl. Instrum. Meth. A462 (2001) 152.

- [16] P. Golonka and Z. Was, *PHOTOS Monte Carlo: A precision tool for QED corrections* in Z and W decays, Eur. Phys. J. **C45** (2006) 97, arXiv:hep-ph/0506026.
- [17] Geant4 collaboration, J. Allison et al., Geant4 developments and applications, IEEE

 Trans. Nucl. Sci. **53** (2006) 270; Geant4 collaboration, S. Agostinelli et al., Geant4:

 A simulation toolkit, Nucl. Instrum. Meth. **A506** (2003) 250.
- [18] M. Clemencic et al., The LHCb simulation application, Gauss: Design, evolution and experience, J. Phys. Conf. Ser. **331** (2011) 032023.
- [19] L. Breiman, J. H. Friedman, R. A. Olshen, and C. J. Stone, *Classification and regression trees*, Wadsworth international group, Belmont, California, USA, 1984.
- [20] R. E. Schapire and Y. Freund, A decision-theoretic generalization of on-line learning
 and an application to boosting, J. Comput. Syst. Sci. 55 (1997) 119.
- [21] A. Feder, Your BibTeX resource, http://www.bibtex.org/.
- [22] E. Majorana, Teoria simmetrica dell'elettrone e del positrone, Nuovo Cim. **14** (1937) 171.
- [23] M. Shell, Mciteplus: Enhanced multicitations, http://www.michaelshell.org/tex/mciteplus/.
- [24] R. N. Mohapatra and G. Senjanovic, Neutrino Mass and Spontaneous Parity Violation, Phys. Rev. Lett. 44 (1980) 912; S. Pascoli and S. T. Petcov, Majorana neutrinos, neutrino mass spectrum and the $|\langle m \rangle| \sim 10^{-3}$ eV frontier in neutrinoless double beta decay, Phys. Rev. D77 (2008) 113003, arXiv:0711.4993.
- [25] LHCb collaboration, Optimization and calibration of the LHCb flavour tagging performance using 2010 data, LHCb-CONF-2011-003.
- [26] J. Dickens, A measurement of the photon efficiency from the 2010 data, LHCb-INT-2011-047; C. Adrover et al., Searches for $B_s^0 \to \mu^+\mu^-$ and $B^0 \to \mu^+\mu^-$ in 370 pb⁻¹ at LHCb, LHCb-ANA-2011-078; P. Owen, Measurement of branching fractions, isospin asymmetries and angular observables in exclusive electroweak penguin decays, CERN-THESIS-2014-057; P. Perret, First Years of Running for the LHCb Calorimeter system, LHCb-PROC-2014-017; U. Egede, Future of heavy flavour physics, LHCb-TALK-2014-257.
- [27] LHCb collaboration, R. Aaij et al., Measurement of charged particle multiplicities and densities in pp collisions at $\sqrt{s} = 7$ TeV in the forward region, Eur. Phys. J. C74 (2014) 2888, arXiv:1402.4430.
- [28] LHCb collaboration, R. Aaij et al., Observation of the decay $B_s^0 \to \overline{D}^0 \phi$, Phys. Lett. B727 (2013) 403, arXiv:1308.4583.

- [29] LHCb collaboration, R. Aaij et al., Observation of $J/\psi p$ resonances consistent with pentaquark states in $\Lambda_b^0 \to J/\psi p K^-$ decays, Phys. Rev. Lett. **115** (2015) 072001, arXiv:1507.03414.
- [30] Heavy Flavor Averaging Group, Y. Amhis et al., Averages of b-hadron, c-hadron,
 and τ-lepton properties as of summer 2014, arXiv:1412.7515, updated results and
 plots available at http://www.slac.stanford.edu/xorg/hfag/.
- [31] A. Tsaregorodtsev et al., DIRAC3: The new generation of the LHCb grid software,
 J. Phys. Conf. Ser. **219** (2010) 062029; R. Graciani Diaz et al., Belle-DIRAC setup
 for using Amazon Elastic Compute Cloud, Journal of Grid Computing **9** (2011) 65.
- [32] T. Skwarnicki, A study of the radiative cascade transitions between the Upsilon-prime and Upsilon resonances, PhD thesis, Institute of Nuclear Physics, Krakow, 1986, DESY-F31-86-02.
- [33] S. S. Wilks, The large-sample distribution of the likelihood ratio for testing composite hypotheses, Ann. Math. Stat. **9** (1938) 60.
- [34] W. D. Hulsbergen, *Decay chain fitting with a Kalman filter*, Nucl. Instrum. Meth. **A552** (2005) 566, arXiv:physics/0503191.
- [35] M. Pivk and F. R. Le Diberder, sPlot: A statistical tool to unfold data distributions, Nucl. Instrum. Meth. **A555** (2005) 356, arXiv:physics/0402083.
- [36] G. Punzi, Sensitivity of searches for new signals and its optimization, in Statistical Problems in Particle Physics, Astrophysics, and Cosmology (L. Lyons, R. Mount, and R. Reitmeyer, eds.), p. 79, 2003. arXiv:physics/0308063.
- [37] LHCb collaboration, R. Aaij et al., Measurement of the fragmentation fraction ratio f_s/f_d and its dependence on B meson kinematics, JHEP **04** (2013) 001, arXiv:1301.5286, f_s/f_d value updated in LHCb-CONF-2013-011.
- [38] R. Aaij et al., Performance of the LHCb calorimeters, LHCb-DP-2013-004, in preparation.
- [39] LHCb collaboration, R. Aaij et al., Measurement of the track reconstruction efficiency at LHCb, JINST 10 (2015) P02007, arXiv:1408.1251.
- [40] F. Archilli et al., Performance of the muon identification at LHCb, JINST 8 (2013) P10020, arXiv:1306.0249.
- [41] A. Affolder et al., Radiation damage in the LHCb vertex locator, JINST 8 (2013) P08002, arXiv:1302.5259.
- [42] D. van Eijk et al., Radiation hardness of the LHCb Outer Tracker, Nucl. Instrum.

 Meth. A685 (2012) 62.

- [43] R. B. Appleby et al., Simulation of machine induced background in the LHCb experiment: methodology and implementation, IEEE Trans. Nucl. Sci. **59** (2012) 1681.
- ⁷⁰² [44] M. Anelli et al., Performance of the LHCb muon system with cosmic rays, JINST 5
 ⁷⁰³ (2010) P10003, arXiv:1009.1963.
- [45] S. Borghi et al., First spatial alignment of the LHCb VELO and analysis of beam absorber collision data, Nucl. Instrum. Meth. A618 (2010) 108.
- ⁷⁰⁶ [46] LHCb collaboration, *LHCb Trigger and Online Technical Design Report*, CERN-⁷⁰⁷ LHCC-2014-016. LHCb-TDR-016.
- ⁷⁰⁸ [47] LHCb collaboration, *LHCb Tracker Upgrade Technical Design Report*, CERN-LHCC-⁷⁰⁹ 2014-001. LHCb-TDR-015.
- ⁷¹⁰ [48] LHCb collaboration, *LHCb PID Upgrade Technical Design Report*, CERN-LHCC-⁷¹¹ 2013-022. LHCb-TDR-014.
- [49] LHCb collaboration, *LHCb VELO Upgrade Technical Design Report*, CERN-LHCC-2013-021. LHCb-TDR-013.
- ⁷¹⁴ [50] LHCb collaboration, Framework TDR for the LHCb Upgrade: Technical Design Report, CERN-LHCC-2012-007. LHCb-TDR-012.
- [51] LHCb collaboration, *LHCb computing: Technical Design Report*, CERN-LHCC-2005-019. LHCb-TDR-011.
- [52] LHCb collaboration, *LHCb trigger system: Technical Design Report*, CERN-LHCC-2003-031. LHCb-TDR-010.
- [53] LHCb collaboration, *LHCb reoptimized detector design and performance: Technical Design Report*, CERN-LHCC-2003-030. LHCb-TDR-009.
- [54] LHCb collaboration, *LHCb inner tracker: Technical Design Report*, CERN-LHCC-2002-029. LHCb-TDR-008.
- [55] LHCb collaboration, *LHCb online system, data acquisition and experiment control:*Technical Design Report, CERN-LHCC-2001-040. LHCb-TDR-007.
- [56] LHCb collaboration, *LHCb outer tracker: Technical Design Report*, CERN-LHCC-227 2001-024. LHCb-TDR-006.
- [57] LHCb collaboration, LHCb VELO (VErtex LOcator): Technical Design Report, CERN-LHCC-2001-011. LHCb-TDR-005.
- [58] LHCb collaboration, *LHCb muon system: Technical Design Report*, CERN-LHCC-2001-010. LHCb-TDR-004.

- [59] LHCb collaboration, LHCb RICH: Technical Design Report, CERN-LHCC-2000-037.
 LHCb-TDR-003.
- [60] LHCb collaboration, *LHCb calorimeters: Technical Design Report*, CERN-LHCC-2000-036. LHCb-TDR-002.
- ⁷³⁶ [61] LHCb collaboration, *LHCb magnet: Technical Design Report*, CERN-LHCC-2000-⁷³⁷ 007. LHCb-TDR-001.
- [62] LHCb collaboration, R. Aaij et al., Search for Higgs-like boson decaying into pair of long-lived particles, LHCb-PAPER-2016-014, in preparation.
- [63] LHCb collaboration, R. Aaij et al., Measurement of the CP violating asymmetry a_{sl}^s , LHCb-PAPER-2016-013, in preparation.
- [64] LHCb collaboration, R. Aaij et al., The differential branching fraction of $B^0 \rightarrow K^*(892)^0 \mu^+ \mu^-$ and S-wave fraction in $B^0 \rightarrow K^+ \pi^- \mu^+ \mu^-$ decays, LHCb-PAPER-2016-012, in preparation.
- [65] LHCb collaboration, R. Aaij et al., Measurement of forward W and Z boson production in association with jets in proton-proton collisions at $\sqrt{s}=8~TeV$, arXiv:1605.00951, submitted to JHEP.
- [66] LHCb collaboration, R. Aaij et al., Measurement of the properties of the Ξ_b^{*0} baryon, arXiv:1604.03896, submitted to JHEP.
- [67] LHCb collaboration, R. Aaij et al., Model-independent evidence for $J/\psi p$ contributions to $\Lambda_b \to J/\psi p K^-$ decays, arXiv:1604.05708, submitted to Phys. Rev. Lett.
- [68] LHCb collaboration, R. Aaij et al., Measurements of the mass and lifetime of the Ω_b^- baryon, arXiv:1603.01412, to appear in Phys. Rev. D.
- [69] LHCb collaboration, R. Aaij et al., Measurement of the CKM angle γ using $B^0 \rightarrow DK^{*0}$ with $D \rightarrow K_S^0 \pi^+ \pi^-$ decays, arXiv:1605.01082, submitted to JHEP.
- [70] LHCb collaboration, R. Aaij et al., Model-independent measurement of the CKM angle γ using $B^0 \to DK^{*0}$ decays with $D \to K_S^0 \pi^+ \pi^-$ and $K_S^0 K^+ K^-$, arXiv:1604.01525, submitted to JHEP.
- [71] LHCb collaboration, R. Aaij et al., Search for violations of Lorentz invariance and CPT symmetry in $B_{(s)}^0$ mixing, arXiv:1603.04804, submitted to Phys. Rev. Lett.
- [72] LHCb collaboration, R. Aaij et al., Observations of $\Lambda_b^0 \to \Lambda K^+\pi^-$ and $\Lambda_b^0 \to \Lambda K^+K^-$ decays and searches for other Λ_b^0 and Ξ_b^0 decays to Λh^+h^- final states, arXiv:1603.00413, to appear in JHEP.

- [73] LHCb collaboration, R. Aaij et al., Measurement of CP observables in $B^{\pm} \rightarrow DK^{\pm}$ and $B^{\pm} \rightarrow D\pi^{\pm}$ with two- and four-body D meson decays, arXiv:1603.08993, submitted to Phys. Lett. B.
- ⁷⁶⁸ [74] LHCb collaboration, R. Aaij *et al.*, Observation of the $\Lambda_b \to \Lambda \phi$ decay, arXiv:1603.02870, submitted to Phys. Lett. B.
- [75] LHCb collaboration, R. Aaij et al., Search for B_c decays to the $p\bar{p}\pi$ final state, arXiv:1603.07037, submitted to Phys. Lett. B.
- [76] LHCb collaboration, R. Aaij et al., Observation of the $\Lambda_b^0 \to \psi(2S)pK^-$ decay, arXiv:1603.06961, submitted to JHEP.
- [77] LHCb collaboration, R. Aaij et al., Constraints on the unitarity triangle angle γ from Dalitz plot analysis of $B^0 \to DK^+\pi^-$ decays, arXiv:1602.03455, submitted to Phys. Rev. D.
- [78] LHCb collaboration, R. Aaij et al., Study of $\psi(2S)$ production cross-sections and cold nuclear matter effects in pPb collisions at $\sqrt{s_{\scriptscriptstyle NN}}=5~TeV$, arXiv:1601.07878, to appear in JHEP.
- [79] LHCb collaboration, R. Aaij et al., First observation of $D^0 \bar{D}^0$ oscillations in $D^0 \to K^+\pi^+\pi^-\pi^-$ decays and a measurement of the associated coherence parameters, arXiv:1602.07224, submitted to Phys. Rev. Lett.
- [80] LHCb collaboration, R. Aaij et al., Neural-network-based same side kaon tagging algorithm calibrated with $B_s^0 \to D_s^- \pi^+$ and $B_{s2}^* (5840)^0 \to B^+ K^-$ decays, arXiv:1602.07252, submitted to J. Inst.
- [81] LHCb collaboration, R. Aaij et al., Measurement of the difference of time-integrated asymmetries in $D^0 \to K^-K^+$ and $D^0 \to \pi^-\pi^+$ decays, arXiv:1602.03160, to appear in Phys. Rev. Lett.
- ⁷⁸⁹ [82] LHCb collaboration, R. Aaij et al., First observation of the rare $B^+ \to D^+ K^+ \pi^-$ ⁷⁹⁰ decay, Phys. Rev. **D93** (2015) 051101(R), arXiv:1512.02494.
- [83] LHCb collaboration, R. Aaij et al., Measurement of the inclusive $B_s^0 \to D_s^{(*)+} D_s^{(*)-}$ branching fraction, arXiv:1602.07543, to appear in Phys. Rev. D.
- ⁷⁹³ [84] LHCb collaboration, R. Aaij et al., Study of D_{sJ}^+ mesons decaying to $D^{*+}K_S^0$ and $D^{*0}K^+$ final states, JHEP **02** (2015) 133, arXiv:1601.01495.
- ⁷⁹⁵ [85] LHCb collaboration, R. Aaij et al., Angular analysis of the $B^0 \to K^{*0}\mu^+\mu^-$ decay using 3 fb⁻¹ of integrated luminosity, JHEP **02** (2016) 104, arXiv:1512.04442.
- [86] LHCb collaboration, R. Aaij et al., Observation of $B^0_s \to \overline{D}^0 K^0_S$ and evidence for $B^0_s \to \overline{D}^{*0} K^0_S$ decays, Phys. Rev. Lett. **116** (2016) 161802, arXiv:1603.02408.

- [87] LHCb collaboration, R. Aaij et al., Measurement of forward W and Z boson production in pp collisions at $\sqrt{s} = 8$ TeV, JHEP **01** (2015) 155, arXiv:1511.08039.
- [88] LHCb collaboration, R. Aaij et al., Search for the lepton-flavour violating decay $D^0 \to e^{\pm}\mu^{\mp}$, Phys. Lett. **B754** (2015) 167, arXiv:1512.00322.
- [89] LHCb collaboration, R. Aaij et al., Evidence for the strangeness-changing weak decay $\Xi_b^- \to \Lambda_b^0 \pi^-$, Phys. Rev. Lett. **115** (2015) 241801, arXiv:1510.03829.
- [90] LHCb collaboration, R. Aaij et al., Production of associated Υ and open charm hadrons in pp collisions at $\sqrt{s}=7$ and 8 TeV via double parton scattering, arXiv:1510.05949, submitted to JHEP.
- [91] LHCb collaboration, R. Aaij et al., Forward production of Υ mesons in pp collisions at $\sqrt{s} = 7$ and 8 TeV, JHEP 11 (2015) 103, arXiv:1509.02372.
- 810 [92] LHCb collaboration, R. Aaij et al., Search for the decays $B^0 \to J/\psi \gamma$ and $B_s^0 \to J/\psi \gamma$ 811 , Phys. Rev. **D92** (2015) 112002, arXiv:1510.04866.
- [93] LHCb collaboration, R. Aaij et al., First observation of the decay $D^0 \to K^-\pi^+\mu^+\mu^$ in the $\rho^0 - \omega$ region of the dimuon mass spectrum, arXiv:1510.08367, submitted to Phys.Lett.B.
- Elso [94] LHCb collaboration, R. Aaij et al., Model-independent measurement of mixing parameters in $D^0 \to K_S^0 \pi^+ \pi^-$ decays, JHEP **04** (2016) 033, arXiv:1510.01664.
- Elso [95] LHCb collaboration, R. Aaij et al., Measurements of prompt charm production cross-sections in pp collisions at $\sqrt{s}=13~TeV$, JHEP **03** (2016) 159, arXiv:1510.01707.
- [96] LHCb collaboration, R. Aaij et al., Measurement of two-particle correlations in proton-ion collisions at $\sqrt{s_{\scriptscriptstyle NN}}=5~TeV$, arXiv:1512.00439, submitted to Phys.Lett.B.
- [97] LHCb collaboration, R. Aaij et al., Measurement of the forward-backward asymmetry in $Z/\gamma^* \to \mu^+\mu^-$ decays and determination of the effective weak mixing angle, JHEP 11 (2015) 190, arXiv:1509.07645.
- [98] LHCb collaboration, R. Aaij et al., A model-independent confirmation of the $Z(4430)^-$ state, Phys. Rev. **D92** (2015) 112009, arXiv:1510.01951.
- [99] LHCb collaboration, R. Aaij et al., Measurement of forward J/ψ production cross-sections in pp collisions at $\sqrt{s}=13$ TeV, JHEP **10** (2015) 172, arXiv:1509.00771.
- [100] LHCb collaboration, R. Aaij et al., Search for hidden-sector bosons in $B^0 \to K^{*0} \mu^+ \mu^$ decays, Phys. Rev. Lett. **115** (2015) 161802, arXiv:1508.04094.

- [101] LHCb collaboration, R. Aaij et al., First measurement of the differential branching fraction and CP asymmetry of the $B^+ \to \pi^+ \mu^+ \mu^-$ decay, JHEP **10** (2015) 034, arXiv:1509.00414.
- [102] LHCb collaboration, R. Aaij et al., Measurement of CP violation parameters and polarisation fractions in $B_s^0 \to J/\psi \overline{K}^{*0}$ decays, JHEP **11** (2015) 082, arXiv:1509.00400.
- [103] LHCb collaboration, R. Aaij et al., Observation of the $B_s \to J/\psi \phi \phi$ decay, JHEP 03 (2016) 040, arXiv:1601.05284.
- [104] LHCb collaboration, R. Aaij et al., Study of the productions of Λ_b^0 and \bar{B}^0 hadrons in pp collisions and first measurement of the $\Lambda_b^0 \to J/\psi p K^-$ branching fraction, Chin. Phys. C **40** (2015) 011001, arXiv:1509.00292.
- ⁸⁴² [105] LHCb collaboration, R. Aaij et al., Measurement of the B^0 oscillation frequency Δm_d with $B^0 \to D^{(*)-} \mu^+ \nu_\mu$, LHCb-PAPER-2015-031, in preparation.
- [106] LHCb collaboration, R. Aaij et al., Measurement of the time-integrated CP asymmetry in $D^0 \to K_S^0 K_S^0$ decays, JHEP 10 (2015) 055, arXiv:1508.06087.
- LHCb collaboration, R. Aaij et al., Measurement of the $B_s^0 \to \phi \phi$ branching fraction and search for the decay $B^0 \to \phi \phi$, JHEP 10 (2015) 053, arXiv:1508.00788.
- ⁸⁴⁸ [108] LHCb collaboration, R. Aaij et al., B flavour tagging using charm decays at the LHCb experiment, JINST **10** (2015) P10005, arXiv:1507.07892.
- [109] LHCb collaboration, R. Aaij et al., Studies of the resonance structure in $D^0 \rightarrow K_S^0 K^{\pm} \pi^{\mp}$ decays, Phys. Rev. **D93** (2016) 052018, arXiv:1509.06628.
- [110] LHCb collaboration, R. Aaij et al., Measurement of the ratio of branching fractions $\mathcal{B}(\overline{B}^0 \to D^{*+}\tau^-\overline{\nu}_{\tau})/\mathcal{B}(\overline{B}^0 \to D^{*+}\mu^-\overline{\nu}_{\mu})$, Phys. Rev. Lett. **115** (2015) 111803, arXiv:1506.08614.
- ESS [111] LHCb collaboration, R. Aaij et al., Measurement of the branching fraction ratio $\mathcal{B}(B_c^+ \to \psi(2S)\pi^+)/\mathcal{B}(B_c^+ \to J/\psi\pi^+)$, Phys. Rev. **D92** (2015) 057007, arXiv:1507.03516.
- ESS [112] LHCb collaboration, R. Aaij et al., Angular analysis and differential branching fraction of the decay $B_s^0 \to \phi \mu^+ \mu^-$, JHEP **09** (2015) 179, arXiv:1506.08777.
- [113] LHCb collaboration, R. Aaij et al., First observation of top quark production in the forward region, Phys. Rev. Lett. 115 (2015) 112001, arXiv:1506.00903.
- EXECUTE: 114] LHCb collaboration, R. Aaij et al., Study of W boson production in association with beauty and charm, Phys. Rev. **D92** (2015) 052012, arXiv:1505.04051.

- [115] LHCb collaboration, R. Aaij et al., Study of $B^- \to DK^-\pi^+\pi^-$ and $B^- \to D\pi^-\pi^+\pi^$ decays and determination of the CKM angle γ , Phys. Rev. **D92** (2015) 112005, arXiv:1505.07044.
- [116] LHCb collaboration, R. Aaij et al., Search for the $\Lambda_b^0 \to \Lambda \eta$ and $\Lambda_b^0 \to \Lambda \eta'$ decays with the LHCb detector, JHEP **09** (2015) 006, arXiv:1505.03295.
- [117] LHCb collaboration, R. Aaij et al., First observation of the decay $B_s^0 \to K_S^0 K^*(892)^0$, JHEP **01** (2015) 012, arXiv:1506.08634.
- [118] LHCb collaboration, R. Aaij et al., Amplitude analysis of $B^0 \to \overline{D}^0 K^+ \pi^-$ decays, Phys. Rev. **D92** (2015) 012012, arXiv:1505.01505.
- [119] LHCb collaboration, R. Aaij et al., Identification of beauty and charm quark jets at LHCb, JINST 10 (2015) P06013, arXiv:1504.07670.
- ETS [120] LHCb collaboration, R. Aaij et al., Quantum numbers of the X(3872) state and orbital angular momentum in its $\rho^0 J/\psi$ decays, Phys. Rev. **D92** (2015) 011102(R), arXiv:1504.06339.
- [121] LHCb collaboration, R. Aaij et al., A study of CP violation in $B^{\mp} \to Dh^{\mp}$ ($h = K, \pi$) with the modes $D \to K^{\mp}\pi^{\pm}\pi^{0}$, $D \to \pi^{+}\pi^{-}\pi^{0}$ and $D \to K^{+}K^{-}\pi^{0}$, Phys. Rev. **D91** (2015) 112014, arXiv:1504.05442.
- [122] LHCb collaboration, R. Aaij et al., Determination of the quark coupling strength $|V_{ub}|$ using baryonic decays, Nature Physics 11 (2015) 743, arXiv:1504.01568.
- ⁸⁸³ [123] LHCb collaboration, R. Aaij *et al.*, Search for the decay $B_s^0 \to \overline{D}^0 f_0(980)$, JHEP **08** (2015) 005, arXiv:1505.01654.
- ENSE [124] LHCb collaboration, R. Aaij et al., Measurement of the exclusive $\Upsilon(nS)$ production cross-section in pp collisions at $\sqrt{s}=7$ TeV and 8 TeV, JHEP **09** (2015) 084, arXiv:1505.08139.
- ESSE [125] LHCb collaboration, R. Aaij et al., Observation of the decay $\overline{B}^0_s \to \psi(2S)K^+\pi^-$, Phys. Lett. B747 (2015) 484, arXiv:1503.07112.
- ⁸⁹⁰ [126] LHCb collaboration, R. Aaij et al., Differential branching fraction and angular analysis of $\Lambda_b^0 \to \Lambda \mu^+ \mu^-$ decays, JHEP **06** (2015) 115, arXiv:1503.07138.
- E92 [127] LHCb collaboration, R. Aaij et al., First observation and measurement of the branching fraction for the decay $B_s^0 \to D_s^{*\mp} K^\pm$, JHEP **06** (2015) 130, arXiv:1503.09086.
- Eq. [128] LHCb collaboration, R. Aaij et al., First observation and amplitude analysis of the $B^- \to D^+ K^- \pi^-$ decay, Phys. Rev. **D91** (2015) 092002, arXiv:1503.02995.

- ESSE [129] LHCb collaboration, R. Aaij et al., Observation of the $B^0 \to \rho^0 \rho^0$ decay from an amplitude analysis of $B^0 \to (\pi^+\pi^-)(\pi^+\pi^-)$ decays, Phys. Lett. **B747** (2015) 468, arXiv:1503.07770.
- [130] LHCb collaboration, R. Aaij et al., Measurement of the time-dependent CP asymmetries in $B_s^0 \to J/\psi K_S^0$, JHEP **06** (2015) 131, arXiv:1503.07055.
- [131] LHCb collaboration, R. Aaij et al., Measurement of CP violation in $B^0 \to J/\psi K_S^0$ decays, Phys. Rev. Lett. **115** (2015) 031601, arXiv:1503.07089.
- ⁹⁰³ [132] LHCb collaboration, R. Aaij et al., Measurement of $Z \rightarrow e^+e^-$ production at $\sqrt{s} = 8$ ⁹⁰⁴ TeV, JHEP **05** (2015) 109, arXiv:1503.00963.
- [133] LHCb collaboration, R. Aaij et al., Search for long-lived heavy charged particles
 using a ring-imaging Cherenkov technique at LHCb, Eur. Phys. J. C75 (2015) 595,
 arXiv:1506.09173.
- [134] LHCb collaboration, R. Aaij et al., Measurement of the forward Z boson cross-section in pp collisions at $\sqrt{s} = 7$ TeV, JHEP **08** (2015) 039, arXiv:1505.07024.
- [135] LHCb collaboration, R. Aaij et al., Dalitz plot analysis of $B^0 \to \overline{D}^0 \pi^+ \pi^-$ decays, Phys. Rev. **D92** (2015) 032002, arXiv:1505.01710.
- [136] LHCb collaboration, R. Aaij et al., Measurement of indirect CP asymmetries in $D^0 \to K^-K^+$ and $D^0 \to \pi^-\pi^+$ decays, JHEP **04** (2015) 043, arXiv:1501.06777.
- [137] LHCb collaboration, R. Aaij et al., Measurement of CP asymmetries and polarisation fractions in $B_s^0 \to K^{*0} \overline{K}^{*0}$ decays, JHEP **07** (2015) 166, arXiv:1503.05362.
- ⁹¹⁶ [138] LHCb collaboration, R. Aaij et al., Precise measurements of the properties of the $B_1(5721)^{0,+}$ and $B_2^*(5747)^{0,+}$ states and observation of structure at higher invariant mass in the $B^+\pi^-$ and $B^0\pi^+$ spectra, JHEP **04** (2015) 024, arXiv:1502.02638.
- [139] LHCb collaboration, R. Aaij et al., Angular analysis of the $B^0 \to K^{*0}e^+e^-$ decay in the low-q² region, JHEP **04** (2015) 024, arXiv:1501.03038.
- [140] LHCb collaboration, R. Aaij et al., Observation of the $B_s^0 \to \eta' \eta'$ decay, Phys. Rev. Lett. 115 (2015) 051801, arXiv:1503.07483.
- [141] LHCb collaboration, R. Aaij et al., Determination of the branching fractions of $B_s^0 \to D_s^{\mp} K^{\pm}$ and $B^0 \to D_s^{\mp} K^{+}$, JHEP **02** (2015) 029, arXiv:1412.7654.
- LHCb collaboration, R. Aaij et al., Study of the rare B_s^0 and B^0 decays into the $\pi^+\pi^-\mu^+\mu^-$ final state, Phys. Lett. **B743** (2015) 46, arXiv:1412.6433.
- ⁹²⁷ [143] LHCb collaboration, R. Aaij et al., Search for long-lived particles decaying to jet pairs, Eur. Phys. J. C75 (2014) 152, arXiv:1412.3021.

- [144] LHCb collaboration, R. Aaij et al., Observation of two new Ξ_b^- baryon resonances, Phys. Rev. Lett. **114** (2015) 062004, arXiv:1411.4849.
- [145] LHCb collaboration, R. Aaij et al., Measurement of the lifetime of the B_c^+ meson using the $B_c^+ \to J/\psi \pi^+$ decay mode, Phys. Lett. **B742** (2015) 29, arXiv:1411.6899.
- [146] LHCb collaboration, R. Aaij et al., Precision measurement of CP violation in $B_s^0 \to J/\psi K^+K^-$ decays, Phys. Rev. Lett. **114** (2015) 041801, arXiv:1411.3104.
- [147] LHCb collaboration, R. Aaij et al., Measurement of the CP-violating phase β in $\overline{B}^0 \to J/\psi \pi^+ \pi^-$ decays and limits on penguin effects, Phys. Lett. **B742** (2015) 38, arXiv:1411.1634.
- ⁹³⁸ [148] LHCb collaboration, R. Aaij et al., Measurement of the inelastic pp cross-section at a centre-of-mass energy of $\sqrt{s} = 7$ TeV, JHEP **02** (2014) 029, arXiv:1412.2500.
- [149] LHCb collaboration, R. Aaij et al., Study of η - η' mixing from measurement of $B^0_{(s)} \to J/\psi \eta^{(\prime)}$ decay rates, JHEP **01** (2015) 024, arXiv:1411.0943.
- [150] LHCb collaboration, R. Aaij et al., Measurement of the Z+b-jet cross-section in pp collisions at $\sqrt{s}=7$ TeV in the forward region, JHEP **01** (2015) 064, arXiv:1411.1264.
- ⁹⁴⁵ [151] LHCb collaboration, R. Aaij et al., Search for CP violation in $D^0 \to \pi^-\pi^+\pi^0$ decays with the energy test, Phys. Lett. **B740** (2015) 158, arXiv:1410.4170.
- LHCb collaboration, R. Aaij et al., Measurement of the semileptonic CP asymmetry in $B^0-\overline{B}^0$ mixing, Phys. Rev. Lett. **114** (2015) 041601, arXiv:1409.8586.
- ⁹⁴⁹ [153] LHCb collaboration, R. Aaij et al., Search for the lepton flavour violating decay $\tau^- \to \mu^- \mu^+ \mu^-$, JHEP **02** (2015) 121, arXiv:1409.8548.
- [154] LHCb collaboration, R. Aaij et al., Measurement of the CP-violating phase ϕ_s in $\overline{B}_s^0 \to D_s^+ D_s^-$ decays, Phys. Rev. Lett. 113 (2014) 211801, arXiv:1409.4619.
- ⁹⁵³ [155] LHCb collaboration, R. Aaij et al., Measurement of B_c^+ production at $\sqrt{s}=8$ TeV, Phys. Rev. Lett. **114** (2014) 132001, arXiv:1411.2943.
- ⁹⁵⁵ [156] CMS and LHCb collaborations, V. Khachatryan *et al.*, Observation of the rare $B_s^0 \to \mu^+\mu^-$ decay from the combined analysis of CMS and LHCb data, Nature **522** (2015) 68, arXiv:1411.4413.
- LHCb collaboration, R. Aaij et al., Precision measurement of the mass and lifetime of the Ξ_b^- baryon, Phys. Rev. Lett. **113** (2014) 242002, arXiv:1409.8568.
- [158] LHCb collaboration, R. Aaij et al., Precision luminosity measurements at LHCb,
 JINST 9 (2014) P12005, arXiv:1410.0149.

- ⁹⁶² [159] LHCb collaboration, R. Aaij et al., Search for CP violation using T-odd correlations ⁹⁶³ in $D^0 \to K^+K^-\pi^+\pi^-$ decays, JHEP **10** (2014) 005, arXiv:1408.1299.
- ⁹⁶⁴ [160] LHCb collaboration, R. Aaij et al., Determination of γ and $-2\beta_s$ from charmless two-body decays of beauty mesons, Phys. Lett. **B739** (2014) 1, arXiv:1408.4368.
- [161] LHCb collaboration, R. Aaij et al., Measurement of CP violation in the three-body phase space of charmless B^{\pm} decays, Phys. Rev. **D90** (2014) 112004, arXiv:1408.5373.
- [162] LHCb collaboration, R. Aaij et al., Observation of $B^0_s \to K^{*\pm}K^{\mp}$ and evidence of $B^0_s \to K^{*-}\pi^+$ decays, New J. Phys. **16** (2014) 123001, arXiv:1407.7704.
- [163] LHCb collaboration, R. Aaij et al., Measurement of the \overline{B}^0 - B^0 and \overline{B}^0_s - B^0_s production asymmetries in pp collisions at $\sqrt{s}=7$ TeV, Phys. Lett. **B739** (2014) 218, arXiv:1408.0275.
- [164] LHCb collaboration, R. Aaij et al., Measurement of the CKM angle γ using $B^{\pm} \rightarrow DK^{\pm}$ with $D \rightarrow K_S^0 \pi^+ \pi^-$, $K_S^0 K^+ K^-$ decays, JHEP **10** (2014) 097, arXiv:1408.2748.
- ⁹⁷⁷ [165] LHCb collaboration, R. Aaij et al., Measurement of the $\chi_b(3P)$ mass and of the rela-⁹⁷⁸ tive rate of $\chi_{b1}(1P)$ and $\chi_{b2}(1P)$ production, JHEP **10** (2014) 088, arXiv:1409.1408.
- [166] LHCb collaboration, R. Aaij et al., First observation of a baryonic B_c^+ decay, Phys. Rev. Lett. **113** (2014) 152003, arXiv:1408.0971.
- ⁹⁸¹ [167] LHCb collaboration, R. Aaij et al., Measurement of CP asymmetry in $B_s^0 \to D_s^{\mp} K^{\pm}$ ⁹⁸² decays, JHEP **11** (2014) 060, arXiv:1407.6127.
- ⁹⁸³ [168] LHCb collaboration, R. Aaij et al., Measurement of the \overline{B}_s^0 meson lifetime in $D_s^+\pi^-$ ⁹⁸⁴ decays, Phys. Rev. Lett. **113** (2014) 172001, arXiv:1407.5873.
- ⁹⁸⁵ [169] LHCb collaboration, R. Aaij et al., Dalitz plot analysis of $B_s^0 \to \overline{D}^0 K^- \pi^+$ decays, Phys. Rev. **D90** (2014) 072003, arXiv:1407.7712.
- [170] LHCb collaboration, R. Aaij et al., Observation of overlapping spin-1 and spin-3 \overline{D}^0K^- resonances at mass 2.86 GeV/c^2 , Phys. Rev. Lett. **113** (2014) 162001, arXiv:1407.7574.
- ⁹⁹⁰ [171] LHCb collaboration, R. Aaij et al., Evidence for CP violation in $B^+ \to p\overline{p}K^+$ decays, Phys. Rev. Lett. **113** (2014) 141801, arXiv:1407.5907.
- ⁹⁹² [172] LHCb collaboration, R. Aaij et al., Measurement of the forward W boson production ⁹⁹³ cross-section in pp collisions at $\sqrt{s} = 7$ TeV, JHEP **12** (2014) 079, arXiv:1408.4354.

- ⁹⁹⁴ [173] LHCb collaboration, R. Aaij et al., Measurement of CP asymmetries in the decays $B^0 \to K^{*0} \mu^+ \mu^-$ and $B^+ \to K^+ \mu^+ \mu^-$, JHEP **09** (2014) 177, arXiv:1408.0978.
- [174] LHCb collaboration, R. Aaij et al., Study of χ_b meson production in pp collisions at \sqrt{s} =7 and 8 TeV and observation of the decay $\chi_b \to \Upsilon(3S)\gamma$, Eur. Phys. J. C74 (2014) 3092, arXiv:1407.7734.
- ⁹⁹⁹ [175] LHCb collaboration, R. Aaij et al., First observations of the rare decays $B^+ \rightarrow K^+\pi^+\pi^-\mu^+\mu^-$ and $B^+ \rightarrow \phi K^+\mu^+\mu^-$, JHEP **10** (2014) 064, arXiv:1408.1137.
- 1001 [176] LHCb collaboration, R. Aaij et al., Measurement of the $\eta_c(1S)$ production cross1002 section in proton-proton collisions via the decay $\eta_c(1S) \to p\overline{p}$, Eur. Phys. J. C75
 1003 (2015) 311, arXiv:1409.3612.
- ¹⁰⁰⁴ [177] LHCb collaboration, R. Aaij et al., Measurement of CP violation parameters in $B^0 \to DK^{*0}$ decays, Phys. Rev. **D90** (2014) 112002, arXiv:1407.8136.
- 1006 [178] LHCb collaboration, R. Aaij et al., Observation of charmonium pairs produced exclusively in pp collisions, J. Phys. **G41** (2014) 115002, arXiv:1407.5973.
- 1008 [179] LHCb collaboration, R. Aaij et al., Measurement of CP violation in $B_s^0 \to \phi \phi$ decays, Phys. Rev. **D90** (2014) 052011, arXiv:1407.2222.
- 1010 [180] LHCb collaboration, R. Aaij et al., Measurement of the ratio of B_c^+ branching 1011 fractions to $J/\psi\pi^+$ and $J/\psi\mu^+\nu_\mu$, Phys. Rev. **D90** (2014) 032009, arXiv:1407.2126.
- [181] LHCb collaboration, R. Aaij et al., Test of lepton universality using $B^+ \to K^+ \ell^+ \ell^$ loss decays, Phys. Rev. Lett. **113** (2014) 151601, arXiv:1406.6482.
- 1014 [182] LHCb collaboration, R. Aaij et al., First measurement of the charge asymmetry in beauty-quark pair production, Phys. Rev. Lett. 113 (2014) 082003, arXiv:1406.4789.
- 1016 [183] LHCb collaboration, R. Aaij et al., Observation of Z production in proton-lead collisions at LHCb, JHEP **09** (2014) 030, arXiv:1406.2885.
- [184] LHCb collaboration, R. Aaij et al., Precision measurement of the mass and lifetime of the Ξ_b^0 baryon, Phys. Rev. Lett. 113 (2014) 032001, arXiv:1405.7223.
- 1020 [185] LHCb collaboration, R. Aaij et al., Observation of the $\Lambda_b^0 \to J/\psi p\pi^-$ decay, JHEP 07 (2014) 103, arXiv:1406.0755.
- [186] LHCb collaboration, R. Aaij et al., Measurement of the CP-violating phase ϕ_s in $\overline{B}_s^0 \to J/\psi \pi^+ \pi^-$ decays, Phys. Lett. **B736** (2014) 186, arXiv:1405.4140.
- 1024 [187] LHCb collaboration, R. Aaij et al., Search for CP violation in $D^{\pm} \to K_S^0 K^{\pm}$ and 1025 $D_s^{\pm} \to K_S^0 \pi^{\pm}$ decays, JHEP **10** (2014) 025, arXiv:1406.2624.

- [188] LHCb collaboration, R. Aaij et al., Measurement of CP violation and constraints on the CKM angle γ in $B^{\pm} \rightarrow DK^{\pm}$ with $D \rightarrow K_S^0 \pi^+ \pi^-$ decays, Nucl. Phys. **B888** (2014) 169, arXiv:1407.6211.
- [189] LHCb collaboration, R. Aaij et al., Observation of the $B_s^0 \to J/\psi K_S^0 K^{\pm} \pi^{\mp}$ decay, JHEP **07** (2014) 140, arXiv:1405.3219.
- [190] LHCb collaboration, R. Aaij et al., Study of Υ production and cold nuclear effects in pPb collisions at $\sqrt{s_{NN}} = 5$ TeV, JHEP **07** (2014) 094, arXiv:1405.5152.
- 1033 [191] LHCb collaboration, R. Aaij et al., Observation of the resonant character of the $Z(4430)^-$ state, Phys. Rev. Lett. **112** (2014) 222002, arXiv:1404.1903.
- ¹⁰³⁵ [192] LHCb collaboration, R. Aaij et al., Measurement of CP asymmetry in $D^0 \to K^-K^+$ ¹⁰³⁶ and $D^0 \to \pi^-\pi^+$ decays, JHEP **07** (2014) 041, arXiv:1405.2797.
- [193] LHCb collaboration, R. Aaij et al., Measurement of the resonant and CP components in $\overline{B}^0 \to J/\psi \pi^+ \pi^-$ decays, Phys. Rev. **D90** (2014) 012003, arXiv:1404.5673.
- 1039 [194] LHCb collaboration, R. Aaij et al., Effective lifetime measurements in the $B_s^0 \to K^+K^-$, $B^0 \to K^+\pi^-$ and $B_s^0 \to \pi^+K^-$ decays, Phys. Lett. **B736** (2014) 446, arXiv:1406.7204.
- 1042 [195] LHCb collaboration, R. Aaij et al., Measurement of the Ξ_b^- and Ω_b^- baryon lifetimes, 1043 Phys. Lett. **B736** (2014) 154, arXiv:1405.1543.
- ¹⁰⁴⁴ [196] LHCb collaboration, R. Aaij *et al.*, Evidence for the decay $B_c^+ \to J/\psi 3\pi^+ 2\pi^-$, JHEP **05** (2014) 148, arXiv:1404.0287.
- 1046 [197] LHCb collaboration, R. Aaij et al., Evidence for the decay $X(3872) \rightarrow \psi(2S)\gamma$, Nucl. Phys. **B886** (2014) 665, arXiv:1404.0275.
- 1048 [198] LHCb collaboration, R. Aaij et al., Angular analysis of charged and neutral $B \rightarrow K\mu^+\mu^-$ decays, JHEP **05** (2014) 082, arXiv:1403.8045.
- 1050 [199] LHCb collaboration, R. Aaij et al., Measurement of polarization amplitudes and CP asymmetries in $B^0 \to \phi K^*(892)^0$, JHEP **05** (2014) 069, arXiv:1403.2888.
- [200] LHCb collaboration, R. Aaij et al., Study of the kinematic dependences of Λ_b^0 production in pp collisions and a measurement of the $\Lambda_b^0 \to \Lambda_c^+ \pi^-$ branching fraction, JHEP **08** (2014) 143, arXiv:1405.6842.
- [201] LHCb collaboration, R. Aaij et al., Precision measurement of the ratio of the Λ_b^0 to \overline{B}^0 lifetimes, Phys. Lett. **B734** (2014) 122, arXiv:1402.6242.
- ¹⁰⁵⁷ [202] LHCb collaboration, R. Aaij et al., Study of beauty hadron decays into pairs of charm hadrons, Phys. Rev. Lett. **112** (2014) 202001, arXiv:1403.3606.

- 1059 [203] LHCb collaboration, R. Aaij et al., Observation of photon polarization in the $b \to s\gamma$ 1060 transition, Phys. Rev. Lett. **112** (2014) 161801, arXiv:1402.6852.
- [204] LHCb collaboration, R. Aaij et al., Measurement of resonant and CP components in $\overline{B}_s^0 \to J/\psi \pi^+ \pi^-$ decays, Phys. Rev. **D89** (2014) 092006, arXiv:1402.6248.
- [205] LHCb collaboration, R. Aaij et al., A study of CP violation in $B^{\pm} \to DK^{\pm}$ and $B^{\pm} \to D\pi^{\pm}$ decays with $D \to K_S^0 K^{\pm} \pi^{\mp}$ final states, Phys. Lett. **B733** (2014) 36, arXiv:1402.2982.
- [206] LHCb collaboration, R. Aaij et al., Measurement of $\psi(2S)$ polarisation in pp collisions at $\sqrt{s}=7$ TeV, Eur. Phys. J. C74 (2014) 2872, arXiv:1403.1339.
- 1068 [207] LHCb collaboration, R. Aaij et al., Measurement of Υ production in pp collisions at $\sqrt{s} = 2.76$ TeV, Eur. Phys. J. C74 (2014) 2835, arXiv:1402.2539.
- 1070 [208] LHCb collaboration, R. Aaij et al., Measurements of the B^+ , B^0 , B_s^0 meson and Λ_b^0 baryon lifetimes, JHEP **04** (2014) 114, arXiv:1402.2554.
- [209] LHCb collaboration, R. Aaij et al., Search for Majorana neutrinos in $B^- \to \pi^+ \mu^- \mu^$ decays, Phys. Rev. Lett. **112** (2014) 131802, arXiv:1401.5361.
- 1074 [210] LHCb collaboration, R. Aaij et al., Measurement of the B_c^+ meson lifetime using $B_c^+ \to J/\psi \mu^+ \nu_\mu X$ decays, Eur. Phys. J. C74 (2014) 2839, arXiv:1401.6932.
- 1076 [211] LHCb collaboration, R. Aaij et al., Observation of associated production of a Z boson 1077 with a D meson in the forward region, JHEP **04** (2014) 091, arXiv:1401.3245.
- [212] LHCb collaboration, R. Aaij et al., Searches for Λ_b^0 and Ξ_b^0 decays to $K_S^0 p \pi^-$ and $K_S^0 p K^-$ final states with first observation of the $\Lambda_b^0 \to K_S^0 p \pi^-$ decay, JHEP **04** (2014) 087, arXiv:1402.0770.
- [213] LHCb collaboration, R. Aaij et al., Measurement of the $\overline{B}^0_s \to D_s^- D_s^+$ and $\overline{B}^0_s \to D_s^- D_s^+$ effective lifetimes, Phys. Rev. Lett. **112** (2014) 111802, arXiv:1312.1217.
- [214] LHCb collaboration, R. Aaij et al., Updated measurements of exclusive J/ψ and $\psi(2S)$ production cross-sections in pp collisions at $\sqrt{s}=7$ TeV, J. Phys. **G41** (2014) 055002, arXiv:1401.3288.
- 1086 [215] LHCb collaboration, R. Aaij et al., Study of forward Z+jet production in pp collisions at $\sqrt{s}=7$ TeV, JHEP **01** (2014) 033, arXiv:1310.8197.
- ¹⁰⁸⁸ [216] LHCb collaboration, R. Aaij et al., Search for CP violation in the decay $D^+ \rightarrow \pi^-\pi^+\pi^+$, Phys. Lett. **B728** (2014) 585, arXiv:1310.7953.
- ¹⁰⁹⁰ [217] LHCb collaboration, R. Aaij et al., Study of beauty baryon decays to D^0ph^- and $\Lambda_c^+h^-$ final states, Phys. Rev. **D89** (2014) 032001, arXiv:1311.4823.

- 1092 [218] LHCb collaboration, R. Aaij et al., Observation of $\overline{B}_{(s)}^0 \to J/\psi f_1(1285)$ decays and 1093 measurement of the $f_1(1285)$ mixing angle, Phys. Rev. Lett. **112** (2014) 091802, arXiv:1310.2145.
- [219] LHCb collaboration, R. Aaij et al., Measurements of indirect CP asymmetries in $D^0 \to K^-K^+$ and $D^0 \to \pi^-\pi^+$ decays, Phys. Rev. Lett. **112** (2014) 041801, arXiv:1310.7201.
- [220] LHCb collaboration, R. Aaij et al., Measurement of $D^0-\overline{D}^0$ mixing parameters and search for CP violation using $D^0 \to K^+\pi^-$ decays, Phys. Rev. Lett. **111** (2013) 251801, arXiv:1309.6534.
- 1101 [221] LHCb collaboration, R. Aaij et al., Study of J/ψ production and cold nuclear matter effects in pPb collisions at $\sqrt{s_{\scriptscriptstyle NN}}=5$ TeV, JHEP **02** (2014) 072, arXiv:1308.6729.
- [222] LHCb collaboration, R. Aaij et al., Measurement of CP violation in the phase space of $B^{\pm} \rightarrow K^+K^-\pi^{\pm}$ and $B^{\pm} \rightarrow \pi^+\pi^-\pi^{\pm}$ decays, Phys. Rev. Lett. **112** (2014) 011801, arXiv:1310.4740.
- 1106 [223] LHCb collaboration, R. Aaij *et al.*, Search for the decay $D^0 \to \pi^+\pi^-\mu^+\mu^-$, Phys. Lett. **B728** (2014) 234, arXiv:1310.2535.
- 1108 [224] LHCb collaboration, R. Aaij et al., Search for the doubly charmed baryon Ξ_{cc}^+ , JHEP 1109 12 (2013) 090, arXiv:1310.2538.
- 1110 [225] LHCb collaboration, R. Aaij et al., Measurement of the charge asymmetry in $B^{\pm} \rightarrow \phi K^{\pm}$ and search for $B^{\pm} \rightarrow \phi \pi^{\pm}$ decays, Phys. Lett. **B728** (2014) 85, arXiv:1309.3742.
- 1113 [226] LHCb collaboration, R. Aaij et al., Observation of the decay $B_c^+ \to J/\psi K^+ K^- \pi^+$, 1114 JHEP **11** (2013) 094, arXiv:1309.0587.
- [227] LHCb collaboration, R. Aaij et al., Measurement of the $B_s^0 \to \mu^+\mu^-$ branching fraction and search for $B^0 \to \mu^+\mu^-$ decays at the LHCb experiment, Phys. Rev. Lett. 1117 111 (2013) 101805, arXiv:1307.5024.
- 1118 [228] LHCb collaboration, R. Aaij et al., First observation of $\overline{B}^0 \to J/\psi K^+K^-$ and search for $\overline{B}^0 \to J/\psi \phi$ decays, Phys. Rev. **D88** (2013) 072005, arXiv:1308.5916.
- [229] LHCb collaboration, R. Aaij et al., Observation of the decay $B_c^+ \to B_s^0 \pi^+$, Phys. Rev. Lett. **111** (2013) 181801, arXiv:1308.4544.
- [230] LHCb collaboration, R. Aaij et al., Measurement of the CP asymmetry in $B^+ \rightarrow K^+\mu^+\mu^-$ decays, Phys. Rev. Lett. 111 (2013) 151801, arXiv:1308.1340.
- [231] LHCb collaboration, R. Aaij et al., Study of $B^0_{(s)} \to K^0_S h^+ h'^-$ decays with first observation of $B^0_s \to K^0_S K^\pm \pi^\mp$ and $B^0_s \to K^0_S \pi^+ \pi^-$, JHEP **10** (2013) 143, arXiv:1307.7648.

- [232] LHCb collaboration, R. Aaij et al., Model-independent search for CP violation in $D^0 \to K^-K^+\pi^+\pi^-$ and $D^0 \to \pi^-\pi^+\pi^-\pi^+$ decays, Phys. Lett. **B726** (2013) 623, arXiv:1308.3189.
- 1130 [233] LHCb collaboration, R. Aaij et al., First measurement of time-dependent CP violation 1131 in $B_s^0 \to K^+K^-$ decays, JHEP **10** (2013) 183, arXiv:1308.1428.
- 1132 [234] LHCb collaboration, R. Aaij et al., Observation of a resonance in $B^+ \to K^+ \mu^+ \mu^-$ 1133 decays at low recoil, Phys. Rev. Lett. **111** (2013) 112003, arXiv:1307.7595.
- [235] LHCb collaboration, R. Aaij et al., First evidence for the two-body charmless baryonic decay $B^0 \to p\overline{p}$, JHEP 10 (2013) 005, arXiv:1308.0961.
- [236] LHCb collaboration, R. Aaij et al., Measurement of form-factor-independent observables in the decay $B^0 \to K^{*0} \mu^+ \mu^-$, Phys. Rev. Lett. **111** (2013) 191801, arXiv:1308.1707.
- 1139 [237] LHCb collaboration, R. Aaij et al., Observation of $B_s^0 \overline{B}_s^0$ mixing and measurement of mixing frequencies using semileptonic B decays, Eur. Phys. J. C73 (2013) 2655, arXiv:1308.1302.
- 1142 [238] LHCb collaboration, R. Aaij et al., Branching fraction and CP asymmetry of the decays $B^+ \to K_S^0 \pi^+$ and $B^+ \to K_S^0 K^+$, Phys. Lett. **B726** (2013) 646, arXiv:1308.1277.
- 1145 [239] LHCb collaboration, R. Aaij et al., Measurement of the flavour-specific CP-violating asymmetry $a_{\rm sl}^s$ in B_s^0 decays, Phys. Lett. **B728** (2014) 607, arXiv:1308.1048.
- 1147 [240] LHCb collaboration, R. Aaij et al., Precision measurement of the Λ_b^0 baryon lifetime, 1148 Phys. Rev. Lett. **111** (2013) 102003, arXiv:1307.2476.
- 1149 [241] LHCb collaboration, R. Aaij et al., Studies of the decays $B^+ \to p\overline{p}h^+$ and observation of $B^+ \to \overline{\Lambda}(1520)p$, Phys. Rev. **D88** (2013) 052015, arXiv:1307.6165.
- [242] LHCb collaboration, R. Aaij et al., Search for the lepton-flavour-violating decays $B_s^0 \to e^{\pm}\mu^{\mp}$ and $B^0 \to e^{\pm}\mu^{\mp}$, Phys. Rev. Lett. **111** (2013) 141801, arXiv:1307.4889.
- 1154 [243] LHCb collaboration, R. Aaij et al., Searches for $B^0_{(s)} \to J/\psi p \overline{p}$ and $B^+ \to J/\psi p \overline{p} \pi^+$ 1155 decays, JHEP **09** (2013) 006, arXiv:1306.4489.
- 1156 [244] LHCb collaboration, R. Aaij et al., Measurement of the relative rate of prompt χ_{c0} , χ_{c1} and χ_{c2} production at $\sqrt{s}=7$ TeV, JHEP 10 (2013) 115, arXiv:1307.4285.
- 1158 [245] LHCb collaboration, R. Aaij et al., Measurement of CP violation in the phase space of $B^{\pm} \rightarrow K^{\pm}\pi^{+}\pi^{-}$ and $B^{\pm} \rightarrow K^{\pm}K^{+}K^{-}$ decays, Phys. Rev. Lett. **111** (2013) 101801, arXiv:1306.1246.

- 1161 [246] LHCb collaboration, R. Aaij et al., Study of D_J meson decays to $D^+\pi^-$, $D^0\pi^+$ and $D^{*+}\pi^-$ final states in pp collisions, JHEP **09** (2013) 145, arXiv:1307.4556.
- 1163 [247] LHCb collaboration, R. Aaij et al., Measurement of the differential branching fraction of the decay $\Lambda_b^0 \to \Lambda \mu^+ \mu^-$, Phys. Lett. **B725** (2013) 25, arXiv:1306.2577.
- 1165 [248] LHCb collaboration, R. Aaij et al., Observation of $B_s^0 \to \chi_{c1} \phi$ decay and study of $B^0 \to \chi_{c1,2} K^{*0}$ decays, Nucl. Phys. **B874** (2013) 663, arXiv:1305.6511.
- 1167 [249] LHCb collaboration, R. Aaij et al., Measurement of the polarization amplitudes in $B^0 \to J/\psi K^*(892)^0$ decays, Phys. Rev. **D88** (2013) 052002, arXiv:1307.2782.
- [250] LHCb collaboration, R. Aaij et al., Measurements of the branching fractions of the decays $B_s^0 \to \overline{D}^0 K^- \pi^+$ and $B^0 \to \overline{D}^0 K^+ \pi^-$, Phys. Rev. **D87** (2013) 112009, arXiv:1304.6317.
- 1172 [251] LHCb collaboration, R. Aaij et al., First observation of the decay $B_c^+ \to J/\psi K^+$, JHEP **09** (2013) 075, arXiv:1306.6723.
- 1174 [252] LHCb collaboration, R. Aaij et al., A measurement of the CKM angle γ from a combination of $B^{\pm} \to Dh^{\pm}$ analyses, Phys. Lett. **B726** (2013) 151, arXiv:1305.2050.
- 1176 [253] LHCb collaboration, R. Aaij et al., Differential branching fraction and angular analysis of the decay $B^0 \to K^{*0} \mu^+ \mu^-$, JHEP **08** (2013) 131, arXiv:1304.6325.
- LHCb collaboration, R. Aaij et al., First observation of CP violation in the decays of B_s^0 mesons, Phys. Rev. Lett. **110** (2013) 221601, arXiv:1304.6173.
- 1180 [255] LHCb collaboration, R. Aaij et al., Differential branching fraction and angular analysis of the decay $B_s^0 \to \phi \mu^+ \mu^-$, JHEP 07 (2013) 084, arXiv:1305.2168.
- 1182 [256] LHCb collaboration, R. Aaij et al., Production of J/ψ and Υ mesons in pp collisions at $\sqrt{s}=8$ TeV, JHEP **06** (2013) 064, arXiv:1304.6977.
- 1184 [257] LHCb collaboration, R. Aaij et al., Measurement of the effective $B_s^0 \to J/\psi K_S^0$ 1185 lifetime, Nucl. Phys. **B873** (2013) 275, arXiv:1304.4500.
- 1186 [258] LHCb collaboration, R. Aaij et al., Searches for violation of lepton flavour and baryon 1187 number in tau lepton decays at LHCb, Phys. Lett. **B724** (2013) 36, arXiv:1304.4518.
- 1188 [259] LHCb collaboration, R. Aaij *et al.*, Search for the rare decay $D^0 \to \mu^+\mu^-$, Phys. Lett. **B725** (2013) 15, arXiv:1305.5059.
- 1190 [260] LHCb collaboration, R. Aaij et al., First observation of the decay $B_s^0 \to \phi \overline{K}^{*0}$, JHEP 1191 11 (2013) 092, arXiv:1306.2239.
- [261] LHCb collaboration, R. Aaij et al., Precision measurement of D meson mass differences, JHEP **06** (2013) 065, arXiv:1304.6865.

- 1194 [262] LHCb collaboration, R. Aaij et al., Observation of $B_c^+ \to J/\psi D_s^+$ and $B_c^+ \to J/\psi D_s^{*+}$ 1195 decays, Phys. Rev. **D87** (2013) 112012, arXiv:1304.4530.
- 1196 [263] LHCb collaboration, R. Aaij et al., Limits on neutral Higgs boson production 1197 in the forward region in pp collisions at $\sqrt{s}=7$ TeV, JHEP **05** (2013) 132, 1198 arXiv:1304.2591.
- 1199 [264] LHCb collaboration, R. Aaij et al., Measurement of J/ψ polarization in pp collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C73 (2013) 2631, arXiv:1307.6379.
- [265] LHCb collaboration, R. Aaij et al., First measurement of the CP-violating phase in $B_s^0 \to \phi \phi$ decays, Phys. Rev. Lett. **110** (2013) 241802, arXiv:1303.7125.
- [266] LHCb collaboration, R. Aaij et al., Precision measurement of the $B_s^0 \overline{B}_s^0$ oscillation frequency in the decay $B_s^0 \to D_s^- \pi^+$, New J. Phys. **15** (2013) 053021, arXiv:1304.4741.
- 1206 [267] LHCb collaboration, R. Aaij et al., Measurement of the $B^0 \to K^{*0}e^+e^-$ branching 1207 fraction at low dilepton mass, JHEP **05** (2013) 159, arXiv:1304.3035.
- [268] LHCb collaboration, R. Aaij et al., Measurement of B meson production cross-sections in proton-proton collisions at $\sqrt{s}=7$ TeV, JHEP **08** (2013) 117, arXiv:1306.3663.
- 1211 [269] LHCb collaboration, R. Aaij et al., Search for direct CP violation in $D^0 \to h^-h^+$ 1212 modes using semileptonic B decays, Phys. Lett. **B723** (2013) 33, arXiv:1303.2614.
- [270] LHCb collaboration, R. Aaij et al., Measurement of CP violation and the B_s^0 meson decay width difference with $B_s^0 \to J/\psi K^+K^-$ and $B_s^0 \to J/\psi \pi^+\pi^-$ decays, Phys. Rev. **D87** (2013) 112010, arXiv:1304.2600.
- 1216 [271] LHCb collaboration, R. Aaij et al., Determination of the X(3872) meson quantum numbers, Phys. Rev. Lett. **110** (2013) 222001, arXiv:1302.6269.
- [272] LHCb collaboration, R. Aaij et al., Measurements of the $\Lambda_b^0 \to J/\psi \Lambda$ decay amplitudes and the Λ_b^0 polarisation in pp collisions at $\sqrt{s}=7$ TeV, Phys. Lett. **B724** (2013) 27, arXiv:1302.5578.
- [273] LHCb collaboration, R. Aaij *et al.*, Search for the decay $B_s^0 \to D^{*\mp}\pi^{\pm}$, Phys. Rev. **D87** (2013) 071101(R), arXiv:1302.6446.
- [274] LHCb collaboration, R. Aaij et al., Observation of the suppressed ADS modes $B^{\pm} \to [\pi^{\pm}K^{\mp}\pi^{+}\pi^{-}]_{D}K^{\pm}$ and $B^{\pm} \to [\pi^{\pm}K^{\mp}\pi^{+}\pi^{-}]_{D}\pi^{\pm}$, Phys. Lett. **B723** (2013) 44, arXiv:1303.4646.
- 1226 [275] LHCb collaboration, R. Aaij et al., Observation of the decay $B_c^+ \to \psi(2S)\pi^+$, Phys. Rev. **D87** (2013) 071103(R), arXiv:1303.1737.

- [276] LHCb collaboration, R. Aaij et al., Observations of $B_s^0 \to \psi(2S)\eta$ and $B_{(s)}^0 \to \psi(2S)\pi^+\pi^-$ decays, Nucl. Phys. **B871** (2013) 403, arXiv:1302.6354.
- 1230 [277] LHCb collaboration, R. Aaij et al., Search for CP violation in $D^+ \to \phi \pi^+$ and $D_s^+ \to K_S^0 \pi^+$ decays, JHEP **06** (2013) 112, arXiv:1303.4906.
- 1232 [278] LHCb collaboration, R. Aaij et al., Search for $D^+_{(s)} \to \pi^+ \mu^+ \mu^-$ and $D^+_{(s)} \to \pi^- \mu^+ \mu^+$ 1233 decays, Phys. Lett. **B724** (2013) 203, arXiv:1304.6365.
- 1234 [279] LHCb collaboration, R. Aaij et al., First observations of $\overline{B}^0_s \to D^+D^-$, $D^+_sD^-$ and D $^0\overline{D}^0$ decays, Phys. Rev. **D87** (2013) 092007, arXiv:1302.5854.
- 1236 [280] LHCb collaboration, R. Aaij *et al.*, Search for rare $B^0_{(s)} \to \mu^+ \mu^- \mu^+ \mu^-$ decays, Phys. Rev. Lett. **110** (2013) 211801, arXiv:1303.1092.
- 1238 [281] LHCb collaboration, R. Aaij et al., Measurements of the Λ_b^0 , Ξ_b^- , and Ω_b^- baryon masses, Phys. Rev. Lett. **110** (2013) 182001, arXiv:1302.1072.
- 1240 [282] LHCb collaboration, R. Aaij et al., Measurements of the branching fractions of $B^+ \to p\overline{p}K^+$ decays, Eur. Phys. J. C73 (2013) 2462, arXiv:1303.7133.
- [283] LHCb collaboration, R. Aaij et al., Study of $B^0 \to D^{*-}\pi^+\pi^-\pi^+$ and $B^0 \to D^{*-}K^+\pi^-\pi^+$ decays, Phys. Rev. **D87** (2013) 092001, arXiv:1303.6861.
- [284] LHCb collaboration, R. Aaij et al., Analysis of the resonant components in $\overline{B}^0 \to J/\psi \pi^+ \pi^-$, Phys. Rev. **D87** (2013) 052001, arXiv:1301.5347.
- 1246 [285] LHCb collaboration, R. Aaij et al., Exclusive J/ψ and $\psi(2S)$ production in pp collisions at $\sqrt{s}=7$ TeV, J. Phys. **G40** (2013) 045001, arXiv:1301.7084.
- [286] LHCb collaboration, R. Aaij et al., First evidence for the decay $B_s^0 \to \mu^+\mu^-$, Phys. Rev. Lett. **110** (2013) 021801, arXiv:1211.2674.
- 1250 [287] LHCb collaboration, R. Aaij et al., Measurement of CP observables in $B^0 \to DK^{*0}$ 1251 with $D \to K^+K^-$, JHEP **03** (2013) 067, arXiv:1212.5205.
- 1252 [288] LHCb collaboration, R. Aaij et al., Prompt charm production in pp collisions at $\sqrt{s} = 7$ TeV, Nucl. Phys. **B871** (2013) 1, arXiv:1302.2864.
- 1254 [289] LHCb collaboration, R. Aaij et al., Amplitude analysis and branching fraction measurement of $\overline{B}^0_s \to J/\psi K^+K^-$, Phys. Rev. **D87** (2013) 072004, arXiv:1302.1213.
- 1256 [290] LHCb collaboration, R. Aaij et al., Measurement of J/ψ production in pp collisions at $\sqrt{s}=2.76$ TeV, JHEP **02** (2013) 041, arXiv:1212.1045.
- 1258 [291] LHCb collaboration, R. Aaij et al., Observation of $D^0-\overline{D}^0$ oscillations, Phys. Rev. Lett. **110** (2013) 101802, arXiv:1211.1230.

- 1260 [292] LHCb collaboration, R. Aaij et al., Measurement of the fragmentation fraction ratio f_s/f_d and its dependence on B meson kinematics, JHEP **04** (2013) 001, arXiv:1301.5286.
- 1263 [293] LHCb collaboration, R. Aaij et al., Measurement of the cross-section for $Z \to e^+e^-$ 1264 production in pp collisions at $\sqrt{s} = 7$ TeV, JHEP **02** (2013) 106, arXiv:1212.4620.
- 1265 [294] LHCb collaboration, R. Aaij et al., Measurement of the time-dependent CP asym-1266 metry in $B^0 \to J/\psi K_S^0$ decays, Phys. Lett. **B721** (2013) 24, arXiv:1211.6093.
- [295] LHCb collaboration, R. Aaij et al., Measurement of the forward energy flow in pp collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C73 (2013) 2421, arXiv:1212.4755.
- 1269 [296] LHCb collaboration, R. Aaij *et al.*, First observation of the decays $\overline{B}_{(s)}^{0} \rightarrow D_{s}^{+}K^{-}\pi^{+}\pi^{-}$ and $\overline{B}_{s}^{0} \rightarrow D_{s1}(2536)^{+}\pi^{-}$, Phys. Rev. **D86** (2012) 112005, arXiv:1211.1541.
- 1272 [297] LHCb collaboration, R. Aaij et al., Measurement of the $B^0-\overline{B}^0$ oscillation frequency Δm_d with the decays $B^0 \to D^-\pi^+$ and $B^0 \to J/\psi K^{*0}$, Phys. Lett. **B719** (2013) 318, arXiv:1210.6750.
- 1275 [298] LHCb collaboration, R. Aaij et al., and A. Bharucha et al., Implications of LHCb mea-1276 surements and future prospects, Eur. Phys. J. C73 (2013) 2373, arXiv:1208.3355.
- [299] LHCb collaboration, R. Aaij et al., First observation of the decay $B_{s2}^*(5840)^0 \rightarrow B^{*+}K^-$ and studies of excited B_s^0 mesons, Phys. Rev. Lett. **110** (2013) 151803, arXiv:1211.5994.
- [300] LHCb collaboration, R. Aaij et al., A study of the Z production cross-section in pp collisions at $\sqrt{s} = 7$ TeV using tau final states, JHEP **01** (2013) 111, arXiv:1210.6289.
- [301] LHCb collaboration, R. Aaij et al., Measurements of B_c^+ production and mass with the $B_c^+ \to J/\psi \pi^+$ decay, Phys. Rev. Lett. **109** (2012) 232001, arXiv:1209.5634.
- [302] LHCb collaboration, R. Aaij et al., A model-independent Dalitz plot analysis of $B^{\pm} \to DK^{\pm}$ with $D \to K_S^0 h^+ h^-$ ($h = \pi, K$) decays and constraints on the CKM angle γ , Phys. Lett. **B718** (2012) 43, arXiv:1209.5869.
- 1287 [303] LHCb collaboration, R. Aaij et al., Measurement of the D^{\pm} production asymmetry in 7 TeV pp collisions, Phys. Lett. **B718** (2013) 902, arXiv:1210.4112.
- [304] LHCb collaboration, R. Aaij et al., First evidence for the annihilation decay mode $B^+ \to D_s^+ \phi$, JHEP **02** (2013) 043, arXiv:1210.1089.
- [305] LHCb collaboration, R. Aaij et al., Differential branching fraction and angular analysis of the $B^+ \to K^+ \mu^+ \mu^-$ decay, JHEP **02** (2013) 105, arXiv:1209.4284.

- [306] LHCb collaboration, R. Aaij et al., Search for the rare decay $K_S^0 \to \mu^+\mu^-$, JHEP **01** (2013) 090, arXiv:1209.4029.
- [307] LHCb collaboration, R. Aaij et al., Evidence for the decay $B^0 \to J/\psi\omega$ and measurement of the relative branching fractions of B_s^0 meson decays to $J/\psi\eta$ and $J/\psi\eta'$, Nucl. Phys. **B867** (2013) 547, arXiv:1210.2631.
- [308] LHCb collaboration, R. Aaij et al., Measurement of the CP asymmetry in $B^0 \rightarrow K^{*0}\mu^+\mu^-$ decays, Phys. Rev. Lett. **110** (2013) 031801, arXiv:1210.4492.
- [309] LHCb collaboration, R. Aaij et al., First observation of the decay $B^+ \to \pi^+ \mu^+ \mu^-$, JHEP 12 (2012) 125, arXiv:1210.2645.
- [310] LHCb collaboration, R. Aaij et al., Measurement of the ratio of branching fractions $\mathcal{B}(B^0 \to K^{*0}\gamma)/\mathcal{B}(B_s^0 \to \phi\gamma)$ and the direct CP asymmetry in $B^0 \to K^{*0}\gamma$, Nucl. Phys. **B867** (2013) 1, arXiv:1209.0313.
- [311] LHCb collaboration, R. Aaij et al., Observation of $B^0 \to \overline{D}^0 K^+ K^-$ and evidence for $B^0_s \to \overline{D}^0 K^+ K^-$, Phys. Rev. Lett. **109** (2012) 131801, arXiv:1207.5991.
- 1307 [312] LHCb collaboration, R. Aaij et al., Measurement of the \overline{B}_s^0 effective lifetime in the $J/\psi f_0(980)$ final state, Phys. Rev. Lett. 109 (2012) 152002, arXiv:1207.0878.
- 1309 [313] LHCb collaboration, R. Aaij et al., Study of D_{sJ} decays to $D^+K_S^0$ and D^0K^+ final states in pp collisions, JHEP **10** (2012) 151, arXiv:1207.6016.
- [314] LHCb collaboration, R. Aaij et al., Measurement of the fraction of $\Upsilon(1S)$ originating from $\chi_b(1P)$ decays in pp collisions at $\sqrt{s}=7$ TeV, JHEP 11 (2012) 031, arXiv:1209.0282.
- 1314 [315] LHCb collaboration, R. Aaij et al., Measurement of the $B_s^0 \to J/\psi \overline{K}^{*0}$ branching fraction and angular amplitudes, Phys. Rev. **D86** (2012) 071102(R), arXiv:1208.0738.
- 1316 [316] LHCb collaboration, R. Aaij et al., Measurement of the effective $B_s^0 \to K^+K^-$ 1317 lifetime, Phys. Lett. **B716** (2012) 393, arXiv:1207.5993.
- 1318 [317] LHCb collaboration, R. Aaij et al., Observation of excited Λ_b^0 baryons, Phys. Rev. Lett. 109 (2012) 172003, arXiv:1205.3452.
- 1320 [318] LHCb collaboration, R. Aaij et al., Measurement of the isospin asymmetry in $B \to K^{(*)}\mu^+\mu^-$ decays, JHEP **07** (2012) 133, arXiv:1205.3422.
- 1322 [319] LHCb collaboration, R. Aaij et al., Measurement of relative branching fractions of B decays to $\psi(2S)$ and J/ψ mesons, Eur. Phys. J. C72 (2012) 2118, arXiv:1205.0918.
- 1324 [320] LHCb collaboration, R. Aaij et al., Measurement of the D_s^+ - D_s^- production asymmetry in 7 TeV pp collisions, Phys. Lett. **B713** (2012) 186, arXiv:1205.0897.

- 1326 [321] LHCb collaboration, R. Aaij et al., Inclusive W and Z production in the forward region at $\sqrt{s} = 7$ TeV, JHEP **06** (2012) 058, arXiv:1204.1620.
- 1328 [322] LHCb collaboration, R. Aaij et al., Strong constraints on the rare decays $B_s^0 \to \mu^+\mu^-$ 1329 and $B^0 \to \mu^+\mu^-$, Phys. Rev. Lett. **108** (2012) 231801, arXiv:1203.4493.
- 1330 [323] LHCb collaboration, R. Aaij et al., Measurement of the CP-violating phase ϕ_s in $\overline{B}_s^0 \to J/\psi \pi^+ \pi^-$ decays, Phys. Lett. **B713** (2012) 378, arXiv:1204.5675.
- 1332 [324] LHCb collaboration, R. Aaij et al., Analysis of the resonant components in $\overline{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$, Phys. Rev. **D86** (2012) 052006, arXiv:1204.5643.
- 1334 [325] LHCb collaboration, R. Aaij et al., Measurement of the polarization amplitudes and 1335 triple product asymmetries in the $B_s^0 \to \phi \phi$ decay, Phys. Lett. **B713** (2012) 369, arXiv:1204.2813.
- [326] LHCb collaboration, R. Aaij et al., Observation of double charm production involving open charm in pp collisions at $\sqrt{s} = 7$ TeV, JHEP **06** (2012) 141, Addendum ibid. **03** (2014) 108, arXiv:1205.0975.
- [327] LHCb collaboration, R. Aaij et al., Measurement of b-hadron branching fractions for two-body decays into charmless charged hadrons, JHEP **10** (2012) 037, arXiv:1206.2794.
- 1343 [328] LHCb collaboration, R. Aaij *et al.*, Observation of CP violation in $B^{\pm} \rightarrow DK^{\pm}$ decays, Phys. Lett. **B712** (2012) 203, Erratum ibid. **B713** (2012) 351, arXiv:1203.3662.
- 1346 [329] LHCb collaboration, R. Aaij et al., Measurement of $\psi(2S)$ meson production in pp collisions at $\sqrt{s}=7$ TeV, Eur. Phys. J. C72 (2012) 2100, arXiv:1204.1258.
- 1348 [330] LHCb collaboration, R. Aaij et al., First observation of the decay $B_c^+ \to J/\psi \pi^+ \pi^- \pi^+$, Phys. Rev. Lett. **108** (2012) 251802, arXiv:1204.0079.
- 1350 [331] LHCb collaboration, R. Aaij et al., Measurement of the B^{\pm} production cross-section 1351 in pp collisions at $\sqrt{s} = 7$ TeV, JHEP **04** (2012) 093, arXiv:1202.4812.
- [332] LHCb collaboration, R. Aaij et al., Measurement of the ratio of branching fractions $\mathcal{B}(B^0 \to K^{*0}\gamma)/\mathcal{B}(B^0_s \to \phi\gamma)$, Phys. Rev. **D85** (2012) 112013, arXiv:1202.6267.
- 1354 [333] LHCb collaboration, R. Aaij et al., Measurement of the $B_s^0 \to J/\psi K_S^0$ branching fraction, Phys. Lett. **B713** (2012) 172, arXiv:1205.0934.
- 1356 [334] LHCb collaboration, R. Aaij et al., First observation of the decays $\overline{B}^0 \to D^+K^-\pi^+\pi^-$ 1357 and $B^- \to D^0K^-\pi^+\pi^-$, Phys. Rev. Lett. 108 (2012) 161801, arXiv:1201.4402.
- 1358 [335] LHCb collaboration, R. Aaij et al., Searches for Majorana neutrinos in B^- decays, Phys. Rev. **D85** (2012) 112004, arXiv:1201.5600.

- 1360 [336] LHCb collaboration, R. Aaij et al., Measurement of prompt hadron production ratios in pp collisions at $\sqrt{s}=0.9$ and 7 TeV, Eur. Phys. J. C72 (2012) 2168, arXiv:1206.5160.
- 1363 [337] LHCb collaboration, R. Aaij et al., Measurement of Υ production in pp collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C72 (2012) 2025, arXiv:1202.6579.
- 1365 [338] LHCb collaboration, R. Aaij *et al.*, *Measurement of b-hadron masses*, Phys. Lett. B708 (2012) 241, arXiv:1112.4896.
- [339] LHCb collaboration, R. Aaij et al., Observation of X(3872) production in pp collisions at $\sqrt{s}=7$ TeV, Eur. Phys. J. C72 (2011) 1972, arXiv:1112.5310.
- 1369 [340] LHCb collaboration, R. Aaij et al., Search for the X(4140) state in $B^+ \to J/\psi \phi K^+$ 1370 decays, Phys. Rev. **D85** (2012) 091103(R), arXiv:1202.5087.
- 1371 [341] LHCb collaboration, R. Aaij et al., Measurement of mixing and CP violation param-1372 eters in two-body charm decays, JHEP **04** (2012) 129, arXiv:1112.4698.
- [342] LHCb collaboration, R. Aaij et al., Measurement of the CP violating phase ϕ_s in $\overline{B}_s^0 \to J/\psi f_0(980)$, Phys. Lett. **B707** (2012) 497, arXiv:1112.3056.
- 1375 [343] LHCb collaboration, R. Aaij et al., Measurement of the ratio of prompt χ_c to J/ψ production in pp collisions at $\sqrt{s}=7$ TeV, Phys. Lett. **B718** (2012) 431, arXiv:1204.1462.
- 1378 [344] LHCb collaboration, R. Aaij et al., First evidence of direct CP violation in charmless two-body decays of B_s^0 mesons, Phys. Rev. Lett. **108** (2012) 201601, arXiv:1202.6251.
- 1381 [345] LHCb collaboration, R. Aaij et al., Determination of the sign of the decay width difference in the B_s^0 system, Phys. Rev. Lett. 108 (2012) 241801, arXiv:1202.4717.
- 1383 [346] LHCb collaboration, R. Aaij et al., Opposite-side flavour tagging of B mesons at the LHCb experiment, Eur. Phys. J. C72 (2012) 2022, arXiv:1202.4979.
- 1385 [347] LHCb collaboration, R. Aaij *et al.*, Observation of $\overline{B}_{s}^{0} \to J/\psi f_{2}'(1525)$ in $J/\psi K^{+}K^{-}$ 1386 final states, Phys. Rev. Lett. **108** (2012) 151801, arXiv:1112.4695.
- 1387 [348] LHCb collaboration, R. Aaij *et al.*, Search for the rare decays $B_s^0 \to \mu^+\mu^-$ and $B^0 \to \mu^+\mu^-$, Phys. Lett. **B708** (2012) 55, arXiv:1112.1600.
- [349] LHCb collaboration, R. Aaij et al., Measurements of the branching fractions and CP asymmetries of $B^{\pm} \to J/\psi \pi^{\pm}$ and $B^{\pm} \to \psi(2S)\pi^{\pm}$ decays, Phys. Rev. **D85** (2012) 091105(R), arXiv:1203.3592.
- 1392 [350] LHCb collaboration, R. Aaij et al., Evidence for CP violation in time-integrated $D^0 \rightarrow h^-h^+$ decay rates, Phys. Rev. Lett. 108 (2012) 111602, arXiv:1112.0938.

- 1394 [351] LHCb collaboration, R. Aaij et al., Measurements of the branching fractions of the decays $B_s^0 \to D_s^{\mp} K^{\pm}$ and $B_s^0 \to D_s^{-} \pi^{+}$, JHEP **06** (2012) 115, arXiv:1204.1237.
- 1396 [352] LHCb collaboration, R. Aaij et al., Measurement of the CP-violating phase ϕ_s in the decay $B_s^0 \to J/\psi \phi$, Phys. Rev. Lett. 108 (2012) 101803, arXiv:1112.3183.
- [353] LHCb collaboration, R. Aaij et al., Differential branching fraction and angular analysis of the decay $B^0 \to K^{*0} \mu^+ \mu^-$, Phys. Rev. Lett. **108** (2012) 181806, arXiv:1112.3515.
- [354] LHCb collaboration, R. Aaij et al., Measurement of the cross-section ratio $\sigma(\chi_{c2})/\sigma(\chi_{c1})$ for prompt χ_c production at $\sqrt{s}=7$ TeV, Phys. Lett. **B714** (2012) 215, arXiv:1202.1080.
- 1404 [355] LHCb collaboration, R. Aaij et al., Measurement of b hadron production fractions in 1405 7 TeV pp collisions, Phys. Rev. **D85** (2012) 032008, arXiv:1111.2357.
- 1406 [356] LHCb collaboration, R. Aaij et al., Search for CP violation in $D^+ \to K^-K^+\pi^+$ 1407 decays, Phys. Rev. **D84** (2011) 112008, arXiv:1110.3970.
- [357] LHCb collaboration, R. Aaij et al., Measurements of the branching fractions for $B_{(s)} \to D_{(s)} \pi \pi \pi$ and $\Lambda_b^0 \to \Lambda_c^+ \pi \pi \pi$, Phys. Rev. **D84** (2011) 092001, Erratum ibid. **D85** (2012) 039904, arXiv:1109.6831.
- 1411 [358] LHCb collaboration, R. Aaij et al., Absolute luminosity measurements with the LHCb detector at the LHC, JINST 7 (2012) P01010, arXiv:1110.2866.
- [359] LHCb collaboration, R. Aaij et al., Measurement of the effective $B_s^0 \to K^+K^$ lifetime, Phys. Lett. **B707** (2012) 349, arXiv:1111.0521.
- [360] LHCb collaboration, R. Aaij et al., Observation of J/ψ -pair production in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. **B707** (2012) 52, arXiv:1109.0963.
- [361] LHCb collaboration, R. Aaij et al., First observation of the decay $B_s^0 \to K^{*0} \overline{K}^{*0}$, Phys. Lett. **B709** (2012) 50, arXiv:1111.4183.
- [362] LHCb collaboration, R. Aaij et al., Measurement of charged particle multiplicities in pp collisions at $\sqrt{s}=7$ TeV in the forward region, Eur. Phys. J. C72 (2012) 1947, arXiv:1112.4592.
- 1422 [363] LHCb collaboration, R. Aaij et al., Measurement of the $B_s^0 \overline{B}_s^0$ oscillation frequency 1423 Δm_s in $B_s^0 \to D_s^-(3)\pi$ decays, Phys. Lett. **B709** (2012) 177, arXiv:1112.4311.
- [364] LHCb collaboration, R. Aaij et al., Search for lepton number violating decays $B^+ \to \pi^- \mu^+ \mu^+$ and $B^+ \to K^- \mu^+ \mu^+$, Phys. Rev. Lett. **108** (2012) 101601, arXiv:1110.0730.

- [365] LHCb collaboration, R. Aaij et al., First observation of the decay $\overline{B}_s^0 \to D^0 K^{*0}$ and a measurement of the ratio of branching fractions $\frac{\mathcal{B}(\overline{B}_s^0 \to D^0 K^{*0})}{\mathcal{B}(\overline{B}^0 \to D^0 \rho^0)}$, Phys. Lett. **B706** (2011) 32, arXiv:1110.3676.
- [366] LHCb collaboration, R. Aaij et al., Measurement of the inclusive ϕ cross-section in pp collisions at $\sqrt{s}=7$ TeV, Phys. Lett. **B703** (2011) 267, arXiv:1107.3935.
- [367] LHCb collaboration, R. Aaij et al., Determination of f_s/f_d for 7 TeV pp collisions and measurement of the $B^0 \to D^-K^+$ branching fraction, Phys. Rev. Lett. 107 (2011) 211801, arXiv:1106.4435.
- [368] LHCb collaboration, R. Aaij et al., Measurement of V^0 production ratios in pp collisions at $\sqrt{s} = 0.9$ and 7 TeV, JHEP **08** (2011) 034, arXiv:1107.0882.
- [369] LHCb collaboration, R. Aaij *et al.*, Search for the rare decays $B_s^0 \to \mu^+\mu^-$ and $B^0 \to \mu^+\mu^-$, Phys. Lett. **B699** (2011) 330, arXiv:1103.2465.
- [370] LHCb collaboration, R. Aaij et al., Measurement of J/ψ production in pp collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C71 (2011) 1645, arXiv:1103.0423.
- [371] LHCb collaboration, R. Aaij et al., First observation of $B_s^0 \to J/\psi f_0(980)$ decays, Phys. Lett. **B698** (2011) 115, arXiv:1102.0206.
- 1443 [372] LHCb collaboration, R. Aaij et al., First observation of $\overline{B}_s^0 \to D_{s2}^{*+} X \mu^- \overline{\nu}$ decays, Phys. Lett. **B698** (2011) 14, arXiv:1102.0348.
- 1445 [373] LHCb collaboration, R. Aaij et al., Measurement of $\sigma(pp \to b\bar{b}X)$ at $\sqrt{s} = 7$ TeV in the forward region, Phys. Lett. **B694** (2010) 209, arXiv:1009.2731.
- 1447 [374] LHCb collaboration, R. Aaij et al., Prompt K_S^0 production in pp collisions at $\sqrt{s} = 0.9$ 1448 TeV, Phys. Lett. **B693** (2010) 69, arXiv:1008.3105.
- 1449 [375] LHCb collaboration, Search for structure in the $B_s^0 \pi^{\pm}$ invariant mass spectrum, 1450 LHCb-CONF-2016-004.
- 1451 [376] LHCb collaboration, Study of Cold Nuclear Matter effect with prompt D⁰ meson production in pPb collisions at LHCb, LHCb-CONF-2016-003.
- 1453 [377] LHCb collaboration, Measurement of the $Z \to \mu\mu$ production cross-section at forward 1454 rapidities in pp collisions at $\sqrt{s} = 13$ TeV, LHCb-CONF-2016-002.
- 1455 [378] LHCb collaboration, LHCb γ combination update from $B \to DKX$ decays, LHCb-1456 CONF-2016-001.
- [379] LHCb collaboration, Study of $\psi(2S)$ production and cold nuclear matter effects in pPb collisions at $\sqrt{s_{NN}} = 5$ TeV, LHCb-CONF-2015-005.

- [380] LHCb collaboration, First measurements of long-range near-side angular correlations in $\sqrt{s_{NN}} = 5$ TeV proton-lead collisions in the forward region, LHCb-CONF-2015-004.
- [381] LHCb collaboration, Measurement of the B^0 oscillation frequency Δm_d with $B^0 \rightarrow D^{(*)-}\mu^+\nu_\mu$, LHCb-CONF-2015-003.
- 1464 [382] LHCb collaboration, Angular analysis of the $B_d^0 \to K^{*0} \mu^+ \mu^-$ decay, LHCb-CONF-1465 2015-002.
- 1466 [383] LHCb collaboration, Study of the decay $B^+ \to K^+\pi^0$ at LHCb, LHCb-CONF-2015-1467 001.
- 1468 [384] LHCb collaboration, Improved constraints on γ : CKM2014 update, LHCb-CONF-1469 2014-004.
- [385] ALICE and LHCb collaborations, Reference pp cross-sections for $\Upsilon(1S)$ studies in proton-lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV and comparisons between ALICE and LHCb results, Aug, 2014. LHCb-CONF-2014-003; ALICE-PUBLIC-2014-002.
- [386] LHCb collaboration, Measurement of the forward W boson cross-section in pp collisions at $\sqrt{s} = 7$ TeV, LHCb-CONF-2014-002.
- 1475 [387] LHCb collaboration, A search for heavy long-lived stau pair production in the LHCb detector, LHCb-CONF-2014-001.
- [388] ALICE and LHCb collaborations, Reference pp cross-sections for J/ψ studies in proton-lead collisions at $\sqrt{s_{NN}}=5.02$ TeV and comparisons between ALICE and LHCb results, Dec, 2013. LHCb-CONF-2013-013, ALICE-PUBLIC-2013-002.
- [389] CMS and LHCb collaborations, Combination of results on the rare decays $B_{(s)}^0 \rightarrow \mu^+\mu^-$ from the CMS and LHCb experiments, Jul, 2013. CMS-PAS-BPH-13-007, LHCb-CONF-2013-012.
- 1483 [390] LHCb collaboration, Updated average f_s/f_d b-hadron production fraction ratio for 7 TeV pp collisions, LHCb-CONF-2013-011.
- [391] LHCb collaboration, Search for the $\Lambda_b^0 \to \Lambda \eta'$ decay at LHCb, LHCb-CONF-2013-010.
- [392] LHCb collaboration, *CP* and up-down asymmetries in $B^{\pm} \to K^{\pm}\pi^{\mp}\pi^{\pm}\gamma$ decays, LHCb-CONF-2013-009.
- [393] LHCb collaboration, Study of the J/ψ production cross-section in proton-lead collisions at $\sqrt{s_{NN}} = 5~TeV$, LHCb-CONF-2013-008.
- [394] LHCb collaboration, Measurement of the cross section for $Z \to \mu^+\mu^-$ production with 1.0 fb⁻¹ of pp collisions at $\sqrt{s} = 7$ TeV, LHCb-CONF-2013-007.

- [395] LHCb collaboration, A measurement of γ from a combination of $B^{\pm} \rightarrow DK^{\pm}$ analyses including first results using 2 fb⁻¹ of 2012 data, LHCb-CONF-2013-006.
- [396] LHCb collaboration, Graphical comparison of the LHCb measurements of W and Z boson production with ATLAS and CMS, LHCb-CONF-2013-005.
- [397] LHCb collaboration, Model-independent measurement of CP violation parameters in $B^{\pm} \to (K_S^0 h^+ h^-)_D K^{\pm}$ decays, LHCb-CONF-2013-004.
- [398] LHCb collaboration, A search for time-integrated CP violation in $D^0 \to K^-K^+$ and $D^0 \to \pi^-\pi^+$ decays, LHCb-CONF-2013-003.
- 1501 [399] LHCb collaboration, Measurement of $\sigma(b\bar{b})$ with inclusive final states, LHCb-CONF-1502 2013-002.
- 1503 [400] LHCb collaboration, Measurement of the forward-central $b\bar{b}$ production asymmetry at LHCb, LHCb-CONF-2013-001.
- 1505 [401] LHCb collaboration, First look at the pPb pilot run, LHCb-CONF-2012-034.
- LHCb collaboration, Optimization and calibration of the same-side kaon tagging algorithm using hadronic B_s^0 decays in 2011 data, LHCb-CONF-2012-033.
- [403] LHCb collaboration, A measurement of γ from a combination of $B^+ \to Dh^+$ analyses, LHCb-CONF-2012-032.
- ¹⁵¹⁰ [404] LHCb collaboration, Studies of $\Lambda_b^0 \to J/\psi \Lambda$ production in pp collisions at $\sqrt{s} = 7~TeV$, LHCb-CONF-2012-031.
- 1512 [405] LHCb collaboration, Search for the suppressed ADS modes $B^{\pm} \rightarrow [\pi^{\pm}K^{\mp}\pi^{+}\pi^{-}]_{D}K^{\pm}$ 1513 and $B^{\pm} \rightarrow [\pi^{\pm}K^{\mp}\pi^{+}\pi^{-}]_{D}\pi^{\pm}$, LHCb-CONF-2012-030.
- 1514 [406] LHCb collaboration, Measurement of the time-dependent CP-violation parameters 1515 $in B_s^0 \to D_s^{\mp} K^{\pm}$, LHCb-CONF-2012-029.
- 1516 [407] LHCb collaboration, Evidence for CP violation in $B \to KK\pi$ and $B \to \pi\pi\pi$ decays, LHCb-CONF-2012-028.
- 1518 [408] LHCb collaboration, Search for the lepton flavour violating and baryon number violating decays $\tau^- \to \overline{p}\mu^+\mu^-$ and $\tau^- \to p\mu^-\mu^-$, LHCb-CONF-2012-027.
- [409] LHCb collaboration, Performance of flavor tagging algorithms optimised for the analysis of $B_s^0 \to J/\psi \phi$, LHCb-CONF-2012-026.
- [410] LHCb collaboration, Production of J/ψ and $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$ mesons at $\sqrt{s} = 8~TeV$, LHCb-CONF-2012-025.
- [411] LHCb collaboration, Measurement of CP observables in $B^0 \to DK^{*0}$ with $D \to K^+K^-$, LHCb-CONF-2012-024.

- 1526 [412] LHCb collaboration, Branching fraction measurements of $B_{d,s}^0$ decays to $K_S^0 h h'$ final states, including first observation of $B_s^0 \to K_S K \pi$, LHCb-CONF-2012-023.
- [413] LHCb collaboration, Measurement of the flavour-specific CP violating asymmetry $a_{\rm sl}^s$ in B_s^0 decays, LHCb-CONF-2012-022.
- 1530 [414] LHCb collaboration, First observation of $B^- \to D^0 K^- \pi^+ \pi^-$ decays to CP even final states, LHCb-CONF-2012-021.
- ¹⁵³² [415] LHCb collaboration, Observation of $\chi_b(3P)$ state at LHCb in pp collisions at $\sqrt{s} = 7$ TeV, LHCb-CONF-2012-020.
- 1534 [416] LHCb collaboration, Search for CP violation in $D^0 \to \pi^- \pi^+ \pi^+ \pi^-$ decays, LHCb1535 CONF-2012-019.
- 1536 [417] LHCb collaboration, Evidence for CP violation in $B \to K\pi\pi$ and $B \to KKK$ 1537 decays, LHCb-CONF-2012-018.
- 1538 [418] LHCb collaboration, Search for the rare decays $B_{(s)}^0 \to \mu\mu$ at the LHC with the ATLAS, CMS and LHCb experiments, LHCb-CONF-2012-017.
- 1540 [419] LHCb collaboration, Measurement of jet production in $Z^0/\gamma^* \to \mu^+\mu^-$ events at LHCb in $\sqrt{s} = 7$ TeV pp collisions, LHCb-CONF-2012-016.
- 1542 [420] LHCb collaboration, Search for the lepton flavour violating decay $\tau^- \to \mu^+ \mu^- \mu^-$, LHCb-CONF-2012-015.
- [421] LHCb collaboration, Search for (Higgs-like) bosons decaying into long-lived exotic particles, LHCb-CONF-2012-014.
- 1546 [422] LHCb collaboration, Inclusive low mass Drell-Yan production in the forward region at $\sqrt{s} = 7$ TeV, LHCb-CONF-2012-013.
- 1548 [423] LHCb collaboration, Measurement of the forward energy flow in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ with the LHCb experiment, LHCb-CONF-2012-012.
- 1550 [424] LHCb collaboration, Measurement of the cross-section for $Z^0 \to e^+e^-$ production in pp collisions at $\sqrt{s} = 7$ TeV, LHCb-CONF-2012-011.
- ¹⁵⁵² [425] LHCb collaboration, Search for the rare decays $B_s^0 \to \mu^+ \mu^- \mu^+ \mu^-$ and $B_d^0 \to \mu^+ \mu^- \mu^+ \mu^-$, LHCb-CONF-2012-010.
- 1554 [426] LHCb collaboration, First observations and branching fraction measurements of \overline{B}_s^0 1555 to double-charm final states, LHCb-CONF-2012-009.
- 1556 [427] LHCb collaboration, Differential branching fraction and angular analysis of the 1557 $B^0 \to K^{*0} \mu^+ \mu^- decay$, LHCb-CONF-2012-008.

- 1558 [428] LHCb collaboration, Measurement of time-dependent CP violation in charmless 1559 two-body B decays, LHCb-CONF-2012-007.
- 1560 [429] LHCb collaboration, First observation of $B^+ \to \pi^+ \mu^+ \mu^-$, LHCb-CONF-2012-006.
- 1561 [430] LHCb collaboration, Search for the $D^0 \to \mu^+\mu^-$ decay with 0.9 fb⁻¹ at LHCb, LHCb-1562 CONF-2012-005.
- 1563 [431] LHCb collaboration, Measurement of the direct CP asymmetry in the $B_d^0 \to K^{*0}\gamma$ 1564 decay, LHCb-CONF-2012-004.
- 1565 [432] LHCb collaboration, Measurement of the ratio of branching fractions for $B_s^0 \to \phi \mu \mu$ 1566 and $B_s^0 \to J/\psi \phi$, LHCb-CONF-2012-003.
- 1567 [433] LHCb collaboration, Tagged time-dependent angular analysis of $B_s^0 \to J/\psi \phi$ decays at LHCb, LHCb-CONF-2012-002.
- 1569 [434] LHCb collaboration, Measurement of the effective $B_s^0 \to K^+K^-$ lifetime, LHCb1570 CONF-2012-001.
- 1571 [435] LHCb collaboration, Measurement of the relative cross-section $\sigma(\chi_{c_2})/\sigma(\chi_{c_1})$ of prompt χ_c mesons using at LHCb, LHCb-CONF-2011-062.
- 1573 [436] LHCb collaboration, A search for time-integrated CP violation in $D^0 \to h^- h^+$ decays, 1574 LHCb-CONF-2011-061.
- 1575 [437] LHCb collaboration, Measurement of the masses of the Ξ_b^- and Ω_b^- , LHCb-CONF-1576 2011-060.
- 1577 [438] LHCb collaboration, Relative branching ratio measurements of charmless B^{\pm} decays to three hadrons, LHCb-CONF-2011-059.
- 1579 [439] LHCb collaboration, Measurements of the relative branching fractions of the $B^{\pm} \rightarrow p\bar{p}K^{\pm}$ decay channel including charmonium contributions, LHCb-CONF-2011-058.
- 1581 [440] LHCb collaboration, Measurements of the relative and absolute branching fractions 1582 of the decays $B_s^0 \to D_s^{\mp} K^{\pm}$ and $B_s^0 \to D_s^{-} \pi^{+}$, LHCb-CONF-2011-057.
- 1583 [441] LHCb collaboration, Combination of ϕ_s measurements from $B_s^0 \to J/\psi \phi$ and $B_s^0 \to J/\psi f_0(980)$, LHCb-CONF-2011-056.
- [442] LHCb collaboration, Measurement of the ratio of branching fractions $\mathcal{B}(B_d \to K^{*0}\gamma)/\mathcal{B}(B_s \to \phi\gamma)$ with the LHCb experiment at $\sqrt{s} = 7$ TeV, LHCb-CONF-2011-055.
- 1588 [443] LHCb collaboration, Measurement of the Charm Mixing Parameter y_{CP} in Two-Body Charm Decays, LHCb-CONF-2011-054.

- LHCb collaboration, Observations of Orbitally Excited $B_{(s)}^{**}$ Mesons, LHCb-CONF-2011-053.
- 1592 [445] LHCb collaboration, Study of Triple Product Asymmetries in $B_s \to \phi \phi$ decays, LHCb-CONF-2011-052.
- 1594 [446] LHCb collaboration, Measurement of ϕ_s in $B_s \to J/\psi f_0(980)$, LHCb-CONF-2011-1595 051.
- 1596 [447] LHCb collaboration, Measurement of Δm_s in the decay $B_s^0 \to D_s^-(K^+K^-\pi^-)\pi^+$ 1597 using opposite-side and same-side flavour tagging algorithms, LHCb-CONF-2011-1598 050.
- 1599 [448] LHCb collaboration, Tagged time-dependent angular analysis of $B_s \to J/\psi \phi$ decays with 337 pb⁻¹ at LHCb, LHCb-CONF-2011-049.
- ¹⁶⁰¹ [449] LHCb collaboration, Measurement of the $B_s^0 \to J/\psi K_s^0$ branching fraction, LHCb-¹⁶⁰² CONF-2011-048.
- [450] CMS and LHCb collaborations, Search for the rare decay $B_s^0 \to \mu^+\mu^-$ at the LHC with the CMS and LHCb experiments, LHCb-CONF-2011-047,CMS-PAS-BPH-11-019.
- 1606 [451] LHCb collaboration, Measurement of the CP Violation Parameter \mathcal{A}_{Γ} in Two-Body
 1607 Charm Decays, LHCb-CONF-2011-046.
- ¹⁶⁰⁸ [452] LHCb collaboration, Search for X(4140) in $B^+ \to J/\psi \phi K^+$, LHCb-CONF-2011-¹⁶⁰⁹ 045.
- ¹⁶¹⁰ [453] LHCb collaboration, Evidence for the suppressed decay $B^{\pm} \to (K^{\mp}\pi^{\pm})_D K^{\pm}$, LHCb¹⁶¹¹ CONF-2011-044.
- [454] LHCb collaboration, Inclusive X(3872) production in pp collisions at $\sqrt{s} = 7$ TeV, LHCb-CONF-2011-043.
- ¹⁶¹⁴ [455] LHCb collaboration, Charmless charged two-body B decays at LHCb with 2011 data, LHCb-CONF-2011-042.
- 1616 [456] LHCb collaboration, Z cross-section measurement at $\sqrt{s} = 7$ TeV using the channel $Z \to \tau \tau$, LHCb-CONF-2011-041.
- ¹⁶¹⁸ [457] LHCb collaboration, First observation of $B_c^+ \to J/\psi \pi^+ \pi^- \pi^+$, LHCb-CONF-2011-¹⁶¹⁹ 040.
- [458] LHCb collaboration, Updated measurements of W and Z production at $\sqrt{s} = 7$ TeV with the LHCb experiment, LHCb-CONF-2011-039.
- [459] LHCb collaboration, Angular analysis of $B^0 \to K^{*0} \mu^+ \mu^-$, LHCb-CONF-2011-038.

- [460] LHCb collaboration, Search for the rare decays $B^0_{(s)} \to \mu^+\mu^-$ with 300 pb⁻¹ at LHCb, LHCb-CONF-2011-037.
- [461] LHCb collaboration, Studies of beauty baryons decaying to $D^0p\pi^-$ and D^0pK^- , LHCb-CONF-2011-036.
- 1627 [462] LHCb collaboration, Analysis of $\overline{B}^0_s \to J/\psi (\pi^+\pi^- \text{ and } K^+K^-)$ and the first observation of $J/\psi f_2'(1525)$, LHCb-CONF-2011-035.
- [463] LHCb collaboration, Average f_s/f_d b-hadron production fraction for 7 TeV pp collisions, LHCb-CONF-2011-034.
- [464] LHCb collaboration, Measurement of the B^{\pm} production cross-section at LHCb, LHCb-CONF-2011-033.
- [465] LHCb collaboration, A measurement of the ratio of branching fractions: $\frac{\mathcal{B}(B^{\pm}\to DK^{\pm})}{\mathcal{B}(B^{\pm}\to D\pi^{\pm})}$ for $D\to K\pi$, KK, $K\pi\pi\pi$ and $K_S^0\pi\pi$, LHCb-CONF-2011-031.
- [466] LHCb collaboration, Measurement of the Ratio of Branching Fractions $\mathcal{B}(B^{\pm} \to J/\psi \pi^{\pm})/\mathcal{B}(B^{\pm} \to J/\psi K^{\pm})$ at $\sqrt{s} = 7$ TeV with the LHCb Detector, LHCb-1637 CONF-2011-030.
- 1638 [467] LHCb collaboration, Time integrated ratio of wrong-sign to right-sign $D^0 \to K\pi$ 1639 decays in 2010 data at LHCb, LHCb-CONF-2011-029.
- [468] LHCb collaboration, Measurement of b-hadron production fractions in 7 TeV centreof-mass energy pp collisions, LHCb-CONF-2011-028.
- [469] LHCb collaboration, Measurement of b-hadron masses with exclusive $J/\psi X$ decays in 2010 data, LHCb-CONF-2011-027.
- ¹⁶⁴⁴ [470] LHCb collaboration, Measurement of the $\psi(2S)$ production cross-section at $\sqrt{s}=$ 7 TeV in LHCb, LHCb-CONF-2011-026.
- 1646 [471] LHCb collaboration, Evidence for the decay $B_s^0 \to J/\psi \overline{K}^{*0}$, LHCb-CONF-2011-025.
- ¹⁶⁴⁷ [472] LHCb collaboration, First observations of the Cabibbo-suppressed decays $\overline{B}^0 \to D^+K^-\pi^+\pi^-$ and $B^- \to D^0K^-\pi^+\pi^-$, LHCb-CONF-2011-024.
- 1649 [473] LHCb collaboration, A search for time-integrated CP violation in $D^0 \to h^+h^-$ decays 1650 and a measurement of the D^0 production asymmetry, LHCb-CONF-2011-023.
- 1651 [474] LHCb collaboration, Central exclusive dimuon production at $\sqrt{s}=7$ TeV, LHCb-1652 CONF-2011-022.
- ¹⁶⁵³ [475] LHCb collaboration, Measurement of the X (3872) mass with first LHCb data, LHCb-¹⁶⁵⁴ CONF-2011-021.

- 1655 [476] LHCb collaboration, A measurement of the cross-section ratio $\sigma(\chi_{c2})/\sigma(\chi_{c1})$ for prompt χ_c production at $\sqrt{s}=7$ TeV in LHCb, LHCb-CONF-2011-020.
- ¹⁶⁵⁷ [477] LHCb collaboration, First observation of the decay $B_s^0 \to K^{*0} \overline{K}^{*0}$, LHCb-CONF-¹⁶⁵⁸ 2011-019.
- 1659 [478] LHCb collaboration, Measurement of the effective $B_s^0 \to K^+K^-$ Lifetime, LHCb-1660 CONF-2011-018.
- ¹⁶⁶¹ [479] LHCb collaboration, Measurement of the B_c^+ to B^+ production cross-section ratios at $\sqrt{s}=7$ TeV in LHCb, LHCb-CONF-2011-017.
- [480] LHCb collaboration, Measurement of the $\Upsilon(1S)$ production cross-section at $\sqrt{s} = 7$ TeV in LHCb, LHCb-CONF-2011-016.
- 1665 [481] LHCb collaboration, Inclusive jets and dijets in LHCb, LHCb-CONF-2011-015.
- 1666 [482] LHCb collaboration, $\mathcal{B}(B_s^0 \to \psi(2S)\phi)/\mathcal{B}(B_s^0 \to J/\psi\phi)$, LHCb-CONF-2011-014.
- [483] LHCb collaboration, Measurement of the relative yields of the decay modes $B^0 \rightarrow D^-\pi^+$, $B^0 \rightarrow D^-K^+$, $B^0 \rightarrow D^-\pi^+$, and determination of f_s/f_d for 7 TeV pp collisions, LHCb-CONF-2011-013.
- ¹⁶⁷⁰ [484] LHCb collaboration, W and Z production at $\sqrt{s} = 7$ TeV with the LHCb experiment, LHCb-CONF-2011-012.
- [485] LHCb collaboration, Measurement of direct \mathcal{CP} violation in charmless charged two-body B decays at LHCb, LHCb-CONF-2011-011.
- [486] LHCb collaboration, Measurement of Δm_d in $B^0 \to D^-(K^+\pi^-\pi^-)\pi^+$, LHCb-CONF-2011-010.
- 1676 [487] LHCb collaboration, Observation of double J/ψ production in proton-proton collisions 1677 at a centre-of-mass energy of $\sqrt{s}=7~TeV$, LHCb-CONF-2011-009.
- 1678 [488] LHCb collaboration, First observation of the decay $\overline{B}_s^0 \to D^0 K^{*0}$ and measurement of the ratio of branching fractions $\frac{\mathcal{B}(\overline{B}_s^0 \to D^0 K^{*0})}{\mathcal{B}(\overline{B}_d^0 \to D^0 \rho^0)}$, LHCb-CONF-2011-008.
- [489] LHCb collaboration, Improved Measurements of the Cabibbo Favored Decays $B_{(s)} \rightarrow D_{(s)}\pi\pi\pi$ and $\Lambda_b \rightarrow \Lambda_{c\pi\pi\pi}$ Branching Fractions, LHCb-CONF-2011-007.
- [490] LHCb collaboration, Tagged time-dependent angular analysis of $B_s^0 \to J/\psi \phi$ decays with the 2010 LHCb data, LHCb-CONF-2011-006.
- [491] LHCb collaboration, Measurement of Δm_s in the decay $B_s^0 \to D_s^-(K^+K^-\pi^-)(3)\pi$, LHCb-CONF-2011-005.

- ¹⁶⁸⁶ [492] LHCb collaboration, Search for CP violation in $B^0 \to J/\psi K_S^0$ decays with first LHCb data, LHCb-CONF-2011-004.
- [493] LHCb collaboration, Flavor-untagged angular analysis of $B_d^0 \to J/\psi K^*$ and $B_s^0 \to J/\psi \phi$ decays, LHCb-CONF-2011-002.
- [494] LHCb collaboration, b-hadron lifetime measurements with exclusive $b \to J/\psi X$ decays reconstructed in the 2010 data, LHCb-CONF-2011-001.
- [495] LHCb collaboration, Measurement of the inclusive ϕ cross-section in pp collisions at $\sqrt{s} = 7$ TeV with the LHCb experiment, LHCb-CONF-2010-014.
- [496] LHCb collaboration, Prompt charm production in pp collisions at $\sqrt{s}=7$ TeV, LHCb-CONF-2010-013.
- 1696 [497] LHCb collaboration, Measurements of B^0 mesons production cross-section in pp 1697 collisions at $\sqrt{s} = 7$ TeV using $B^0 \to D^{*-}\mu^+\nu_\mu X$ decays, LHCb-CONF-2010-012.
- [498] LHCb collaboration, Measurement of prompt $\overline{\Lambda}/\Lambda$ and $\overline{\Lambda}/K_{\rm S}^0$ production ratios in inelastic non-diffractive pp collisions at $\sqrt{s}=0.9$ and 7 TeV, LHCb-CONF-2010-1700 011.
- 1701 [499] LHCb collaboration, Measurement of the J/ψ production cross-section at $\sqrt{s} = 7 \text{ TeV}$ in LHCb, LHCb-CONF-2010-010.
- 1703 [500] LHCb collaboration, Measurement of the \bar{p}/p ratio in LHCb at $\sqrt{s} = 900 \,\text{GeV}$ and 7 TeV, LHCb-CONF-2010-009.
- [501] LHCb collaboration, Prompt K_S^0 production in pp collisions at $\sqrt{s} = 900$ GeV, LHCb-CONF-2010-008.

LHCb collaboration

 1707 A. N. Other¹.

 1 University of nowhere

The author list for journal publications is generated from the Membership Database shortly after 'approval to go to paper' has been given. It will be sent to you by email shortly after a paper number has been assigned. The author list should be included already at first circulation, to allow new members of the collaboration to verify that they have been included correctly. Occasionally a misspelled name is corrected, or associated institutions become full members. Therefore an updated author list will be sent to you after the final EB review of the paper. In case line numbering doesn't work well after including the authorlist, try moving the \bigskip after the last author to a separate line.

The authorship for Conference Reports should be "The LHCb collaboration", with a footnote giving the name(s) of the contact author(s), but without the full list of collaboration names.