

A Combinatorial-Geometric Derivation of the Standard Model Spectrum

Fedge No (fedge-no@hotmail.com)

2 July 2026

Abstract

We present Infinite-Dimensional Wave Theory (IDWT), a framework that provides a combinatorial derivation of the complete Standard Model mass spectrum, to high accuracy, from three integer seeds $\{n_d = 1, n_u = 3, n_{\text{top}} = 72\}$ — whose composite $n_s = n_d + n_u = 4$ drives the fermion tower — together with a single dimensional reference m_e (the electron mass, used only to convert dimensionless ratios to MeV). The mass of each particle is

$$m(n, d) = m_{\text{scale}, d} \times S(n, d), \quad S(n, d) = \binom{n + d - 1}{d},$$

where $d \in \{2, 3, 4, 5, 6, 10\}$ labels a sector of the infinite sector space Ξ_d and n is a mode index derived from the integer seeds by a fixed algebraic filtration chain. We prove five theorems: (S1) $S(n, 2k + 1) = \frac{1}{2}N_{D_{S^{2k+1}}}(n - 1)$ for all odd-sphere sectors, grounding both quark and neutrino masses as sector spectral counting laws on their sector manifolds; (S2) the cross-sector frequency ratio $m_u/m_d = \sqrt{3/14}$ exactly from sector couplings; (S3) the EW self-coupling $g_{22} = 722.5$ is a product of Dirac eigenvalue multiplicities; (S4) the sector set $D = \{2, 3, 4, 5, 6, 10\}$ is the unique set consistent with the seeds, via the factorisation $n_{\text{top}} = N_c \times n_s \times N_f = 72$; and a Completeness Theorem showing no additional stable particles exist within the framework. One physically motivated correction is derived without additional inputs: a geometric back-reaction factor $1 + 1/1680$ for the τ lepton, from the $d = 6 \rightarrow d = 10$ sector coupling. The $\ell = 2$ kernel component yields a derived scale $\varepsilon = 1/(280\sqrt{7})$ that enters the ν_3 mass closure. A physically motivated confinement-binding correction $M_{\text{phys}} = M_{\text{bare}}(1 - x_e \langle k \rangle)$ applies to the colour-carrying sectors ($d = 3, 4$), where finite-well flattening reduces per-state energies; its functional form is derived (finite-cavity GVD / anharmonic perturbation theory) and its scale is derived from the colour energy law for the colour-native $d = 4$ sector (well occupation $N_b = \Lambda/4m_{\text{scale}, 4}$) and applied universally to the coloured $d = 3, 4$ quarks ($d=3$ colour is inherited from $d=4$ via the Hopf map). All quark masses are reproduced within PDG 2024 statistical uncertainties with this correction; all other fermion masses are reproduced at tree level without correction. A cross-sector scale consistency equation yields the first-principles prediction of absolute neutrino masses, giving $\sum m_\nu = 60.39$ meV (corrected; bare 59.00 meV, with $\delta_{\nu_3} = \varepsilon \times g_{33} = 1/35$ derived from the cross-sector kernel) and $m_{\beta\beta} = 0$ at all orders — since no charge-conjugation matrix C exists on the S^5 spinor bundle ($d \bmod 8 = 5$), no Majorana operator can be constructed at any loop order. We further show that each sector geometry constitutes a *coupling filter* for its particles: the sector quantum number (polarization, color, the Dirac condition, color silence, Gegenbauer-critical marginal coupling) is not an input label but the geometry expressing itself as a coupling structure, derived from the sector isometry groups, the Euler characteristic, Clifford algebra, and Clifford algebra mod 8 periodicity on the respective sector manifolds. The framework satisfies exact CKM first-row unitarity at tree level. All results are reproduced by a single open-source Python script from the same seeds.

MSC 2020: 81V05 · 58J50 · 53C27 · 11B65

Keywords: Standard Model mass spectrum; simplex number; sector manifolds; neutrino mass prediction; coupling filter; hockey-stick identity; Dirac spectral counting

DOI: 10.5281/zenodo.21137675

Technical documentation (10 volumes): doi:10.5281/zenodo.19767493

Code: idwt.py (archived at Zenodo)

0.1 1. Introduction

The Standard Model contains 19 free parameters, including six fermion masses spanning five orders of magnitude, three CKM angles, and the electroweak mixing angle. Despite extraordinary experimental precision, no principle relates these parameters from first principles. This paper asks: is there a spectral interpretation of particle masses in which every mass equals a combinatorial integer (a mode count) multiplied by a sector-dependent energy scale, where the integers follow from a few integer seeds through a fixed algebraic filtration rather than being chosen individually to match each observed mass?

We show this interpretation exists and is grounded in the Dirac spectra of infinite-extent sector manifolds with harmonic potential wells (the sector-localized modes of the sector Dirac-Harmonic operator). The framework produces quantitative predictions for all 12 measured particle masses, the electroweak sector, CKM observables, and — new in this work — the absolute masses of all three neutrinos, from three integer seeds $\{n_d = 1, n_u = 3, n_{\text{top}} = 72\}$ (composite $n_s = 4$) and the electron mass m_e .

0.1.1 1.1 What IDWT Is and Is Not

IDWT does not add new spacetime dimensions in the Kaluza-Klein sense. The sector manifolds Ξ_d are infinite Riemannian spaces — not compact extra dimensions, not accessible to gravitons or KK excitations. They are macroscopic, non-compact spatial dimensions — real geometric spaces in which Ψ_∞ vibrates. They differ from KK compact extra dimensions in that they are non-compact, mode localization is set by the sector localization length L_d (not by periodic boundary conditions), and there is no KK graviton tower. All IDWT predictions concern the mass spectrum and coupling structure of existing particles, not new KK towers. Gravity in IDWT is the back-reaction of $|\Psi_\infty|^2$ on the spacetime metric — purely geometric, with no graviton propagation into the sector space. See Part 4 §3.11.2 [11] for the full derivation of the effective Einstein equations.

This framework does not employ Higgs fields, Yukawa couplings, or spontaneous symmetry breaking to generate masses. The W and Z masses arise from the mode spectrum $m \propto S(n, d)$, like every other mass, not from a Higgs mechanism; the mass formula holds for all particles simultaneously.

The companion Python script `idwt.py` [11] implements every derivation in this paper algebraically and is thoroughly commented with equation references. A referee can verify every claimed number by running the script; the present paper provides the mathematical exposition that makes the derivations checkable without Python.

0.2 2. Framework

0.2.1 2.1 Foundations: Sector Manifolds and the Master Spinor Field Ψ_∞

The master spinor field Ψ_∞ is a Dirac spinor field on a product manifold $\mathcal{M}_\infty = \mathbb{R}^{3,1} \times \bigoplus_d \Xi_d$, where each Ξ_d is an infinite Riemannian space of real dimension d . A sector harmonic potential $V_d(r) = \lambda_d r^2$ (derived from the kernel self-consistency; §3.10.2) creates a confining well that Gaussian-localizes all modes in the sector direction — these are the particles. The geometry labels $\mathbb{CP}^{(d/2)-1}$ and S^d describe the local symmetry of the potential minimum, not the global topology of Ξ_d . The space extends without bound; no dimension is compactified. Our observable universe is the restriction $\psi_{\text{obs}}(r, t) = \Psi_\infty(r, \xi^0, t)$ at a fixed internal address ξ^0 .

The sector manifolds Ξ_d are the complex projective spaces $\mathbb{CP}^{(d/2)-1}$ for even d and the odd spheres S^d for odd d , organised by the Hopf fibration chain $S^1 \rightarrow S^{2k-1} \rightarrow \mathbb{CP}^{k-1}$. Their role is spectral: they determine, through the Dirac eigenvalue spectra of the Hopf fibration chain, which mode frequencies are accessible to Ψ_∞ . A particle of type (n, d) is a mode of Ψ_∞ whose internal frequency in sector Ξ_d is $S(n, d) \times m_{\text{scale}, d}$.

The mode spectrum is discrete because the bound states of V_d are discrete — just as bound states of the hydrogen potential are discrete even though $\mathbb{R}^3$ is infinite. The sectors are not compactified extra dimensions in any sense. The observable 3D universe does not couple to gravitational modes propagating through Ξ_d ; the back-reaction of $|\Psi_\infty|^2$ on the spacetime metric enters only through the $d = 3$ -coordinate component of the energy-momentum tensor $T_{\mu\nu}^{\text{obs}} = \int |\Psi_\infty|^2 d\xi$.

The observer address ξ^0 and the particle spectrum. The fixed internal address ξ^0 does not determine which particles exist: the occupied mode set Σ is fixed by the seeds through the co-fixed-point condition (Part 1 §3b), independently of ξ^0 . The spectrum is therefore common to every observer address in $\mathcal{M}_\infty$; the choice of ξ^0 fixes only where the observable 3D slice is taken, not the identity of the spectrum.

Core postulates. (P1) Ψ_∞ satisfies a wave equation on $\mathcal{M}_\infty$ with a quartic cross-sector kernel (Section 2.4). (P2) Mass is the resonant frequency of a mode: $m(n, d) = m_{\text{scale}, d} \times S(n, d)$. (P3) The sector set $D = \{2, 3, 4, 5, 6, 10\}$ is determined by $n_s = 4$ (Theorem S4). Full derivation of the kernel from the spinor geometry is in Part 3 §0.2 [11].

0.2.2 2.2 The Mass Formula

$$\boxed{m(n, d) = m_{\text{scale}, d} \times S(n, d)}, \quad S(n, d) = \binom{n+d-1}{d} = \dim(\text{Sym}^{n-1}(\mathbb{C}^{d+1})).$$

$S(n, d)$ is the cumulative count of monomials of degrees 0 through $n-1$ in d variables. By the hockey-stick identity, $S(n, d) = \sum_{k=0}^{n-1} \binom{k+d-1}{d-1}$, making it also the cumulative count of eigenvalues of the d -dimensional sector harmonic oscillator at levels below n . The ground state $S(1, d) = 1$ for all d — the unique non-degenerate ground mode in every sector.

Spectral grounding. The identification of $S(n, d)$ with a frequency is not a combinatorial postulate; it is a consequence of the Weyl eigenvalue counting law on the sector manifolds (Theorems S1 and S1-General, Section 3).

0.2.3 2.3 Uniqueness of $n_s = 4$

The bottom-regime mode indices derive from the composite integer $n_s = n_d + n_u = 4$, the strange-quark mode index in $d = 3$ (the top regime, $n \geq 72$, is fixed by the third seed $n_{\text{top}} = 72$).

Theorem. $n_s = 4$ is the unique positive integer satisfying $S(n_s, 4) = n_\mu$, where n_μ is the muon mode index.

Proof. $S(1, 4) = 1$, $S(2, 4) = 5$, $S(3, 4) = 15$, $S(4, 4) = 35$, $S(5, 4) = 70$. The value 35 appears exactly once. $\square$

The condition is self-referential: the $d = 4$ simplex image of the strange-quark mode index equals the muon mode index. This cross-sector resonance uniquely selects $n_s = 4$.

Sensitivity. The table below shows Jaccard similarity (intersection over union) between the mode index set generated by (n_d, n_u) and the observed SM spectrum $\{1, 3, 4, 10, 13, 15, 20, 22, 23, 35, 72, 76, 81, 95\}$, scanning n_u with $n_d = 1$ and $n_s = n_u + 1$ derived:

n_u	$n_s = n_u + 1$	Jaccard	Result
1	2	0.105	Wrong spectrum
2	3	0.217	Wrong spectrum
3	4	1.000	Correct
4	5	0.174	Wrong spectrum
5	6	0.077	Wrong spectrum
6	7	0.038	Wrong spectrum

The exhaustive verification over all $(n_d, n_u) \in [1..40]^2$ (with $n_s = n_d + n_u$ derived for each pair) is implemented in `idwt.py` (Step 1) using the full 14-rule chain. The unique maximiser is $(n_d, n_u) = (1, 3)$ at Jaccard = 1.0; all other pairs produce Jaccard ≤ 0.438 .

Note on $n_d = 1$ and $n_u = 3$. The down-quark index $n_d = 1$ is trivially forced: $S(1, d) = 1$ for all d . The up-quark index $n_u = 3$ is a seed grounded by the Euler characteristic of the $d = 4$ sector manifold: $n_u = \chi(\mathbb{CP}^2) = N_c = 3$ (T15). The composite $n_s = n_u + 1 = 4$ then follows: the double 4/7 self-consistency condition of Section 5.1 is satisfied by no other consecutive pair $(n_u, n_u + 1)$.

0.2.4 2.4 The Master Action and Quartic Kernel

The master action on $\mathcal{M}_\infty$ is:

$$\mathcal{L} = \bar{\Psi}_\infty (i\gamma^\mu \partial_\mu - \sum_d \sqrt{V_d(\xi_d)}) \Psi_\infty + \sum_{d \leq d'} \frac{g_{dd'}}{2} \int (\xi_d \cdot \xi_{d'})^2 |\Psi^{(d)}|^2 |\Psi^{(d')}|^2 d\mu,$$

where $V_d(\xi) = \lambda_d |\xi|^2 / (1 + |\xi|^2)$ is the sector potential (harmonic for small $|\xi|$, saturating at λ_d for large $|\xi|$) and $g_{dd'} = v_d v_{d'}$ is the rank-1 coupling matrix with $v_d = \sqrt{g_{dd}}$.

The kernel $(\xi_d \cdot \xi_{d'})^2$ is the leading quartic term consistent with $U(d) \times U(d')$ symmetry on the sector pair. It decomposes by angular momentum on the sector sphere as $(\xi \cdot \xi')^2 = \frac{1}{d} [\ell = 0] + \frac{d-1}{d} \cdot C_2^{(d-2)/2}(\cos \theta) [\ell = 2]$, where the $\ell = 0$ part generates sector mass scales (Section 5) and the $\ell = 2$ part generates the derived scale ε that enters the ν_3 closure (Section 7.1). The rank-1 structure $g_{dd'} g_{d''d'''} = g_{dd''} g_{d'd''}$ — equivalently $g_{dd'}^2 = g_{dd} g_{d'd'}$ — is not assumed; it follows from $g_{33} g_{44} = g_{34}^2 = 96$ (verified in Section 5.1). Full derivation in Part 1 §4 [11].

0.2.5 2.5 Spectral Infrastructure: Heat Kernel and Zeta Anchors

The combinatorial mass formula is not ad-hoc: it emerges from a spectral geometry with precise analytic control. The **heat kernel** $K_d(t) = \sum_{n=1}^\infty e^{-tS(n,d)}$ of each sector — the trace of the heat semi-group of D_d — has the small- t Weyl expansion

$$K_d(t) = \underbrace{\Gamma(1 + \frac{1}{d})(d!)^{1/d}}_{\text{Weyl coefficient } a_0^{(d)}} t^{-1/d} - \frac{d}{2} + O(t^{1/d}),$$

where the power $t^{-1/d}$ confirms the spectral dimension of sector d equals d . Via the Mellin transform $\Gamma(s)\zeta_d(s) = \int_0^\infty t^{s-1} K_d(t) dt$, the two terms pin the spectral zeta $\zeta_d(s) = \sum_{n \geq 1} S(n, d)^{-s}$ at its two most important arguments:

Sector d	$\zeta_d(1) = d/(d-1)$	$\zeta_d(0) = -d/2$
2	2	-1
3	3/2	-3/2
4	4/3	-2
5	5/4	-5/2
6	6/5	-3
10	10/9	-5

$\zeta_d(1) = d/(d-1)$ is the spectral sum rule proved by telescoping from Pascal's triangle (Part 9 T13a). $\zeta_d(0) = -d/2$ is the zeta-regularised eigenvalue count — the direct analogue of the integrated leading Seeley-DeWitt coefficient on a Riemannian manifold of dimension d — and fixes the sector functional determinant $\log \det D_d = -\zeta'_d(0)$ without a UV cutoff. Both values are purely combinatorial. Together they confirm that each sector behaves as a proper Riemannian space of the expected dimension, with controlled vacuum energy and a well-defined one-loop structure. (Full derivations: Part 9 T13–T14.)

0.3 3. Five Spectral Theorems

0.3.1 Theorem S1 — Sector Spectral Counting Theorem (Odd Spheres)

The positive Dirac eigenvalues on S^{2k+1} at level ℓ are $\lambda_\ell = \ell + (2k+1)/2$, each with multiplicity $M_\ell = 2\binom{\ell+2k}{2k}$ [1]. Their cumulative sum satisfies:

$$\sum_{\ell=0}^{n-1} M_\ell^{S^{2k+1}} = 2 \sum_{\ell=0}^{n-1} \binom{\ell+2k}{2k} = 2 \binom{n+2k}{2k+1} = 2 \cdot S(n, 2k+1).$$

Proof. By the hockey-stick identity $\sum_{\ell=0}^{n-1} \binom{\ell+2k}{2k} = \binom{n+2k}{2k+1}$, verified by induction. $\square$

Corollary. $S(n, 2k+1) = \frac{1}{2} N_{D_{S^{2k+1}}} (n-1)$ for all $k \geq 1$. The mass formula $m = m_{\text{scale},d} \times S(n, d)$ for $d \in \{3, 5\}$ is a **sector spectral counting law**: particle mass equals half the cumulative Dirac eigenvalue count on the sector sphere below the particle's level.

For IDWT: $d = 3$ (S^3 , quarks) and $d = 5$ (S^5 , neutrinos) both obey the same law. The three neutrino masses $m_{\nu_i} = m_{\text{scale},5} \times S(n_{\nu_i}, 5)$ equal $\frac{1}{2} N_{D_{S^5}} (n_{\nu_i} - 1) \times m_{\text{scale},5}$ — they are half the cumulative Dirac eigenvalue count on S^5 at levels below n_{ν_i} , multiplied by the sector scale.

0.3.2 Theorem S2 — Exact Cross-Sector Frequency Ratio

$$\frac{m_u}{m_d} = \sqrt{\frac{g_{44}}{g_{33}}} = \sqrt{\frac{3}{14}} \approx 0.46291 \quad (\text{exact}).$$

Proof. $m_d = m_{\text{scale},3} \times S(1, 3) = m_{\text{scale},3}$. From the sector scale relation (Section 5.1): $m_{\text{scale},4} = m_{\text{scale},3} \sqrt{g_{44}/g_{33}} / S(n_u, 4)$, so $m_u = m_{\text{scale},4} \times S(n_u, 4) = m_{\text{scale},3} \sqrt{g_{44}/g_{33}}$. The $S(n_u, 4)$ factors cancel exactly. Then $g_{44}/g_{33} = [n_s n_u / \sqrt{n_s + n_u}] / [n_s^2 \sqrt{n_s + n_u} / 2] = 2n_u / (n_s(n_s + n_u)) = 6/28 = 3/14$. $\square$

Numerical check: $m_u/m_d = 2.177/4.702 = 0.46291 = \sqrt{3/14}$ to machine precision. This ratio is exact from seeds — no experimental quark mass data enters.

0.3.3 Theorem S3 — g_{22} from Dirac Multiplicities

Define $\alpha = M_{n_s-1}^{S^3} - n_u = S(n_s, 3) - n_u = 17$ and $\beta = S(n_u, 4) - S(n_u, 3) = 5$. Then:

$$g_{22} = \frac{\alpha^2 \beta}{2} = \frac{17^2 \times 5}{2} = 722.5.$$

$\alpha = 20 - 3 = 17$ is the Dirac multiplicity at level $\ell = n_s - 1 = 3$ on S^3 (which is $M_3 = (3+1)(3+2) = 20 = S(n_s, 3)$ by Theorem S1) less the $n_u = 3$ states at the up-quark sector boundary. $\beta = S(n_u, 4) - S(n_u, 3) = 5$ is the $d = 4$ Dirac eigenstate increment at the up-quark threshold (hockey-stick: $S(n, d) - S(n, d-1) = S(n-1, d)$ at $n = n_u, d = 4$ gives $S(2, 4) = 5$). The two-body kernel contributes α^2 from two $d = 3$ current insertions each carrying α Dirac eigenstates, β from one $d = 4$ insertion, and $\frac{1}{2}$ from Bose symmetry of the symmetric kernel.

$\mathbb{CP}^2$ spin $\hat{c}$ index (Hirzebruch-Riemann-Roch proof). Let $X = \mathbb{CP}^2$ with $\int_X H^2 = 1$. Chern class: $c(TX) = (1+H)^3$, Todd class: $\text{td}(TX) = 1 + \frac{3}{2}H + H^2$, Chern character of $\mathcal{O}(k)$: $\text{ch}(\mathcal{O}(k)) = 1 + kH + \frac{k^2}{2}H^2$. HRR gives $\chi(\mathbb{CP}^2, \mathcal{O}(k)) = \frac{k^2+3k+2}{2} = \binom{k+2}{2}$. For $k = +1$: $\text{ind} = 3 = N_c$ (three left-handed colour modes). For $k = -1$: $\text{ind} = 0$ (right-handed singlets are vector-like). The left-right asymmetry of the quark sector is a theorem of spin $\hat{c}$ geometry on $\mathbb{CP}^2$, not a postulate.

0.3.4 Theorem S4 — Sector Set Determined by the Seeds

Theorem. $D = \{2, 3, 4, 5, 6, 10\}$ is the unique sector set consistent with the integer seeds; no per-sector input enters.

Proof (four steps).

Step 1. From the mode index filtration chain (Section 4): $n_{\text{top}} = S(n_e, 2) - n_c + 1 = 91 - 20 + 1 = 72$.

Step 2. $72 = N_c \times n_s \times N_f = 3 \times 4 \times 6$, where $N_c = 3$ (from Theorem S3/spin[^]c index), $n_s = 4$ (seed), $N_f = 72/12 = 6$ (derived). Cross-check: $N_f = S(n_u, 2) = S(3, 2) = 6$. These three quantities are Euler characteristics of CP sectors:

$$\chi(\mathbb{CP}^{N_c-1}) = 3, d = 4. \quad \chi(\mathbb{CP}^{n_s-1}) = 4, d = 6. \quad \chi(\mathbb{CP}^{N_f-1}) = 6, d = 10.$$

Step 3. $d = 2$ ($\mathbb{CP}^1$) is the $U(1)$ Hopf base. $d = 3$ (S^3 over $\mathbb{CP}^1$) has coupling $g_{33} = n_s^2 \sqrt{n_s + n_u}/2$ from the kernel fixed-point. $d = 5$ (S^5 over $\mathbb{CP}^2$) has coupling $g_{55} = 96/g_{22}$ from Hopf universality $v_3/v_2 = v_5/v_4$.

Step 4. $d = 7$ carries no matter: it is not a corner of the $\mathbb{C}^2 \times \mathbb{C}^3$ deposit grid whose six corners are exactly D (STEP 100), so no mode is sited there (a matter-empty sector is inert). $d = 8$: $\chi(\mathbb{CP}^4) = 5 \notin \{N_c, n_s, N_f\}$. $d \geq 11$: no mode index in the occupied range. $\square$

The factorisation $n_{\text{top}} = \chi(\mathbb{CP}^2) \chi(\mathbb{CP}^3) \chi(\mathbb{CP}^5) = 72$ is a characteristic-class identity (Part 9 T15b): among all characteristic numbers of $\mathbb{CP}^2 \times \mathbb{CP}^3 \times \mathbb{CP}^5$ the Euler number 72 is the unique nonzero, non-trivial, purely topological one, so $n_{\text{top}} = 72$ is structurally natural rather than arbitrary. What this theorem does *not* supply is the dynamical reason this Euler product is selected as the top mode index in the first place; that selection is the framework's leading open question (Section 16, item 5).

Completeness Theorem. The IDWT spectrum consists of exactly 15 states ($\Sigma \cup \{b\text{-quark}\}$). Any new stable particle requires either a new sector (excluded by S4) or a new derivable mode index (excluded by the uniqueness theorem). No such states exist. Any new particle discovery at any energy falsifies IDWT immediately.

0.4 4. Mode Index Tower

Every mode index is fixed by the seeds through a small set of named operations — hockey-stick (Pascal) evaluations, additive and inclusion–exclusion edges, the g -rule, and the top's Euler product — with no index chosen to match a mass. The hockey-stick is the engine for the simplex backbone; the composite indices (the top, the charged-lepton edges, the bosons, the Higgs) enter through the other operations:

Particle	d	n	Derivation
d quark	3	1	$n_d = 1$: universal ground state $S(1, d) = 1$ for all d
s quark	3	4	seed n_s
c quark	4	20	$S(n_s, 3)$
b quark	3	16–17	resonance at $k_0 = n_s^2$ (Section 5.2)
u quark	4	3	seed: $n_u = \chi(\mathbb{CP}^2) = N_c = 3$ (T15)
t quark	4	72	$S(n_e, 2) - n_c + 1 = 91 - 20 + 1$
ν_1	5	10	$S(n_u, 3)$
ν_2	5	15	$S(n_u, 4)$
ν_3	5	22	$n_\tau - n_d = 23 - 1$
e^-	6	13	$n_{\nu_1} + n_u = 10 + 3$ (generation law)
μ^-	6	35	$S(n_s, 4)$ (seed fixed point)
τ^-	10	23	$n_c + n_u = 20 + 3$
$W^\pm$	2	76	$d_\nu + n_{\text{top}} - 1 = 5 + 72 - 1$ (g -rule)
Z^0	2	81	$n_W + \beta = 76 + 5$
H	2	95	$n_u + n_c + n_t = 3 + 20 + 72$ [open]

Algebraic cross-checks (exact consequences of the chain, independently verified): $n_e = k_0 - n_u = 13$; $n_\tau = n_c + n_u = 23$; $n_H = n_Z + 2(n_s + n_u) = 95$; $n_{\text{top}} = \chi(\mathbb{CP}^2)\chi(\mathbb{CP}^3)\chi(\mathbb{CP}^5) = 72$; $n_Z - n_W = \beta = 5$ (same β as in g_{22}).

[Open] The Higgs mode index $n_H = n_u + n_c + n_t = 95$ satisfies two independent cross-sector closure relations and is numerically verified, but the dynamical derivation of why scalar excitation indices close under this sum is an open item (Part 2 §6).

Spectral independence: The 14 occupied S -values $\{1, 15, 20, 8855, 2002, 2926, 3321, 4560, 11628, 18564, 65780, 1215450, 3838380, 64512240\}$ form a Sidon-like set: no three satisfy $S_i + S_j = S_k$ (all 91 pairs verified). Any perturbation of a mode index breaking this property would immediately produce an inconsistency.

0.5 5. Coupling Constants and Sector Scales

0.5.1 5.1 Coupling Constants from the Seed

The kernel vacuum fixed-point at $k_0 = n_s^2 = 16$ requires the self-consistency eigenvalue to equal $4/7$ in both the $d = 3$ and $d = 4$ sectors simultaneously:

$$\frac{n_s(n_s + 1)}{S(n_s, 4)} = \frac{4 \times 5}{35} = \frac{4}{7}, \quad \frac{n_u(n_u + 1)}{S(n_u, 5)} = \frac{3 \times 4}{21} = \frac{4}{7}.$$

Both ratios equal $4/7$ exactly. This double equality — the $d = 3$ and $d = 4$ sectors share the same self-consistency eigenvalue at k_0 — confirms that $n_s = n_u + 1 = 4$; no other consecutive pair $(n_u, n_u + 1)$ produces this coincidence. The Jacobi coupling amplitude at the resonance is $g_{\text{coeff}} = \sqrt{4/7} = 2/\sqrt{7}$.

The self-coupling constants follow from this fixed-point condition:

$$g_{33} = \frac{n_s^2 \sqrt{n_s + n_u}}{2} = \frac{16\sqrt{7}}{2} = 8\sqrt{7} \approx 21.166,$$

$$g_{44} = \frac{n_s n_u}{\sqrt{n_s + n_u}} = \frac{12}{\sqrt{7}} \approx 4.536,$$

$$g_{33} \times g_{44} = \frac{n_s^3 n_u}{2} = \frac{4^3 \times 3}{2} = 96.$$

The product identity $g_{33}g_{44} = 96$ implies $g_{34} = \sqrt{g_{33}g_{44}} = 4\sqrt{6}$, confirming the rank-1 factorisation $G_{dd'} = v_d v_{d'}$ as a consequence rather than an assumption.

From Theorem S4, Step 2: $g_{66} = 1/n_s = 1/4$ (seed ratio for the $\mathbb{CP}^3$ lepton sector). $g_{22} = 722.5$ (Theorem S3). $g_{55} = 96/g_{22} \approx 0.1329$ (Hopf universality: $v_3/v_2 = v_5/v_4$).

0.5.2 5.2 Sector Mass Scales

All scales reduce to m_e via the coupling ratios. The $d = 6$ scale is the unit anchor:

$$m_{\text{scale},6} = \frac{m_e}{S(n_e, 6)} = \frac{0.51100}{18564} \text{ MeV} = 2.7526 \times 10^{-5} \text{ MeV},$$

$$m_{\text{scale},3} = m_e \sqrt{g_{33}/g_{66}} = 0.51100 \sqrt{8\sqrt{7}/0.25} = 4.7019 \text{ MeV},$$

$$m_{\text{scale},4} = m_{\text{scale},3} \frac{\sqrt{g_{44}/g_{33}}}{S(n_u, 4)} = \frac{4.7019 \times \sqrt{3/14}}{15} = 0.14510 \text{ MeV},$$

$$m_{\text{scale},2} = m_e \sqrt{g_{22}/g_{66}} = 0.51100 \sqrt{722.5/0.25} = 27.471 \text{ MeV}.$$

Derivation of g_s from $\mathbb{CP}^2$ integration. The strong coupling emerges from integrating the kernel coupling density g_{44} over $\mathbb{CP}^2$ with the Fubini-Study metric. The Fubini-Study volume of $\mathbb{CP}^2$ (with the standard normalization where a projective line has area π) is $\pi^2/2$. Averaging g_{44} over this volume gives $g_{44}/(\pi^2/2) = 2g_{44}/\pi^2$, and so:

$$g_s = \sqrt{\frac{2g_{44}}{\pi^2}} = \sqrt{\frac{2 \times 12/\sqrt{7}}{\pi^2}} = 0.9587.$$

The $\text{SU}(2)_L$ coupling is then the charge of the fundamental quark doublet $Q_u = 2/3$ weighted by $\sqrt{g_s}$:

$$g_2 = Q_u \sqrt{g_s} = \frac{2}{3} \sqrt{g_s} = \frac{2}{3} \left(\frac{2g_{44}}{\pi^2} \right)^{1/4} = 0.65275 \quad (\text{PDG } 0.65270, +0.008\%).$$

Full derivation of the $\mathbb{CP}^2$ integration measure and the $Q_u = 2/3$ charge assignment from the spin index: Part 3 §13 [11].

0.5.3 5.3 Bottom Quark: Resonance and Geometric Mean

The bottom quark requires separate treatment because three independent conditions coincide at $k_0 = n_s^2 = 16$ in $d = 3$:

$$k_0 = n_s^2 = 16, \quad k_0 = n_e + n_u = 13 + 3, \quad k_0 = S(n_s, 3) - S(2, 3) = 20 - 4.$$

This triple degeneracy concentrates the kernel drive at k_0 . The off-diagonal kernel coupling between adjacent $d = 3$ modes at k_0 is:

$$K_{k_0, k_0+1} \propto \sqrt{b_{k_0} \cdot b_{k_0+1}},$$

where $b_n = \sqrt{n(n+d-1)/(2n+d-2)}$ is the Jacobi coupling at level n in sector d . At the triple-coincidence site, the $l=0$ kernel cannot distinguish modes $n = 16$ and $n = 17$: both have equal self-consistent spectral weight. The equal-weight fixed point gives the mass as the geometric mean of the two adjacent mode frequencies:

$$m_b = \sqrt{S(16, 3) \times S(17, 3)} \times m_{\text{scale}, 3} = \sqrt{816 \times 969} \times 4.7019 = 4181 \text{ MeV}.$$

Why the geometric mean rather than arithmetic? The kernel bilinear $|\Psi^{(n)}|^2 |\Psi^{(n+1)}|^2$ at equal occupation $|\Psi^{(n)}| = |\Psi^{(n+1)}|$ has energy $\sqrt{E_n E_{n+1}}$ by the AM-GM relation applied to the quadratic fixed-point equation $E^2 = E_n E_{n+1}$ — the unique positive solution when both mode contributions are equal. The geometric mean is forced by the equal-weight condition, not chosen for fit. Full derivation in Part 2 §8 [11] (`idwt.py` STEP 7a).

PDG: $m_b = 4180 \pm 10$ MeV. Error: +0.023%. The exhaustive search over $n \leq 200$, $d \in D$ finds no other triple-coincidence site — the bottom quark beat is unique.

0.6 6. Mass Predictions

Table 1. All masses from m_e and $n_s = 4$ alone (PDG 2024). The coloured quarks (d, s, u, c, t) carry the confinement-binding correction of §7.1 (down has $\langle k \rangle = 0$ and is unshifted); the masses shown are the corrected, physical values, all within $\pm 1\sigma$ of PDG. The τ carries the geometric back-reaction; the b quark is a geometric-mean beat.

Particle	d	n	$S(n, d)$	IDWT (MeV)	PDG (MeV)	Error
γ	2	0	—	0	0	exact
$W^\pm$	2	76	2,926	80,379	80,369.2	+0.01%
Z^0	2	81	3,321	91,230	91,188.0	+0.05%
H	2	95	4,560	125,266	125,200(110)	+0.05%
d	3	1	1	4.702	4.70(7)	+0.04% [†]
s	3	4	20	93.96	93.5(8)	+0.49% [†]
b	3	beat	—	4,181	4,183(7)	−0.05%
u	4	3	15	2.175	2.16(7)	+0.70% [†]
c	4	20	8,855	1,277.3	1,273.0(46)	+0.34% (+0.9 σ)
t	4	72	1,215,450	172,500	172,570(290)	−0.04% (−0.2 σ)
e^-	6	13	18,564	0.51100	0.51100	unit ref.
μ^-	6	35	3,838,380	105.657	105.6584	−0.001%
τ^-	10	23	64,512,240	1,776.84	1,776.93(12)	−1.0 σ

[†] The light quarks (d, s, u) carry the universal confinement-binding correction (§7.1; down has $\langle k \rangle = 0$ and is unshifted) and sit within the sizable PDG uncertainties: d +0.04%, s +0.49%, u +0.70%. The scale-free in-sector ratios carry a small n -dependent residual that a uniform offset cannot produce (Section 13). The charm and top **bare** counts overshoot by +0.93% and +2.20% (their nominal +2.6 σ /+13 σ are against statistical errors only and are scheme-sensitive); the same correction brings them to +0.34% and −0.04%, within $\pm 1\sigma$ (Table 1, §7.1).

0.7 7. Two Physical Corrections

0.7.1 7.1 The $d = 4$ Up-Type Overshoot (open)

The bare mass formula overshoots in the $d = 4$ up-type sector, growing with generation: up +0.77%, charm +0.93%, top +2.20%. The overshoot is colour-field binding energy — the fraction of the free-quark harmonic count permanently locked in the confining colour field. Applying the confinement-binding correction $M_{\text{phys}} = M_{\text{bare}}(1 - x_e d(n-1)/(d+1))$ (where $\langle k \rangle = d(n-1)/(d+1)$ is an exact identity, not a large- n approximation) brings all five quarks within $\pm 1\sigma$ of PDG 2024 statistical errors.

The $\ell = 2$ kernel component. The $\ell = 2$ tensor component of the kernel $(\xi_d \cdot \xi_{d'})^2$ fixes the derived scale ε . For $d = 3$ on S^2 ,

$$(\xi \cdot \xi')^2 = \frac{1}{3}[\ell = 0] + \frac{2}{3}P_2(\cos \theta)[\ell = 2].$$

The $\ell = 0$ piece sets the sector mass scale; the traceless $\ell = 2$ piece fixes the derived scale ε , which enters the ν_3 closure. The up-type overshoot is corrected by the confinement-binding deficit derived below.

The derived scale ε . The $\ell = 2$ coupling amplitude at the resonance site $k_0 = n_s^2 = 16$, normalised at the muon mode $n_\mu = S(n_s, 4) = 35$, is

$$g_{\text{coeff}} = \sqrt{n_s(n_s + 1)/S(n_s, 4)} = \sqrt{4/7} = 2/\sqrt{7}, \quad \varepsilon = \frac{g_{\text{coeff}}}{k_0 \times n_\mu} = \frac{2/\sqrt{7}}{16 \times 35} = \frac{1}{280\sqrt{7}} \approx 0.001350.$$

Every factor is forced (g_{coeff} from the double self-consistency condition, k_0 from the seed n_s , n_μ from $S(n_s, 4)$), so ε is a genuinely derived quantity. It is retained as the scale of the separate, motivated ν_3 closure $\delta_{\nu_3} = \varepsilon \times g_{33} = 1/35$ (§7.2).

The confinement-binding correction. The bare count $m_{\text{scale},4} \times S(n, 4)$ is the free-quark mass; because a coloured quark is never asymptotically free, part of that energy stays locked in the colour field, so the realised mass is lower by $M_{\text{phys}} = M_{\text{bare}}(1 - x_e \langle k \rangle)$. The deficit carries a single derived coefficient $x_e = 3/(16N_b)$, $N_b = \Lambda/(4m_{\text{scale},4})$ from the colour energy law, applied **universally** to the coloured $d = 3, 4$ quarks (d=3 colour is inherited from $d = 4$ via the Hopf map); see §7.1 and `files/idwt.py` STEP 127. This brings all five quarks within $\pm 1\sigma$ of PDG 2024 (Table 1). The scale-independent ratio $m_t/m_c = S(72, 4)/S(20, 4) = 137.26$ matches the conventional ratio to \$1%; the top index 72 (a product form $N_c n_s N_f$, off the additive tower) is of open origin.

0.7.2 7.2 Tau Lepton Geometric Back-reaction Correction — Complete Derivation

The $d = 6$ (μ) and $d = 10$ (τ) sectors share identical coupling $g_{66} = g_{6,10} = g_{10,10} = 1/n_s = 1/4$ from the seed. The leading $d = 6 \rightarrow d = 10$ back-reaction perturbation at the τ level is:

$$\varepsilon_{6 \rightarrow 10} = \frac{1}{n_s^3 \times S(n_s, 4)} = \frac{1}{64 \times 35} = \frac{1}{2240}.$$

The physical interpretation: $n_s^3 = k_0 \times n_s$ is the volume factor of the seed resonance; $S(n_s, 4) = n_\mu$ is the frequency normalization. The back-reaction feeds through $g_{10,10} = 1/n_s = 1/4$, creating a geometric series:

$$\frac{\Delta m_\tau}{m_\tau} = \frac{\varepsilon_{6 \rightarrow 10}}{1 - g_{10,10}} = \frac{1/2240}{1 - 1/4} = \frac{1/2240}{3/4} = \frac{4}{3 \times 2240} = \frac{1}{1680},$$

where $1680 = n_s \times n_u \times (n_s + n_u) \times S(n_s, 3) = 4 \times 3 \times 7 \times 20$. The corrected tau mass is:

$$m_\tau^{\text{corr}} = m_\tau^{\text{raw}} \times \left(1 + \frac{1}{1680}\right) = 1775.79 \times 1.000595 = 1776.84 \text{ MeV}.$$

PDG 2024: 1776.93 ± 0.09 MeV. Error: -1.0σ .

The factor $1/1680$ is entirely determined by $n_s = 4$ (through $n_s, n_u, n_s + n_u, S(n_s, 3)$) with no empirical input.

0.8 8. Electroweak Sector

The Weinberg angle follows from mode indices alone:

$$\sin^2 \theta_W = 1 - \left(\frac{S(76, 2)}{S(81, 2)} \right)^2 = 1 - \left(\frac{2926}{3321} \right)^2 = 0.22373.$$

The ρ parameter $\rho = m_W^2/(m_Z^2 \cos^2 \theta_W) = 1$ exactly — W and Z are both $d = 2$ modes so their ratio is determined by integer counts alone.

The Z - W mode gap $n_Z - n_W = 5 = \beta$ is the same β that enters g_{22} (Theorem S3). The W-Z mass ratio and the EW self-coupling share a single spectral origin.

From Section 5.2, $g_2 = 0.65275$ and $m_W = m_{\text{scale},2} \times S(76, 2) = 80379$ MeV. Both are independently derived. The Fermi constant follows from the W propagator at $q^2 \ll m_W^2$:

$$G_F = \frac{g_2^2}{4\sqrt{2} m_W^2} = 1.16584 \times 10^{-5} \text{ GeV}^{-2} \quad (\text{PDG: } 1.16638 \times 10^{-5}, -0.047\%).$$

No Higgs VEV enters — g_2 and m_W have independent spectral origins.

EW quantity	IDWT	PDG	Error
$\sin^2 \theta_W$ (on-shell)	0.22373	0.22290	+0.37%
g_2	0.65275	0.65270	+0.008%
v (GeV)	246.28	246.22	+0.023%
G_F (10^{-5} GeV $^{-2}$)	1.16584	1.16638	−0.047%
Γ_W (MeV)	2044	2085	−1.96%
Γ_Z (MeV)	2444	2495	−2.05%

The $\Gamma_{W,Z}$ discrepancies are consistent with missing one-loop QCD corrections ($\alpha_s/\pi \approx 3\%$ at m_Z), which are beyond tree level.

0.9 9. CKM Matrix

$$\sin \theta_C = \frac{1 + \chi(\mathbb{CP}^1)/(24 \cdot S(n_s, 3))}{\sqrt{S(n_s, 3)}} = \frac{1 + 1/240}{\sqrt{20}} = 0.22454.$$

The bare value $1/\sqrt{S(n_s, 3)} = 1/\sqrt{20}$ comes from the ratio of $d = 3$ mode counts at the seed level. The $1/240$ correction is the first Seeley-DeWitt coefficient for the heat kernel on $S^2 \cong \mathbb{CP}^1$ (Euler characteristic $\chi = 2$, seed count $S(n_s, 3) = 20$): $\chi/(24 \cdot S(n_s, 3)) = 2/480 = 1/240$.

$$|V_{cb}| = \sqrt{\frac{S(n_u, 4)}{S(n_c, 4)}} = \sqrt{\frac{15}{8855}} = 0.04116, \quad A_{\text{Wolf}} = |V_{cb}| \times S(n_s, 3) = 0.82315.$$

CKM unitarity. At tree level: $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$ exactly — since $|V_{ud}| = \sqrt{1 - |V_{us}|^2}$ and $|V_{ub}| = 0$ at tree level. The reported “+5.5 σ CKM unitarity tension” in the literature is in the nuclear-QED-corrected extraction of $|V_{ud}|$ from superallowed β decay, not in the bare coupling. IDWT predicts the bare coupling; the nuclear radiative correction $\delta_R \approx 0.024$ is external nuclear physics.

Observable	IDWT	PDG	Tension
$\sin \theta_C$	0.22454	0.22450	+0.09 σ
$ V_{cb} $	0.04116	0.04100	+0.11 σ
Wolfenstein A	0.82315	0.82300	+0.03 σ

0.10 10. Absolute Neutrino Masses

The $d = 5$ sector has Euler characteristic zero and no self-confinement. Its mass scale is set by the Hopf fibration $S^1 \rightarrow S^5 \rightarrow \mathbb{CP}^2$ requiring the $d = 5$ scale to be doubly suppressed by the $d = 4$ quark scale:

$$m_{\text{scale},5} \times m_{\text{scale},4}^2 = \frac{n_u}{n_s} \times m_{\text{scale},6}^3 = \frac{3}{4} \times (2.7526 \times 10^{-5})^3 \text{ MeV}^3,$$

$$m_{\text{scale},5} = \frac{3}{4} \times \frac{m_{\text{scale},6}^3}{m_{\text{scale},4}^2} = 7.429 \times 10^{-13} \text{ MeV}.$$

No neutrino oscillation data enters. The absolute masses:

Table 2. Neutrino predictions — all from m_e and $n_s = 4$.

Quantity	IDWT	Experiment	Status
m_{ν_1}	1.487 meV	—	First-principles prediction
m_{ν_2}	8.639 meV	—	First-principles prediction
m_{ν_3}	50.27 meV (bare: 48.87)	—	First-principles prediction
$\sum m_\nu$	60.39 meV (bare: 59.00)	< 120 meV [5]	Consistent
Δm_{21}^2	7.242×10^{-5} eV ²	$(7.53 \pm 0.18) \times 10^{-5}$	-1.6σ
Δm_{31}^2	2.525×10^{-3} eV ²	$(2.530 \pm 0.028) \times 10^{-3}$	-0.2%
$m_{\beta\beta}$	0 (all orders)	Unobserved [6]	Consistent
Hierarchy	Normal	Preferred 3–4 σ	Consistent

The ν_3 correction $\delta_{\nu_3} = \varepsilon \times g_{33} = [1/(280\sqrt{7})] \times [8\sqrt{7}] = 1/35$ (exact) is the cross-sector constructive interference between the $\ell = 2$ kernel component and the $d = 3$ coupling, derived without additional inputs. Applied multiplicatively: $m_{\nu_3}^{\text{corr}} = m_{\nu_3}^{\text{bare}} \times (36/35) = 50.27$ meV. This closes the Δm_{31}^2 discrepancy from -5.7% (bare) to -0.2% (corrected) against PDG 2024 (2.530×10^{-3} eV², NO).

The $m_{\beta\beta} = 0$ prediction holds at all orders: the $d = 5$ sector has $d \bmod 8 = 5$, which means no charge-conjugation matrix C exists on the S^5 spinor bundle. Since no C exists globally, cross-sector couplings cannot construct $\psi^T C \psi$ at any loop order — the Majorana operator is geometrically absent, not merely suppressed. Any $0\nu\beta\beta$ signal immediately falsifies IDWT. $\sum m_\nu = 60.39$ meV is within reach of CMB-S4 (~ 14 meV sensitivity) and the Simons Observatory (~ 30 – 40 meV); IDWT will be confirmed or excluded within this decade.

0.11 11. Dynamical Picture: How Sector Geometry Determines Observable Physics

The $\ell = 0$ part of the kernel sets the sector scales (mass formula). The $\ell = 2$ part generates the derived scale ε (used for the ν_3 closure) and, at loop level, contributes to boson self-energies. The emergent gauge symmetries arise from the isometry groups of the sector manifolds: $\text{SU}(3)_c$ from the isometry $\text{SU}(3)$ of $\mathbb{CP}^2$, and the electroweak $\text{SU}(2)_L \times \text{U}(1)_Y$ from the $\text{U}(2) \subset \text{SU}(3)$ holonomy of $\mathbb{CP}^2$ (hypercharge Y from its T_8).

The physical particle spectrum is the co-fixed-point spectrum of the seed-driven filtration chain — the modes that are simultaneous self-consistent solutions of the coupled sector equations (Part 1 §3b; Part 7 §1.1). A mode that is not a co-fixed point does not form a stable particle: the $d = 3$ modes at $n = 2$ and $n = 3$ are such non-fixed points, consistent with the Completeness Theorem and with observation.

Coordinate containment: whether a force can reach a given particle is fixed by the nesting of their sectors. The sector coordinates form a single chain, $\Xi_2 \subset \Xi_3 \subset \dots$, so any two sectors share the coordinates of the smaller, and a force couples to a mode on those shared coordinates. The $d = 2$ photon’s coordinates lie inside every sector, so it reaches every charged mode; a force seated in a larger sector reaches a mode only through whatever coordinates the two share. This fixes which interactions are possible; the coupling structure of those interactions is the subject of Section 12.

Gravity: variation of the master action with respect to $g_{\mu\nu}$ yields the observer’s Einstein equations $G_{\mu\nu} = 8\pi G_N T_{\mu\nu}^{\text{eff}}$, where $T_{\mu\nu}^{\text{eff}} = \int_{\Xi} T_{\mu\nu}^{\text{Dirac}} d\mu_{\xi}$ is the sector-space integral of the matter stress-energy. G_N is a measured constant of spacetime; the sector geometry Ξ is a fixed background and contributes no gravitational degrees of freedom. Full treatment in Part 4 §3.11.2 [11].

0.12 12. Sector Geometry as Coupling Filter

The coordinate containment principle (Section 11) answers which forces can reach a given particle's sector coordinates. A complementary principle governs the coupling structure of that interaction: **the particle's own sector geometry determines what coupling handles it presents to the world and what entire classes of interaction are geometrically forbidden to it.** The sector quantum number — polarization, color, the Dirac condition, color neutrality — is not an input label assigned to the particle. It is the geometry of the sector expressing itself as a coupling structure.

$d = 2$ ($\mathbb{CP}^1$, $\mathbf{U}(1)$) — **orientation filter.** The photon's two helicity states are the complete internal geometry of $\mathbb{CP}^1$. The coupling to a target goes as $\varepsilon_\mu j^\mu$ — the polarization vector contracted with the current. Perpendicular currents receive zero coupling, not suppression. Polarization is not a property of the photon in addition to its geometry; it is the $\mathbf{U}(1)$ fiber structure of $\mathbb{CP}^1$ expressed as what the photon can and cannot couple to.

$d = 3$ (S^3 , $\mathbf{SO}(4)$) — **weak isospin filter.** The isometry $\mathbf{SO}(4) = \mathbf{SU}(2)_L \times \mathbf{SU}(2)_R$ gives down-type quarks their left-handed weak coupling. The right-handed component does not participate in the weak interaction — $\mathbf{SU}(2)_R$ is latent in the geometry. Color coupling is inherited derivatively via coordinate containment inside Ξ_4 , not from the S^3 structure itself.

$d = 4$ ($\mathbb{CP}^2$, $\mathbf{SU}(3)$) — **color filter.** Color originates here. $\chi(\mathbb{CP}^2) = 3$ — three color coupling handles as a theorem of $\mathbb{CP}^2$ topology, not a parameter. Color conservation is not a dynamical rule; isolated color-nonsinglet asymptotic states are geometrically impossible. Confinement is this filter at the level of which states can be constructed.

$d = 5$ (S^5 , $\mathbf{SO}(6)$) — **Majorana/LNV filter.** The Clifford algebra on S^5 ($d \bmod 8 = 5$) admits no charge-conjugation matrix C on the spinor bundle — C does not exist globally on S^5 . Since no C exists, cross-sector couplings cannot construct $\psi^T C \psi$ at any loop order, and the Majorana mass operator is geometrically forbidden to all orders, not merely at leading order. This is the basis for the $m_{\beta\beta} = 0$ prediction (Section 10). The S^5 Hopf fibration $S^1 \rightarrow S^5 \rightarrow \mathbb{CP}^2$ additionally projects the $\mathbb{CP}^2$ color representation onto its singlet, giving color-neutral neutrinos despite their coordinate support inside Ξ_4 . The $\mathbf{SO}(6) \cong \mathbf{SU}(4)$ structure gives neutrinos their $B - L$ charge.

$d = 6$ ($\mathbb{CP}^3$, $\mathbf{SU}(4)$) — **colour silence filter.** $\chi(\mathbb{CP}^3) = 4$, not 3; colour contributions cancel in the $\mathbf{SU}(4)$ representation. Zero strong coupling at any energy — geometrically absent, not suppressed. The $d = 2$ photon sector ($\mathbb{CP}^1$) sits inside $d = 6$ via the coordinate nesting $\Xi_2 \subset \Xi_6$; the electron-photon coupling follows from coordinate containment, giving pure $\mathbf{U}(1)$ EM with coupling structure fixed by the $\mathbb{CP}^3$ isometry.

$d = 10$ ($\mathbb{CP}^5$, $\mathbf{SU}(6)$) — **Gegenbauer-critical coupling.** At the Gegenbauer critical endpoint, modes sit at the Jacobi coupling boundary — neither freely sector-delocalized nor robustly sector-bound. The tau couples in principle to every sector (its coordinates span all of Ξ_{10}), but coupling weight is distributed with no dominant channel. No single decay mode is selected; every channel is accessible; none carries concentrated weight. This explains the tau's decay pattern: short lifetime (no gap protecting it) but broad distribution across many channels. The geometric back-reaction correction $\delta_\tau = 1/1680$ (Section 7.2) is required because the naive perturbation series does not converge at the critical point.

Relationship to coordinate containment. Coordinate containment is necessary but not sufficient. A particle may have coordinate support in a force's sector and still have zero coupling if its own sector geometry projects the relevant representation to zero. The S^5 Hopf structure projects the $\mathbb{CP}^2$ color representation onto the singlet; the $\mathbb{CP}^3$ index cancellation gives zero strong coupling despite the lepton's coordinates containing Ξ_4 . Both principles together determine the full coupling structure of any particle.

Significance. The Standard Model takes quantum numbers as axioms: quarks have three colors by measurement; neutrinos are (possibly) Dirac by measurement; leptons are color-neutral by measurement. IDWT derives each from the Euler characteristic, Clifford algebra mod 8 periodicity, or Clifford algebra on the respective sector manifold. The coupling structure of every particle in the Standard Model is a theorem of each particle's sector manifold geometry, not a postulate.

0.13 13. Light-Quark Scale Offset and In-Sector Structure

PDG reports light-quark masses in $\overline{\text{MS}}$ at $\mu = 2$ GeV; IDWT computes them from the sector scale $m_{\text{scale},d}$, fixed at the confinement scale $\Lambda_{\text{QCD}} \approx N_c f_\pi = 3 \times 94 = 282$ MeV. The bare-count predictions already sit within the sizable PDG $\overline{\text{MS}}$ uncertainties ($d +0.04\%$, $s +0.57\%$, $u +0.77\%$; $b -0.05\%$) even before the §7.1 confinement-binding correction (which brings s , u to $+0.49\%$, $+0.70\%$), so the overall sector scale is compatible with the data up to a possible scheme/scale offset between the IDWT confinement scale and $\overline{\text{MS}}$ at $\mu = 2$ GeV.

That offset does not account for the whole residual, and the residual is not uniform within a sector. A scheme or scale shift multiplies $m_{\text{scale},d}$ by one common factor for every mode in sector d (the kernel is rank-1), so it cancels identically in the scale-free **in-sector ratio** $S(n_{\text{hi}}, d)/S(n_{\text{lo}}, d)$. Those ratios are not exact: $s/d = 20.000$ against 19.894 ($+0.54\%$), $c/u = 590.33$ against 589.35 ($+0.17\%$), and $t/c = 137.26$ against 135.56 ($+1.25\%$), whereas the boson (Z/W , H/W) and charged-lepton (μ/e) in-sector ratios agree to a few parts in 10^4 . The residual therefore grows with the level gap and is confined to the quark sectors $d = 3$ and $d = 4$: it is n -dependent structure in the combinatorial count $S(n, d)$, not a uniform sector scale error. It is reported as measured accuracy and not absorbed as a correction (§7.1).

The analytic derivation of the scheme conversion from g_s to $\overline{\text{MS}}$ at $\mu = 2$ GeV requires computing the QCD running from Λ_{QCD} to the PDG renormalization point. This is a well-defined calculation (involving the two-loop QCD β -function with $N_f = 3$ active flavors and matching at m_c) that remains to be performed within the IDWT coupling framework. It is flagged as an open item; the current offsets are not in tension with experiment given the large PDG uncertainties on light-quark masses.

0.14 14. Comparison with Other Approaches

Framework	Mass parameters	Gauge structure	Neutrino masses
Standard Model	9 fermion masses free	Given	Free (Yukawa)
Koide relation [3]	Lepton masses only; 1 relation	Not derived	—
Noncommutative geometry [4]	Some constraints	Derived	Not fixed
String landscape	Statistical distribution	In principle	Statistical
IDWT (this work)	All 15 from $n_s = 4$	Derived from sector isometries	60.39 meV predicted

IDWT is distinctive in providing: (a) masses for all 15 predicted particles, (b) a derivation of the sector set from the integer seeds, (c) a spectrum-completeness theorem (no stable mode outside the selected set), and (d) parameter-free neutrino mass predictions falsifiable within this decade.

0.15 15. Falsifiable Predictions

1. $\sum m_\nu = 60.39$ **meV**. A measurement outside $[55, 65]$ meV falsifies the cross-sector scale equation.
2. $m_{\beta\beta} = 0$ **exactly**. Any positive $0\nu\beta\beta$ detection falsifies IDWT.
3. $m_u/m_d = \sqrt{3/14} = 0.46291$ **exactly (Theorem S2)**. A ratio outside $[0.40, 0.55]$ falsifies the coupling derivation.
4. **No new stable particles at any energy (Completeness Theorem)**. Any new particle discovery falsifies IDWT.
5. **Normal neutrino hierarchy**. Inverted hierarchy is excluded by mode ordering.
6. **No stable states at 18.807 MeV or 47.019 MeV**. These $d = 3$ modes ($n = 2$, $n = 3$) are not co-fixed points, so no stable particle forms; they may appear as broad resonances only.

0.16 16. Open Items

1. **CP-violating phase δ .** Requires loop-level computation of the Hopf Chern-Simons integral.
 2. **PMNS mixing angles.** Derived via spectral geometry (Part 9 T6): $\sin^2 \theta_{23} = 0.5590$ (PDG 2024: 0.553, +1.07%), $\sin^2 \theta_{12} = 0.3086$ (PDG 0.307, +0.51%), $\sin^2 \theta_{13} = 0.0211$ (PDG 0.0219, +0.96%). All three from $g_{55} = 96/g_{22}$ and four mode indices. CP phase δ remains open (requires spectral phase integral).
 3. **Δm_{31}^2 discrepancy.** Closed by $\delta_{\nu_3} = \varepsilon \times g_{33} = 1/35$ (cross-sector constructive interference, exact). Corrected $\Delta m_{31}^2 = 2.525 \times 10^{-3} \text{ eV}^2$ (PDG 2024: 2.530×10^{-3} , -0.2%).
 4. **g_1 residual -1.88% .** After 1-loop $U(1)_Y$ running from m_W to m_Z . Remaining gap consistent with 2-loop QED threshold matching between IDWT fiber scheme and $\overline{\text{MS}}$.
 5. **Top index $n_{\text{top}} = 72$.** The product form $\chi(\mathbb{CP}^2) \times \chi(\mathbb{CP}^3) \times \chi(\mathbb{CP}^5) = 3 \times 4 \times 6$ is structurally natural but the selecting condition — why this Euler product and not another combination — has no derivation. This is the leading open dynamical question of the framework.
 6. **The gravitational input G_∞ .** Gravity in IDWT is the back-reaction of $|\Psi_\infty|^2$ on the spacetime metric (Part 4 §3.12.2 [11]); the in-sector Green's function fixes the per-dimension coupling $G_d = G_\infty / [(d-2)S_{d-1}]$, equal to the measured G_N at $d = 3$. G_∞ (equivalently G_N) is a second dimensional input alongside m_e and is not derived from the sector combinatorics. It is not commensurable with the active-force couplings: gravity in IDWT is the curvature of M_∞ sourced by $|\Psi_\infty|^2$, whereas the active forces are the wave's kernel self-coupling on shared sector coordinates — geometry versus coupling, not two forces of the same kind. There is therefore no ratio between gravity and the active forces to fine-tune, and the dimensionless $G_\infty m_e^2 \approx 2.2 \times 10^{-44}$ is simply an input, not a prediction. No fine-tuning is hidden anywhere else either: IDWT masses are discrete spectral ranks $S(n, d)$, not action coefficients, so they receive no additive UV renormalisation and the Higgs-style hierarchy problem does not arise (Part 8 §3a [11]). The genuinely open question is narrower — whether G_∞ could instead be fixed geometrically, reducing the two dimensional inputs to one — which the present construction does not achieve.
 7. **Spectral action EW scale.** The spectral action gives $\sqrt{\text{Tr}(D^2)} = 248.3 \text{ GeV}$, within $+0.82\%$ of v_{Higgs} ; both are spectral quantities built from the same seeds, so the residual is a spectral-normalisation artefact (Part 9 §T7 [11]), not an independent confirmation. G_N enters the Einstein–Hilbert term as a measured input.
-

0.17 17. Conclusion

Starting from three integer seeds $\{n_d = 1, n_u = 3, n_{\text{top}} = 72\}$ — whose composite $n_s = n_d + n_u = 4$ is independently forced by the topological fixed point $S(4, 4) = 35$ — and the dimensional reference m_e , IDWT provides a combinatorial derivation of the complete Standard Model mass spectrum, the electroweak sector, CKM observables, and absolute neutrino masses to high accuracy. Five theorems ground the framework in spectral geometry, with complete proofs supplied above. The Completeness Theorem proves no additional particles exist within the framework — a consequence, not a claim. The τ back-reaction and ν_3 closure corrections are derived from the kernel structure without additional inputs; the confinement-binding correction for the colour sectors ($d = 3, 4$) has a derived functional form, with its $d = 4$ scale derived from the colour energy law and its $d = 3$ scale anchored.

A complementary structural result (Section 12) shows that each sector geometry constitutes a coupling filter for its particles. The sector quantum number — polarization, color, the Dirac condition, total color silence, Gegenbauer-critical marginal coupling — is not an input label but the geometry expressing itself as a coupling structure, derived from the Euler characteristic, Clifford algebra mod 8 periodicity, and Clifford algebra on the sector manifold. Coordinate containment determines whether coupling is possible; the sector geometry determines the structure and forbidden classes of that coupling. These are independent principles that together give a complete geometric account of why each particle interacts the way it does.

The framework makes five concrete predictions falsifiable with near-term experiments, most critically $\sum m_\nu = 60.39$ meV (corrected; bare 59.00 meV, with $\delta_{\nu_3} = 1/35$ derived from the cross-sector kernel; CMB-S4 target: ~ 14 meV) and $m_{\beta\beta} = 0$ at all orders (current KamLAND-Zen bound: $< 28\text{--}122$ meV, NME-dependent [6]). These will confirm or exclude IDWT within this decade.

0.18 Acknowledgements

The author thanks Claude, GPT, Meta, Grok, Perplexity, Z.ai, and Mistral Le Chat.

0.19 Appendix A: Quick-Reference Summary

All results from $n_s = 4$ and m_e alone.

Identity	Statement	Source
Hockey-stick	$S(n, d) = S(n, d-1) + S(n-1, d)$	Combinatorics
Weyl law	$S(n, 2k+1) = \frac{1}{2} N_{D_{S_{2k+1}}} (n-1)$	Theorem S1
m_u/m_d	$\sqrt{g_{44}/g_{33}} = \sqrt{3/14}$ (exact)	Theorem S2
g_{22}	$(S(n_s, 3) - n_u)^2 \times (S(n_u, 4) - S(n_u, 3))/2 = 722.5$	Theorem S3
Sector set	$D = \{2, 3, 4, 5, 6, 10\}$ from $72 = N_c \times n_s \times N_f$	Theorem S4
$\ell = 2$ scale	$\varepsilon = 1/(280\sqrt{7}) = 0.001350$ (enters δ_{ν_3} ; not applied to quarks)	$\ell = 2$ kernel
Back-reaction	$1/1680 = 1/(n_s n_u (n_s + n_u) S(n_s, 3))$	$d = 6 \rightarrow d = 10$
$\sum m_\nu$	60.39 meV (corrected; bare: 59.00 meV, $\delta_{\nu_3} = 1/35$)	§10
$m_{\beta\beta}$	0 (all orders; no C on S^5 bundle, $d \bmod 8 = 5$)	§10, §12
CKM unitarity	$ V_{ud} ^2 + V_{us} ^2 + V_{ub} ^2 = 1$ (exact)	§9

All 15 mode indices ($n_s = 4$ forces every entry):

Particle	d	n	Particle	d	n
d quark	3	1	ν_1	5	10
u quark	4	3	e^-	6	13
s quark	3	4	ν_2	5	15
c quark	4	20	ν_3	5	22
t quark	4	72	τ^-	10	23
b quark	3	beat	μ^-	6	35
$W^\pm$	2	76	Z^0	2	81
H	2	95			

All 6 coupling constants (exact from $n_s = 4$):

Coupling	Value	Sector
g_{33}	$8\sqrt{7} \approx 21.166$	$d = 3$ quarks
g_{44}	$12/\sqrt{7} \approx 4.536$	$d = 4$ quarks
g_{22}	722.5 (Theorem S3)	$d = 2$ gauge
g_{55}	$96/722.5 \approx 0.1329$	$d = 5$ neutrinos
g_{66}	1/4 (seed)	$d = 6$ leptons
$g_{10,10}$	1/4 (seed, shared with $d = 6$)	$d = 10$ tau

0.20 References

1. C. Bär, *Geom. Funct. Anal.* **6**, 899 (1996). (*Dirac spectrum on S^n*)
2. H. B. Lawson and M.-L. Michelsohn, *Spin Geometry* (Princeton University Press, 1989). (*Clifford algebra mod 8 periodicity, Majorana conditions*)
3. Y. Koide, *Lett. Nuovo Cim.* **34**, 201 (1983).
4. A. Chamseddine, A. Connes, and V. Mukhanov, *JHEP* **2015**, 104 (2015).
5. Planck Collaboration (N. Aghanim et al.), “Planck 2018 results. VI. Cosmological parameters,” *A&A* **641**, A6 (2020), arXiv:1807.06209.
6. KamLAND-Zen Collaboration, “Search for Majorana Neutrinos with the Complete KamLAND-Zen Dataset,” *Phys. Rev. Lett.* (2024), arXiv:2406.11438.
7. CMB-S4 Collaboration, arXiv:1610.02743 (2016).
8. Simons Observatory Collaboration (P. Ade et al.), *JCAP* **2019**, 056 (2019).
9. KATRIN Collaboration (M. Aker et al.), *Nature Phys.* **18**, 160 (2022).
10. Particle Data Group (R. L. Workman et al.), *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022).
11. Fedge No, *Infinite-Dimensional Wave Theory: Technical Documentation* (10 volumes), Zenodo, doi:10.5281/zenodo.19767493 (2026).