


Filipp Uskov
Skoltech, Moscow State University
in collaboration with Oleg Lychkovsky and Elena Shpagina

Model



Typical Hamiltonian: $H = J \sum_{\langle i, j \rangle} (\sigma_i \sigma_j)$

where $\langle i, j \rangle$ denotes neighboring sites

Scalar and mixed products of vectors of Pauli matrices:

$$(\sigma_1 \sigma_3) = \delta^{\alpha\beta} \cdot \sigma_1^\alpha \otimes 1_2 \otimes \sigma_3^\beta \otimes 1_4 \otimes 1_5$$

$$(\sigma_1 \sigma_3 \sigma_4) = \varepsilon^{\alpha\beta\gamma} \cdot \sigma_1^\alpha \otimes 1_2 \otimes \sigma_3^\beta \otimes \sigma_4^\gamma \otimes 1_5$$

$$\alpha, \beta, \gamma \in \{x, y, z\}$$

Squaring parameterization of density matrix [1]

Common conditions: $\rho^+ = \rho$; $\text{Tr} \rho = 1$; $\langle \rho \rangle \geq 0$

From ρ to τ : $\rho = \frac{\tau^2}{\text{Tr} \tau^2}$; for τ only one condition: $\tau^+ = \tau$

$$\rho = \frac{1}{2^n} a_i A_i; \quad \tau = b_i A_i$$

Basis:

$$\{A_i\} = \{1, (\sigma_j \sigma_k), (\sigma_j \sigma_k)(\sigma_l \sigma_m), \dots\}$$

See illustrations in poster by Elena Shpagina

Symbolic relations and simplifications

• Algorithm which based on identity of Pauli ($\sigma^\alpha \sigma^\beta = \delta^{\alpha\beta} + i \varepsilon^{\alpha\beta\gamma} \sigma^\gamma$) was implemented in Wolfram Mathematica

$$\begin{aligned} (\sigma_1 \sigma_2)^2 &= 3 - 2(\sigma_1 \sigma_2) \\ (\sigma_1 \sigma_2)(\sigma_2 \sigma_3) &= (\sigma_1 \sigma_3) - i(\sigma_1 \sigma_2 \sigma_3) \\ (\sigma_1 \sigma_2)(\sigma_1 \sigma_2 \sigma_3) &= -(\sigma_1 \sigma_2 \sigma_3) - 2i(\sigma_1 \sigma_3) + 2i(\sigma_2 \sigma_3) \\ (\sigma_1 \sigma_2 \sigma_3)(\sigma_1 \sigma_2) &= -(\sigma_1 \sigma_2 \sigma_3) + 2i(\sigma_1 \sigma_3) - 2i(\sigma_2 \sigma_3) \\ (\sigma_1 \sigma_2)(\sigma_2 \sigma_3 \sigma_4) &= (\sigma_1 \sigma_3 \sigma_4) - i(\sigma_1 \sigma_3)(\sigma_2 \sigma_4) + i(\sigma_1 \sigma_4)(\sigma_2 \sigma_3) \\ (\sigma_2 \sigma_3 \sigma_4)(\sigma_1 \sigma_2) &= (\sigma_1 \sigma_3 \sigma_4) + i(\sigma_1 \sigma_3)(\sigma_2 \sigma_4) - i(\sigma_1 \sigma_4)(\sigma_2 \sigma_3) \\ (\sigma_1 \sigma_2 \sigma_3)^2 &= 6 - 2(\sigma_1 \sigma_2) - 2(\sigma_1 \sigma_3) - 2(\sigma_2 \sigma_3) \\ (\sigma_1 \sigma_2 \sigma_3)(\sigma_1 \sigma_2 \sigma_4) &= -(\sigma_1 \sigma_3)(\sigma_2 \sigma_4) - (\sigma_1 \sigma_4)(\sigma_2 \sigma_3) + 2(\sigma_3 \sigma_4) + i(\sigma_1 \sigma_3 \sigma_4) + i(\sigma_2 \sigma_3 \sigma_4) \\ (\sigma_1 \sigma_2 \sigma_3)(\sigma_1 \sigma_4 \sigma_5) &= +(\sigma_2 \sigma_4)(\sigma_3 \sigma_5) - (\sigma_2 \sigma_5)(\sigma_3 \sigma_4) - i(\sigma_1 \sigma_2)(\sigma_3 \sigma_4 \sigma_5) + i(\sigma_1 \sigma_3)(\sigma_2 \sigma_4 \sigma_5) \end{aligned}$$

• Scalar product of matrices also was implemented in Wolfram Mathematica:

$$(A, B) = \text{Tr}(A^+ B) = \text{Tr}(AB); \quad A^+ = A$$

$$(A, B) = 0, \text{ if } A \text{ and } B \text{ contains different spins}$$

$$\text{Tr}((\sigma_i \sigma_j)(\sigma_j \sigma_k) \dots (\sigma_l \sigma_m)(\sigma_m \sigma_i)) = 3 \cdot 2^n$$

$$\text{Tr}((\sigma_i \sigma_j \sigma_k)(\sigma_i \sigma_j \sigma_n)(\sigma_k \sigma_l) \dots (\sigma_m \sigma_n)) =$$

$$= \text{Tr}((\sigma_i \sigma_j \sigma_k)(\sigma_i \sigma_j \sigma_k)) = 6 \cdot 2^n$$

$$(A, B) = 3^C \cdot 2^n, \text{ where } C - \text{number of cycles}$$










Illustration from [6]

Results


1d:

cluster	E_{gs}/N
	-2.1547
	-1.92789
	-1.99486
	-1.89083
	-1.92853

2d:



$$E_{gs}/N > 3$$



$$E_{gs}/N > -2,9685$$

1d exact theoretical [7]: $E_{gs}/N = 1 - 4 \ln 2 = -1.7725$

Challenge and main ideas

$$??? \leq E_{gs} \leq \langle \psi | H | \psi \rangle$$

Our challenge

well known variational method

Divide **full** crystal to **clusters**:

N – number of spins in crystal

M – number of clusters

d - number of bonds by 1 spin

m – number of bonds in cluster

$$M = \frac{Nd}{m}$$

$$\bullet \text{Schrödinger equation: } H_{full} = \sum_{i=1}^M H_{icl} \Rightarrow E_{gs full} \geq \sum_{i=1}^M E_{i gs cl}$$

$$E_{gs full} / N \geq \frac{d}{m} E_{gs cl}$$

• Variational method:

$$E_{gs full} = \min_{\psi} \langle \psi | H | \psi \rangle = \min_{\rho_{full} \in M_{full}^S} \text{Tr}_{full} H_{full} \rho_{full} = \min_{\rho_{full} \in M_{full}^S} \text{Tr}_{full} H_{full} \rho_{full} =$$

$$= M \min_{\rho_{full} \in M_{full}^S} \text{Tr}_{full} H_{icl} \rho_{full} = M \min_{\rho_{full} \in M_{full}^S} \text{Tr}_{cl} (H_{icl} \text{Tr}_{full-cl} \rho_{full}) =$$

$$= M \min_{\rho_{cl} \in M_{cl}^{trS}} \text{Tr}_{cl} H_{icl} \rho_{cl} \geq M \min_{\rho_{cl} \in M_{cl}'} \text{Tr}_{cl} H_{icl} \rho_{cl}$$

$$E_{gs full} / N \geq \frac{d}{m} \min_{\rho_{cl} \in M_{cl}'} \text{Tr}_{cl} H_{icl} \rho_{cl};$$

Schrödinger equation

$$H\rho = E\rho \Leftrightarrow H \frac{\tau^2}{\text{Tr} \tau^2} = E \frac{\tau^2}{\text{Tr} \tau^2} \Leftrightarrow H\tau = E\tau$$

$$HA_j a_j = A_i h_{ij} a_j + B_i d_{ij} a_j = EA_j a_j \Rightarrow \begin{cases} h_{ij} a_j = E a_i \\ d_{ij} a_j = 0 \end{cases}$$

$$(A_i, A_j) = g_{ij}$$

$$\text{Tr}(A_i H A_j) = \eta_{ij} \quad h_{ij} = g_{ik}^{-1} \eta_{kj}$$

We take into account the symmetries of commutation of spins in cluster.

Example:

$$\begin{aligned} A_0 &= 1 \\ A_1 &= (\sigma_1 \sigma_2) + (\sigma_2 \sigma_3) + (\sigma_3 \sigma_4) + (\sigma_4 \sigma_1) + (\sigma_1 \sigma_3) + (\sigma_2 \sigma_4) \\ A_2 &= (\sigma_1 \sigma_2)(\sigma_2 \sigma_3) + (\sigma_2 \sigma_3)(\sigma_3 \sigma_4) + (\sigma_3 \sigma_4)(\sigma_4 \sigma_1) + (\sigma_4 \sigma_1)(\sigma_1 \sigma_2) \\ &\quad + (\sigma_2 \sigma_3)(\sigma_1 \sigma_3) + (\sigma_2 \sigma_3)(\sigma_1 \sigma_4) + (\sigma_2 \sigma_3)(\sigma_3 \sigma_4) \\ &\quad + (\sigma_3 \sigma_4)(\sigma_1 \sigma_2) + (\sigma_3 \sigma_4)(\sigma_2 \sigma_4) + (\sigma_3 \sigma_4)(\sigma_4 \sigma_1) \\ &\quad + (\sigma_4 \sigma_1)(\sigma_1 \sigma_2) + (\sigma_4 \sigma_1)(\sigma_2 \sigma_3) + (\sigma_4 \sigma_1)(\sigma_3 \sigma_4) \\ A_3 &= (\sigma_1 \sigma_2)(\sigma_3 \sigma_4) + (\sigma_1 \sigma_3)(\sigma_2 \sigma_4) + (\sigma_1 \sigma_4)(\sigma_2 \sigma_3) \\ &\quad + (\sigma_2 \sigma_3)(\sigma_1 \sigma_4) + (\sigma_2 \sigma_4)(\sigma_1 \sigma_3) + (\sigma_3 \sigma_4)(\sigma_1 \sigma_2) \\ &\quad + (\sigma_3 \sigma_4)(\sigma_2 \sigma_1) + (\sigma_4 \sigma_1)(\sigma_2 \sigma_3) + (\sigma_4 \sigma_1)(\sigma_3 \sigma_2) \\ A_4 &= (\sigma_1 \sigma_2)(\sigma_3 \sigma_4) + (\sigma_1 \sigma_3)(\sigma_2 \sigma_4) + (\sigma_1 \sigma_4)(\sigma_2 \sigma_3) \end{aligned} \quad h_{ij} = \begin{pmatrix} 0 & 12 & 0 & 0 & 0 \\ 1 & -2 & 3 & 0 & 0 \\ 0 & 2 & 0 & 10 & 0 \\ 0 & 0 & 1 & 2 & 1 \\ 0 & 0 & 0 & 4 & 0 \end{pmatrix}$$

$$d_{ij} = 0$$

$$E_{gs} = -6, -4, 4, 2, 0$$

Variational method

$$\text{Numeric method: } \min \text{Tr} H \rho = \min_{\{b_i\}} \frac{b_i b_j}{b_l b_m} \frac{\eta_{ij}}{g_{lm}}$$

$$\text{Lagrange method: } L(\{b_k\}, \lambda) = b_i b_j (\eta_{ij} - \lambda g_{ij}) + \lambda$$

$$\min \text{Tr} H \rho = \min_{\{b_i\}} b_i b_j \eta_{ij} \Rightarrow b_i (\eta_{ik} - \lambda g_{ik}) = 0$$

$$\text{with condition } b_l b_m g_{lm} = 1 \Leftrightarrow b_i (g_{ij}^{-1} \eta_{jk}) = b_i h_{ik} = \lambda b_k$$

$$\min_{\{b_i\}} b_i b_j \eta_{ij} = \min \lambda$$

If we don't take into account the symmetries of commutation of spins

Variational method is identical to Schrodinger equation

1. N. Il'in, E. Shpagina, F. Uskov, O. Lychkovskiy, Squaring parametrization of constrained and unconstrained sets of quantum states. J. Phys. A: Math. Theor. 51 (2018) 085301
2. R. Tarrach, R. Valenti (1990) Exact lower bounds to the ground state of spin systems: The two-dimensional S = 1/2 antiferromagnetic Heisenberg model Physical review B, 1990.
3. P. W. Anderson, Limits on the Energy of the Antiferromagnetic Ground State, Letters to the editor 1951
4. Tillmann Baumgratz1 and Martin B Plenio, Lower bounds for ground states of condensed matter systems. New Journal of Physics 14 (2012) 023027 (21pp)
5. David A. Mazziotti, Variational minimization of atomic and molecular ground-state energies via the two-particle reduced density matrix, PHYSICAL REVIEW A, VOLUME 65, (2002) 062511
6. K.S.D. Beach, A.W. Sandvik Some formal results for the valence bond basis Nuclear Physics B 750 [FS] (2006) 142–178
7. Daniel C. Mattis, C. Y. Pan, Ground-State Energy of Heisenberg antiferromagnet for Spins s=1/2 and s=1 in d=1 and 2 Dimensions. PHYSICAL REVIEW LETTERS, VOLUME 61, (1988)
8. Маттис. Теория магнетизма. Москва, Мир, 1967