

Theoretical Models of Power Spectrum

The papers for this code:

V12: <https://arxiv.org/pdf/1207.0839>

V13: <https://arxiv.org/pdf/1308.6294>

O14: <https://arxiv.org/pdf/1312.4214v2>

H19: <https://arxiv.org/pdf/1906.02875>

Q is D , in this software 'D' is a function, so we instead use Q.

PL is the linear power spectrum.

$\mathcal{H} = a H$ where a is the scale factor, H is the Hubble parameter.

Pmnc denotes the loop terms defined in C.Howlett 2019 (H19).

Pmn denotes the loop terms defined in Vlah12 (V12).

We want to check if there is any bugs in Pmnc of H19.

Sec.1: Comparing the expressions of the power spectrum of H19 to those of V12 and O14

1.1: The density power spectrum

This is the density power spectrum of H19, see the Eq.A2 of H19:

$$In[*]:= P\delta c = p00c + \mu^2 (2 p01c + p02c + p11c) + \mu^4 \left(p03c + p04c + p12c + p13c + \frac{1}{4} \times p22c \right); (* Eq.A2 of H19 *)$$

ExpandAll[Pδc]

[展开全部](#)

Out[*]=

$$p00c + 2 p01c \mu^2 + p02c \mu^2 + p11c \mu^2 + p03c \mu^4 + p04c \mu^4 + p12c \mu^4 + p13c \mu^4 + \frac{p22c \mu^4}{4}$$

On the other hand, the expression of density power of V12 is given by :

$$\begin{aligned} \text{In}[*]:= \mathbf{P}\delta = & \mathbf{p00} + \left(\frac{\mathbf{k}\mu}{\mathcal{H}}\right)^2 \mathbf{p11} + \frac{1}{4} \times \left(\frac{\mathbf{k}\mu}{\mathcal{H}}\right)^4 \mathbf{p22} + \\ & 2 \left(\frac{-\mathbf{i}\mathbf{k}\mu}{\mathcal{H}} \mathbf{p01} + \left(-\frac{1}{2} \times \left(\frac{\mathbf{k}\mu}{\mathcal{H}}\right)^2\right) \mathbf{p02} + \frac{\mathbf{i}}{6} \times \left(\frac{\mathbf{k}\mu}{\mathcal{H}}\right)^3 \mathbf{p03} + \left(-\frac{\mathbf{i}}{2} \times \left(\frac{\mathbf{k}\mu}{\mathcal{H}}\right)^3\right) \mathbf{p12} + \right. \\ & \left. \left(-\frac{1}{6} \times \left(\frac{\mathbf{k}\mu}{\mathcal{H}}\right)^4 \mathbf{p13}\right) + \frac{1}{24} \times \left(\frac{\mathbf{k}\mu}{\mathcal{H}}\right)^4 \mathbf{p04} \right); (* \text{ Eq.2.23 of V12 } *) \end{aligned}$$

ExpandAll[Pδ]

展开全部

Out[*]=

$$\begin{aligned} \mathbf{p00} - \frac{2 \mathbf{i} \mathbf{k} \mathbf{p01} \mu}{\mathcal{H}} - \frac{\mathbf{k}^2 \mathbf{p02} \mu^2}{\mathcal{H}^2} + \frac{\mathbf{k}^2 \mathbf{p11} \mu^2}{\mathcal{H}^2} + \\ \frac{\mathbf{i} \mathbf{k}^3 \mathbf{p03} \mu^3}{3 \mathcal{H}^3} - \frac{\mathbf{i} \mathbf{k}^3 \mathbf{p12} \mu^3}{\mathcal{H}^3} + \frac{\mathbf{k}^4 \mathbf{p04} \mu^4}{12 \mathcal{H}^4} - \frac{\mathbf{k}^4 \mathbf{p13} \mu^4}{3 \mathcal{H}^4} + \frac{\mathbf{k}^4 \mathbf{p22} \mu^4}{4 \mathcal{H}^4} \end{aligned}$$

Comparing the above $\mathbf{P}\delta$ to $\mathbf{P}\delta\mathbf{c}$, we can find the relations between \mathbf{P}_{mn} of H19 and \mathbf{P}_{mn} of V12, which are given by :

In[*]:= Solve[p00c == p00, p00c]

解方程

$$\text{Solve}\left[2 \mathbf{p01c} \mu^2 == -\frac{2 \mathbf{i} \mathbf{k} \mathbf{p01} \mu}{\mathcal{H}}, \mathbf{p01c}\right]$$

解方程

$$\text{Solve}\left[\mathbf{p02c} \mu^2 == -\frac{\mathbf{k}^2 \mathbf{p02} \mu^2}{\mathcal{H}^2}, \mathbf{p02c}\right]$$

解方程

$$\text{Solve}\left[\mathbf{p11c} \mu^2 == \frac{\mathbf{k}^2 \mathbf{p11} \mu^2}{\mathcal{H}^2}, \mathbf{p11c}\right]$$

解方程

$$\text{Solve}\left[\mathbf{p03c} \mu^4 == \frac{\mathbf{i} \mathbf{k}^3 \mathbf{p03} \mu^3}{3 \mathcal{H}^3}, \mathbf{p03c}\right]$$

解方程

$$\text{Solve}\left[\mathbf{p04c} \mu^4 == \frac{\mathbf{k}^4 \mathbf{p04} \mu^4}{12 \mathcal{H}^4}, \mathbf{p04c}\right]$$

解方程

$$\text{Solve}\left[\mathbf{p12c} \mu^4 == -\frac{\mathbf{i} \mathbf{k}^3 \mathbf{p12} \mu^3}{\mathcal{H}^3}, \mathbf{p12c}\right]$$

解方程

$$\text{Solve}\left[\mathbf{p13c} \mu^4 == -\frac{\mathbf{k}^4 \mathbf{p13} \mu^4}{3 \mathcal{H}^4}, \mathbf{p13c}\right]$$

解方程

$$\text{Solve}\left[\frac{\mathbf{p22c} \mu^4}{4} == \frac{\mathbf{k}^4 \mathbf{p22} \mu^4}{4 \mathcal{H}^4}, \mathbf{p22c}\right]$$

解方程

Clear["Global`*"]

清除

Out[*]=

$$\{\{\mathbf{p00c} \rightarrow \mathbf{p00}\}\}$$

Out[*]=

$$\left\{\left\{\mathbf{p01c} \rightarrow -\frac{\mathbf{i} \mathbf{k} \mathbf{p01}}{\mathcal{H} \mu}\right\}\right\}$$

Out[*]=

$$\left\{ \left\{ p_{02c} \rightarrow -\frac{k^2 p_{02}}{\mathcal{H}^2} \right\} \right\}$$

Out[*]=

$$\left\{ \left\{ p_{11c} \rightarrow \frac{k^2 p_{11}}{\mathcal{H}^2} \right\} \right\}$$

Out[*]=

$$\left\{ \left\{ p_{03c} \rightarrow \frac{i k^3 p_{03}}{3 \mathcal{H}^3 \mu} \right\} \right\}$$

Out[*]=

$$\left\{ \left\{ p_{04c} \rightarrow \frac{k^4 p_{04}}{12 \mathcal{H}^4} \right\} \right\}$$

Out[*]=

$$\left\{ \left\{ p_{12c} \rightarrow -\frac{i k^3 p_{12}}{\mathcal{H}^3 \mu} \right\} \right\}$$

Out[*]=

$$\left\{ \left\{ p_{13c} \rightarrow -\frac{k^4 p_{13}}{3 \mathcal{H}^4} \right\} \right\}$$

Out[*]=

$$\left\{ \left\{ p_{22c} \rightarrow \frac{k^4 p_{22}}{\mathcal{H}^4} \right\} \right\}$$

1.2: The momentum power spectrum

This is the momentum power spectrum of H19, see Eq.A3 of H19:

$$In[*]:= \text{Pmc} = \frac{\mathcal{H}^2}{k^2} \times (p_{11c} + \mu^2 \times (2 p_{12c} + 3 p_{13c} + p_{22c})); (* \text{ Eq.A3 of H19 } *)$$

ExpandAll[Pmc]

[展开全部](#)

Out[*]=

$$\frac{p_{11c} \mathcal{H}^2}{k^2} + \frac{2 p_{12c} \mathcal{H}^2 \mu^2}{k^2} + \frac{3 p_{13c} \mathcal{H}^2 \mu^2}{k^2} + \frac{p_{22c} \mathcal{H}^2 \mu^2}{k^2}$$

On the other hand, the expression of momentum power of O14 is given by :

$$In[*]:= \text{Solve}\left[\left(\frac{k \mu}{\mathcal{H}}\right)^2 \text{Pm} == \left(\frac{k \mu}{\mathcal{H}}\right)^2 p_{11} - 2 i \left(\frac{k \mu}{\mathcal{H}}\right)^3 p_{12} - \left(\frac{k \mu}{\mathcal{H}}\right)^4 p_{13} + \left(\frac{k \mu}{\mathcal{H}}\right)^4 p_{22}, \text{Pm}\right]$$

[解方程](#)

(* Eq.4.2 of O14 *)

Out[*]=

$$\left\{ \left\{ \text{Pm} \rightarrow \frac{p_{11} \mathcal{H}^2 - 2 i k p_{12} \mathcal{H} \mu - k^2 p_{13} \mu^2 + k^2 p_{22} \mu^2}{\mathcal{H}^2} \right\} \right\}$$

$$\text{In}[*]:= \text{Pm} = \frac{\text{p11} \mathcal{H}^2 - 2 \text{i} k \text{p12} \mathcal{H} \mu - k^2 \text{p13} \mu^2 + k^2 \text{p22} \mu^2}{\mathcal{H}^2};$$

(*it should be taken from the above eq*)

ExpandAll[Pm]

[展开全部](#)

Out[*]=

$$\text{p11} - \frac{2 \text{i} k \text{p12} \mu}{\mathcal{H}} - \frac{k^2 \text{p13} \mu^2}{\mathcal{H}^2} + \frac{k^2 \text{p22} \mu^2}{\mathcal{H}^2}$$

Comparing the above Pm to Pmc, we can find the relation between Pmn of H19 and Pmn of V12 :

$$\text{In}[*]:= \text{Solve}\left[\frac{\text{p11c} \mathcal{H}^2}{k^2} == \text{p11}, \text{p11c}\right]$$

[解方程](#)

$$\text{Solve}\left[\frac{2 \text{p12c} \mathcal{H}^2 \mu^2}{k^2} == -\frac{2 \text{i} k \text{p12} \mu}{\mathcal{H}}, \text{p12c}\right]$$

[解方程](#)

$$\text{Solve}\left[\frac{3 \text{p13c} \mathcal{H}^2 \mu^2}{k^2} == -\frac{k^2 \text{p13} \mu^2}{\mathcal{H}^2}, \text{p13c}\right]$$

[解方程](#)

$$\text{Solve}\left[\frac{\text{p22c} \mathcal{H}^2 \mu^2}{k^2} == \frac{k^2 \text{p22} \mu^2}{\mathcal{H}^2}, \text{p22c}\right]$$

[解方程](#)

Clear["Global`*"]

[清除](#)

Out[*]=

$$\left\{\left\{\text{p11c} \rightarrow \frac{k^2 \text{p11}}{\mathcal{H}^2}\right\}\right\}$$

Out[*]=

$$\left\{\left\{\text{p12c} \rightarrow -\frac{\text{i} k^3 \text{p12}}{\mathcal{H}^3 \mu}\right\}\right\}$$

Out[*]=

$$\left\{\left\{\text{p13c} \rightarrow -\frac{k^4 \text{p13}}{3 \mathcal{H}^4}\right\}\right\}$$

Out[*]=

$$\left\{\left\{\text{p22c} \rightarrow \frac{k^4 \text{p22}}{\mathcal{H}^4}\right\}\right\}$$

The above equations are the same as the Eqs of Sec 1.1 .

1.3: The cross power spectrum

This is the cross power spectrum of H19:

$$\text{In}[*]:= \text{P}\delta\text{mc} = \frac{\mathcal{H}}{k} \times \mu \times \left(\text{p01c} + \text{p02c} + \text{p11c} + \mu^2 \times \left(\frac{3}{2} \times \text{p03c} + 2 \text{p04c} + 2 \text{p12c} + 3 \text{p13c} + \frac{1}{2} \times \text{p22c} \right) \right);$$

ExpandAll[Pδmc]

[展开全部](#)

Out[*]=

$$\frac{\frac{\mathcal{H}}{k} \text{p01c}}{k} + \frac{\frac{\mathcal{H}}{k} \text{p02c}}{k} + \frac{\frac{\mathcal{H}}{k} \text{p11c}}{k} + \frac{3 \frac{\mathcal{H}}{k} \text{p03c}}{2k} + \frac{2 \frac{\mathcal{H}}{k} \text{p04c}}{k} + \frac{2 \frac{\mathcal{H}}{k} \text{p12c}}{k} + \frac{3 \frac{\mathcal{H}}{k} \text{p13c}}{k} + \frac{\frac{\mathcal{H}}{k} \text{p22c}}{2k}$$

In practice, $\text{P}\delta\text{mc}$ should be the imaginary part of the above equation, i.e.

$$\text{P}\delta\text{mc} = \frac{\mathcal{H}\mu}{k} \left(\text{p01c} + \text{p02c} + \text{p11c} + \mu^2 \left(\frac{3}{2} \text{p03c} + 2 \text{p04c} + 2 \text{p12c} + 3 \text{p13c} + \frac{1}{2} \text{p22c} \right) \right).$$

On the other hand, the expression of cross power of O14 is given by :

$$\text{In}[*]:= \text{Solve}\left[\left(-\frac{\mathcal{H}k\mu}{\mathcal{H}}\right) \text{P}\delta\text{m} == \right.$$

[解方程](#)

$$\left. -\frac{\mathcal{H}k\mu}{\mathcal{H}} \text{p01} + \left(\frac{k\mu}{\mathcal{H}}\right)^2 \text{p11} - \left(\frac{k\mu}{\mathcal{H}}\right)^2 \text{p02} + \frac{\mathcal{H}}{2} \left(\frac{k\mu}{\mathcal{H}}\right)^3 \text{p03} - 2 \frac{\mathcal{H}}{2} \left(\frac{k\mu}{\mathcal{H}}\right)^3 \text{p12} + \frac{1}{6} \left(\frac{k\mu}{\mathcal{H}}\right)^4 \text{p04} - \left(\frac{k\mu}{\mathcal{H}}\right)^4 \text{p13} + \frac{1}{2} \left(\frac{k\mu}{\mathcal{H}}\right)^4 \text{p22}, \text{P}\delta\text{m} \right] (* \text{ Eq.4.1 of O14 } *)$$

Out[*]=

$$\left\{ \left\{ \text{P}\delta\text{m} \rightarrow \frac{1}{6 \mathcal{H}^3} \left(3 \left(2 \text{p01} \mathcal{H}^3 - k^2 \text{p03} \mathcal{H} \mu^2 + 4 k^2 \text{p12} \mathcal{H} \mu^2 \right) + \mathcal{H} \left(-6 k \text{p02} \mathcal{H}^2 \mu + 6 k \text{p11} \mathcal{H}^2 \mu + k^3 \text{p04} \mu^3 - 6 k^3 \text{p13} \mu^3 + 3 k^3 \text{p22} \mu^3 \right) \right) \right\} \right\}$$

$$\text{In}[*]:= \text{P}\delta\text{m} = \frac{1}{6 \mathcal{H}^3} \left(3 \left(2 \text{p01} \mathcal{H}^3 - k^2 \text{p03} \mathcal{H} \mu^2 + 4 k^2 \text{p12} \mathcal{H} \mu^2 \right) + \mathcal{H} \left(-6 k \text{p02} \mathcal{H}^2 \mu + 6 k \text{p11} \mathcal{H}^2 \mu + k^3 \text{p04} \mu^3 - 6 k^3 \text{p13} \mu^3 + 3 k^3 \text{p22} \mu^3 \right) \right);$$

(*this should be taken from the above eq.*)

ExpandAll[Pδm]

[展开全部](#)

Out[*]=

$$\text{p01} - \frac{\mathcal{H}k \text{p02} \mu}{\mathcal{H}} + \frac{\mathcal{H}k \text{p11} \mu}{\mathcal{H}} - \frac{k^2 \text{p03} \mu^2}{2 \mathcal{H}^2} + \frac{2 k^2 \text{p12} \mu^2}{\mathcal{H}^2} + \frac{\mathcal{H}k^3 \text{p04} \mu^3}{6 \mathcal{H}^3} - \frac{\mathcal{H}k^3 \text{p13} \mu^3}{\mathcal{H}^3} + \frac{\mathcal{H}k^3 \text{p22} \mu^3}{2 \mathcal{H}^3}$$

Comparing the above $\text{P}\delta\text{m}$ to $\text{P}\delta\text{mc}$ we can find the the relation between Pmn of H19 and Pmn of V12:

In[*]:= **Solve** $\left[\frac{\mathfrak{i} \text{p01c } \mathcal{H} \mu}{k} == \text{p01}, \text{p01c}\right]$
 解方程

Solve $\left[\frac{\mathfrak{i} \text{p02c } \mathcal{H} \mu}{k} == -\frac{\mathfrak{i} k \text{p02 } \mu}{\mathcal{H}}, \text{p02c}\right]$
 解方程

Solve $\left[\frac{\mathfrak{i} \text{p11c } \mathcal{H} \mu}{k} == \frac{\mathfrak{i} k \text{p11 } \mu}{\mathcal{H}}, \text{p11c}\right]$
 解方程

Solve $\left[\frac{3 \mathfrak{i} \text{p03c } \mathcal{H} \mu^3}{2 k} == -\frac{k^2 \text{p03 } \mu^2}{2 \mathcal{H}^2}, \text{p03c}\right]$
 解方程

Solve $\left[\frac{2 \mathfrak{i} \text{p04c } \mathcal{H} \mu^3}{k} == \frac{\mathfrak{i} k^3 \text{p04 } \mu^3}{6 \mathcal{H}^3}, \text{p04c}\right]$
 解方程

Solve $\left[\frac{2 \mathfrak{i} \text{p12c } \mathcal{H} \mu^3}{k} == \frac{2 k^2 \text{p12 } \mu^2}{\mathcal{H}^2}, \text{p12c}\right]$
 解方程

Solve $\left[\frac{3 \mathfrak{i} \text{p13c } \mathcal{H} \mu^3}{k} == -\frac{\mathfrak{i} k^3 \text{p13 } \mu^3}{\mathcal{H}^3}, \text{p13c}\right]$
 解方程

Solve $\left[\frac{\mathfrak{i} \text{p22c } \mathcal{H} \mu^3}{2 k} == \frac{\mathfrak{i} k^3 \text{p22 } \mu^3}{2 \mathcal{H}^3}, \text{p22c}\right]$
 解方程

Clear["Global`*"]
 清除

Out[*]=

$$\left\{\left\{\text{p01c} \rightarrow -\frac{\mathfrak{i} k \text{p01}}{\mathcal{H} \mu}\right\}\right\}$$

Out[*]=

$$\left\{\left\{\text{p02c} \rightarrow -\frac{k^2 \text{p02}}{\mathcal{H}^2}\right\}\right\}$$

Out[*]=

$$\left\{\left\{\text{p11c} \rightarrow \frac{k^2 \text{p11}}{\mathcal{H}^2}\right\}\right\}$$

Out[*]=

$$\left\{\left\{\text{p03c} \rightarrow \frac{\mathfrak{i} k^3 \text{p03}}{3 \mathcal{H}^3 \mu}\right\}\right\}$$

Out[*]=

$$\left\{\left\{\text{p04c} \rightarrow \frac{k^4 \text{p04}}{12 \mathcal{H}^4}\right\}\right\}$$

Out[*]=

$$\left\{\left\{\text{p12c} \rightarrow -\frac{\mathfrak{i} k^3 \text{p12}}{\mathcal{H}^3 \mu}\right\}\right\}$$

Out[*]=

$$\left\{\left\{\text{p13c} \rightarrow -\frac{k^4 \text{p13}}{3 \mathcal{H}^4}\right\}\right\}$$

Out[*] =

$$\left\{ \left\{ p_{22c} \rightarrow \frac{k^4 p_{22}}{\mathcal{H}^4} \right\} \right\}$$

The above equations are the same as the Eqs of Sec1.1.

1.4: Over all the relation between Pmnc and Pmn are:

We finally have the following relations between Pmnc of H19 and Pmn of Vlah12 :

$$\begin{aligned} \text{In[*]} := & \quad p_{00c} = p_{00}; \quad p_{01c} = -\frac{i k p_{01}}{\mathcal{H} \mu}; \quad p_{02c} = -\frac{k^2 p_{02}}{\mathcal{H}^2}; \quad p_{03c} = \frac{i k^3 p_{03}}{3 \mathcal{H}^3 \mu}; \quad p_{04c} = \frac{k^4 p_{04}}{12 \mathcal{H}^4}; \\ & \quad p_{11c} = \frac{k^2 p_{11}}{\mathcal{H}^2}; \quad p_{12c} = -\frac{i k^3 p_{12}}{\mathcal{H}^3 \mu}; \quad p_{13c} = -\frac{k^4 p_{13}}{3 \mathcal{H}^4}; \quad p_{22c} = \frac{k^4 p_{22}}{\mathcal{H}^4}; \\ & \quad \text{Clear["Global`*"]} \\ & \quad \text{清除} \end{aligned}$$

Sec.2: Validating the loop terms Pmn

2.1: P00

$$\begin{aligned} \text{In[*]} := & \quad P_{11dd} = PL; (* \text{ linear power spectrum in V12*}) \\ & \quad P_{22dd} = 2 I_{00}; (* \text{ Eq3.2 of V12*}) \\ & \quad P_{13dd} = 3 k^2 PL J_{00}; (* \text{ Eq3.2 of V12*}) \\ & \quad P_{00} = Q^2 P_{11dd} + Q^4 (P_{22dd} + 2 P_{13dd}); (* \text{ Eq3.1 of V12*}) \\ & \quad Phh_{00} = b_1^2 P_{00} + Q^4 \times \left(2 b_1 (b_2 K_{00} + b_s K_{s00}) + \right. \\ & \quad \left. \frac{1}{2} (b_2^2 K_{01} + b_s^2 K_{s01}) + b_2 b_s K_{s02} + 2 b_3 n_l \sigma_3^2 PL \right); (* \text{ Eq2.28 of V13*}) \end{aligned}$$

Important Notice : In the above equation, do not fogort the prefactor 'Q⁴' for the term of $2 b_1 (b_2 K_{00} + b_s K_{s00}) + \frac{1}{2} (b_2^2 K_{01} + b_s^2 K_{s01}) + b_2 b_s K_{s02} + 2 b_3 n_l \sigma_3^2 PL$

$$\begin{aligned} \text{In[*]} := & \quad Phh_{00} = \text{FullSimplify}[\text{ExpandAll}[Phh_{00}]]; \\ & \quad \text{完全简化} \quad \text{展开全部} \\ & \quad P_{00c} = \text{FullSimplify}[Phh_{00}] \\ & \quad \text{完全简化} \end{aligned}$$

Out[*] =

$$\begin{aligned} & 2 b_1 (b_2 K_{00} + b_s K_{s00}) Q^4 + b_1^2 Q^2 (PL + 2 (I_{00} + 3 J_{00} k^2 PL) Q^2) + \\ & \quad \frac{1}{2} Q^4 (b_2^2 K_{01} + b_s^2 K_{s01} + 2 b_2 b_s K_{s02} + 4 b_3 n_l PL \sigma_3^2) \end{aligned}$$

The corrected P00 of H19 should be:

In[*]:= P00cor =

$$b1^2 Q^2 (PL + 2 Q^2 (I00 + 3 k^2 PL J00)) + 2 b1 Q^4 \left(b2 K00 + bs Ks00 + \frac{b3nl PL \sigma^2}{b1} \right) + Q^4 \left(\frac{1}{2} b2^2 K01 + \frac{1}{2} bs^2 Ks01 + b2 bs Ks02 \right);$$

ExpandAll[P00cor] - ExpandAll[P00c]

[展开全部](#)

[展开全部](#)

Clear["Global`*"]

[清除](#)

Out[*]=

0

2.2: P01

In[*]:= P11δθ = -f ℋ PL; (*Eq.3.7 of V12 . *)

P22δθ = -2 f ℋ I01; (*Eq.3.7 of V12*)

P13δθ = -3 f ℋ k^2 PL J01; (*Eq.3.7 of V12*)

Pδθ = Q^2 P11δθ + Q^4 (P22δθ + 2 P13δθ); (*Eq.3.6 of V12*)

AA211 = -2 f ℋ $\frac{\mu}{k}$ I10; (*Eq.3.9 of V12*)

AA121 = -2 f ℋ μ k PL $\left(3 J10 + \frac{1}{2} (\sigma^2 + \sigma^2) \right)$; (*Eq.3.9 of V12*)

AA112 = f ℋ μ k PL $(\sigma^2 + \sigma^2)$; (*Eq.3.9 of V12*)

A01 = Q^4 (AA211 + AA112 + AA121); (*Eq.3.8 of V12*)

α = - μ $\dot{\imath}$ / k ;

P01 = - $\dot{\imath}$ $\frac{\mu}{k}$ Pδθ - $\dot{\imath}$ A01; (*Eq.3.4 of V12*)

Phh01 = b1^2 P01 + b1 (1 - b1) α Pδθ +

f ℋ Q^4 × (α (b2 K10 + bs Ks10) + α b1 (b2 K11 + bs Ks11) + α b3nl σ^2 PL);

(*Eq.2.29 of V13 , Again, in this equation,

do not fogort the pre-factor 'fℋQ^4' for the term of 'α(b2K10+bsKs10)+αb1(b2K11+bsKs11)+αb3nlσ^2PL'. *)

P01c = FullSimplify[ExpandAll[$-\frac{\dot{\imath} k Phh01}{\mathcal{H} \mu}$]]

[完全简化](#)

[展开全部](#)

Out[*]=

$$f Q^2 (2 b1^2 (I10 + 3 J10 k^2 PL) Q^2 + b1 (PL + (2 I01 - b2 K11 - bs Ks11 + 6 J01 k^2 PL) Q^2) - Q^2 (b2 K10 + bs Ks10 + b3nl PL \sigma^2))$$

The P01 of H19 is correct:


```

In[*]:= P01cor = f b1 Q^2
          (PL + 2 Q^2 (I01 + b1 I10 + 3 k^2 PL (J01 + b1 J10)) - b2 Q^2 K11 - bs Q^2 Ks11) -
          f Q^4 (b2 K10 + bs Ks10 + b3nl σ3^2 PL);
ExpandAll[P01cor] - ExpandAll[P01c]
|展开全部 |展开全部
Clear["Global`*"]
|清除

Out[*]=
0

```

2.3: P02

```

In[*]:= A211 =  $\left(\frac{f \mathcal{H}}{k}\right)^2 (I02 + \mu^2 I20); (*Eq.3.34 of V12*)$ 
A121 = (f \mathcal{H})^2 PL (J02 + \mu^2 J20); (*Eq.3.34 of V12*)
A112 = A121; (*Eq.3.34 of V12*)
A02 = Q^4 (A211 + A121 + A112); (*Eq.3.33 of V12*)
B1111 = - (f \mathcal{H})^2 PL σv^2; (*Eq.3.34 of V12*)
B02 = Q^4 B1111; (*Eq.3.33 of V12, do not fogort Q^4*)
P02 = - (A02 + B02) (*Eq.3.31 of V12. I moved Q^4 to A02 and B02. *);
|虚数单位

P02 = P02 - f^2 PL Q^4 \mathcal{H}^2 σv^2; (*see the texts below Eq. 3.38 of V12,
omitting the velocity dispersion part from P02*)
Phh02 = b1 P02 + phh00 σv^2 - (k^-2 \mathcal{H}^2 f^2 Q^4) × (b2 K20 + bs Ks20);
(*Eq.2.31 of V13. phh00=p00c. In this equation,
|输入行

do not fogort the pre-factor 'k^-2 \mathcal{H}^2 f^2 Q^4' for '(b2K20+bsKs20)'. *)
phh00 = p00c;
Phh02 = FullSimplify[ExpandAll[Phh02]];
|完全简化 |展开全部

P02c = -  $\frac{k^2 Phh02}{\mathcal{H}^2}$ ;
P02c = FullSimplify[ExpandAll[P02c]]
|完全简化 |展开全部

Out[*]=
f^2 Q^4 (b2 K20 + bs Ks20 + b1 (I02 + I20 μ^2 + 2 k^2 PL (J02 + J20 μ^2))) -  $\frac{k^2 p00c σv^2}{\mathcal{H}^2}$ 

```

The corrected P02 of H19 should be

```

In[*]:= P02cor = f^2 b1 Q^4 (I02 + μ^2 I20 + 2 k^2 PL (J02 + μ^2 J20)) -
           $\frac{1}{\mathcal{H}^2 f^2} f^2 k^2 σv^2 p00c - (-1) × f^2 Q^4 (b2 K20 + bs Ks20); (* phh00=p00c *)$ 
ExpandAll[P02cor] - ExpandAll[P02c]
|展开全部 |展开全部

Out[*]=
0

```

Important Notice : If we define: $σ_v^{vlah} \equiv \mathcal{H} f σ_v^{Howlett}$ for the equations in Vlah's paper, then we have:

```

In[*]:= σv = ℋ f σvc; (*σv is Vlah, σvc is Howlett.*)
P02cnew = FullSimplify[ExpandAll[P02c]];
          完全简化      展开全部
P02cor = f^2 b1 Q^4 (I02 + μ^2 I20 + 2 k^2 PL (J02 + μ^2 J20))
        - f^2 k^2 σvc^2 p00c - (-1) × f^2 Q^4 (b2 K20 + bs Ks20); (*H19*)
ExpandAll[P02cor] - ExpandAll[P02cnew]
          展开全部      展开全部
Clear["Global`*"]
          清除
Out[*]=
0

```

So, the above is the corrected P02 of H19 is correct.

2.4: P03

```

In[*]:= phh01 =  $\frac{p01c}{-\frac{\hbar k}{\hbar \mu}}$ ;
Phh03 = 3 phh01 σv^2; (*Eq.2.40 of V13 *)
(* ON the other hand, we also have: *)
Phh03 =  $\frac{p03c}{\frac{\hbar k^3}{3 \hbar^3 \mu}}$ ;
(*So, we can solve the equation: *)
Solve[ $\frac{p03c}{\frac{\hbar k^3}{3 \hbar^3 \mu}} = 3 \frac{p01c}{-\frac{\hbar k}{\hbar \mu}} \sigma v^2$ , p03c]
          解方程
Out[*]=
{ {p03c → -  $\frac{k^2 p01c \sigma v^2}{\hbar^2}$  } }

```

Again, if we define: $\sigma_v^{\text{Vlah}} \equiv \mathcal{H} f \sigma_v^{\text{Howlett}}$ for the equations in Vlah's paper, then we have:

```

In[*]:= σv = ℋ f σvc;
P03c = ExpandAll[ -  $\frac{k^2 p01c \sigma v^2}{\hbar^2}$  ]
          展开全部
Out[*]=
- f^2 k^2 p01c σvc^2

```

The P03 of H19 is correct:

```

In[*]:= P03cor = - f^2 k^2 σvc^2 p01c;
ExpandAll[P03cor] - ExpandAll[P03c]
          展开全部      展开全部
Clear["Global`*"]
          清除
Out[*]=
0

```

2.5: P04

```

In[*]:= A211 =  $\left(\frac{f \mathcal{H}}{k}\right)^2 (I02 + \mu^2 I20);$  (*Eq.3.34 of V12*)
A121 =  $(f \mathcal{H})^2 \text{PL} (J02 + \mu^2 J20);$  (*Eq.3.34 of V12*)
A112 = A121; (*Eq.3.34 of V12*)
A02 =  $Q^4 (A211 + A121 + A112);$  (*Eq.3.33 of V12*)
 $\sigma v = \mathcal{H} f \sigma v c;$  (* Again, we re-define  $\sigma_v$  use  $\mathcal{H} f \sigma_v$  *)
 $\sigma 4 = \mathcal{H} f \sigma 4 c;$ 
B1111 =  $-(f \mathcal{H})^2 \text{PL} \sigma v^2;$  (*Eq.3.34 of V12*)
B02 =  $Q^4 B1111;$  (*Eq.3.33 of V12*)
P02 =  $-(A02 + B02)$  (*Eq.3.31 of V12. I moved  $Q^4$  to A02 and B02. *);
|虚数单位
 $\overline{P02} = P02 - f^2 \text{PL} Q^4 \mathcal{H}^2 \sigma v^2;$  (*see the texts below Eq. 3.38 of V12,
omitting the velocity dispersion part from P02*)
Phh04 =  $6 b1 \overline{P02} \sigma v^2 + b1 b1 \text{phh00} (3 \sigma v^4 + \sigma 4^4);$  (* Eq.2.47 of V13*)
phh00 = p00c;
P04c = FullSimplify[ExpandAll[ $\frac{k^4 \text{Phh04}}{12 \mathcal{H}^4}$ ]]
|完全简化 |展开全部
Out[*]=
 $\frac{1}{12} b1 f^4 k^2 (-6 Q^4 (I02 + I20 \mu^2 + 2 k^2 \text{PL} (J02 + J20 \mu^2)) \sigma v c^2 + b1 k^2 p00c (\sigma 4 c^4 + 3 \sigma v c^4))$ 

```

The corrected P04 of H19 should be:

```

In[*]:= P04cor =  $\left(-\frac{1}{2}\right) f^4 b1 k^2 \sigma v c^2 Q^4 (I02 + \mu^2 I20 + 2 k^2 \text{PL} (J02 + \mu^2 J20))$ 
+  $\frac{1}{4} f^4 b1^2 k^4 p00c \left(\sigma v c^4 + \sigma 4 c^4 \times \frac{1}{3}\right);$  (* phh00=p00c *)
ExpandAll[P04cor] - ExpandAll[P04c] (* phh00=p00c *)
|展开全部 |展开全部
Clear["Global`*"]
|清除
Out[*]=
0

```

In the $\sigma 4 c^4 \times \frac{1}{3}$, $\frac{1}{3}$ can be absorbed into $\sigma 4 c$, so P04 of H19 is correct.

2.6: P11

```

In[*]:= P11tt =  $\mathcal{H}^2 f^2 \text{PL}$ ; (*Eq.3.21 of V12*)
P22tt = 2  $\mathcal{H}^2 f^2 \text{I11}$ ; (*Eq.3.21 of V12*)
P13tt = 3  $\mathcal{H}^2 f^2 k^2 \text{PL J11}$ ; (*Eq.3.21 of V12*)
Ptt =  $Q^2 \text{P11tt} + Q^4 (\text{P22tt} + 2 \text{P13tt})$ ; (*Eq.3.20 of V12*)
Bs211 = 2  $\mathcal{H}^2 f^2 \mu k \text{PL} (3 \text{J10} + 1/2 (\sigma v^2 + \sigma 3^2))$ ; (*Eq.3.23 of V12*)
Bs121 = -  $\mathcal{H}^2 f^2 \mu k \text{PL} (\sigma v^2 + \sigma 3^2)$ ; (*Eq.3.23 of V12*)
Bs112 = 2  $\mathcal{H}^2 f^2 \frac{\mu}{k} \text{I22}$ ; (*Eq.3.23 of V12*)
B11 =  $Q^4 (\text{Bs211} + \text{Bs112} + \text{Bs121})$ ; (*Eq.3.22 of V12*)
C1111 =  $\mathcal{H}^2 f^2 k^{-2} (\text{I31} + \text{I13} \mu^2)$ ; (*Eq.3.24 of V12*)
C11 =  $Q^4 \text{C1111}$ ; (* do not fogort  $Q^4$  *)
P11 =  $\frac{\mu^2}{k^2} \text{Ptt} + \frac{2 \mu}{k} \text{B11} + \text{C11}$ ; (*Eq.3.18 of V12*)
Phh11 =  $\text{P11} + ((b1 - 1) + (b1 - 1)) \frac{\mu}{k} \text{B11} + (b1 b1 - 1) \text{C11}$ ; (*Eq.2.35 of V13*)
P11c = FullSimplify[ExpandAll[ $\frac{k^2 \text{Phh11}}{\mathcal{H}^2}$ ]]

```

```

Out[*]=  $f^2 Q^2 (\text{PL} \mu^2 + Q^2 (b1^2 \text{I31} + (2 \text{I11} + b1 (b1 \text{I13} + 4 \text{I22}) + 6 (2 b1 \text{J10} + \text{J11}) k^2 \text{PL}) \mu^2)$ 

```

The P11 of H19 is correct:

```

In[*]:= P11cor =  $f^2 Q^2 (\mu^2 (\text{PL} + Q^2 (2 \text{I11} + 4 b1 \text{I22} + b1^2 \text{I13} + 6 k^2 \text{PL} (\text{J11} + 2 b1 \text{J10}))) +$ 
 $b1^2 Q^2 \text{I31})$ ;
ExpandAll[P11cor] - Expand[P11c]

```

清除

```

Out[*]=

```

0

2.7: P12

```

In[ ]:= (*we need to firstly calculate P01 of V12*)
P11δθ = -f ℋ PL; (*Eq.3.7 of V12, the Hubble parameter aH is ignored. *)
P22δθ = -2 f ℋ I01; (*Eq.3.7 of V12*)
P13δθ = -3 f ℋ k² PL J01; (*Eq.3.7 of V12*)
Pδθ = Q² P11δθ + Q⁴ (P22δθ + 2 P13δθ); (*Eq.3.6 of V12*)
AA211 = -2 f ℋ  $\frac{\mu}{k}$  I10; (*Eq.3.9 of V12*)
AA121 = -2 f ℋ μ k PL  $\left( 3 J10 + \frac{1}{2} (\sigma_{vc}^2 + \sigma_3^2) \right)$ ;
(*Eq.3.9 of V12, These should be σvc,
because the growth rate is taken outside sigma_v in V12. *)
AA112 = f ℋ μ k PL (σvc² + σ3²); (*Eq.3.9 of V12. These should be σvc,
because the growth rate is taken outside sigma_v in V12. *)
A01 = Q⁴ (AA211 + AA112 + AA121); (*Eq.3.8 of V12*)
α = - μ i / k;
P01 = FullSimplify[ExpandAll[- i  $\frac{\mu}{k}$  Pδθ - i A01]]; (*Eq.3.4 of V12*)
完全简化 展开全部

(*Then, we calculate P02*)
k11 = k μ;
As211 = -  $\frac{(f \mathcal{H})^3}{k^2}$  (I12 + μ² I21); (*Eq.3.44 of V12*)
As121 = - (f ℋ)³ PL (J02 + μ² J20); (*Eq.3.44 of V12*)
As112 = As121; (*Eq.3.44 of V12*)
A12 = Q⁴ (As211 + As121 + As112); (*Eq.3.43 of V12*)
B12 = Q⁴ (f ℋ)³ μ k⁻³ (I03 + μ² I30); (*Eq.3.44 of V12, do not forgot Q⁴ *)
C12 = Q⁴ (f ℋ)³ PL σvc²; (*Eq.3.44 of V12. Should be σvc as well*)
P12 = -  $\frac{i}{k^2}$  (k11 A12 + k² B12 + k11 C12);
(*Eq.3.41 of V12*)
phh01 =  $\frac{p01c}{-\frac{i k}{\mathcal{H} \mu}}$ ;
Phh12 = P12 - i (b1 - 1) B12 - (phh01 - Q² P01) σv²;
(*Eq.2.38 of V13. For consistency,
For循环

this should have a Q². Because in C12 we are affectively fitting sigma_v at z=
因为

0, then rescaling to z=z. But here we have rescaled the P01 contains P_L(z=0),
while the sigma_v is still being evaluated at z=z. *)
σv = f ℋ σvc;
P12c = FullSimplify[ExpandAll[-  $\frac{i k^3 Phh12}{\mathcal{H}^3 \mu}$ ]];
完全简化 展开全部

Out[ ]:=
f² (f Q⁴ (I12 + I21 μ² - b1 (I03 + I30 μ²) + 2 k² PL (J02 + J20 μ²)) +
k² (-p01c + 2 f (I01 + I10 + 3 (J01 + J10) k² PL) Q⁶) σvc²)

```

The corrected P12 of H19 should be:

```
In[*]:= P12cor = f^3 Q^4 (I12 + I21 μ^2 - b1 (I03 + I30 μ^2) + 2 k^2 PL (J02 + J20 μ^2))
          - f^2 k^2 σvc^2 p01c + 2 f^3 k^2 (I01 + I10 + 3 (J01 + J10) k^2 PL) Q^4 (Q σvc)^2;
ExpandAll[P12cor] - Expand[P12c]
|展开全部 |展开
Clear["Global`*"]
|清除

Out[*]=
0
```

2.8: P13

```
In[*]:= phh11 =  $\frac{p11c}{\mathcal{H}^2}$ ;

Phh13 = 3 phh11 σv^2; (* Eq.2.43 of V13 *)
(* ON the other hand, we also have: *)

Phh13 =  $\frac{p13c}{-\frac{k^4}{3 \mathcal{H}^4}}$ ;

(*So, we can solve the equation: *)

Solve[ $\frac{p13c}{-\frac{k^4}{3 \mathcal{H}^4}} == 3 \frac{p11c}{\mathcal{H}^2} \sigma v^2$ , p13c]
|解方程

Out[*]=
{{p13c → - $\frac{k^2 p11c \sigma v^2}{\mathcal{H}^2}$ }}
```

```
In[*]:= σv = f H σvc;

P13c = ExpandAll[ $-\frac{k^2 p11c \sigma v^2}{\mathcal{H}^2}$ ]
|展开全部

Out[*]=
-f^2 k^2 p11c σvc^2
```

The P13 of H19 is correct:

```
In[*]:= P13cor = - f^2 k^2 σvc^2 p11c;
ExpandAll[P13cor] - ExpandAll[P13c]
|展开全部 |展开全部
Clear["Global`*"]
|清除

Out[*]=
0
```

2.9: P22

```

In[*]:= A211 =  $\left(\frac{f \mathcal{H}}{k}\right)^2 (I02 + \mu^2 I20)$ ; (*Eq.3.34 of V12*)
A121 =  $(f \mathcal{H})^2 PL (J02 + \mu^2 J20)$ ; (*Eq.3.34 of V12*)
A112 = A121; (*Eq.3.34 of V12*)
A02 =  $Q^4 (A211 + A121 + A112)$ ; (*Eq.3.33 of V12*)
 $\sigma v = f \mathcal{H} \sigma v c$ ;
B1111 =  $-(f \mathcal{H})^2 PL \sigma v^2$ ; (*Eq.3.34 of V12*)
B02 =  $Q^4 B1111$ ; (*Eq.3.33 of V12, do not forget  $Q^4$ *)
P02 =  $-(A02 + B02)$  (*Eq.3.31 of V12. I moved  $Q^4$  to A02 and B02. *);
 $\overline{P02}$  = Simplify[ExpandAll[P02 -  $f^2 PL Q^4 \mathcal{H}^2 \sigma v^2$ ]];
(*Finally, calculate P22*)
 $\overline{P22} = \frac{\frac{1}{16} f^4 Q^4 \mu^4 (I23 + 2 \mu^2 I32 + \mu^4 I33)}{\frac{1}{4} \left(\frac{k \mu}{\mathcal{H}}\right)^4}$ ; (*Eq.3.48 of V12*)
Phh00Pb22 = 2 phh00  $\sigma v^4$ ; (*texts below Eq.3.50 of V12. *)
phh00 = p00c;
phh02 =  $\frac{p02c}{-\frac{k^2}{\mathcal{H}^2}}$ ;
Phh22 =  $\overline{P22} + b1 \sigma v^2 \overline{P02} + phh02 \sigma v^2 + Phh00Pb22$ ; (*Eq.2.45 of V13 *)
(* The following p02c is taken from Sec2.3. Please check
it if the calculation of P02 of Sec2.3 has been updated. *)
p02c =  $f^2 b1 Q^4 (I02 + \mu^2 I20 + 2 k^2 PL (J02 + \mu^2 J20))$ 
-  $f^2 k^2 \sigma v c^2 p00c - (-1) \times f^2 Q^4 (b2 K20 + bs Ks20)$ ;
P22c = FullSimplify[ExpandAll[ $\frac{k^4 Phh22}{\mathcal{H}^4}$ ]];

```

Out[*]=

$$\frac{1}{4} f^4 \left(Q^4 (I23 + 2 I32 \mu^2 + I33 \mu^4) - 4 k^2 Q^4 (b2 K20 + bs Ks20 + 2 b1 (I02 + I20 \mu^2 + 2 k^2 PL (J02 + J20 \mu^2))) \right) \sigma v c^2 + 12 k^4 p00c \sigma v c^4$$

The P22 of H19 is correct:

```

In[*]:= P22cor =  $\frac{1}{4} f^4 Q^4 (I23 + 2 \mu^2 I32 + \mu^4 I33) + f^4 k^4 \sigma v c^4 p00c -$ 
 $f^2 k^2 \sigma v c^2 (2 p02c - f^2 Q^4 (b2 K20 + bs Ks20))$ ; (* phh00=p00c *)
ExpandAll[P22cor] - ExpandAll[P22c]
Clear["Global`*"]

```

Out[*]=

0

Sec.3: Summary

The corrected Pmn are:

$$\begin{aligned} P00cor = & b1^2 Q^2 (PL + 2 Q^2 (I00 + 3 k^2 PL J00)) + \\ & 2 b1 Q^4 (b2 K00 + bs Ks00 + \frac{b3n1 PL \sigma^2}{b1}) + \\ & Q^4 (\frac{1}{2} b2^2 K01 + \frac{1}{2} bs^2 Ks01 + b2 bs Ks02) \end{aligned}$$

$$\begin{aligned} P02cor = & f^2 b1 Q^4 (I02 + \mu^2 I20 + 2 k^2 PL (J02 + \mu^2 J20)) - \\ & f^2 k^2 \sigma^2 p00c - (-1) \times f^2 Q^4 (b2 K20 + bs Ks20) \end{aligned}$$

$$\begin{aligned} P12cor = & f^3 Q^4 (I12 + I21 \mu^2 - b1 (I03 + I30 \mu^2) + 2 k^2 PL (J02 + J20 \mu^2)) \\ & - f^2 k^2 \sigma^2 p01c + 2 f^3 k^2 (I01 + I10 + 3 (J01 + J10) k^2 PL) Q^4 (Q \sigma^2)^2 ; \end{aligned}$$

Sec.4: Cross power spectrum for multi-surveys

4.1: P00

```
In[1]:=  $\delta h \delta h b = b1 b1c \delta \delta + \frac{b1 b2c + b2 b1c}{2} \delta \delta 2 +$ 
 $\frac{b1 bsc + bs b1c}{2} \delta s2 + \frac{1}{3} \frac{b1 b3c + b3 b1c}{2} \delta \delta 3 + \frac{1}{4} b2 b2c \delta 2 \delta 2 +$ 
 $\frac{1}{4} bs bsc s2s2 + \frac{1}{2} \frac{b2 bsc + bs b2c}{2} \delta 2s2 ; (*Eq.2.13 of V13*)$ 
 $\delta \delta = P00 ; (*Eq.2.15 of V13*)$ 
 $\delta \delta 2 = 2 K00 + 2 \frac{34}{21} P00 \sigma^2 ; (*Eq.2.15 and 2.17 of V13*)$ 
 $\delta \delta 3 = 3 P00 \sigma^2 ; (*Eq.2.15 of V13*)$ 
 $\delta s2 = 2 Ks00 ; (*Eq.2.15 and 2.17 of V13*)$ 
 $\delta 2 \delta 2 = 2 K01 ; (*Eq.2.20 of V13*)$ 
 $s2s2 = 2 Ks01 ; (*Eq.2.20 of V13*)$ 
 $\delta 2s2 = 2 Ks02 ; (*Eq.2.20 of V13*)$ 
 $Phh00 = \delta h \delta h b ; (*Eq.2.15 and 2.17 of V13*)$ 
Collect[Phh00, P00]
```

[合并同类项](#)

Out[10]=

$$\begin{aligned} & (b1c b2 + b1 b2c) K00 + \frac{b2 b2c K01}{2} + (b1c bs + b1 bsc) Ks00 + \frac{bs bsc Ks01}{2} + \\ & \frac{1}{2} (b2c bs + b2 bsc) Ks02 + P00 \left(b1 b1c + \frac{34}{21} (b1c b2 + b1 b2c) \sigma^2 + \frac{1}{2} (b1c b3 + b1 b3c) \sigma^2 \right) \end{aligned}$$

Following V13 Eq2.18, We re-difine the linear bais parameter b1 to make:

$$b1 b1c \rightarrow b1 b1c + \frac{34}{21} (b1c b2 + b1 b2c) \sigma^2 + \frac{1}{2} (b1c b3 + b1 b3c) \sigma^2$$

$$\text{Phh00} = (b1c \, b2 + b1 \, b2c) \, K00 + \frac{b2 \, b2c \, K01}{2} + (b1c \, bs + b1 \, bsc) \, Ks00 + \frac{bs \, bsc \, Ks01}{2} + \frac{1}{2} (b2c \, bs + b2 \, bsc) \, Ks02 + P00 \times b1 \, b1c$$

Then, we have:

$$\text{In[11]:= Phh00} = b1 \, b1c \, P00 + Q^4 \times \left((b1c \, b2 + b1 \, b2c) \, K00 + \frac{b2 \, b2c \, K01}{2} + (b1c \, bs + b1 \, bsc) \, Ks00 + \frac{bs \, bsc \, Ks01}{2} + \frac{1}{2} (b2c \, bs + b2 \, bsc) \, Ks02 + 2 \, b3nl \, \sigma^3^2 \, PL \right);$$

P11dd = PL;

P22dd = 2 I00;

P13dd = 3 k² PL J00;

P00 = Q² P11dd + Q⁴ (P22dd + 2 P13dd);

Phh00 = FullSimplify[ExpandAll[Phh00]];
 完全简化 展开全部

P00c = FullSimplify[Phh00]
 完全简化

Out[17]=

$$b1 \, b1c \, PL \, Q^2 + b1 \, (b2c \, K00 + bsc \, Ks00 + 2 \, b1c \, (I00 + 3 \, J00 \, k^2 \, PL)) \, Q^4 + \frac{1}{2} \, Q^4 \, (b2 \, b2c \, K01 + 2 \, b1c \, (b2 \, K00 + bs \, Ks00) + bs \, bsc \, Ks01 + b2c \, bs \, Ks02 + b2 \, bsc \, Ks02 + 4 \, b3nl \, PL \, \sigma^3^2)$$

If set:

b1c=b1; b2c=b2; bsc=bs;

we should get auto-PS:

In[18]:= b1c = b1;

b2c = b2;

bsc = bs;

ExpandAll[P00c - (2 b1 (b2 K00 + bs Ks00) Q⁴ + b1² Q² (PL + 2 (I00 + 3 J00 k² PL) Q²) +
 展开全部

$$\frac{1}{2} \, Q^4 \, (b2^2 \, K01 + bs^2 \, Ks01 + 2 \, b2 \, bs \, Ks02 + 4 \, b3nl \, PL \, \sigma^3^2))]$$

Clear["Global`*"]

清除

Out[21]=

0

4.2: P01

```

In[23]:= P11δθ = -f ℋ PL;
P22δθ = -2 f ℋ I01;
P13δθ = -3 f ℋ k² PL J01;
Pδθ = Q² P11δθ + Q⁴ (P22δθ + 2 P13δθ);
AA211 = -2 f ℋ  $\frac{\mu}{k}$  I10;
AA121 = -2 f ℋ μ k PL  $\left( 3 J10 + \frac{1}{2} (\sigma v^2 + \sigma^3) \right)$ ;
AA112 = f ℋ μ k PL  $(\sigma v^2 + \sigma^3)$ ;
A01 = Q⁴ (AA211 + AA112 + AA121);
α = - μ  $\dot{\imath}$  / k;
P01 = -  $\dot{\imath}$   $\frac{\mu}{k}$  Pδθ -  $\dot{\imath}$  A01;
Phh01 = b1 b1c P01 + b1 (1 - b1c) α Pδθ +
  f ℋ Q⁴ × ( α (b2 K10 + bs Ks10) + α b1c (b2 K11 + bs Ks11) + α b3nl σ³ PL );
(*Eq.2.26 of V13 *)
P01c = FullSimplify[ExpandAll[ $-\frac{\dot{\imath} k \text{Phh01}}{\mathcal{H} \mu}$ ]]
完全简化 展开全部

Out[34]=
f Q² (b1 (PL + 2 (I01 + b1c I10 + 3 (J01 + b1c J10) k² PL) Q²) -
  Q² (b2 (K10 + b1c K11) + bs Ks10 + b1c bs Ks11 + b3nl PL σ³))

If set: b1c=b1; b2c=b2; bsc=bs; we should get auto-PS:

In[35]:= b1c = b1;
b2c = b2;
bsc = bs;
ExpandAll[P01c - ( f Q²
展开全部
  (2 b1² (I10 + 3 J10 k² PL) Q² + b1 (PL + (2 I01 - b2 K11 - bs Ks11 + 6 J01 k² PL) Q²) -
    Q² (b2 K10 + bs Ks10 + b3nl PL σ³))
)]
Clear["Global`*"]
清除

Out[38]=
0

```

4.3: P11

```
In[40]:= P11tt =  $\mathcal{H}^2 f^2$  PL;
P22tt = 2  $\mathcal{H}^2 f^2$  I11;
P13tt = 3  $\mathcal{H}^2 f^2 k^2$  PL J11;
Ptt =  $Q^2$  P11tt +  $Q^4$  ( P22tt + 2 P13tt ) ;
Bs211 = 2  $\mathcal{H}^2 f^2 \mu k$  PL ( 3 J10 + 1 / 2 (  $\sigma v^2$  +  $\sigma 3^2$  ) ) ;
Bs121 = -  $\mathcal{H}^2 f^2 \mu k$  PL (  $\sigma v^2$  +  $\sigma 3^2$  ) ;
Bs112 = 2  $\mathcal{H}^2 f^2 \frac{\mu}{k}$  I22;
B11 =  $Q^4$  ( Bs211 + Bs112 + Bs121 ) ;
C1111 =  $\mathcal{H}^2 f^2 k^{-2}$  ( I31 + I13  $\mu^2$  ) ;
C11 =  $Q^4$  C1111;
P11 =  $\frac{\mu^2}{k^2}$  Ptt +  $\frac{2 \mu}{k}$  B11 + C11;
Phh11 = P11 + ( (b1 - 1) + (b1c - 1) )  $\frac{\mu}{k}$  B11 + (b1 b1c - 1) C11 ;
(*Eq.2.35 of V13*)
P11c = FullSimplify[ExpandAll[ $\frac{k^2 \text{Phh11}}{\mathcal{H}^2}$ ]]
|完全简化 |展开全部
```

```
Out[52]=  $f^2 Q^2 \left( \text{PL} \mu^2 + Q^2 \left( b1 b1c I31 + \right. \right.$ 
 $\left. \left( 2 I11 + b1 b1c I13 + 2 b1 I22 + 2 b1c I22 + 6 \left( (b1 + b1c) J10 + J11 \right) k^2 \text{PL} \right) \mu^2 \right)$ 
```

If set : b1c=b1; b2c=b2; bsc=bs; we should get auto-PS:

```
In[53]:= b1c = b1;
b2c = b2;
bsc = bs;
ExpandAll[P11c - (  $f^2 Q^2$ 
|展开全部
 $\left( \text{PL} \mu^2 + Q^2 \left( b1^2 I31 + \left( 2 I11 + b1 (b1 I13 + 4 I22) + 6 (2 b1 J10 + J11) k^2 \text{PL} \right) \mu^2 \right) \right) ]$ 
Clear["Global`*"]
|清除
```

```
Out[56]= 0
```

4.4: P02, P03 and P04, P12 P13 and P22:

P02:

We do not need to modify P02 for cross-PS. The expression of P02 of cross-PS is the same as auto-PS. In V13, Eq. 2.13, the $\overline{P02}$ is calculated from dark matter PS (see the first term of Eq.2.30 $\langle \delta | u_{\parallel}^2 \rangle$),

it is purely the contribution from dark matter and should be an auto-like term.

The P^{hh}_{00} of Eq. 2.13 should be auto-like term rather than cross-like term P^{hhbar}_{00} . From the second term of Eq.2.30, we can see that

$$\langle \delta | \delta u_{\parallel}^2 \rangle = \langle \delta | \delta \rangle \langle u_{\parallel}^2 \rangle = \langle \delta | \delta \rangle \sigma_v^2$$

where $\langle \delta | \delta \rangle$ is P^{hh}_{00} which is an auto-like term.

P03: $P03_{cor} = -f^2 k^2 \sigma_{vc}^2 p01c$: use cross-like p01c of Sec 4.2.

P04: In Eq.2.47 of V13, $\overline{P02}$ is auto-like term, in H19 Eq.A8, just replace b_1^2 with $b_1\overline{b_1}$ and use P00 of Sec 4.1 .

P12: use cross-like p01 of Sec 4.2.

P13: $p_{13cor} = -f^2 k^2 \sigma_{vc}^2 p_{11c}$; use cross-like p_{11c} of Sec 4.3.

P22: use cross-like P00 of Sec 4.1 . Do not need to modify p02.

The end of the code . ## ## ## ## ## ## ## ## ## ## ## ## ## ## ## ##
##

Figure 9-7: The end of the code.

Shi-Fan Chen's cross-power

Wolfram Cloud

(unnamed) - Wolfram Cloud

www.wolframcloud.com/env/da696323-333e-4670-a95c-ae6bb4868516

WOLFRAM CLOUD

Plan: Basic **UPGRADE** Documentation Language Intro Quick Links Cloud Files

(unnamed)

File Edit Format Insert Evaluation View Help Share Publish

In[1]:= $\text{ExpandAll}[\text{Solve}[(-((i k \mu)/\mathcal{H})) P\delta m == -((i k \mu)/\mathcal{H}) p_{01} + ((k \mu)/\mathcal{H})^2 p_{11} - ((k \mu)/\mathcal{H})^2 p_{02} + i/2 ((k \mu)/\mathcal{H})^3 p_{03} - 3/2 i ((k \mu)/\mathcal{H})^3 p_{12} + 1/6 ((k \mu)/\mathcal{H})^4 p_{04} - 2/3 ((k \mu)/\mathcal{H})^4 p_{13} + 1/2 ((k \mu)/\mathcal{H})^4 p_{22}, P\delta m]]$

Out[1]= $\left\{\left\{P\delta m \rightarrow p_{01} - \frac{i k p_{02} \mu}{\mathcal{H}} + \frac{i k p_{11} \mu}{\mathcal{H}} - \frac{k^2 p_{03} \mu^2}{2 \mathcal{H}^2} + \frac{3 k^2 p_{12} \mu^2}{2 \mathcal{H}^2} + \frac{i k^3 p_{04} \mu^3}{6 \mathcal{H}^3} - \frac{2 i k^3 p_{13} \mu^3}{3 \mathcal{H}^3} + \frac{i k^3 p_{22} \mu^3}{2 \mathcal{H}^3}\right\}\right\}$

In[2]:= $P\delta mc = i \mathcal{H} / k * \mu * (p_{01c} + p_{02c} + p_{11c} + \mu^2 * (3/2 * p_{03c} + 2 p_{04c} + 3/2 p_{12c} + 2 p_{13c} + 1/2 * p_{22c}));$
 $\text{ExpandAll}[P\delta mc]$

Out[3]= $\frac{i p_{01c} \mathcal{H} \mu}{k} + \frac{i p_{02c} \mathcal{H} \mu}{k} + \frac{i p_{11c} \mathcal{H} \mu}{k} + \frac{3 i p_{03c} \mathcal{H} \mu^3}{2 k} + \frac{2 i p_{04c} \mathcal{H} \mu^3}{k} + \frac{3 i p_{12c} \mathcal{H} \mu^3}{2 k} + \frac{2 i p_{13c} \mathcal{H} \mu^3}{k} + \frac{i p_{22c} \mathcal{H} \mu^3}{2 k}$

In[4]:= $\text{Solve}[3 i p_{12c} \mathcal{H} \mu^3 / (2 k) == (3 k^2 p_{12} \mu^2) / (2 \mathcal{H}^2), p_{12c}]$
 $\text{Solve}[(2 i p_{13c} \mathcal{H} \mu^3) / k == -((2 i k^3 p_{13} \mu^3) / (3 \mathcal{H}^3)), p_{13c}]$

Out[4]= $\left\{\left\{p_{12c} \rightarrow -\frac{i k^3 p_{12}}{\mathcal{H}^3 \mu}\right\}\right\}$

Out[5]= $\left\{\left\{p_{13c} \rightarrow -\frac{k^4 p_{13}}{3 \mathcal{H}^4}\right\}\right\}$