


Statistical Methods for Quantitative MS-Based Proteomics:

1. Identification & False discovery rate

Lieven Clement

Statistics and Genomics Seminar, UC Berkeley, California

Challenges in Label Free MS-based Quantitative Proteomics



Quantification Identification

Challenges in Label Free MS-based Quantitative Proteomics






Table of Outcomes

	Called Bad	Called Correct	
Bad hit	TN	FP	m_0
Correct hit	FN	TP	m_1
Total	NR	R	m

- TN: number of true negatives
- FP: number of false positives
- FN: number of false negatives
- TP: number of true positives
- NR: number of non-rejections, R: number of rejections

Table of Outcomes

	Called Bad	Called Correct	
Bad hit	TN	FP	m_0
Correct hit	FN	TP	m_1
Total	NR	R	m

Random Variables

Table of Outcomes

		Called Bad	Called Correct	
Unobservable	Bad hit	TN	FP	m_0
	Correct hit	FN	TP	m_1
Observable	Total	NR	R	m

Table of Outcomes

		Called Bad	Called Correct	
Unobservable	Bad hit	TN	FP	m_0
	Correct hit	FN	TP	m_1
Observable	Total	NR	R	m


$FDP = \frac{FP}{FP+TP}$. But is unknown! (FDP: false discovery proportion)

Table of Outcomes

		Called Bad	Called Correct	
Unobservable	Bad hit	TN	FP	m_0
	Correct hit	FN	TP	m_1
Observable	Total	NR	R	m


$$FDR = E \left[\frac{FP}{FP+TP} \right]. \text{ (FDR: false discovery rate)}$$

Search engines return score that discriminates good from bad matches




Search engines return score that discriminates good from bad matches

Score threshold t ?




Search engines return score that discriminates good from bad matches



Score threshold t ?

$$f(x) = \pi_0 f_0(x) + (1 - \pi_0) f_1(x)$$

Search engines return score that discriminates good from bad matches




Score threshold t ?

$$f(x) = \pi_0 f_0(x) + (1 - \pi_0) f_1(x)$$

$$\text{FDR}(t) = E \left[\frac{FP}{FP + TP} \right]$$

Search engines return score that discriminates good from bad matches




Score threshold t ?

$$f(x) = \pi_0 f_0(x) + (1 - \pi_0) f_1(x)$$

$$\text{FDR}(t) = E \left[\frac{FP}{FP + TP} \right]$$

$$\text{FDR}(t) = \frac{mP[FP]P[x \geq t | FP]}{mP[x \geq t]}$$

Search engines return score that discriminates good from bad matches



Score threshold t ?


$$f(x) = \pi_0 f_0(x) + (1 - \pi_0) f_1(x)$$

$$\text{FDR}(t) = E \left[\frac{FP}{FP + TP} \right]$$

$$\text{FDR}(t) = \frac{mP[FP]P[x \geq t | FP]}{mP[x \geq t]}$$

$$\text{FDR}(t) = \frac{\pi_0 P_0[x \geq t]}{P[x \geq t]}$$

Search engines return score that discriminates good from bad matches



Score threshold t ?

$$f(x) = \pi_0 f_0(x) + (1 - \pi_0) f_1(x)$$


$$\text{FDR}(t) = E \left[\frac{FP}{FP + TP} \right]$$

$$\text{FDR}(t) = \frac{mP[FP]P[x \geq t | FP]}{mP[x \geq t]}$$

$$\text{FDR}(t) = \frac{\pi_0 P_0[x \geq t]}{P[x \geq t]}$$

$$P[x \geq t] = \int_t^{\infty} f(x)$$


How to estimate FDR?



$$P.[x \geq t] = \int_t^{\infty} f.(x)$$

$$\text{FDR}(t) = E \left[\frac{FP}{FP + TP} \right] = \frac{m\pi_0 P[x \geq t | FP]}{mP[x \geq t]}$$

How to estimate FDR?




$$P.[x \geq t] = \int_t^{\infty} f.(x)$$

$$P.[x \geq t] \approx \frac{\#x \geq t}{m}$$

$$\text{FDR}(t) = E \left[\frac{FP}{FP + TP} \right] = \frac{m\pi_0 P[x \geq t|FP]}{mP[x \geq t]}$$


$$\widehat{\text{FDR}}(t) = \frac{m\pi_0 P[x \geq t|FP]}{\#x \geq t}$$

- $F(t) = \int_{-\infty}^t f(x)$ using the Empirical cumulative distribution function (ECDF): $\bar{F}(t)$



$$\rightarrow \widehat{FDR}(t) = \frac{m\pi_0 P[x \geq t|FP]}{\#x \geq t} = \frac{\pi_0 P[x \geq t|FP]}{\frac{\#x \geq t}{m}} = \frac{\pi_0 [1 - F(t)]}{1 - \bar{F}(t)}$$


- $F(t) = \int_{-\infty}^t f(x)$ using the Empirical cumulative distribution function (ECDF): $\bar{F}(t)$



$$\rightarrow \widehat{FDR}(t) = \frac{m\pi_0 P[x \geq t|FP]}{\#x \geq t} = \frac{\pi_0 P[x \geq t|FP]}{\frac{\#x \geq t}{m}} = \frac{\pi_0 [1 - F(t)]}{1 - \bar{F}(t)}$$


- How to characterize $F_0(t)$ and π_0 in proteomics?

Target-Decoy approach to establish null distribution




- Searching against decoy databases to generate representative bad hits
- Reversed databases are a popular choice

Target-Decoy approach to establish null distribution



- Searching against decoy databases to generate representative bad hits
- Reversed databases are a popular choice
- Concatenated search


Target-Decoy approach to establish null distribution



- Searching against decoy databases to generate representative bad hits
- Reversed databases are a popular choice
- Concatenated search
- Assumption that bad hits have an equal probability to map on forward (target) and reverse database (decoy)

$$\hat{\pi}_0 = \frac{\#decoys}{\#targets}$$


Target-Decoy approach to establish null distribution



- Searching against decoy databases to generate representative bad hits
- Reversed databases are a popular choice
- Concatenated search
- Assumption that bad hits have an equal probability to map on forward (target) and reverse database (decoy)

$$\hat{\pi}_0 = \frac{\#decoys}{\#targets}$$

Target-Decoy approach to establish null distribution




- Score cutoff?

$$\text{FDR}(x) = E \left[\frac{FP}{FP + TP} \right]$$

- Competitive Target - decoy:

$$\widehat{\text{FDR}}(x) = \frac{\# \text{decoys} | X \geq x}{\# \text{targets} | X \geq x}$$

Target-Decoy approach to establish null distribution



- Score cutoff?

$$\text{FDR}(x) = E \left[\frac{FP}{FP + TP} \right]$$


- Competitive Target - decoy:

$$\widehat{\text{FDR}}(x) = \frac{\# \text{decoys} | X \geq x}{\# \text{targets} | X \geq x}$$

$$\widehat{\text{FDR}}(x) = \frac{\# \text{decoys}}{\# \text{targets}} \frac{\frac{\# \text{decoys} | X \geq x}{\# \text{decoys}}}{\frac{\# \text{targets} | X \geq x}{\# \text{targets}}}$$

$$\widehat{\text{FDR}}(x) = \hat{\pi}_0 \frac{1 - \bar{F}_0(x)}{1 - \bar{F}(x)}$$

Target-Decoy approach to establish null distribution



- Score cutoff?

$$\text{FDR}(x) = E \left[\frac{FP}{FP + TP} \right]$$


- Competitive Target - decoy:

$$\widehat{\text{FDR}}(x) = \frac{\# \text{decoys} | X \geq x}{\# \text{targets} | X \geq x}$$

$$\widehat{\text{FDR}}(x) = \frac{\# \text{decoys}}{\# \text{targets}} \frac{\frac{\# \text{decoys} | X \geq x}{\# \text{decoys}}}{\frac{\# \text{targets} | X \geq x}{\# \text{targets}}}$$

$$\widehat{\text{FDR}}(x) = \hat{\pi}_0 \frac{1 - \bar{F}_0(x)}{1 - \bar{F}(x)}$$

Target-Decoy approach to establish null distribution



- Score cutoff?

$$\text{FDR}(x) = E \left[\frac{FP}{FP + TP} \right]$$

- Competitive Target - decoy:

$$\widehat{\text{FDR}}(x) = \frac{\# \text{decoys} | X \geq x}{\# \text{targets} | X \geq x}$$

$$\widehat{\text{FDR}}(x) = \frac{\# \text{decoys}}{\# \text{targets}} \frac{\frac{\# \text{decoys} | X \geq x}{\# \text{decoys}}}{\frac{\# \text{targets} | X \geq x}{\# \text{targets}}}$$




$$\widehat{\text{FDR}}(x) = \hat{\pi}_0 \frac{1 - \bar{F}_0(x)}{1 - \bar{F}(x)}$$

We have to evaluate that

- The decoys are good simulations of the targets: compare $\bar{F}_0(x)$ with $\bar{F}(x)$
- $\hat{\pi}_0 = \frac{\#decoys}{\#targets}$ is a good estimator for π_0 .
- We will use Probability-Probability-plots for this purpose.
- They plot the ECDFs from two samples in function of each other.




PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot




PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot






PP-plot






PP-plot






PP-plot






PP-plot






PP-plot






PP-plot






PP-plot






PP-plot





PP-plot





PP-plot



PP-plot: pyrococcus





PP-plot: pyrococcus





What about $\hat{\pi}_0$?



PP-plot: pyrococcus





PP-plot: pyrococcus





PP-plot: pyrococcus





PP-plot: pyrococcus





PP-plot: pyrococcus





PP-plot: pyrococcus





PP-plot: pyrococcus





PP-plot: pyrococcus





PP-plot: pyrococcus



PP-plot: pyrococcus



PP-plot: pyrococcus



PP-plot: pyrococcus

