



Statistical Methods for Quantitative MS-Based Proteomics:


1. Identification & False discovery rate

Lieven Clement

Proteomics Data Analysis Shortcourse




Challenges in Label Free MS-based Quantitative Proteomics




Quantification Identification

Challenges in Label Free MS-based Quantitative Proteomics




Identification




E-values

Probability that a random hit produces a higher score than the observed PSM score.




E-values

Probability that a random hit produces a higher score than the observed PSM score.




E-values

Probability that a random hit produces a higher score than the observed PSM score.




E-values

Probability that a random hit produces a higher score than the observed PSM score.



E-values

Probability that a random hit produces a higher score than the observed PSM score.



E-values

Probability that a random hit produces a higher score than the observed PSM score.

- A bad hit is the random hit with the best score so it is also bound to have a low E-value.

E-values

Probability that a random hit produces a higher score than the observed PSM score.

- A bad hit is the random hit with the best score so it is also bound to have a low E-value.
- If we look at E-values for all PSMs they are only useful as a score.

E-values

Probability that a random hit produces a higher score than the observed PSM score.

- A bad hit is the random hit with the best score so it is also bound to have a low E-value.
- If we look at E-values for all PSMs they are only useful as a score.
- We should know the distribution of the maximum score when we want to do the statistics.

Table of Outcomes

	Called Bad	Called Correct	
Bad hit	TN	FP	m_0
Correct hit	FN	TP	m_1
Total	NR	R	m

- TN: number of true negatives
- FP: number of false positives
- FN: number of false negatives
- TP: number of true positives
- NR: number of non-rejections, R: number of rejections

Table of Outcomes

		Called Bad	Called Correct	
Unobservable	Bad hit	TN	FP	m_0
	Correct hit	FN	TP	m_1
Observable	Total	NR	R	m


$FDP = \frac{FP}{FP+TP}$. But is unknown! (FDP: false discovery proportion)

Table of Outcomes

		Called Bad	Called Correct	
Unobservable	Bad hit	TN	FP	m_0
	Correct hit	FN	TP	m_1
Observable	Total	NR	R	m


$$FDR = E \left[\frac{FP}{FP+TP} \right]. \text{ (FDR: false discovery rate)}$$

Search engines return score that discriminates good from bad matches




Search engines return score that discriminates good from bad matches

Score threshold t ?




Search engines return score that discriminates good from bad matches



Score threshold t ?

$$f(x) = \pi_0 f_0(x) + (1 - \pi_0) f_1(x)$$

Search engines return score that discriminates good from bad matches




Score threshold t ?

$$f(x) = \pi_0 f_0(x) + (1 - \pi_0) f_1(x)$$

$$\text{FDR}(t) = E \left[\frac{FP}{FP + TP} \right]$$

Search engines return score that discriminates good from bad matches




Score threshold t ?

$$f(x) = \pi_0 f_0(x) + (1 - \pi_0) f_1(x)$$

$$\text{FDR}(t) = E \left[\frac{FP}{FP + TP} \right]$$

$$\text{FDR}(t) = \frac{mP[FP]P[x \geq t | FP]}{mP[x \geq t]}$$

Search engines return score that discriminates good from bad matches



Score threshold t ?


$$f(x) = \pi_0 f_0(x) + (1 - \pi_0) f_1(x)$$

$$\text{FDR}(t) = E \left[\frac{\text{FP}}{\text{FP} + \text{TP}} \right]$$

$$\text{FDR}(t) = \frac{mP[\text{FP}]P[x \geq t | \text{FP}]}{mP[x \geq t]}$$

$$\text{FDR}(t) = \frac{\pi_0 P_0[x \geq t]}{P[x \geq t]}$$

Search engines return score that discriminates good from bad matches



Score threshold t ?

$$f(x) = \pi_0 f_0(x) + (1 - \pi_0) f_1(x)$$


$$\text{FDR}(t) = E \left[\frac{FP}{FP + TP} \right]$$

$$\text{FDR}(t) = \frac{mP[FP]P[x \geq t|FP]}{mP[x \geq t]}$$

$$\text{FDR}(t) = \frac{\pi_0 P_0[x \geq t]}{P[x \geq t]}$$

$$P[x \geq t] = \int_t^{\infty} f(x) dx$$

How to estimate FDR?




$$\text{FDR}(t) = E \left[\frac{FP}{FP+TP} \right]$$

$$= \frac{\pi_0 P_0[x \geq t]}{P[x \geq t]}$$

$$P[x \geq t] = \int_t^{\infty} f(x) dx$$

How to estimate FDR?



$$\hat{P}[x \geq t] = \frac{\#x \geq t}{m} \Rightarrow$$


$$\text{FDR}(t) = E \left[\frac{FP}{FP+TP} \right]$$

$$= \frac{\pi_0 P_0[x \geq t]}{P[x \geq t]}$$

$$P[x \geq t] = \int_t^{\infty} f(x) dx$$

$$\widehat{\text{FDR}}(t) = \frac{\pi_0 P_0[x \geq t]}{\frac{\#x \geq t}{m}}$$

How to estimate FDR?



$$\text{FDR}(t) = E \left[\frac{FP}{FP+TP} \right]$$


$$= \frac{\pi_0 P_0[x \geq t]}{P[x \geq t]}$$

$$P.[x \geq t] = \int_t^{\infty} f.(x)dx$$

$$\hat{P}[x \geq t] = \frac{\#x \geq t}{m} \Rightarrow \widehat{\text{FDR}}(t) = \frac{\pi_0 P_0[x \geq t]}{\frac{\#x \geq t}{m}}$$


How to characterize $f_0(t)$ and π_0 in proteomics?

Target-Decoy approach to establish null distribution




- Search against decoy database to generate representative bad hits
- Reversed databases are popular

Target-Decoy approach to establish null distribution




- Search against decoy database to generate representative bad hits
- Reversed databases are popular
- Concatenated search

Target-Decoy approach to establish null distribution



- Search against decoy database to generate representative bad hits
- Reversed databases are popular
- Concatenated search


Target-Decoy approach to establish null distribution



- Search against decoy database to generate representative bad hits
- Reversed databases are popular
- Concatenated search
- Assumption: bad hits has equal probability to map on target and decoy

$$\hat{\pi}_0 = \frac{\#decoys}{\#targets}$$

Target-Decoy approach to establish null distribution



- Search against decoy database to generate representative bad hits
- Reversed databases are popular
- Concatenated search
- Assumption: bad hits has equal probability to map on target and decoy

$$\hat{\pi}_0 = \frac{\#decoys}{\#targets}$$


- Score cutoff:

$$FDR(x) = E \left[\frac{FP}{FP+TP} \right]$$


Target-Decoy approach to establish null distribution

- Competitive Target - decoy:

$$\widehat{\text{FDR}}(x) = \frac{\# \text{decoys} | X \geq x}{\# \text{targets} | X \geq x}$$



Target-Decoy approach to establish null distribution



- Competitive Target - decoy:


$$\widehat{\text{FDR}}(x) = \frac{\# \text{decoys} | X \geq x}{\# \text{targets} | X \geq x}$$

$$\widehat{\text{FDR}}(x) = \frac{\# \text{decoys}}{\# \text{targets}} \frac{\frac{\# \text{decoys} | X \geq x}{\# \text{decoys}}}{\frac{\# \text{targets} | X \geq x}{\# \text{targets}}}$$

$$\widehat{\text{FDR}}(x) = \hat{\pi}_0 \frac{\int_t^{+\infty} f_0(x) dx}{\int_t^{+\infty} f(x) dx}$$

$$\widehat{\text{FDR}}(x) = \hat{\pi}_0 \frac{\hat{P}_0[X \geq x]}{\hat{P}[X \geq x]}$$

Target-Decoy approach to establish null distribution



- Competitive Target - decoy:


$$\widehat{\text{FDR}}(x) = \frac{\# \text{decoys} | X \geq x}{\# \text{targets} | X \geq x}$$

$$\widehat{\text{FDR}}(x) = \frac{\# \text{decoys}}{\# \text{targets}} \frac{\frac{\# \text{decoys} | X \geq x}{\# \text{decoys}}}{\frac{\# \text{targets} | X \geq x}{\# \text{targets}}}$$

$$\widehat{\text{FDR}}(x) = \hat{\pi}_0 \frac{\int_t^{+\infty} f_0(x) dx}{\int_t^{+\infty} f(x) dx}$$

$$\widehat{\text{FDR}}(x) = \hat{\pi}_0 \frac{\hat{P}_0[X \geq x]}{\hat{P}[X \geq x]}$$

Target-Decoy approach to establish null distribution



- Competitive Target - decoy:

$$\widehat{\text{FDR}}(x) = \frac{\# \text{decoys} | X \geq x}{\# \text{targets} | X \geq x}$$

$$\widehat{\text{FDR}}(x) = \frac{\# \text{decoys}}{\# \text{targets}} \frac{\frac{\# \text{decoys} | X \geq x}{\# \text{decoys}}}{\frac{\# \text{targets} | X \geq x}{\# \text{targets}}}$$

$$\widehat{\text{FDR}}(x) = \hat{\pi}_0 \frac{\int_t^{+\infty} f_0(x) dx}{\int_t^{+\infty} f(x) dx}$$

$$\widehat{\text{FDR}}(x) = \hat{\pi}_0 \frac{\hat{P}_0[X \geq x]}{\hat{P}[X \geq x]}$$

Assess TDA assumptions


We have to evaluate that

- The decoys are good simulations of the bad target hits: compare distributions $F_0(x)$ with $F(x)$

$$F_0(x) = \int_{-\infty}^x f_0(x)dx \quad \leftrightarrow \quad F(x) = \int_{-\infty}^x f(x)dx$$

- $\hat{\pi}_0 = \frac{\#decoys}{\#targets}$ is a good estimator for π_0 .
- We will use Probability-Probability-plots (PP-plot) for this purpose.




- To make PP-plots we need estimates for $F_0(x)$ and $F(x)$.
- The empirical cumulative distribution (ECDF) is used for that purpose



$$\hat{F}_0(x) = \frac{\#decoys | X \leq x}{\#decoys}, \quad \hat{F}(x) = \frac{\#targets | X \leq x}{\#targets}$$




PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot

PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot




PP-plots have the property that they show a straight 45 degree line through the origin if and only if both distributions are equivalent.






PP-plot






PP-plot






PP-plot






PP-plot






PP-plot






PP-plot






PP-plot






PP-plot






PP-plot





PP-plot





PP-plot



PP-plot: pyrococcus





PP-plot: pyrococcus





What about $\hat{\pi}_0$?



PP-plot: pyrococcus





PP-plot: pyrococcus





PP-plot: pyrococcus





PP-plot: pyrococcus





PP-plot: pyrococcus





PP-plot: pyrococcus





PP-plot: pyrococcus





PP-plot: pyrococcus





PP-plot: pyrococcus



PP-plot: pyrococcus



PP-plot: pyrococcus



PP-plot: pyrococcus

