

# **UPDATE: DOWNSCALING OF THE SEAS5 TO THE GREATER REGION OF LUXEMBOURG**

VERIFICATION

FELICITAS PAIXAO

LUXEMBOURG INSTITUTE OF SCIENCE AND TECHNOLOGY  
BELVAUX LUXEMBOURG

28 08 2019

# **SUMMARY OF WORK UP TO MEETING IN TRIER**

## **Creating a literature pool to acquire knowledge about task**

Literature on:

- seasonal forecasting and products therefor
  - ▶ decided on the Seasonal Forecast System 5 (SEAS5) provided by the ECMWF
- dynamical downscaling and products therefor
  - ▶ decided on the COSMO-CLM provided by a range of meteorological services and maintained by the CLM community
- ensemble forecasting
- verification methods and measures
  - ▶ how to create a reference dataset
  - ▶ decided on E-OBS provided by European Climate Assessment and Dataset

# SUMMARY OF WORK UP TO MEETING IN TRIER

## **First requirements concerning the necessary datasets**

SEAS5:

- getting familiar with dataset in general
  - ▶ spatial resolution and driving atmospheric/ocean models
  - ▶ range of variables and temporal resolution of them
  - ▶ which data is available (slightly pre-processed (monthly means/ensemble mean/etc.), completely raw (continuous, probabilistic values) and much more
- getting familiar with technicalities surrounding the dataset
  - ▶ "streams" through which different, above mentioned data is provided
  - ▶ getting familiar with Meteorological Archival and Retrieval System (MARS)
  - ▶ getting familiar with the specific dataset descriptions via "confluence", the wiki and collaboration platform of ECMWF
- collecting more literature on this specific product

## **First requirements concerning the necessary datasets**

E-OBS:

- getting familiar with dataset in general
  - ▶ spatial and temporal resolution
  - ▶ range of variables
  - ▶ how is the dataset produced
- getting familiar with technicalities surrounding the dataset
  - ▶ downloading the E-OBS manually, as it is just 5 available variables
- collecting more literature on this specific product

## Acquirement of SEAS5 data with MARS

- MARS uses either the Web User Interface on the ECMWF website or can be used via a Web-API service interface which uses a python script
- Web-API client has to be installed (description and all steps until it's usable can be found in confluence wiki<sup>1</sup>)
- to not download/overwrite already existing files script also checks for filename and variations of it
- the whole looping procedure, as well as the identification as an authorized user is done in bash commands
- all needed variables are then redirected to an outputfile (request\_date\_em.mars) that can be read by the mars.py script that was installed above

---

<sup>1</sup><https://confluence.ecmwf.int/display/WEBAPI/Access+MARS>

# SUMMARY OF WORK UP TO MEETING IN TRIER

## SEAS5 data

- 6 hrly continuous forecast (fc)
- 51 ensemble members
- 7 months of data per initialization (ini) date
- 4 ini dates (lead times)
- global latlon grid
- variables: t2m in K, total precipitation in m

## E-OBS data

- daily mean values
- ensemble mean
- data from 2011 until 2019
- european latlon grid (
- variables: air temperature, thickness of rainfall amount in mm
- no values over ocean domains

## Regridding

- Both datasets were regridded according to a reference file for the EU-CORDEX domain (here called geogrid)
  - ▶ geogrid is a "generic" grid type
  - ▶ created grid file for geogrid to regrid the other datasets
  - ▶ "curvilinear" as grid type = rectangular shape, not following lon/lat lines
- regridded with the cdo command remapbil
  - ▶ bilinear interpolation remapping
  - ▶ detailed description of the interpolation scheme can be found in the SCRIP<sup>2</sup> description

---

<sup>2</sup><https://github.com/SCRIP-Project/SCRIP/blob/master/SCRIP/doc/SCRIPusers.pdf>



## Tasks of matching

- creating daily means for SEAS5
- masking SEAS5 with E-OBS data to get the same NA values at the same grid points
- inter comparing timeline of datasets and cutting them where ever the other set had no values
- changing units of E-OBS dataset to match the SEAS5 dataset
- create intuitive names to loop through for E-OBS
- creating script that sets up a directory tree from base directory

## **Datasets were now**

- on the same grid
- at the same temporal resolution
- within the same time range
- matching in terms of NA values
- matching in terms of units

## First attempt at verification

- as JJA is of interest datasets were reduced to it
- creating RMSE, MAE and bias with the same scripting method
  - ▶ looping through each time step of grid points of SEAS5 ensemble mean, creating the mean, then standard deviation and then RMSE/MAE/Bias

# **OUTCOME OF MEETING**

# OUTCOME OF MEETING

## Reference dataset

E-OBS dataset is insufficient, as there is not enough observations and therefore the interpolated results are lacking.

ERA5 would be a reasonable choice for substitution, as it is a well described and verified reanalysis. The data can be retrieved from the Copernicus<sup>3</sup> server. Details of dataset on later slides.

## Verification metrics

Equations should be clearly stated.

As there were problems in description of the used RMSE in the metrics it is very important to explain and document way of using different verification measures.

---

<sup>3</sup><https://cds.climate.copernicus.eu/>

# OUTCOME OF MEETING

## Forecast dataset

Temporal range of operational forecast not big enough, re-forecast for a bigger span would be more valuable.

The re-forecast of the SEAS5 dataset is also available on the Copernicus<sup>4</sup> server. It has a set of 25 ensemble members and a resolution of  $1^{\circ} \times 1^{\circ}$ . More details of dataset on later slides.

## Literature reference

Identification of papers with similar problem of verification.

I go forward mainly but not only with the paper by Johnson et al. [12] on the verification of specifically the SEAS5 as I use the re-forecast dataset now.

---

<sup>4</sup><https://cds.climate.copernicus.eu/>

# **SETUP AFTER MEETING**

## Forecast

SEAS5 operational forecast

- initial dates of 01.03., 01.04., 01.05., 01.06. for different lead times (for JJA)
- 51 ensemble members
- total precipitation and 2 meter temperature
- $0.4^{\circ}$

Problem:

Not enough temporal cover (just 3 months of actual forecast). A span of ~30 years would be much better for a sufficient verification.



## Reference

### E-OBS dataset

- daily values for JJA
- ensemble mean
- precipitation and 2 meter temperature
- $0.25^\circ$

### Problem:

Measurement stations are sparse, which makes the interpolation very inaccurate. A reanalysis would be better for a sufficient verification.

# **PROGRESS SINCE MEETING**

Creation of automated script to download SEAS5 re-forecast dataset from Copernicus server.

```

1  #!/usr/bin/env python
2  import cdsapi
3  import os.path
4
5  c = cdsapi.Client()
6
7  pwd = os.getcwd()
8  dirout = [str(pwd) + '/reforecast/']
9  if not os.path.exists(dirout[0]):
10     os.mkdir(dirout[0])
11
12  os.chdir(dirout[0])
13
14  mon=[ "%-2d" % i for i in range(3,7)]
15
16  steps = []
17  for i in range(6,5161,6):
18     steps.append(i)
19
20  for y in range(1993, 2017):
21     for m in mon:
22         if not
os.path.exists('refc_{year}_{month}.grib'.format(year=y,month=m))
23         and
os.path.exists('refc_{year}_{month}.nc'.format(year=y,month=m))
24         and
os.path.exists('BJ_refc_{year}_{month}_JJA.nc'.format(year=y,month=m))
25         )) and
os.path.exists('refc_{year}_{month}_JJA.nc'.format(year=y,month=m)):
26
c.retrieve('seasonal-original-single-levels',
27
{
'originating_centre': 'ecmwf',
28
'system': '5',
29
'variable': '2m_temperature',
30
'year': y,
31
'month': m,
32
'day': '01',
33
'leadtime_hour': steps,
34
'format': 'grib'
35
},
'refc_{year}_{month}.grib'.format(year=y,month=m)

```

## Re-forecast download script

- creates output directory, if not yet created
- loops over months for ini dates (03/04/05/06) for years from 1993 until 2017
- 25 ensemble members
- 6 hourly temporal resolution
- 1° spatial resolution
- works similar to first retrieval script but retrieves directly, not creating a temporal file (request.mars above)

Creation of automated script to download ERA5 dataset from Copernicus server.

```
1  #!/usr/bin/env python
2  import cdsapi
3  from datetime import datetime, timedelta
4  from dateutil.relativedelta import relativedelta
5  import os.path
6
7  c = cdsapi.Client()
8
9  pwd = os.getcwd()
10 dirout = [str(pwd) + '/ERA5/']
11 if not os.path.exists(dirout[0]):
12     os.mkdir(dirout[0])
13
14 os.chdir(dirout[0])
15
16 def days_of_month(y, m):
17     d0 = datetime(y, m, 1)
18     d1 = d0 + relativedelta(months=1)
19     out = list()
20     while d0 < d1:
21         out.append(d0.strftime('%Y-%m-%d'))
22         d0 += timedelta(days=1)
23     return out
24
25 year=range(1993, 2019)
26 mon=range(6,9)
27
28 for y in year:
29     for m in mon:
30         timerange=days_of_month(y, m)
31         file=[str(dirout[0]+'ERA5_{yyyy}_{0mm}.grib'.format(yyyy=y,mm=m))]
32         if not os.path.exists(file[0]):
33             c.retrieve('reanalysis-era5-single-levels',
34                 {
35                     'product_type': 'reanalysis',
36                     'format': 'grib',
37                     'variable': ['2m_temperature', 'total_precipitation'],
38                     'date': timerange,
39                     'time': ['00:00', '06:00', '12:00', '18:00']
40                 },
41                 file[0]
42             )
43
```

### **ERA5 download script**

- creates output directory, if not yet created
- loops over months JJA for years from 1993 until 2017
- reanalysis without members (HRES reference data)
- 6 hourly temporal resolution
- 0.28° spatial resolution
- works similar to first retrieval script but retrieves directly, not creating a temporal file (request.mars above)

## Tasks of matching

- split SEAS5 re-forecast into ensemble members, as they are not treated as a 4th dimension in raw dataset but integrated into "timeline"
- the remapping procedure remains the same as with previous datasets
- reducing SEAS5 to JJA only (ERA5 was already downloaded matching only these months)
- as they are forced by the same atmospheric model matching is much easier to do
- both are also downloaded in GRIB format and have to be transformed

**Mainly after Johnson et al. (2019) [12], Jolliffe (2012) [13] and Wilks (2011) [31]**

- Deterministic measures (ensemble mean)
  - ▶ RMSE
  - ▶ MAE
  - ▶ ME
- Probabilistic measures (ensemble members)
  - ▶ Reliability diagrams
  - ▶ ACC
  - ▶ CRPSS
  - ▶ Brier score



## Equations after which deterministic metrics is calculated:

- Root Mean Squared Error (RMSE)[13]

$$RMSE = \frac{1}{N} \sum_{t=1}^N \sqrt{\frac{1}{M} \sum_{i=1}^M (f_{i,t} - o_{i,t})^2}$$

where M is the number of points in the grid, N the years over which values are taken for JJA,  $f/o_{i,t}$  the value at time t and location i and answers the question "What is the mean grid-based RMSE, averaged across all times (JJA)?"

## Equations after which determinitsic metrics is calculated:

- Mean Absolute Error (MAE)[13]

- Mean Error (ME) - Bias

# VERIFICATION METRICS

## Equations after which probabilistic metrics is calculated:

- Anomaly correlation coefficient (ACC) [13]


$$ACC_t = \frac{\sum_{i=1}^M (f'_i - \bar{f}') (o'_i - \bar{o}')}{M_{s_{f'} s_{o'}}}$$




**To Do NEXT:**

IMPLEMENTING OF VERIFICATION  
METRICS ON NEW DATASET





# REFERENCES I

 HANNAH C. ARNOLD.  
**SHOULD WEATHER AND CLIMATE PREDICTION MODELS BE DETERMINISTIC OR STOCHASTIC?**  
*Weather*, 68, October 2013.

 PHILIP E. BETT, HAZEL E. THOMTON, ALBERTO TROCCOLI, MATTEO DE FELICE, EMMA SUCKLING, LAURENT DUBUS, YVES-MARIE SAINT-DRENAN, AND DAVID J. BRAYSHAW.  
**A SIMPLIFIED SEASONAL FORECASTING STRATEGY, APPLIED TO WIND AND SOLAR POWER IN EUROPE.**  
*Climate Services*, 24:3, 2019.

 JONAS BHEND, IRINA MAHLSTEIN, AND MARK A. LINIGER.  
**PREDICTIVE SKILL OF CLIMATE INDICES COMPARED TO MEAN QUANTITIES IN SEASONAL FORECASTS.**  
*Quarterly Journal of the Royal Meteorological Society*, 143(702):184–194, January 2017.

# REFERENCES II

-  GLENN W. BRIER AND ROGER A. ALLEN.  
**VERIFICATION OF WEATHER FORECASTS, PAGES 841–848.**  
American Meteorological Society, Boston, MA, 1951.
-  CLM-COMMUNITY.  
**CLIMATE LIMITED-AREA MODELLING COMMUNITY.**  
<https://www.clm-community.eu/index.php?menuid=220>.
-  ALEJANDRO DI LUCA, RAMON DE ELIA, AND RENE LAPRISE.  
**CHALLENGES IN THE QUEST FOR ADDED VALUE OF REGIONAL CLIMATE DYNAMICAL DOWNSCALING.**  
*Current Climate Change Reports*, 2015.
-  E. DIEZ, B. ORFILA, M. D. FRIAS, J. FERNANDEZ, A. S. COFINO, AND J. M. GUTIERREZ.  
**DOWNSCALING ECMWF SEASONAL PRECIPITATION FORECASTS IN EUROPE USING THE RCA MODEL.**  
*Tellus A: Dynamic Meteorology and Oceanography*, 63(4):757–762, 2011.

# REFERENCES III



ECMWF.

**SEAS5 USER GUIDE.**

ECMWF, November 2017.



ROBERT FAWCETT.

**VERIFICATION TECHNIQUES AND SIMPLE THEORETICAL FORECAST MODELS.**

*Weather and Forecasting*, 2008.



MARKEL GARCIA-DIEZ, JESUS FERNANDEZ, AND ROBERT VAUTARD.

**AN RCM MULTI-PHYSICS ENSEMBLE OVER EUROPE:  
MULTI-VARIABLE EVALUATION TO AVOID ERROR COMPENSATION.**

*Climate Dynamics*, 2015.




FILIPPO GIORGI AND WILLIAM J. GUTOWSKI JR.

**REGIONAL DYNAMICAL DOWNSCALING AND THE CORDEX INITIATIVE.**

*Annual Review of Environment and Resources*, 2015.

# REFERENCES IV

 S. J. JOHNSON, T. N. STOCKDALE, L. FERRANTI, M. A. BALMASEDA, F. MOLteni, L. MAGNUSSON, S. TIETSCH, D. DECREMER, A. WEISHEIMER, G. BALSAMO, S. P. E. KEELEY, K. MOGENSEN, H. ZUO, AND B. M. MONGE-SANZ.


**SEAS5: THE NEW ECMWF SEASONAL FORECAST SYSTEM.**

*Geoscientific Model Development*, 12(3):1087–1117, 2019.

 IAN T. JOLLIFFE AND DAVID B. STEPHENSON.

**FORECAST VERIFICATION - A PRACTITIONER'S GUIDE IN ATMOSPHERIC SCIENCE.**

John Wiley and Sons, Ltd., second edition, 2012.

 MACIEJ KRYZA, KINGA WAIASZEK, HANNA OJRZYNSKA, MARIUSZ SZYMANOWSKI, MAIGORZATA WERNER, AND ANTHONY J. DORE.  
**HIGH-RESOLUTION DYNAMICAL DOWNSCALING OF ERA-INTERIM USING THE WRF REGIONAL CLIMATE MODEL FOR THE AREA OF POLAND. PART 1: MODEL CONFIGURATION AND STATISTICAL EVALUATION FOR THE 1981–2010 PERIOD.**

*Pure and Applied Geophysics*, 2016.



# REFERENCES V



ERIC M. LAFLAMME, ERNST LINDER, AND YIBIN PAN.  
**STATISTICAL DOWNSCALING OF REGIONAL CLIMATE MODEL OUTPUT TO  
ACHIEVE PROJECTIONS OF PRECIPITATION EXTREMES.**  
*Weather and Climate Extremes*, 2015.






YOAV LEVI AND ITZHAK CARMONA.  
**SEASONAL FORECAST VERIFICATION AND APPLICATION IN TIMES OF  
CHANGE.**  
*Earth System Dynamics*, 2016.



DELEI LI, BAOSHU YIN, JIANLONG FENG, ALESSANDRO DOSIO, BEATE  
GEYER, JIFENG QI, HONGYUAN SHI, AND ZHENHUA XU.  
**PRESENT CLIMATE EVALUATION AND ADDED VALUE ANALYSIS OF  
DYNAMICALLY DOWNSCALED SIMULATIONS OF CORDEX-EAST ASIA.**  
*Journal of Applied Meteorology and Climatology*, 2018.

# REFERENCES VI

-  IRINA MAHLSTEIN, CHRISTOPH SPIRIG, MARK A. LINIGER, AND CHRISTOF APPENZELLER.  
**ESTIMATING DAILY CLIMATOLOGIES FOR CLIMATE INDICES DERIVED FROM CLIMATE MODEL DATA AND OBSERVATIONS.**  
*Journal of Geophysical Research: Atmospheres*, 120:2808–2818, April 2015.
-  R. MANZANAS, J. M. GUTIERREZ, J. FERNANDEZ, E. MEIJGAARD VAN, S. CALMANTI, M. E. MAGARINO, A. S. COFINO, AND S. HERRERA.  
**DYNAMICAL AND STATISTICAL DOWNSCALING OF SEASONAL TEMPERATURE FORECASTS IN EUROPE: ADDED VALUE FOR USER APPLICATION.**  
*Climate Services*, 9:44–56, 2017.
-  MARION MITTERMAIER, NIGEL ROBERTS, AND SIMON A. THOMPSON.  
**A LONG-TERM ASSESSMENT OF PRECIPITATION FORECAST SKILL USING THE FRACTIONS SKILL SCORE.**  
*Meteorological Applications*, 20:176–186, 2013.

## REFERENCES VII



ALLAN H. MURPHY.

**WHAT IS A GOOD FORECAST? AN ESSAY ON THE NATURE OF GOODNESS IN WEATHER FORECASTING.**

*American Meteorological Society*, pages 281–293, 1993.



GRIGORY NIKULIN, SHAKEEL ASHARAF, MARËA EUGENIA MAGARIËO, SANDRO CALMANTI, RITA M. CARDOSO, JONAS BHEND, JESËS FERNËNDEZ, MARËA DOLORES FRËNAS, KRISTINA FRËHLLICH, BARBARA FRËIJH, SIXTO HERRERA GARCËA, RODRIGO MANZANAS, JOSËL MANUEL GUTIËLRREZ, ULF HANSSON, MICHAEL KOLAX, MARK A. LINIGER, PEDRO M. M. SOARES, CHRISTOPH SPIRIG, RICARDO TOME, AND KLAUS WYSER.  
**DYNAMICAL AND STATISTICAL DOWNSCALING OF A GLOBAL SEASONAL HINDCAST IN EASTERN AFRICA.**

*Climate Services*, 9:72–85, 2018.




TIM PALMER.

**THE ECMWF ENSEMBLE PREDICTION SYSTEM: LOOKING BACK (MORE THAN) 25 YEARS AND PROJECTING FORWARD 25 YEARS.**

*Quarterly Journal of the Royal Meteorological Society*, 2018.

## REFERENCES VIII


-  MARKKU RUMMUKAINEN.  
**ADDED VALUE IN REGIONAL CLIMATE MODELING.**  
*WIRES Climate Change*, 7:145–159, 2016.
-  SILJE LUND SORLAND, CHRISTOPH SCHÄDR, DANIEL LÄIJTHI, AND ERIK KJELLSTROM.  
**BIAS PATTERNS AND CLIMATE CHANGE SIGNALS IN GCM-RCM MODEL CHAINS.**  
*Environmental Research Letters*, 13, 2018.
-  PIET TERMONIA, BERT VAN SCHAEYBROECK, LESLEY DE CRUZ, ROZEMIEN DE TROCH, OLIVIER GIOT, RAFIQ HAMDİ, STÄLPHANE VANNITSEM, FRANCOIS DUCHENE, PATRICK WILLEMS, HOSSEIN TABARI, ELS VAN UYTVEN, PARISA HOSSEINZADEHTALAEI, NICOLE VAN LIPZIG, HENDRIK WOUTERS, SAM VANDEN BROUCKE, MATTHIAS DEMUZERE, JEAN-PASCAL VAN YPERSELE, PHILIPPE MARBAIX, CECILLE VILLANUEVA-BIRRIEL, XAVIER FETTWEIS, CORALINE WYARD, CHLOE SCHOLZEN, SEBASTIEN DOUTRELOUP, KOEN DE RIDDER, ANNE GOBIN, DIRK LAUWAET, TRISSEVGENI STAVRAKOU,

## REFERENCES IX

MAITE BAUWENS, JEAN-FRANCOIS M/UELLER, PATRICK LUYTEN,  
STEPHANIE PONSAR, DRIES VAN DEN EYNDE, AND ERIC POTTIAUX.


**COMBINING REGIONAL DOWNSCALING EXPERTISE IN BELGIUM: CORDEX  
AND BEYOND.**

*Belgian Research Action through Interdisciplinary Networks*, 2018.

 P. R. TIWARI, S. C. KAR, U. C. MOHANTY, S. DEY, P. SINHA, M. S. SHEKHAR,  
AND R. S. SOKHI.

**COMPARISON OF STATISTICAL AND DYNAMICAL DOWNSCALING METHODS  
FOR SEASONAL SCALE WINTER PRECIPITATION PREDICTIONS OVER NORTH  
INDIA.**

*International Journal of Climatology*, 2018.

 F. VITART, C. ARDILOUZE, A. BONET, A. BROOKSHAW, M. CHEN,  
C. CODOREAN, M. DÉQUÉ, L. FERRANTI, E. FUCILE, M. FUENTES,  
H. HENDON, J. HODGSON, H.-S. KANG, A. KUMAR, H. LIN, G. LIU, X. LIU,  
P. MALGUZZI, I. MALLAS, M. MANOUSSAKIS, D. MASTRANGELO,  
C. MAC LACHLAN, C. MC LEAN, A. MINAMI, R. MLADEK, T. NAKAZAWA,  
S. NAJM, Y. NIE, M. RIXEN, A. W. ROBERTSON, P. RUTI, C. SUN, Y. TAKAYA,

# REFERENCES X

M. TOLSTYKH, F. VENUTI, D. WALISER, S. WOOLNOUGH, T. WU, D.-J. WON, H. XIAO, R. ZARIPOV, AND L. ZHANG.

**THE SUBSEASONAL TO SEASONAL (S2S) PREDICTION PROJECT DATABASE.**  
*Bulletin of the American Meteorological Society*, 2017.



A. WEISHEIMER AND T. N. PALMER.

**ON THE RELIABILITY OF SEASONAL CLIMATE FORECASTS.**  
*Journal of the Royal Society Interface*, 11, 2014.



CHRISTOPHER J. WHITE, HENRIK CARLSEN, ANDREW W. ROBERTSON, RICHARD J. T. KLEIN, JEFFREY K. LAZO, ARUN KUMAR, FREDERIC VITART, ERIN COUGHLAN DE PEREZ, ANDREA J. RAY, VIRGINIA MURRAY, SUKAINA BHARWANI, DAVE MACLEOD, RACHEL JAMES, LORA FLEMING, ANDREW P. MORSE, BERND EGGEN, RICHARD GRAHAM, ERIK KJELLSTRÅM, EMILY BECKER, KATHLEEN V. PEGION, NEIL J. HOLBROOK, DARRYN McEVOY, MICHAEL DEPLEDGE, SARAH PERKINS-KIRKPATRICK, TIMOTHY J. BROWN, ROGER STREET, LINDSEY JONES, TOMAS A. REMENYI, INDI HODGSON-JOHNSTON, CARLO BUONTEMPO, ROB LAMB, HOLGER MEINKE, BERIT ARHEIMER, AND STEPHEN E. ZEBIAK.

# REFERENCES XI

## **POTENTIAL APPLICATIONS OF SUBSEASONAL-TO-SEASONAL (S2S) PREDICTIONS.**

*Royal Meteorological Society*, 2017.



DANIEL S. WILKS.

## **STATISTICAL METHODS IN THE ATMOSPHERIC SCIENCES, VOLUME 100 OF INTERNATIONAL GEOPHYSICS.**

Elsevier, third edition, 2011.




YONGKANG XUE, ZAVISA JAJNIC, JIMY DUDHIA, RATKO VASIC, AND FERNANDO DE SALES.

## **A REVIEW ON REGIONAL DYNAMICAL DOWNSCALING IN INTRA-SEASONAL TO SEASONAL SIMULATION/PREDICTION AND MAJOR FACTORS THAT AFFECT DOWNSCALING ABILITY.**

*Atmospheric Research*, 2014.

## REFERENCES XII

 DEJIAN YANG, XIU-QUN YANG, DAN YE, XUGUANG SUN, JIABEI FANG, CUJIAO CHU, TAO FENG, YIQUN JIANG, JIN LIANG, XUEJUAN REN, YAOCUN ZANG, AND YUJIN TANG.

**ON THE RELATIONSHIP BETWEEN PROBABILISTIC AND DETERMINISTIC SKILLS IN DYNAMICAL SEASONAL CLIMATE PREDICTION.**

*Journal of Geophysical Research: Atmospheres*, 123:5261–5283, May 2018.