

THEORY

- Pham (finance)

NUMERIC

- ? (mainly the numerical appendix), Moll's website (more examples)
- ? (good slides)

Problem of Household

$$\begin{aligned} \max_{\{c_t\}_{t \geq 0}} \mathbb{E}_0 \int_0^{\infty} e^{-\rho t} u(c_t) dt \\ \text{S.t } da_t = \{r_t a_t + z_t - c_t\} dt \\ z_t = (w - b) dq_{\mu} - (w - b) dq_s, \quad z_t \in \{b, w\} \\ a_t \geq \underline{a} \end{aligned}$$

HJB equation

$$\rho v_i(a, t) = \max_c \left\{ u(c) + \partial_a v_i(a, t) [r_t a + z_i - c] \right\} + \lambda_i \left[v_{-i}(a, t) - v_i(a, t) \right] + \partial_t v_i(a, t) \quad (1)$$

$$\rho v_i(a) = \max_c \left\{ u(c) + v'_i(a) [ra + z_i - c] \right\} + \lambda_i \left[v_{-i}(a) - v_i(a) \right] \quad (2)$$

Where did borrowing constraint go?

- In continuous time, an undistorted foc holds everywhere in the state space
- Borrowing constraint shows only as *state constraint boundary condition*

$$u'(c_i(\underline{a}, t)) = \partial_{\underline{a}} v_i(\underline{a}, t) \geq u'(r_t \underline{a} + z_i) \quad (3)$$

which ensures $c_i(\underline{a}, t) \leq r_t \underline{a} + z_i$ so that the borrowing constraint is never violated.

In general form, the HJB equation is an equation of the form

$$0 = F(\mathbf{x}, V, DV, D^2V) \quad (4)$$

where $\mathbf{x} := (x, \tau)$.

Finite difference methods: replace derivatives by differences.

Suppose we define a grid $\{x_0, x_1, \dots, x_i, \dots\}$ and a set of timesteps $\{i\Delta : i \in \mathbb{N}\}$. Let $V_i^n \approx V(x_i, \tau_n)$ be the approximate value of the solution at node x_i time $\tau^n := T - t$. Then we can write a general discretization of the HJB equation at node (sx_i, τ^{n+1})

$$0 = S_i^{n+1}((\Delta, \Delta x), V_i^{n+1}, \{V_j^m\}) \quad (5)$$

Condition (Monotonicity) . — The numerical scheme (5) is monotone if

$$S_i^{n+1}(\cdot, V_i^{n+1}, \{Y_j^m\}) \leq S_i^{n+1}(\cdot, V_i^{n+1}, \{Z_j^m\})$$

for all $Y \geq Z$.

Condition (Stability) . — The numerical scheme (5) is stable if for every $\tilde{\Delta} > 0$ it has a solution which is uniformly bounded independently of $\tilde{\Delta}$.

Condition (Consistency) . — The numerical scheme (5) is consistent if for every smooth function ϕ with bounded derivatives we have

$$S_i^{n+1}(\tilde{\Delta}, \phi(\mathbf{x}_i^{n+1}), \{\phi(\mathbf{x}_j^m)\}) \rightarrow F(\mathbf{x}, \phi, D\phi, D^2\phi)$$

as $\tilde{\Delta} \rightarrow 0$ and $\mathbf{x}_i^{n+1} \rightarrow \mathbf{x}$.

Theorem ? . *If the numerical scheme (5) satisfies monotonicity, stability and consistency conditions, then its solution converges locally uniformly to the unique viscosity solution of (4).*

Discretization

Back to our example

Recall our *time-dependent* HJB equation as

$$\partial_\tau v_k(a, \tau) + \rho v_k(a, \tau) - \sup_{c \in \Gamma_k(a)} \left\{ u(c) + \mathcal{D}_{\tau-\tau}^c v_k(a, \tau) \right\} = 0 \quad (6)$$

where

$$\mathcal{D}_t^c \phi_k(a) = \partial_a \phi_k(a) [r_t a + z_k - c] + \lambda_k [\phi_{-k}(a) - \phi_k(a)]$$

Define a grid $\{a_1, a_2, \dots, a_i, \dots\}$ and let $v_k^n = (v_k(a_1, \tau^n), \dots, v_k(a_i, \tau^n), \dots)'$. Discretizing this equation requires deciding upon

- which fd approximation to use: forward/backward differencing

$$\partial_a v_k(a) \approx \frac{v_{k,i+1} - v_{k,i}}{a_{i+1} - a_i}, \quad \partial_a v_k(a) \approx \frac{v_{k,i} - v_{k,i-1}}{a_i - a_{i-1}},$$

- Implicit/explicit timestepping

Discretization

Back to our example

Let \mathcal{D}^c be the discrete form of the differential operator \mathcal{D}^c , so that

$$(\mathcal{D}^c v)_{k,i} = \alpha_{k,i}(c)v_{k,i-1} + \beta_{k,i}(c)v_{k,i+1} - (\alpha_{k,i}(c) + \beta_{k,i}(c) + \lambda_i)v_{k,i} + \lambda_i v_{-k,i}$$

and the discretization

$$\frac{v_{k,i}^{n+1} - v_{k,i}^n}{\Delta} - \sup_{c \in \Gamma_{k,i}} \left\{ u(c) + (\mathcal{D}^c v^{n(+1)})_{k,i} \right\} = 0 \quad (7)$$

where discretization can use forward, backward or central discretization. If

$$\alpha_{k,i}^{n+1} \geq 0, \quad \beta_{k,i}^{n+1} \geq 0$$

we say that (7) is *positive coefficient discretization*.

Why do we care?

We care because a *positive coefficient discretization* is also *monotone*. To see it check that

$$S_{k,i}^{n+1}(\tilde{\Delta}, v_{k,i}^{n+1}, v_{k,i+1}^{n(+1)}, v_{k,i-1}^{n(+1)}, v_{k,i}^n, v_{-k,i}^{n(+1)})$$

is a nonincreasing function of the neighbor nodes $\{v_{\ell,j}^m\}$. Check a example!

Upwind scheme

In order to ensure a *positive coefficient discretization* our choice of central/forward/backward differencing will depend, in general, on the control c . A useful rule for this problem is to use the so-called *upwind scheme*.

IDEA: *Use forward difference whenever drift is positive, and use backward whenever it is negative.*

Suppose that we have the value of consumption $c_{k,i}$ at a particular node. Let $s_{k,i} = ra_i + z_k - c_{k,i}$. In this case, the derivatives are approximated

$$\dots \frac{v_{k,i+1} - v_{k,i}}{a_{i+1} - a_i} s_{k,i}^+ + \frac{v_{k,i} - v_{k,i-1}}{a_i - a_{i-1}} s_{k,i}^- + \dots$$

which in terms of our α, β

$$\alpha_{k,i}^{up} = -\frac{s_{k,i}^-}{a_i - a_{i-1}} \geq 0, \quad \beta_{k,i}^{up} = \frac{s_{k,i}^+}{a_{i+1} - a_i} \geq 0$$

But we don't know $c_{k,i}$!!! HJB equation is highly nonlinear, so we need an iterative method to solve it.

Start with a vector v^n and update v^{n+1} according to

$$\frac{v_{k,i}^{n+1} - v_{k,i}^n}{\Delta} + \rho v_{k,i}^{n+1} = u(c_{k,i}^n) + \frac{v_{k,i+1}^{n+1} - v_{k,i}^{n+1}}{a_{i+1} - a_i} [s_{k,i}^{F,n}]^+ + \frac{v_{k,i}^{n+1} - v_{k,i-1}^{n+1}}{a_i - a_{i-1}} [s_{k,i}^{B,n}]^- + \lambda_k [v_{-k,i}^{n+1} - v_{k,i}^{n+1}] \quad (8)$$

- Compute the policy from the foc $\left(u'(c_{k,i}^n) = \partial_a v_{k,i}^n \right)$ for the backward AND forward derivative of the value function.
- Define $s_{k,i}^{B,n} = ra_i + z_k - c_{k,i}^{B,n}$, $s_{k,i}^{F,n} = ra_i + z_k - c_{k,i}^{F,n}$. Set

$$c_{k,i}^n = \mathbb{1} \left\{ s_{k,i}^{B,n} \leq 0 \right\} \times c_{k,i}^{B,n} + \mathbb{1} \left\{ s_{k,i}^{F,n} \geq 0 \right\} \times c_{k,i}^{F,n} + \mathbb{1} \left\{ s_{k,i}^{F,n} \leq 0 \leq s_{k,i}^{B,n} \right\} \times (ra_i + z_k)$$

- Collecting terms with the same subscripts on the right-hand side

$$\frac{v_{k,i}^{n+1} - v_{k,i}^n}{\Delta} + \rho v_{k,i}^{n+1} = u(c_{k,i}^n) + \alpha_{k,i} v_{k,i-1}^{n+1} + \beta_{k,i} v_{k,i+1}^{n+1} - \left(\alpha_{k,i} + \beta_{k,i} + \lambda_k \right) v_{k,i}^{n+1} + \lambda_i v_{-k,i}^{n+1} \quad (9)$$

where

$$\alpha_{k,i}^{up} = -\frac{[s_{k,i}^{B,n}]^-}{a_i - a_{i-1}} \geq 0, \quad \beta_{k,i}^{up} = \frac{[s_{k,i}^{F,n}]^+}{a_{i+1} - a_i} \geq 0$$

- Equation (9) is just a system of linear equations!!

Implicit Timestepping

Equation (9) can be written in matrix notation as

$$\frac{1}{\Delta}(v^{n+1} - v^n) + \rho v^{n+1} = u(c^n) + \mathbf{A}^n v^{n+1}$$

where the sparse matrix \mathbf{A} looks like

$$\mathbf{A}^n = \begin{bmatrix} \gamma_{1,1} & \beta_{1,1} & 0 & 0 & \dots & \lambda_1 & 0 & \dots & & 0 \\ \alpha_{1,2} & \gamma_{1,2} & \beta_{1,2} & 0 & \dots & 0 & \lambda_1 & 0 & \dots & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & \dots & \alpha_{1,l} & \gamma_{1,l} & 0 & 0 & \dots & \dots & 0 & \lambda_1 \\ \lambda_2 & 0 & \dots & \dots & \dots & \gamma_{2,1} & \beta_{2,1} & 0 & \dots & \dots & 0 \\ 0 & \lambda_2 & 0 & \dots & \dots & \alpha_{2,2} & \gamma_{2,2} & \beta_{2,2} & 0 & \dots & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & \dots & \lambda_2 & 0 & 0 & \dots & \dots & \alpha_{2,l} & \gamma_{2,l} \end{bmatrix}_{l+2 \times l+2}$$

entries of row i

$$\left[\underbrace{\alpha_{k,i}}_{\text{inflow } i-1} \quad \underbrace{-(\alpha_{k,i} + \beta_{k,i} + \lambda_k)}_{\text{outflow}} \quad \underbrace{\beta_{k,i}}_{\text{inflow } i+1} \right] \begin{bmatrix} v_{k,i-1} \\ v_{k,i} \\ v_{k,i+1} \end{bmatrix}$$

Kolmogorov Forward equation

Consider the KFE

$$0 = -\frac{d}{da} \left[s(a, z_k) g(a, z_k) \right] - \lambda_k g(a, z_k) + \lambda_{-k} g(a, z_{-k})$$

with the following discretization

$$0 = -\frac{(s_{k,i}^F)^+ g_{k,i} - (s_{k,i-1}^F)^+ g_{k,i-1}}{\Delta a} - \frac{(s_{k,i+1}^B)^- g_{k,i+1} - (s_{k,i}^B)^- g_{k,i}}{\Delta a} - \lambda_k g_{k,i} + \lambda_{-k} g_{-k,i} \quad (10)$$

Collecting terms with the same subscripts on the right-hand side

$$0 = \underbrace{\frac{(s_{k,i-1}^F)}{\Delta a}}_{\beta_{k,i-1}} g_{k,i-1} + \underbrace{\left(\frac{(s_{k,i}^B)}{\Delta a} - \frac{(s_{k,i}^F)}{\Delta a} - \lambda_k \right)}_{\gamma_{k,i}} g_{k,i} + \underbrace{\left(-\frac{(s_{k,i+1}^B)}{\Delta a} \right)}_{\alpha_{k,i+1}} g_{k,i+1} + \lambda_{-k} g_{-k,i}$$

which in matrix notation reads $\mathbf{A}^T \mathbf{g} = \mathbf{0}$.

This makes sense: the operation is exactly the same as that used for finding the stationary distribution of a discrete Poisson process (continuous-time Markov chain). The matrix \mathbf{A} captures the evolution of the stochastic process and to find the stationary distribution, one solves the eigenvalue problem $\mathbf{A}^T \mathbf{g} = \mathbf{0}$.

Here is some content on the slide

$$f(x) = ax^2 + bx + c$$

Some more content

