

Lecture 6

2/2/26
JRA

Resolution

(Ch. 4 in text)

There were two big pioneers in the field of diffraction-limited resolution

Lord Rayleigh - astronomer, studied self-luminous objects
Abbe - microscopist, discusses later

Microscopes have lots of apertures, including lens openings.

Apertures diffract light, causing it to spread.

Ray optics - just trace rays, which land at image point, no spreading

Wave optics - light spreads, degrading images

Objects that are too close together, or viewed w/ a system having apertures that are too small, can't be distinguished due to diffraction

Will see that wavelength also affects amount of diffraction, i.e. thus resolution, w/ longer wavelengths (red) worse than shorter wavelengths (blue)

Why not just use blue light in microscopy?

- Sensors more sensitive to red (& even more sensitive to infrared)
- Blue fluorophores bleach faster
- Multicolor imaging needs several colors (typically red, green, blue)
- Don't normally use UV because it kills cells

Key implications of diffraction

(1) Resolution improves as $\begin{cases} D \uparrow & \text{Aperture diameter increases} \\ \lambda \downarrow & \text{Wavelength decreases} \end{cases}$

(2) Point objects are imaged as a diffraction pattern

Small objects are distorted in size & appear too large

Limit on resolution $\approx 200 \text{ nm} = 0.2 \mu\text{m}$
(w/ blue light)


Objects that are smaller than about $0.2 \mu\text{m}$ in radius appear enlarged to $0.2 \mu\text{m}$ in radius

Example: synaptic vesicles have a radius of $0.05 \mu\text{m}$ but appear to be at least $0.2 \mu\text{m}$ in light microscope

(3) Objects more closely spaced than $\sim 0.2\mu\text{m}$ cannot be distinguished as two objects but instead blur into one object

DEMO

Diffraction




Results:

- No slit $\rightarrow$ spot
- Wide slit $\rightarrow$ spot
-  Narrow slit $\rightarrow$ elongated diffraction pattern

Rayleigh Theory

Hard optics, focus on results

Quantifies angular resolution



How close in angle can two objects get & still be resolved?

$$\tan \Theta_{\min} = \frac{Y}{L} = \boxed{\frac{1.22 \lambda_0}{D}} \approx \Theta_{\min}$$

wavelength in vacuum (or air)
 Minimum resolvable angular separation (in radians)
 small angle approximation

This is what happens w/ eyes


Want $\Theta_{\min}$ to be as small as possible

Key takeaways from equation

$\lambda \downarrow$	$\Theta_{\min} \downarrow$
$D \uparrow$	$\Theta_{\min} \downarrow$

Two other things are important


- What is the minimum resolvable spatial separation $y_{\min}$ between two objects?
- What is the maximum distance $L_{\max}$ from sensor (e.g., eye) two objects can be & still be resolvable?



- Solve for $y_{\min}$ if L is fixed & known
- Solve for $L_{\max}$ if y is fixed & known

ACTIVITY 2

Diffraction-Limited Resolution



- Calculate θ_{min} from λ & D

$$\theta_{\text{min}} = \frac{1.22\lambda}{D} = \frac{1.22 \cdot 550 \cdot 10^{-9} \text{ m}}{2.4 \text{ m}} = 2.8 \cdot 10^{-7} \text{ rad}$$

- Calculate L_{max} from θ_{min} & y

$$\theta_{\text{min}} = \frac{y}{L_{\text{max}}} \Rightarrow L_{\text{max}} = \frac{y}{\theta_{\text{min}}} = \frac{2.5 \cdot 10^3 \text{ m}}{2.8 \cdot 10^{-7} \text{ rad}} = \boxed{8.9 \cdot 10^9 \text{ m}}$$


$= 5.56 \cdot 10^6 \text{ miles}$

The Hubble aperture is large $\left\{ \begin{array}{l} \text{Increased resolution} \\ \text{Increased light detection} \end{array} \right.$

Hubble distinguishes two objects about $1\frac{1}{2}$ miles apart from about $5\frac{1}{2}$ million miles away

Focus to this point on "aperture"

Abbe, who worked for Zeiss, First to focus on "numerical aperture" (NA)



$$NA = n \sin \Theta_{\max}$$

n = refractive index of medium
between specimen & objective

$\Theta_{\max}$ = half-angle of cone of collected light

The medium between the specimen & objective is typically one of the following:

↑ Increasing n	Air	$n = 1$
	Water	$n = 1.33$
↓	Oil	$n = 1.52$

A large NA is good in μ scopy { More light collected
Better resolution
(next lecture)


NA increases as n increases & $\Theta_{\max}$ increases

$$\begin{array}{l} n \uparrow \quad NA \uparrow \\ \Theta_{\max} \uparrow \quad NA \uparrow \end{array}$$

SIMULATION

Mol Expressions

Immersion Oil &
Refractive Index



n_{oil} generally matches n_{glass} & so maximizes light collection

For a microscope, the minimum resolvable separation, $d_{\min}$, is given by

$$d_{\min} = \frac{0.61 \lambda_0}{NA}$$

Want $d_{\min}$ small

$$d_{\min} \downarrow \begin{cases} NA \uparrow & \text{Want large NA} \\ \lambda_0 \downarrow & \text{Want short wavelength} \end{cases}$$

This equation can be used to derive a useful rule of thumb about microscope resolution

$$\text{Assume } \begin{cases} NA \approx 1 & (\text{NAs can get up to } \sim 1.5) \\ 0.61 \approx 0.5 \\ \lambda = 400 \text{ nm} & (\text{use lowest } \lambda \text{ visible light}) \end{cases}$$

$$d_{\min} = \frac{0.5 \cdot 400 \text{ nm}}{1} \approx 200 \text{ nm}$$

Abbe theory will confirm results
Start next lecture

Name _____


Physics 390 In-Class Activity 2

Diffraction-limited Resolution – Rayleigh's Angular Version

As discussed in class, Lord Rayleigh suggested that Airy diffraction patterns from two objects be regarded as barely resolvable if the angular separation of the objects is at least as big as $\theta_{\min}$, given by

$$\tan\theta_{\min} = \frac{y}{L} = \frac{1.22\lambda}{D} \sim \theta_{\min}$$

Here D is the diameter of the diffracting aperture, y is the linear separation of the objects, and L is the distance of the objects from the aperture (see image below).



The asteroid Ida is orbited by the moon Dactyl. If the separation between Ida and Dactyl is 2.5 km, what is the maximum distance at which they can be resolved by the Hubble Space telescope (diameter = 2.4 m) using 550 nm light?