

Figure accessibility and alt text

The aim of alt text is to provide concise and informative descriptions of your figure so that all readers have access to the same level of information and understanding, and that all can engage with and benefit from the visual elements integral to scholarly content.

Including alt text demonstrates a commitment to accessibility and enhances the overall impact and reach of your work. Good alt text ensures that individuals with visual impairments or those using screen readers can comprehend the content and context of your figures.




ACCESSIBLE

ALT TEXT: Graphs and data on *trem2* expression in brain tumours, with subfigures labelled from a to f, illustrating statistical analyses and gene expression profiles.

Figure 1. Brain tumors exhibit significantly higher levels of *TREM2* expression, predominantly in tumor-associated myeloid cells. (A) Top 6 types of tumors with elevated *TREM2* expression in a descending order from left to right, which were glioblastoma (GBM), brain lower grade glioma (LGG), kidney renal clear cell carcinoma (KIRC), kidney renal papillary cell carcinoma (KIRP), pancreatic adenocarcinoma (PAAD), and breast cancer (BRCA). (B) UMAP plots displaying the immune cells in patients with newly diagnosed GBM. (C, D) *TREM2* transcription in different immune cell populations. (E) *TREM2* is not detected in brain tumor cells. The data were shown as mean $\pm$ SEM and analyzed by 1-way ANOVA. (F) The correlation between *TREM2* expression and signature genes related to myeloid cell phagocytosis, immunosuppression, and antigen presentation was evaluated in LGG and GBM patients. The Spearman's correlation test produces both *P*-values and correlation coefficients (R^2). All listed genes had a *P*-value of less than 0.05. Genes with an R^2 value greater than 0.25 were considered to have a correlation (either positive or negative) with *TREM2* expression.

Alt Text Example 1




SOURCE: Zheng, J., Wang, L., Zhao, S., Zhang, W., Chang, Y., Bosco, D. B., Huang, T., Dheer, A., Gao, S., Xu, S., Ayasoufi, K., Al-Kharboosh, R., Qi, F., Xie, M., Johnson, A. J., Dong, H., Quiñones-Hinojosa, A., & Wu, L.-J. (2023). Trem2 mediates MHCII-associated CD4+ T-cell response against gliomas. *Neuro-Oncology*. <https://doi.org/10.1093/neuonc/noad214>

Good alt text should be objective and:

- **CONCISE:** Given the time-consuming nature of screen reader use, strive for alt text under 100 words, and preferably around 25 to 30 words.
- **CONSISTENT:** Maintain language consistency with the main body of text.
- **UNIQUE:** Avoid repeating information found in captions or surrounding text. For images entirely explained by captions or text, consider labeling them as decorative indicating that their purpose is primarily ornamental rather than conveying additional information.
- **CLEAR:** Spell out contractions, abbreviations, numbers, and non-Latin characters. Present information in a logical and consistent sequence.
- **RELEVANT:** Align descriptions with the image's context, intent, and the focus of the text, and title. Different purposes within a work may warrant distinct alt text descriptions.
- **SIMPLE:** Because screen readers don't interpret formatting, refrain from using formatting (e.g., bullet points) in alt text.
- **INCLUSIVE:** Ensure your text doesn't contain additional information that a sighted person would miss.
- **STANDALONE:** Screen readers indicate that alt text replaces an image, so avoid phrases like "Image of..."
- **COMPLETE:** Conclude alt text with a full stop/period, allowing for a pause before the screen reader continues.

Alt Text Example 2



ACCESSIBLE

ALT TEXT: Graphical representation of three phases—completely confined, partially deconfined, and completely deconfined—depicting Polyakov line phases with corresponding transition profiles.

SOURCE: Hanada, M., & Watanabe, H. (2024). On thermal transition in QCD. *Progress of Theoretical and Experimental Physics*, 2024(4). <https://doi.org/10.1093/ptep/ptae033>

Additional considerations:

Alt Text Example 3

- **Grouped images and multi-panel figures**

When images are grouped, provide alt text for individual figures rather than a collective description.

- **Punctuation**


Screen readers are unable to interpret all types of punctuation, so only use the following: periods, commas, semi-colons, colons, parentheses, brackets, quotation mark, and sashes.

- **Art works**

Although starting with a generic “picture of” or “image of” is redundant, it can be beneficial to specify a particular type of image in art works (sculpture, painting, drawing, etc.).

- **Graphs**

Summarize the graph by emphasizing key details, trends, and relationships in the data. If significant, explicitly list underlying values in a logical order. Avoid redundancy if key values have already been discussed in the text. Use full units like “kilometers” for accurate pronunciation by screen readers.



ACCESSIBLE

ALT TEXT: Graphical abstract showing study results where therapeutic inertia in blood pressure management differs by age group in older adults, with no significant association after age 75.

SOURCE: Kristen John, Jan Stenum, Cheng-Chuan Chiang, Margaret A French, Christopher Kim, John Manor, Matthew A Statton, Kendra M Cherry-Allen, Ryan T Roemmich, Accuracy of Video-Based Gait Analysis Using Pose Estimation During Treadmill Walking Versus Overground Walking in Persons After Stroke, Physical Therapy, Volume 104, Issue 2, February 2024, pzad121, <https://doi.org/10.1093/ptj/pzad121>

To Submit Alt Text

To submit alt text with your article, include it alongside figure captions as in the example below, related to the image in Example 4.

Figure 1 caption: L'Exposition Manet au Musée de l'Orangerie in 'Une Rétrospective d'Édouard Manet a Été Inaugurée Hier au Musée de l'Orangerie', L'Excelsior, 19 June 1932. (Photo: Bibliothèque Nationale de France.).

Alt text: Decorative photograph of a vintage French journal cover featuring an article from 1932.

Alt Text Example 4



SOURCE: Published in Michela Passini, A Microhistory of Heritage Creation Processes: The Impressionists Exhibited at the Orangerie (1930-1937), Oxford Art Journal, Volume 46, Issue 2, August 2023, Pages 241-259, <https://doi.org/10.1093/oxartj/kcad023>




ACCESSIBLE

ALT TEXT: Graphs comparing biomechanical data from treadmill vs. overground walking, showing steps, device use, and stride length correlations, with detailed annotations and statistical values.


Alt Text Example 5

(A) Relationships between video-based and motion capture measurements of step length (left) and step time (right) for all steps recorded. The parameters that correspond to the paretic leg are shown in blue, and those that correspond to the nonparetic leg are shown in red. The diamond shape indicates that the participant used an assistive device, and the circle indicates that the participant did not. (B) Relationships between video-based and motion capture measurements of step length (left) and step time (right) for participant-level data, where all steps for each participant were averaged into a single value. Color and shape conventions are the same as those for panel A. (C) Raw step length error (left) and raw step time error (right) for the analyses resulting from sagittal treadmill, sagittal overground, and frontal overground video measurements relative to motion capture measurements. (D) Absolute step length error (left) and absolute step time error (right) for the analyses resulting from sagittal treadmill, sagittal overground, and frontal overground video measurements relative to motion capture measurements. $P < .05$. OG = overground.

SOURCE: Kristen John, Jan Stenum, Cheng-Chuan Chiang, Margaret A French, Christopher Kim, John Manor, Matthew A Statton, Kendra M Cherry-Allen, Ryan T Roemmich, Accuracy of Video-Based Gait Analysis Using Pose Estimation During Treadmill Walking Versus Overground Walking in Persons After Stroke, *Physical Therapy*, Volume 104, Issue 2, February 2024, pzad121, <https://doi.org/10.1093/ptj/pzad121>



Alt Text Example 6

[Open in new tab](#)[Download slide](#)

Adjusted probability of therapeutic inertia by age group and by number of comorbidities during 22,665 primary care visits with uncontrolled BP defined as SBP ≥ 140 mm Hg and/or DBP ≥ 90 mm Hg among 7,415 patients with a hypertension diagnosis. Probabilities were calculated using marginal effects after using generalized linear mixed-effects models with a binomial distribution stratified by age group (65–74 and ≥ 75 years) and adjusted for self-reported race and ethnicity, body mass index, smoking status, clinic site, average time between visits, insurance status, marital status, classes of antihypertensive medications, hypertension severity at the clinic visit, and patient random effects. Abbreviations: BP, blood pressure; DBP, diastolic blood pressure; SBP, systolic blood pressure.




ACCESSIBLE

ALT TEXT: Chart depicting the percentage of therapeutic inertia in hypertension by age and number of comorbidities, showing higher inertia in older age groups and with more comorbidities.

SOURCE: Grant T Hiura, Talar W Markossian, Beatrice D Probst, Mohammad Samie Tootooni, Gregory Wozniak, Michael Rakotz, Holly J Kramer, Age and Comorbidities Are Associated With Therapeutic Inertia Among Older Adults With Uncontrolled Blood Pressure, *American Journal of Hypertension*, Volume 37, Issue 4, April 2024, Pages 280–289, <https://doi.org/10.1093/ajh/hpad108>

Alt Text Example 7




ACCESSIBLE

ALT TEXT: An intricately decorated antique chest with floral and animal motifs, featuring rich colors and a curved lid.

SOURCE: Catalina Ospina, Indigenous Knowledge and Women's Cosmetics: Mopa Mopa in the Colonial Northern Andes, *Art History*, Volume 46, Issue 5, November 2023, Pages 918–945, <https://doi.org/10.1111/1467-8365.12756>

Alt Text Example 8




ACCESSIBLE

ALT TEXT: Educational diagram showing a function $f(x)$ graph with a tangent line to help find inflection points, accompanied by an interactive calculation tool on left.

SOURCE: Anatoli Kouropatov, Regina Ovodenko, An explorative digital tool as a pathway to meaning: the case of the inflection point, *Teaching Mathematics and its Applications: An International Journal of the IMA*, Volume 41, Issue 2, June 2022, Pages 142–166, <https://doi.org/10.1093/teamat/hrac007>

Alt Text Example 9




ACCESSIBLE

ALT TEXT: Bar graph showing concentrations of 11-deoxycortisol and cortisol in follicular fluid at different time points, with significance markers.

SOURCE: Hanada, M., & Watanabe, H. (2024). On thermal transition in QCD. Progress of Theoretical and Experimental Physics, 2024(4). <https://doi.org/10.1093/ptep/ptae033>


Alt Text Example 10



ALT TEXT: Graph depicting the relationship between fls coverage levels and number of romosozumab prescriptions in 2022, illustrated with box plots and individual data points.

SOURCE: Muhammad K Javaid, Tim Jones, Stepehn Hepworth, P078 United Kingdom rates and determinants of romosozumab prescriptions, a need for greater action, Rheumatology, Volume 63, Issue Supplement_1, April 2024, keae163.119, <https://doi.org/10.1093/rheumatology/keae163.119>

Alt Text Example 11



ACCESSIBLE

ALT TEXT: Graphs and microscopic images labelled A to F. A is graph plotting particle count vs. diameter, microscopy image of exosomes, and a western blot analysis for various markers. B is a heatmap showing data across 14 plasma samples with red, green, and black color coding, indicating varying levels of measured variables. C is a Dot plot displaying pathway enrichment differences, with pathways labeled on the y-axis and logp values on the x-axis, using a color gradient to denote significance levels. D is Bar graph comparing levels of exosomal mirnas between pic-m0 and pic-m1 groups, with statistical significance indicators. E is Graph showing roc curves for exosomal mirnas, with lines representing different mirnas annotated with auc values, and a diagonal reference line indicating chance-level performance. F is fluorescence microscopy images showing association of exosomal mir-519c-5p with circulating tumor cells (ctcs), labeled with dapi, cd45, and epcam markers.

Circulating exosomal miR-519e-5p was derived from tumor cells. A, The expression levels of plasma exosomal miR-519e-5p were significantly decreased post operation and significantly higher in papillary thyroid carcinoma (PTC) patients than that in healthy individuals. B, miR-519e-5p was abundantly detected in the primary tumor of PTC patients in fluorescence in situ hybridization test. C, Tissue exosomes were validated by nanoparticle tracking analysis (NTA), transmission electron microscopy (TEM), and Western blot. D, The tissue exosomal miR-519e-5p in PTC tissues was significantly higher than that in the paired paracancerous and normal tissues. E, Exosomal miR-519e-5p expression levels in the culture medium of PTC cells increased with increasing incubation time. F and G, The reduction of fluorescein isothiocyanate-labeled exogenous miR-519e-5p after the secretion of exosomes.

SOURCE: Genpeng Li, Wenjie Chen, Ke Jiang, Jing Huang, Jinjing Zhong, Xiaowei Liu, Tao Wei, Rixiang Gong, Zhihui Li, Jingqiang Zhu, Hubing Shi, Jianyong Lei, Exosome-mediated Delivery of miR-519e-5p Promotes Malignant Tumor Phenotype and CD8+ T-Cell Exhaustion in Metastatic PTC, The Journal of Clinical Endocrinology & Metabolism, 2023;, dgad725, <https://doi.org/10.1210/clinem/dgad725>