

IMPROVED KISSING NUMBERS IN NINETEEN AND TWENTY-ONE DIMENSIONS

ABSTRACT. We prove that the kissing numbers in 19 and 21 dimensions are at least 12270 and 30779. The best previously published lower bounds were 11948, due to Ho, and 29768, due to Cohn and Li, so the gains are 322 and 1011. Both constructions start from the odd-sign kissing configurations of Cohn and Li, delete a small block of their points, and add more new points than were deleted. In 19 dimensions we cut the 20-dimensional configuration by the hyperplane orthogonal to the indicator vector of a tetrad, project 20-dimensional sign vectors into that hyperplane, and give their tails one of two lengths. This gives 12268 points, with a proof by case analysis. We then restore two deleted points by replacing the new points with a table of 112 integer vectors under a group of 16 sign changes that fixes both restored points; this step is checked by computer. In 21 dimensions the new points are weighted sign vectors whose sign patterns meet all eight cosets of a $[21, 9, 6]$ code. They are given by an explicit table of 192 integer vectors and a group of 16 sign changes, and the proof is an exact computation, reduced by that symmetry. We have also formalized both results in Lean 4 with Mathlib.

1. INTRODUCTION

The kissing problem in $\mathbb{R}^n$ asks how many unit spheres can touch a central unit sphere without overlapping [2, Ch. 1, §2.2]. In terms of the points of tangency, it asks for the largest set X of unit vectors in $\mathbb{R}^n$ with $\langle x, y \rangle \leq 1/2$ for all distinct $x, y \in X$. We write τ_n for this maximum, as in [2, Ch. 13]. It is known only for $n \in \{1, 2, 3, 4, 8, 24\}$; see [1, Table 1.1] for references.

For $17 \leq n \leq 23$ the best constructions were Leech’s cross sections of the Leech lattice minimal vectors [7] until Cohn and Li [1] improved dimensions 17 through 21. In 19, 20 and 21 dimensions their idea is short. Take the classical configuration built from a constant-weight code [8], and put an odd number of minus signs on each codeword instead of an even number. Every sign vector of a code orthogonal to the constant-weight code then has inner product at most 6 with the old points, while the kissing condition allows $\frac{1}{2}\sqrt{8n}$, which is larger for $n \geq 19$. So these sign vectors can be added, one per word of a code with minimum distance at least $n/4$. Cohn and Li obtained $\tau_{19} \geq 11692$ in this way, and Ho [6] found a larger code of 1280 words for the same purpose, reaching $\tau_{19} \geq 11948$.

Theorem 1.1. *We have $\tau_{19} \geq 12270$ and $\tau_{21} \geq 30779$.*

Table 1 compares these values with the best published lower bounds and the known upper bounds.

The mechanism is the slack between 6 and $\frac{1}{2}\sqrt{8n}$. Cohn and Li’s added points are flat: each is a sign vector scaled to squared norm 8, and none of them touches an old point. We spend this unused room by changing the shape of the added points.

- In 19 dimensions (Section 3) the added points come from 20 dimensions. Projecting them into a hyperplane leaves a short three-dimensional tail, and we multiply it by 9/11 or 7/5 according to its direction. This lets two added points differ in only four head coordinates when their tails differ, which a code of minimum distance at least 5, such as Ho’s, must avoid. We add 1792 points where Ho adds 1280. The long tails come

TABLE 1. Kissing number bounds in dimensions 19 and 21, as of October 2026 [4]. The previous lower bounds are due to Ho [6] in dimension 19 and to Cohn and Li [1] in dimension 21. The upper bounds are the floors of numerical semidefinite programming values [3] and are not interval-certified.

dimension	previous lower bound	new lower bound	upper bound
19	11948 [6]	12270	24417 [3]
21	29768 [1]	30779	53524 [3]

too close to 192 old points, which we delete, for a net gain of $1792 - 192 - 1280 = 320$. Two of the 192 can be put back. The 1792 points keep their sign patterns but get new, numerically found weights, and a group of 16 sign changes that fixes the two restored points cuts the check to 112 representatives. The net gain becomes $1792 - 190 - 1280 = 322$.

- In 21 dimensions (Section 4) each added point carries its own positive weight on each coordinate. That freedom allows sign patterns from the even cosets of a $[21, 9, 6]$ code as well as from the odd ones, where Cohn and Li use only the four odd cosets. The added points are 192 integer vectors and their images under a group of 16 sign changes. The 192 vectors are half of an earlier table; the other half is replaced by their images under one extra sign change that fixes the octad where deletions happen, which cuts the deletions from 29 to 13. The weights overshoot that octad, and we delete 13 points on it, for a net gain of $3072 - 13 - 2048 = 1011$.

The proofs differ in character. The 12268-point configuration has a case analysis that can be checked by hand, given the finite facts about a code of size 4096 stated in Lemma 3.1. The step to 12270 and the 21-dimensional proof depend on computer calculation in an essential way, unlike the proofs in [1]. Their added points are integer vectors with entries up to $5.8 \cdot 10^7$ in 19 dimensions and between $1.5 \cdot 10^8$ and $2.5 \cdot 10^8$ in absolute value in 21 dimensions. They were obtained by numerical optimization and rounding, not by hand. The theorems are statements about these explicit tables, and their proofs are lists of about 1.3 million and 5.6 million exact integer inequalities. One program checks these in an equivalent form; a second, written independently, checks all 75270315 and 473658031 pairs without the reduction, and we also formalized both results in Lean 4 with Mathlib (Section 5). The formal proof does not verify our case analysis. It verifies the conclusion, that explicit sets of 12270 and 30779 unit vectors with pairwise inner products at most $1/2$ exist.

2. THE ODD-SIGN CONFIGURATIONS

We identify a binary word with its support. For $c \in \mathbb{F}_2^n$ we write $\text{wt}(c)$ for its weight, $(-1)^c \in \{\pm 1\}^n$ for its sign vector, and $\lambda \circ q$ for the coordinatewise product of two vectors.

Let $\mathcal{O}$ be a set of words of length n with weight 8 and pairwise intersections of size at most 4. The *odd configuration* $B(\mathcal{O})$ consists of the $4\binom{n}{2}$ roots $\pm 2e_i \pm 2e_j$ and, for each $o \in \mathcal{O}$, the 128 vectors with entries ± 1 on o , zeros elsewhere and an odd number of minus signs [1, §2]. All have squared norm 8.

Lemma 2.1 (Cohn and Li). *Distinct $b, b' \in B(\mathcal{O})$ satisfy $\langle b, b' \rangle \leq 4$.*

Proof. Two distinct roots have inner product in $\{0, \pm 4, -8\}$. A root has two nonzero coordinates, each contributing at most 2 against an octad vector, so that inner product is at most 4. Two vectors on the same octad have odd sign patterns, so they differ in an even and positive

number $d \geq 2$ of places and have inner product $8 - 2d \leq 4$. Two vectors on different octads share at most four coordinates. $\square$

The next lemma is the reason for odd signs. It is the weighted form of the argument in [1, §2].

Lemma 2.2. *Let $c \in \mathbb{F}_2^n$ meet every word of $\mathcal{O}$ in an even number of coordinates, let $q = (-1)^c$, and let $\lambda \in \mathbb{R}_{>0}^n$. Let b be an octad vector on $o \in \mathcal{O}$. Then q and b disagree in an odd number of coordinates of o , and*

$$\langle \lambda \circ q, b \rangle \leq \sum_{j \in o} \lambda_j - 2 \min_{j \in o} \lambda_j,$$

with equality for some b on o . In particular, for $\lambda = \mathbf{1}$ we have $\langle q, b \rangle = 6$ if q and b disagree in exactly one coordinate and $\langle q, b \rangle \leq 2$ otherwise. For a root $b = \pm 2e_i \pm 2e_j$ the inner product is at most $2(\lambda_i + \lambda_j)$.

Proof. The sign vector q has an even number of minus signs on o , and b has an odd number. So they disagree in an odd number of coordinates of o , in particular in at least one, and the bound follows. Equality holds for the odd pattern that agrees with q except at one coordinate where λ is smallest. For $\lambda = \mathbf{1}$, d disagreements give $\langle q, b \rangle = 8 - 2d$ with d odd. $\square$

Throughout we test compatibility in integers. For nonzero $x, y \in \mathbb{Z}^n$ the angle between them is at least 60° if and only if

$$(1) \quad \langle x, y \rangle \leq 0 \quad \text{or} \quad 4\langle x, y \rangle^2 \leq |x|^2 |y|^2.$$

We call such a pair *compatible*; the same test applies to rational vectors after scaling. If every pair from a finite set of nonzero rays is compatible, then the rays normalized to unit length form a kissing configuration, and distinct rays give distinct points because (1) excludes cosine 1. With $\lambda = \mathbf{1}$, Lemma 2.2 shows that the sign vector of a word that meets every word of $\mathcal{O}$ evenly is compatible with $B(\mathcal{O})$ when $6 \leq \frac{1}{2}\sqrt{8n}$, that is, for $n \geq 18$, with contacts when $n = 18$.

3. DIMENSION 19

3.1. The codes. Coordinates are $1, \dots, 20$. Let $E \subseteq \mathbb{F}_2^{20}$ be the span of the twelve words

$$\begin{aligned} &\{1, 8, 9, 12, 16, 17, 18, 19\}, \{2, 10, 11, 14, 15, 17, 18, 20\}, \{3, 7, 9, 13, 15, 16, 17, 19\}, \\ &\{4, 7, 8, 10, 12, 15, 16, 19\}, \{5, 10, 12, 13, 15, 16, 17, 18\}, \{6, 7, 8, 9, 10, 13, 16, 18\}, \\ &\{1, 4, 7, 9\}, \{1, 5, 6, 18, 20\}, \{1, 3, 12, 15, 20\}, \{1, 10, 13, 19, 20\}, \\ &\{1, 3, 5, 6, 7, 13, 14, 15, 18, 20\}, \{2, 4, 6, 7, 8, 13, 14, 16, 17, 18\}. \end{aligned}$$

On the first 19 coordinates these are Ho's generators $m_1, \dots, m_6, s_1, \dots, s_4, r_1, r_2$ [6, §2]. Our coordinate 20 is Ho's coordinate 24, the last column of his extension matrix P , so we have added it to the words where that column has a 1. Ho shows that the full extension to length 24 is the extended binary Golay code [6, Prop. 2]. So E is the Golay code punctured at Ho's coordinates 20 to 23, and its dual $E^\perp$ is the Golay code shortened at the same four coordinates [9]. The 130 words of weight 8 in $E^\perp$ are the *octads*. They form the shortened Steiner system $S(5, 8, 24)$ with $n = 20$ of [1, Table 2.1], so $B(\text{octads})$ is Cohn and Li's 17400-point configuration C_{20} in $\mathbb{R}^{20}$.

The five octads of the Golay code that contain four given coordinates partition the other twenty coordinates into five tetrads, and these six tetrads form a *sextet* [2, Ch. 10]. Puncturing at the four given coordinates turns those five octads into five words of weight 4 in E .

The following facts are finite computations on the 4096 words of E .

Lemma 3.1. (1) *The weight enumerator of E is*

$$1 + 5z^4 + 64z^5 + 240z^6 + 320z^7 + 250z^8 + 640z^9 + 1056z^{10} \\ + 640z^{11} + 250z^{12} + 320z^{13} + 240z^{14} + 64z^{15} + 5z^{16} + z^{20}.$$

The five words of weight 4 are pairwise disjoint and cover all 20 coordinates.

- (2) *Let $T = \{1, 4, 7, 9\}$, one of the five, and let H be the other 16 coordinates, the head. For $e \in E$ write e_T and e_H for its parts on T and H . The nonzero words with $\text{wt}(e) \leq 7$ have $(\text{wt}(e_T), \text{wt}(e_H))$ equal to $(4, 0)$ for 1 word, $(0, 4)$ for 4, $(1, 4)$ for 64, $(0, 6)$ for 96, $(2, 4)$ for 144, $(1, 6)$ for 256 and $(3, 4)$ for 64.*
- (3) *Two distinct octads meet in at most 4 coordinates. The octads meet T in 0, 2 or 4 coordinates, for 30, 96 and 4 octads respectively.*

The intersections in (3) are even because $T \in E$ and the octads lie in $E^\perp$. By (2), two distinct words $c, c' \in E$ with $d_T = \text{wt}(c_T + c'_T)$ and $d_H = \text{wt}(c_H + c'_H)$ satisfy the following. If $d_T = 0$, then $d_H = 4$ (and $c + c'$ is a tetrad inside H) or $d_H \geq 6$. If $1 \leq d_T \leq 3$, then $d_H \geq 4$. If $d_T = 4$, then $d_H = 0$ (and $c' = c + \mathbf{1}_T$), $d_H = 4$ or $d_H \geq 6$; this is the case $d_T = 0$ applied to c and $c' + \mathbf{1}_T \in E$.

3.2. The configuration. Let

$$R = \begin{pmatrix} 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix},$$

acting on the coordinates $(1, 4, 7, 9)$ of T . The rows of $R/2$ are an orthonormal basis of $\mathbf{1}_T^\perp \cap \mathbb{R}^T$, so $(R/2)^\top(R/2) = I - J/4$. The map $\varphi(v) = (v_H, Rv_T/2) \in \mathbb{R}^{16} \times \mathbb{R}^3$ therefore satisfies

$$(2) \quad \langle \varphi(u), \varphi(v) \rangle = \langle u, v \rangle - \frac{1}{4} \left(\sum_{j \in T} u_j \right) \left(\sum_{j \in T} v_j \right),$$

and it is an isometry from the hyperplane $\mathbf{1}_T^\perp$ onto $\mathbb{R}^{19}$. The configuration in $\mathbb{R}^{19}$ has two parts.

Base. There are 10668 points of C_{20} orthogonal to $\mathbf{1}_T$, as many as in Leech's 19-dimensional configuration. We delete from them the 192 points that lie on the four octads containing T ; these are the points with two plus and two minus signs on T . The remaining 10476 points b enter as $\varphi(b)$. They are the 480 roots inside H , the 12 roots $\pm 2(e_i - e_j)$ with $i, j \in T$, the 3840 odd vectors on the 30 octads disjoint from T , and the 6144 odd vectors on the 96 octads meeting T twice whose two T -entries have opposite signs.

Additions. Let $c \in E$ with $\text{wt}(c_T) \in \{1, 2, 3\}$ and $q = (-1)^c$. The projected tail $t_0 = Rq_T/2$ satisfies $|t_0|^2 = 4 - \frac{1}{4}(\sum_T q_j)^2$ by (2). When $\text{wt}(c_T)$ is odd, $t_0 \in \{\pm 1\}^3$ and $|t_0|^2 = 3$; we call c *diagonal*. When $\text{wt}(c_T) = 2$, $t_0 = \pm 2e_k$ and $|t_0|^2 = 4$; we call c an *axis* word. The added point is

$$a(c) = (q_H, \alpha t_0), \quad \alpha = \begin{cases} 9/11, & c \text{ diagonal,} \\ 7/5, & c \text{ axis,} \end{cases}$$

with $|a(c)|^2 = 2179/121$ or $596/25$. Multiplying by 22 or 10 makes it an integer vector, and the base points $\varphi(b)$ become integer vectors after multiplying by 2.

Which words. Let W be the span of the five tetrads, a code of size 32. The 3584 words with $\text{wt}(c_T) \in \{1, 2, 3\}$ fall into 112 cosets of W . Each coset $c + W$ is a 5-cube: its vertices are the 32 words, and each of the five directions adds one tetrad. We keep one of the two bipartition classes of each cube, 16 words. Lemma 3.2 below holds for either class, so any choice gives a valid configuration of the same size. For definiteness, and to match `data/D19_12268_rays.txt`, we

TABLE 2. The blocks of the 19-dimensional configuration.

block	size	squared norm
roots in H	480	8
roots in T	12	8
octad vectors, octad disjoint from T	3840	8
octad vectors, octad meeting T twice	6144	8
diagonal additions	1024	2179/121
axis additions	768	596/25
total	12268	

keep the words c for which $c + c_{\min}$ is a sum of an even number of tetrads, where $c_{\min}$ is the smallest word of the coset read as the integer $\sum_{j \in c} 2^{j-1}$. This gives 1024 diagonal and 768 axis words, 1792 in all.

The total is $10668 - 192 + 1792 = 12268$. Table 2 lists the blocks.

3.3. Proof. Some restriction inside each cube is needed, because two words of one cube that differ by a tetrad inside H have the same T -pattern and $d_H = 4$. For two diagonal words this gives inner product $8 + 3(9/11)^2 \approx 10.01$ against the bound $\frac{1}{2}|a|^2 \approx 9.00$.

Lemma 3.2. *Two distinct kept words c, c' satisfy $c_H \neq c'_H$, and $d_H \geq 6$ if $c_T = c'_T$.*

Proof. If $c_H = c'_H$, then $c + c'$ is a word supported in T , so it is $\mathbf{1}_T$ by Lemma 3.1(2). Then c and c' lie in the same coset of W and differ by one tetrad, so they lie in opposite classes of the 5-cube and are not both kept. If $c_T = c'_T$ and $d_H = 4$, then $c + c'$ is a tetrad in H , and the same argument applies. $\square$

We now check every pair, using the unnormalized vectors.

Base with base. By (2) the map φ preserves inner products on $\mathbf{1}_T^\perp$, so Lemma 2.1 gives $\langle \varphi(b), \varphi(b') \rangle \leq 4 = \frac{1}{2} \cdot 8$.

Base with additions. Since $\sum_T b_j = 0$, equation (2) gives

$$(3) \quad \langle a(c), \varphi(b) \rangle = \langle q_H, b_H \rangle + \alpha \langle q_T, b_T \rangle = \langle q, b \rangle + (\alpha - 1) \langle q_T, b_T \rangle.$$

The word c lies in E , so it meets every octad evenly, and Lemma 2.2 applies: for an octad vector b , $\langle q, b \rangle = 6$ if q and b disagree in exactly one place of the octad and $\langle q, b \rangle \leq 2$ otherwise. The bound to meet is $\frac{1}{2}|a|\sqrt{8} = \sqrt{2|a|^2}$, which is $\sqrt{4358/121} \approx 6.0014$ for diagonal and $\sqrt{1192/25} \approx 6.9051$ for axis additions.

For the 6144 points on octads meeting T twice, $b_T = \pm(e_i - e_j)$ with i, j the two coordinates of the octad in T , so $\langle q_T, b_T \rangle \in \{-2, 0, 2\}$. If it is 0, the inner product is $\langle q, b \rangle \leq 6$. If it is 2, the inner product is at most $6 + 2(\alpha - 1)$, which is $62/11$ or $34/5$. If it is -2 , then q and b disagree at both coordinates of the octad in T . The number of disagreements is odd, so it is at least 3, $\langle q, b \rangle \leq 2$, and the inner product is at most $2 - 2(\alpha - 1)$. Table 3 collects the maxima.

All maxima are below the bounds, since $36 \cdot 121 = 4356 < 4358$ and $(34/5)^2 = 1156/25 < 1192/25$.

Additions with additions. For kept words $c \neq c'$,

$$\langle a(c), a(c') \rangle = 16 - 2d_H + \alpha \alpha' \langle t_0, t'_0 \rangle, \quad \langle t_0, t'_0 \rangle = \langle q_T, q'_T \rangle - \frac{1}{4} \left(\sum_T q_j \right) \left(\sum_T q'_j \right).$$

TABLE 3. Largest inner product between a base point and an addition in dimension 19.

base points	count	reason	diagonal	axis
roots in H	480	$b_T = 0$	4	4
roots in T	12	$\alpha \langle q_T, b_T \rangle \leq 4\alpha$	36/11	28/5
octads disjoint from T	3840	$b_T = 0$, Lemma 2.2	6	6
octads meeting T twice	6144	three cases above	6	34/5
bound $\sqrt{2 a ^2}$			6.0014	6.9051

TABLE 4. Largest inner products between two additions in dimension 19, for the cases with nonnegative tail product. The bound is $\frac{1}{2}|a||a'|$.

types	case	$\langle a, a' \rangle$	bound	cosine
diagonal, diagonal	$d_T = 0, d_H = 6$	727/121	2179/242	0.3337
	$d_T = 2, \langle t_0, t'_0 \rangle = 1, d_H = 4$	1049/121	2179/242	0.4814
axis, axis	$d_T = 0, d_H = 6$	296/25	298/25	0.4966
	$d_T = 2, d_H = 4$	8	298/25	0.3356
diagonal, axis	$d_T = 1, d_H = 4$	566/55	$\sqrt{324671/3025}$	0.4967

Here $d_H \geq 4$ in every case, and $d_H \geq 6$ when $c_T = c'_T$, by Lemma 3.1(2) and Lemma 3.2. The tail product depends only on the two patterns on T :

- two diagonal words have $d_T \in \{0, 2, 4\}$ and $\langle t_0, t'_0 \rangle = 3, \pm 1, -3$ respectively;
- two axis words have $d_T \in \{0, 2, 4\}$ and $\langle t_0, t'_0 \rangle = 4, 0, -4$ respectively;
- a diagonal and an axis word have $d_T \in \{1, 3\}$ and $\langle t_0, t'_0 \rangle = 2$ or -2 respectively.

The cases with $\langle t_0, t'_0 \rangle < 0$ give at most 8, below every bound. Table 4 lists the others at the smallest allowed d_H .

Every entry is strictly below its bound. For the diagonal–axis row we compare $(566/55)^2 = 320356/3025$ with $324671/3025$. So all pairs are compatible. The roots inside H and inside T already span $\mathbb{R}^{19}$. This proves $\tau_{19} \geq 12268$.

3.4. Remarks. The tail lengths have little room. A diagonal addition against an octad vector on an octad disjoint from T needs $36 \leq 2(16 + 3\alpha^2)$, so $\alpha \geq \sqrt{2/3} \approx 0.8165$ for diagonal words. Two axis additions with the same pattern at $d_H = 6$ need $4 + 4\alpha^2 \leq \frac{1}{2}(16 + 4\alpha^2)$, so $\alpha \leq \sqrt{2} \approx 1.4142$ for axis words. The diagonal–axis pairs at $d_T = 1, d_H = 4$ couple the two factors. Numerically, the conditions behind Tables 3 and 4 allow the diagonal factor in about $[0.8165, 0.877]$ and the axis factor in about $[1.304, 1.4142]$, in a thin region where a larger axis factor permits a larger diagonal factor. We chose 9/11 and 7/5 as simple fractions inside it. With plain projection, both factors equal to 1, the diagonal–axis pairs have inner product 10 against $\frac{1}{2}\sqrt{19 \cdot 20} \approx 9.75$, so plain projection fails.

Axis factors above 1 are what the 192 deleted points cannot tolerate. A deleted point has $b_T \in \{\pm 1\}^4$ with two signs of each kind. An axis word with $q_T = b_T$ whose sign vector disagrees with b in exactly one place of the octad gives inner product $6 + 4 \cdot \frac{2}{5} = 38/5 > 6.9051$. A computer check shows that each deleted point conflicts in this way with some kept axis addition.

We have not determined the largest configuration of this shape. One numerical observation bears on it. The 3584 candidate words have exactly 1792 distinct heads, so both bipartition classes use the same heads. A numerical search over all 2^{16} sign vectors on H , looking for a tail compatible with this 10476-point base by local optimization from 48 starting points, found one only for these 1792 heads. A local search can miss a feasible tail, so this is not a proof.

The additions touch nothing. By direct computation, all 2386584 contacts (pairs with cosine exactly $1/2$) are between base points. The closest pairs involving an addition have cosine $\sqrt{1089/4358} \approx 0.49989$, a diagonal addition against an octad vector on an octad disjoint from T .

3.5. Restoring two deleted points. Two of the 192 deleted points can be put back. This step is due to an AI agent outside this project, which was given the 12268-point configuration of Section 3 and returned a 12270-point configuration in other coordinates. We found a coordinate permutation and sign change that carries its base onto ours, so we state the construction in the coordinates of this section. The new points keep the sign patterns of the 1792 additions, but their weights are new.

Base. Let $h = -e_3 + e_6 + e_8 + e_{19}$ and

$$p_{\pm} = h \pm (e_1 + e_4 - e_7 - e_9).$$

Both are odd vectors on the octad $T \cup \{3, 6, 8, 19\}$ with two plus and two minus signs on T , so they are among the 192 deleted points. Their images are $\varphi(p_{\pm}) = (h_H, \pm 2e_1)$, and $\langle p_+, p_- \rangle = 0$. The base is now the 10476 points of Table 2 together with $\varphi(p_+)$ and $\varphi(p_-)$, 10478 points.

Sign group. For a word $g \in E$ that misses T , let S_g change the signs of the coordinates in g . It acts on $\mathbf{1}_T^{\perp}$ and commutes with φ , and on $\mathbb{R}^{19}$ it changes the signs of head coordinates only. Let $G \subseteq E$ be the span of

$$\{2, 5, 10, 11, 13, 15\}, \quad \{2, 5, 10, 16, 17, 20\}, \quad \{2, 5, 12, 14, 15, 17\}, \quad \{2, 10, 13, 14, 16, 18\}.$$

It has 16 words: 0, twelve words of weight 6, and the three sums of two of the tetrads $\{2, 11, 14, 20\}$, $\{5, 12, 13, 16\}$, $\{10, 15, 17, 18\}$. Every word of G misses $T \cup \{3, 6, 8, 19\}$. The words of E that miss this octad form a group of order 32, and G is one of the four subgroups of index 2 in it that contain none of the three tetrads.

Lemma 3.3. *Every S_g , $g \in G$, maps the 10478-point base onto itself and fixes $\varphi(p_+)$ and $\varphi(p_-)$. On the 48 deleted points of each of the other three octads containing T , the group $\{S_g\}$ has six orbits of size 8.*

Proof. A word $g \in E$ meets every octad evenly, so S_g maps roots to roots and the odd vectors on an octad to odd vectors on the same octad. Since g misses T , S_g preserves $\mathbf{1}_T^{\perp}$. So S_g maps the points of each octad that are orthogonal to $\mathbf{1}_T$ onto themselves, and it maps the 10476 base points of Section 3 onto themselves. It fixes every vector supported on $T \cup \{3, 6, 8, 19\}$, including $p_{\pm}$. On an octad $T \cup u$ with $u \neq \{3, 6, 8, 19\}$, S_g acts through $g \cap u$. The map $g \mapsto g \cap u$ on G has kernel $\{0, u' + u''\}$, where u' and u'' are the other two tetrads, so it takes 8 values. The corresponding sign changes on u are distinct, and no vector with entries ± 1 on u is fixed by a nontrivial one, so every orbit has size 8. $\square$

New points. Let $r_1, \dots, r_{112} \in \mathbb{Z}^{19}$ be the vectors in the **representatives** field of **data/D19_12270_certificate.json** (see Data); Appendix A gives three of them. They have no zero entries, their entries have absolute values at most 58006795, and their squared norms lie between $0.99999998 \cdot 10^{16}$ and $1.00000002 \cdot 10^{16}$. The new points are the $112 \cdot 16 = 1792$ vectors $S_g(r_i)$, $g \in G$.

They are reweighted additions. For each new point x there is a kept word c such that x has the sign of $q = (-1)^c$ on every head coordinate and its tail is closest in angle to the tail $t_0 = Rq_T/2$, and every kept word occurs exactly once. So 64 of the representatives are diagonal and 48 are axis words, and G permutes the kept words. Define the tail factor of a representative as the length of its tail divided by $|t_0|$ and by the root mean square of its head entries. It is between 0.817 and 0.871 for diagonal and between 1.233 and 1.424 for axis words, where Section 3 uses 9/11 and 7/5. Remark (b) below says where the weights change most.

We do not have a closed form for these weights. The outside agent found them by numerical optimization with G imposed: it restored the two points one at a time, minimized a penalty on pairs with cosine above $1/2$ over all 19 coordinates of the representatives, and rounded to integers at squared norm about 10^{16} . The proof does not depend on how they were found.

The total is $10476 + 2 + 1792 = 12270$, that is, $10668 - 190 + 1792$.

Proof. Each S_g is orthogonal and $S_a S_b = S_{a+b}$. With Lemma 3.3, every pair involving a new point is determined by one of these finite lists:

- (1) the $\binom{112}{2} \cdot 16 + 112 \cdot 15 = 101136$ products $\langle r_i, S_g r_j \rangle$ with $i \leq j$, excluding $i = j$, $g = 0$;
- (2) the $112 \cdot 10478 = 1173536$ products $\langle r_i, b \rangle$ with b in the base.

Proposition 3.4. *Let $L = 2 \cdot 10^7$. Every product in lists (1) and (2), with x, y the two vectors, satisfies $\langle x, y \rangle \leq (\frac{1}{2} - \frac{1}{L})|x||y|$.*

Proof. This is a finite computation in exact integers. For $g = \langle x, y \rangle > 0$ the inequality is equivalent to $(2Lg)^2 \leq (L - 2)^2|x|^2|y|^2$. The program `scripts/verify_d19_orbits.py` checks it with Python integers in about 2 seconds. It rebuilds the base from the generators of E , checks Lemma 3.3 and the sign patterns stated above, and then computes both lists. The program `scripts/check_pairs.py` also tests all 75270315 pairs of the configuration directly, in about 7 seconds, and so does the Lean formalization of Section 5. $\square$

The base consists of points of C_{20} orthogonal to $\mathbf{1}_T$, so base pairs are compatible by Lemma 2.1 and (2). So all 12270 points are pairwise compatible, and they span $\mathbb{R}^{19}$ because the base contains all 492 roots. This proves Theorem 1.1 for $n = 19$.

Remarks.

- (a) The restoration spends the margin of the 12268-point configuration, whose largest cosine involving an addition was 0.49989. The smallest value of $\frac{1}{4} - \cos^2$ is now $852356203/888888874369808 \approx 9.59 \cdot 10^{-8}$ over base–new pairs and about $8.95 \cdot 10^{-8}$ over pairs of new points. There are 6736 base–new and 7632 new–new pairs with cosine within 10^{-6} of $1/2$. All 2387420 contacts are still between base points; the 836 more than before involve $p_{\pm}$.
- (b) In Section 3 each deleted point conflicts with axis additions whose sign vector agrees with it on T and disagrees with it in exactly one place of its octad (inner product $38/5$). For $p_{\pm}$ these are the 128 axis points with tail along $\pm e_1$ whose head differs from h in exactly one of the coordinates 3, 6, 8, 19. In the new table exactly these have a shorter tail, with factor about 1.233 instead of 1.40 to 1.42, and entries on 3, 6, 8, 19 about 0.82 times their other head entries. Their cosine with p_+ or p_- is just below $1/2$. The head entries of every other representative are within 5% of each other.
- (c) Each of the 190 points still deleted has cosine between 0.5489 and 0.5563 with some new point. Since G fixes the fiber on $T \cup \{3, 6, 8, 19\}$ pointwise, a point restored there costs no orbit of new points, and the other 46 points of that fiber are the natural candidates. With G kept, the outside agent tried to add 12 of them, one at a time, and one orbit of 8 on another fiber, and found no valid table in any of these cases. This is

a numerical failure, not a proof. A point on one of the other three fibers comes with its orbit of 8.

- (d) A group that fixes two of the four fibers pointwise must be much smaller. The octads span $E^\perp$, so a coordinate sign change that maps C_{20} to itself is S_g for some $g \in E$. The words of E that miss T and two further tetrads of the sextet are only 0, the remaining two tetrads and their sum. So such a group has order at most 4, and the table would need at least 448 representatives.

These facts were computed numerically, except (d), which is a finite computation on E .

4. DIMENSION 21

4.1. The codes. Coordinates are $1, \dots, 24$, and the construction lives on the first 21. Let G be the extended binary Golay code: the cyclic code of length 23 generated by $g(t) = 1 + t + t^5 + t^6 + t^7 + t^9 + t^{11}$, with the coefficient of t^k at coordinate $k + 1$, extended by an overall parity bit at coordinate 24 [2, Ch. 3, §2.8]. Let $\mathcal{O}$ be the 210 octads of G that vanish on coordinates 22, 23 and 24. Then $B = B(\mathcal{O})$ is Cohn and Li's 27720-point configuration in $\mathbb{R}^{21}$.

Let D be G punctured to the first 21 coordinates, a $[21, 12, 5]$ code. Because G is self-dual and the octads in $\mathcal{O}$ vanish on the punctured coordinates, every word of D meets every octad of $\mathcal{O}$ evenly. We call the 21 words of weight 5 in D *lines*. Two sign vectors of words of D at distance d have cosine $1 - 2d/21$, which exceeds $1/2$ only for $d = 5$, where it is $11/21$. Cohn and Li use the 2048 odd-weight words of D . The sum of two odd words is even, so no two of them are at distance 5.

Let $C \subseteq D$ be the span of the nine words (bit j of the binary expansion is coordinate $j + 1$)

1559917, 15163, 774651, 1540761, 485602, 408336, 678478, 1748752, 1231514.

It is a $[21, 9, 6]$ code with only even weights, and D/C has eight cosets. Four are even, with representatives 0, 1393, 2139, 3370, and contain no lines. Four are odd, with representatives 696, 1993, 2787, 3986, and contain 6, 3, 6 and 6 lines. These are finite computations on the 4096 words of D . The code C first appeared in an unpublished 30280-point certificate produced by an AI agent outside this project.

The following lemma is not used in the proof. It explains why the new points must be weighted.

Lemma 4.1. *Let c, c' be distinct words of D . If they lie in the same coset of C , or in two cosets of the same parity, then $\text{wt}(c + c') \geq 6$. If they lie in cosets of opposite parity, then $c + c'$ ranges over a whole odd coset as c and c' range over their cosets, so $\text{wt}(c + c') = 5$ occurs.*

Proof. In the first case $c + c'$ lies in C or in an even coset, and the even words of D have weight at least 6. In the second case $c + c'$ lies in an odd coset, and every odd coset contains a line. $\square$

So flat sign vectors from an even coset and an odd coset conflict at cosine $11/21$.

4.2. Weighted sign vectors. Here is the simplest way around the conflict. Let ℓ be a line, and put $\lambda_j = \sqrt{2/5}$ for $j \in \ell$ and $\lambda_j = 1$ otherwise, so $|\lambda|^2 = 18$. For $c \in D$ the point $\lambda \circ (-1)^c$ has, by Lemma 2.2, inner product at most $\sum_{j \in o} \lambda_j - 2 \min_{j \in o} \lambda_j$ with the octad vectors on o . An octad meets ℓ in 0, 2 or 4 coordinates, because $\ell \in D$, and the bound is then $8 - 2 = 6$, $6 + 2\sqrt{2/5} - 2\sqrt{2/5} = 6$, or $4 + 2\sqrt{2/5} < 6$. This is at most $\frac{1}{2}\sqrt{8}|\lambda| = 6$. So among weight vectors equal to 1 off a line and constant on it, $\sqrt{2/5}$ is the smallest weight on the line that the base allows. A small weight on ℓ weakens the conflicts at distance 5 that run through ℓ .

In general, let $c \in D$, $q = (-1)^c$ and $\lambda \in \mathbb{R}_{>0}^{21}$. Lemma 2.2 shows that $\lambda \circ q$ is compatible with every point of B exactly when

$$(4) \quad \sum_{j \in o} \lambda_j - 2 \min_{j \in o} \lambda_j \leq \sqrt{2} |\lambda| \quad \text{for all } o \in \mathcal{O}, \quad \lambda_i + \lambda_j \leq \frac{1}{\sqrt{2}} |\lambda| \quad \text{for all } i \neq j.$$

The hypothesis $c \in D$ is what makes the first condition exact. Two points $\lambda \circ q$ and $\mu \circ q'$ have inner product $\sum_j \lambda_j \mu_j (-1)^{(c+c')_j}$, which depends only on the weights and on $c + c'$.

4.3. The configuration. *Base.* Let $Q = \{3, 6, 10, 11, 16, 18, 20, 21\} \in \mathcal{O}$. Delete from B the 13 odd sign patterns on Q whose signs, in the coordinate order of Q , are

$$\begin{aligned} &+---+----+, +---+---+-, +-+---+---, +-+---++++, +-+++++---, \\ &+-+++++---, +++-+---+-, +++-++++-, +-+++++---, +-+++++---, \\ &+++++---, ++++++---, ++++++---. \end{aligned}$$

The certificate file lists them in this order as `removed_base_representatives`. We know no closed rule for them; they are exactly the odd patterns on Q that have cosine above $1/2$ with some new point (Section 4.5). The rest, B' , has $840 + 209 \cdot 128 + 115 = 27707$ points.

Sign group. Let $\Sigma \subseteq D$ be the span of 92361, 20689, 346115 and $w = 14723$, with supports

$$\{1, 4, 7, 8, 12, 14, 15, 17\}, \quad \{1, 5, 7, 8, 13, 15\}, \quad \{1, 2, 12, 15, 17, 19\}, \quad \{1, 2, 8, 9, 12, 13, 14\}.$$

For $h \in \Sigma$ let S_h change the signs of the coordinates in h . The group $\{S_h\}$ has order 16. The first three generators span a subgroup $\Sigma_0 \subseteq C$ of order 8; the word w has odd weight, so $w \notin C$, but $w \in D$. Section 4.5 explains where w comes from. All four generators miss Q , so every S_h fixes every vector supported on Q , including the 13 deleted points.

New points. Let $r_1, \dots, r_{192} \in \mathbb{Z}^{21}$ be the first 192 vectors in the `representatives` field of `data/D21_30779_certificate.json` (see Data). (That field has 384 entries; the last 192 are $S_w r_1, \dots, S_w r_{192}$, so that the file also describes the configuration as 384 representatives under Σ_0 . Both readings give the same set of points.) They have no zero entries, their entries have absolute values between 149071270 and 249507763, and their squared norms lie between $0.999999998 \cdot 10^{18}$ and $1.000000002 \cdot 10^{18}$. Appendix A gives one of them for each coset below. The new points are the $192 \cdot 16 = 3072$ vectors $S_h(r_i)$.

The sign words of $r_1, \dots, r_{192}$ and their Σ_0 -images (the words c with $\text{sign}(r) = (-1)^c$), 1536 words in all, fill half of each of the six cosets $c_0 + C$ with

$$c_0 \in \{696, 1993, 2787\} \text{ (odd)}, \quad c_0 \in \{1393, 2139, 3370\} \text{ (even)},$$

with 32 representatives and 256 words on each. Adding w swaps the cosets in the pairs $0 \leftrightarrow 696$, $1393 \leftrightarrow 1993$, $2139 \leftrightarrow 2787$ and $3370 \leftrightarrow 3986$. So the 3072 sign words of all new points meet every coset of C : 512 words on each of 1393, 1993, 2139, 2787 and 256 on each of 0, 696, 3370, 3986. Cohn and Li's four odd cosets are all present, together with all four even ones. On the cosets of 1993 and 2139 the weights of r_i follow the line model, with the lines $\{8, 15, 16, 18, 19\}$ and $\{1, 3, 5, 17, 18\}$: after scaling each representative to $|\lambda|^2 = 18$, every entry is within 10^{-6} of the model. On the coset of 3370 they follow it only approximately, with the line $\{4, 5, 6, 8, 10\}$ and every entry within $7.6 \cdot 10^{-3}$. The cosets of 2139 and 3370 are even and contain no line. Their lines lie in the odd cosets of 1993 and 696, and adding the line to a word of either even coset gives a word in the coset of 3986, at distance 5. So the reduced weights weaken exactly the conflicts with the coset of 3986. On the other three cosets the weights are irregular, and each representative's entries lie between 0.82 and 1.15 times their root mean square. The weights also vary slightly between the 32 orbits of Σ_0 inside a coset.

We do not have a closed form for the irregular weights. They were found by numerical optimization of the conditions of the previous subsection, with a target cosine slightly below $1/2$, and then rounded to integers at scale about $2 \cdot 10^8$. The proof does not depend on how they were found. Replacing the three line-model cosets by their exact limits would not shorten the proof, because the irregular cosets would still need a computer check.

The total is $27707 + 3072 = 30779$.

4.4. Proof.

Lemma 4.2. *Every S_h , $h \in \Sigma$, maps B' onto itself.*

Proof. Every word of Σ lies in D , so it meets every octad of $\mathcal{O}$ in an even number of coordinates. (Membership of the four generators in D is a finite check: the first three lie in $C \subseteq D$, and w lies in D but not in C .) A sign change S_h maps roots to roots. On an octad vector with support o it changes an even number of signs, so the number of minus signs on o stays odd. So $S_h(B) = B$. Since S_h fixes the 13 deleted points, $S_h(B') = B'$. $\square$

Each S_h is orthogonal and $S_a S_b = S_{a+b}$. Hence

$$\langle S_a r_i, S_b r_j \rangle = \langle r_i, S_{a+b} r_j \rangle, \quad \langle S_a r_i, b \rangle = \langle r_i, S_a b \rangle.$$

With Lemma 4.2, every pair involving a new point is determined by one of these finite lists:

- (1) the $\binom{192}{2} \cdot 16 + 192 \cdot 15 = 296256$ products $\langle r_i, S_h r_j \rangle$ with $i \leq j$, excluding $i = j$, $h = 0$;
- (2) the $192 \cdot 27707 = 5319744$ products $\langle r_i, b \rangle$ with $b \in B'$.

Proposition 4.3. *Let $L = 2 \cdot 10^8$. Every product in lists (1) and (2), with x, y the two vectors, satisfies $\langle x, y \rangle \leq (\frac{1}{2} - \frac{1}{L})|x||y|$. In particular every such pair satisfies (1).*

Proof. This is a finite computation in exact integers. For $g = \langle x, y \rangle > 0$ the inequality is equivalent to $(2Lg)^2 \leq (L-2)^2|x|^2|y|^2$. For list (2), Lemma 2.2 gives the maximum of $\langle r_i, b \rangle$ over the full fiber of odd vectors on each octad other than Q and over the roots in closed form, and the 115 retained points on Q are checked one by one. The program `scripts/verify_d21_orbits.py` checks this with Python integers in about 2 seconds. It first checks that representatives 193 to 384 of the certificate file are $S_w r_1, \dots, S_w r_{192}$, and then runs the equivalent computation for the 384 representatives under Σ_0 : 590976 orbit products and the products with B' . A second program, `scripts/check_pairs.py`, written independently, does not use the reduction and tests all 473658031 pairs of the configuration directly, in about one minute. The Lean formalization of Section 5 also checks all pairs directly. $\square$

Base pairs are compatible by Lemma 2.1. So all 30779 points are pairwise compatible, and they span $\mathbb{R}^{21}$ because B' contains all 840 roots. This proves Theorem 1.1 for $n = 21$.

4.5. Remarks. The smallest value of $\frac{1}{4} - \cos^2$ over pairs involving a new point is

$$\frac{105355791297}{799999998594867688} \approx 1.32 \cdot 10^{-8},$$

attained by a new point and an octad vector on an octad other than Q . The tightest pair of two new points has $\frac{1}{4} - \cos^2 \approx 1.98 \cdot 10^{-8}$; their sign vectors are at Hamming distance 5. All 12975933 contacts are between base points.

The deletion is necessary for these representatives. A computer check shows that each of the 13 deleted points has cosine between 0.50444 and 0.50496 with some new point, and that each of the other 115 odd patterns on Q has cosine at most 0.49999998 with every new point. The group $\{S_h\}$ fixes Q pointwise, so its orbits are free to carry different weights near Q .

Where the second half of the table comes from. The table came in two steps, and the second step rests on the following observations; the proof above checks the final table and nothing else.

- (a) For every $h \in D$ that vanishes on Q , the proof of Lemma 4.2 shows that S_h maps B' onto itself and fixes each of the 13 deleted points.
- (b) So if a set A of new points is compatible with B' and with itself, then so is $S_h(A)$, and $A \cup S_h(A)$ needs no further deletions. The only new conditions are the pairs between A and $S_h(A)$.
- (c) Exactly 32 words of D vanish on Q , and 8 of them form Σ_0 . So up to Σ_0 there are three choices of h besides Σ_0 itself. For $A = \{S_g r_i\}$, $g \in \Sigma_0$, the largest cosine between A and $S_h(A)$ is 0.4746 on the class of w , but 0.5117 and 0.9794 on the other two classes, which therefore fail.
- (d) An earlier configuration of 30763 points used 384 representatives under Σ_0 and deleted 29 points on Q . In it, the graph on the 384 orbits that joins two orbits when some pair between them has cosine above 0.4999 has two components of 192 orbits, and no pair across the components has cosine above 0.491. The 29 deleted patterns split disjointly, 13 conflicting with one component and 16 with the other. Keeping the first component, $r_1, \dots, r_{192}$, needs only 13 deletions, and (b) with $h = w$ refills the other 1536 points. This is the whole gain over 30763.
- (e) S_w moves the sign words of A from the cosets of 696, 1993, 2787, 1393, 2139, 3370 to those of 0, 1393, 2139, 1993, 2787, 3986, in that order. This is why all eight cosets appear. By Lemma 4.1, flat sign vectors from cosets of opposite parity conflict at cosine $11/21$; with these weights the largest cross cosine is 0.4746.

These facts were observed numerically.

With the base held fixed, each new point may be moved independently to any unit vector within chord distance $1/(8 \cdot 10^8)$ of its position, and the result is again a kissing configuration, because each inner product then changes by at most $2/(8 \cdot 10^8) < 1/L$. So with B' fixed these configurations form an open set of dimension $3072 \cdot 20 = 61440$. This freedom does not seem to produce room for a 30780th point, and we have not tried to prove that it cannot.

5. FORMAL VERIFICATION

We formalized Theorem 1.1 in Lean 4 [5] with Mathlib [10], using Lean 4.33.1 and Mathlib at tag v4.33.1. The main theorems `kissing_d19` and `kissing_d21` state, for $(n, N) = (19, 12270)$ and $(21, 30779)$, that there is a finite set S in `EuclideanSpace ℝ (Fin n)` with

$$|S| = N, \quad \|x\| = 1 \quad (x \in S), \quad \langle x, y \rangle \leq \frac{1}{2} \quad (x \neq y \in S).$$

A further lemma restates this in terms of spheres: the unit balls centred at $2x$, $x \in S$, touch the unit ball at the origin, and their centres are at pairwise distance at least 2. We do not define τ_n in Lean. The theorems are the existence statements that the bounds $\tau_n \geq N$ mean.

The development has two layers.

- (1) A plain Lean library, `lean/KissingCheck`, builds each configuration as an array of integer rays from its compact description: Ho's twelve generators, the base rules of Section 3, the two restored points, the four generators of G and the 112 representatives, or the Golay polynomial, the 13 deleted sign patterns, the four generators of Σ and the 192 representatives. Each ray is a positive multiple of a point in the text; for instance, the 19-dimensional base rays are $(2b_H, Rb_T)$. One Boolean function tests (1) on every pair, and another tests that every ray has n coordinates and is nonzero.

- (2) A Mathlib library, `lean/KissingD19D21` (the bridge is in `Bridge.lean`), proves, with kernel-checked proofs, that if both tests return true then the normalized rays are N distinct unit vectors with pairwise inner products at most $1/2$. It identifies the Euclidean inner product of two rays with their integer dot product, derives cosine at most $1/2$ from (1), and uses cosine less than 1 to show that the normalized rays are distinct.

The tests are evaluated with `native_decide`. This tactic compiles the Boolean function, runs it, and if the result is true adds an axiom asserting that it evaluates to true. Each use appears in `#print axioms` as its own auxiliary axiom. Each main theorem depends on `propext`, `Classical.choice`, `Quot.sound` and three such axioms, for the size of the array, the nonzero rays and the pairs. Trusting them means trusting that Lean’s compiler, its C code generator and its arbitrary-precision integer runtime compute what the definitions say. A kernel evaluation of 473 million pairs is out of reach. Running both tests natively with `lake exe checkall 19 21` takes about one minute on one core. Once Mathlib is built, the whole project builds in about two minutes on a laptop, about one minute of it for the 21-dimensional module.

Only the conclusion is formally verified. The theorems depend on the three tests and on the bridge in layer (2), not on the construction code. Lean does not verify that the arrays it builds match the descriptions in Sections 3 and 4, and it does not verify our case analysis. We checked the match separately. The script `scripts/compare_with_witness.py` reduces each ray built in Lean (written to `lean/out/` by `lake exe dump rays`) and each ray in the files `data/D19_12270_rays.txt` and `data/D21_30779_rays.txt` to its primitive integer vector, and finds the two sets equal, with 12270 and 30779 directions.

The Lean code names some objects differently from this paper: there K is Σ , H is Σ_0 and T is the 21-dimensional octad Q . In dimension 19 the 10476 base points of Table 2 are called `core`.

The development also contains sanity checks. Kernel-checked `decide` proofs show that the pair test accepts cosine exactly $1/2$ and rejects a pair with $4\langle x, y \rangle^2 - |x|^2|y|^2 = 1$ at size $4 \cdot 10^{18}$. Further checks in `lean/KissingD19D21/Controls.lean`, again by `native_decide`, show that each of the 190 points still deleted in dimension 19 and the 13 deleted in dimension 21 conflicts with some new point.

6. CONCLUSION

We have shown that $\tau_{19} \geq 12270$ and $\tau_{21} \geq 30779$, improving the best published lower bounds of 11948 and 29768. Two questions about the 19-dimensional construction are open to us. Can more of the 46 deleted points on the fiber that G fixes be restored with a new table under the same group? And can a larger table, under a sign group of order at most 4 that fixes two deleted fibers pointwise (Section 3.5, remark (d)), restore points on both fibers? A closed form for the reweighted table would help with both. In 21 dimensions two questions remain. Can some of the 13 deleted points be restored by moving the 192 representatives? And does the step of Section 4.5, keeping one of two decoupled halves of a table and refilling the other with a sign change that fixes the deleted octad, work in 20 or 22 dimensions?

USE OF ARTIFICIAL INTELLIGENCE

GPT-6-Astra Pro and Claude Opus 5.5 were used to find these constructions, write the Lean formalization, and draft this paper.

DATA

The accompanying GitHub repository is

<https://github.com/Felpix-Studios/D19-D21-Kissing>.

It contains short plain-text descriptions of both constructions (`constructions/`), the coordinates of both configurations and of the 12268-point configuration of Section 3, the certificate files with the 112 representatives of Section 3.5 and the 384 representatives of Section 4 (`data/`), the verification programs (`scripts/`) and the Lean project (`lean/`).

APPENDIX A. SAMPLE REPRESENTATIVES

Table 5 lists three of the 112 representatives of Section 3.5, numbered as in `data/D19_12270_certificate.json`: r_1 , an axis word with tail along $-e_1$; r_2 , one of the eight representatives of remark (b), with smaller entries on the coordinates 3, 6, 8, 19; and r_{49} , the first diagonal word. Each is written as three rows of seven, seven and five entries. The first 16 entries are on the head coordinates 2, 3, 5, 6, 8, 10, 11, $\dots$, 20 and the last three form the tail.

TABLE 5. Three representatives r_i in dimension 19.

r_1 , axis	20481341	20313490	-20456771	-20564075	-20601849	20472517	20437129
	-20449319	20482519	20441450	-20457606	20457239	-20463365	20453283
	20368158	20492357	-57452775	-2578	20672		
r_2 , axis	-22223587	-18161323	-22213549	-18167077	18190542	-22211405	-22202628
	-22192123	22213874	22202593	-22213820	22192160	22213558	22211400
	18160057	22223974	-52538238	217	-8099		
r_{49} , diagonal	-23571249	-23567224	23569769	-23569568	-23569515	23569882	23567203
	-23403275	23569946	23569578	23402176	23401749	-23570743	23402033
	23568297	23566505	-19407839	-19578265	-19578365		

Table 6 lists, for dimension 21, the first representative on each of the six cosets met by $r_1, \dots, r_{192}$, in the order in which they appear in `data/D21_30779_certificate.json`. Each is written as three rows of seven entries, coordinates 1 to 21.

TABLE 6. One representative r_i per coset of C .

r_1 , coset 696	246234381	219980925	209785187	−202482891	−224617629	−218092629	216543425
	−219980925	217704730	−239855454	239957324	216009092	220121191	204061050
	236622028	201715279	231423123	216136122	224959675	180951525	181953903
r_{33} , coset 1993	235702242	−235702246	235702245	235702249	−235702248	−235702244	−235702246
	−149071276	235702243	235702244	−235702242	−235702243	−235702244	−235702248
	149071272	149071276	235702246	149071274	149071274	235702248	235702248
r_{65} , coset 2787	225351549	182591213	247281647	−228539751	−187676135	235642930	−216731283
	−207740975	−197342048	231905242	225679867	188757907	−222180761	−210430962
	204605411	249283059	215561908	215633807	221449791	227383590	225778340
r_{97} , coset 1393	214926831	−190063656	233096365	−230585945	200371893	241359405	−235160970
	205103483	182738919	−215303842	−210489005	−184471317	−226272166	−204197249
	239084407	234925920	224054811	221607207	221607197	229029592	224055305
r_{129} , coset 2139	149071274	235702247	149071275	235702251	149071276	−235702243	−235702244
	235702247	−235702244	−235702243	235702242	235702244	−235702243	−235702248
	235702242	235702248	149071275	149071274	235702242	235702247	235702249
r_{161} , coset 3370	235368146	−235367792	235367792	−149956790	149584899	−150128199	235368272
	149956667	−235221677	150184393	−235795937	−235221410	235740042	235392962
	235917546	235399007	235796162	235739740	235942608	235368197	235399408

REFERENCES

- [1] H. Cohn and A. Li, *Improved kissing numbers in seventeen through twenty-one dimensions*, preprint, 2024, version 3, 2026. arXiv:2411.04916v3.
- [2] J. H. Conway and N. J. A. Sloane, *Sphere packings, lattices and groups*, third edition, Grundlehren der mathematischen Wissenschaften 290, Springer, New York, 1999.
- [3] N. Leijenhorst and D. de Laat, *Solving clustered low-rank semidefinite programs arising from polynomial optimization*, Math. Program. Comput. **16** (2024), no. 3, 503–534.
- [4] H. Cohn, *Kissing numbers*, table, <https://cohn.mit.edu/kissing-numbers/>, accessed 4 October 2026.
- [5] L. de Moura and S. Ullrich, *The Lean 4 theorem prover and programming language*, in Automated Deduction, CADE 28, Lecture Notes in Comput. Sci. 12699, Springer, 2021, 625–635.
- [6] B. S. Ho, *A new lower bound for the kissing number in 19 dimensions*, preprint, 2026. arXiv:2603.10425v2.
- [7] J. Leech, *Notes on sphere packings*, Canadian J. Math. **19** (1967), 251–267.
- [8] J. Leech and N. J. A. Sloane, *Sphere packings and error-correcting codes*, Canadian J. Math. **23** (1971), no. 4, 718–745.
- [9] F. J. MacWilliams and N. J. A. Sloane, *The theory of error-correcting codes*, North-Holland, Amsterdam, 1977.
- [10] The mathlib Community, *The Lean mathematical library*, in Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020), ACM, 2020, 367–381.