## Worksheet Week 21

## **Problems**

- **Q1.** X and Y are two (binary) random variables. If X and Y are independent, then P(X,Y) = P(X)P(Y)
  - (a) Give an example of two random variables that are independent.

For example - two flips of a coin. Suppose we have X = coin 1 and Y = coin 2. Then P(X = head, Y = head) = P(X = head)P(Y = head)

(b) Complete the probability table below in such way that the variables X and Y are independent.

	X = 0	X = 1
Y = 0		
Y=1		

Solution:

For example,

Marginalising over X, we get P(X=0)=0.4, P(X=1)=0.6. Marginalising over Y we get P(Y=0)=P(Y=1)=0.5. We see that P(X,Y)=P(X)P(Y) for each combination of X and Y.

(c) Determine the missing entries (a, b) of the joint distribution in such a way that the variables X and Y are again independent.


$$P(Y = 0, X = 0) = 0.1$$
  
 $P(Y = 0, X = 1) = 0.3$   
 $P(Y = 1, X = 0) = a$ 

P(Y = 1, X = 1) = b

Solution:

a = 0.15 and b = 0.45

Q2. Consider the following Bayesian network:



(a) Which random variables are independent of  $X_{3,1}$ ? Solution:

None. Recall that two random variables X and Y are independent if P(X,Y) = P(X)P(Y). In a diagram, this equates to having two disconnected nodes. In the diagram above, consider writing out the joint distribution and then marginalising to obtain e.g.  $P(X_{3,1}, X_{2,2})$ . As we marginalise, we can eliminate some variables, e.g.  $X_{3,3}, X_{2,3}, X_{1,3}, X_{3,2}$ , but we will not be able to end up with simply  $P(X_{3,1}, X_{2,2}) = P(X_{3,1})P(X_{2,2})$ 


A comprehensive algorithm for determining independence is called d-separation, and you can see this at http://web.mit.edu/jmn/www/6.034/d-separation.pdf

(b) Which random variables are independent of  $X_{3,1}$  given  $X_{1,1}$ ? Solution:

Both  $X_{1,2}$  and  $X_{1,3}$  become independent of  $X_{3,1}$  given  $X_{1,1}$ .

Q3. Solve the questions on slides 42 and 44 of the lecture slides. What is the marginal probability, P(S=1), that the sprinkler is on, given that the grass is wet?

$$P(S = 1|W = 1) = \frac{P(S = 1, W = 1)}{P(W = 1)}$$
 
$$P(W = 1) = \sum_{R} \sum_{C} \sum_{S} P(W = 1, S, C, R)$$
 
$$P(S = 1, W = 1) = \sum_{R} \sum_{C} P(W = 1, S = 1, C, R)$$



The joint probability P(W, S, C, R) is given by

$$P(W, S, C, R) = P(C)P(S|C)P(R|C)P(W|S, R)$$

$$\tag{1}$$

So,

$$P(W=1) = \sum_{C} \sum_{S} \sum_{R} P(C)P(S|C)P(R|C)P(W=1|S,R)$$
 (2)

$$= 0.5 * 0.1 * 0.8 * 0.99 + 0.5 * 0.1 * 0.2 * 0.9$$
(3)

$$+0.5*0.9*0.8*0.9+0.5*0.9*0.2*0$$
 (4)

$$+0.5*0.5*0.2*0.99+0.5*0.5*0.8*0.9$$
 (5)

$$+0.5*0.5*0.2*0.9+0.5*0.5*0.8*0$$
 (6)

$$=0.6471$$
 (7)

$$P(W=1, S=1) = \sum_{R} \sum_{C} P(C)P(S=1|C)P(R|C)P(W=1|S=1, R)$$
 (8)

$$= 0.5 * 0.1 * 0.8 * 0.99 + 0.5 * 0.1 * 0.2 * 0.9$$

$$\tag{9}$$

$$+0.5*0.5*0.2*0.99+0.5*0.5*0.8*0.9$$
 (10)

$$=0.2781$$
 (11)

So 
$$P(S=1|W=1) = \frac{P(S=1,W=1)}{P(W=1)} = 0.2781/0.6471 = 0.4297$$

What is the posterior probability, P(S = 1|W = 1, R = 1), that the sprinkler is on given that the grass is wet and it is raining?

$$P(S=1|W=1,R=1) = \frac{P(S=1,W=1,R=1)}{P(W=1,R=1)}$$

$$P(W = 1, R = 1) = \sum_{S} \sum_{C} P(S, W = 1, R = 1, C)$$

$$P(S = 1, W = 1, R = 1) = \sum_{C} P(S = 1, W = 1, R = 1, C)$$

$$P(W=1, R=1) = \sum_{C} \sum_{S} P(C)P(S|C)P(R=1|C)P(W=1|S, R=1)$$
 (12)

$$= 0.5 * 0.1 * 0.8 * 0.99 + 0.5 * 0.9 * 0.8 * 0.9$$

$$(13)$$

$$+0.5*0.5*0.2*0.99+0.5*0.5*0.2*0.9$$
 (14)

$$=0.4581$$
 (15)

$$P(W = 1, R = 1, S = 1) = \sum_{C} P(C)P(S = 1|C)P(R = 1|C)P(W = 1|S = 1, R = 1)$$

$$= 0.5 * 0.1 * 0.8 * 0.99 + 0.5 * 0.5 * 0.2 * 0.99$$


$$(17)$$

$$=0.0846$$
 (18)

So 
$$P(S = 1|W = 1, R = 1) = 0.0846/0.4581 = 0.1846$$

I.e., if we know it is raining, this decreases the probability that the sprinkler is on, given that the ground is wet.

**Q4.** A patient can have a symptom, S, that is caused by two different diseases, A and B. It is known that the presence of a gene G is important in the manifestation of disease A. The Bayes net and conditional probability tables are shown in Figure 2.



P(G)		
g	0.1	
$\neg g$	0.9	

P(A G)				
g	а	1.0		
g	¬a	0.0		
$\neg g$	а	0.1		
$\neg g$	¬a	0.9		

P(B)			
b	0.4		
$\neg b$	0.6		

P(S A,B)				
а	b	S	1.0	
а	b	٦S	0.0	
а	$\neg b$	S	0.9	
а	$\neg b$	٦S	0.1	
¬a	b	S	0.8	
¬a	b	٦S	0.2	
¬a	$\neg b$	S	0.1	
¬a	$\neg b$	٦S	0.9	

Figure 1: Bayes net and probability tables for Q5

(a) What is the probability that a patient has disease A?

$$P(a) = P(a|g)P(g) + P(a|\neg g)P(\neg g)$$
(19)

$$= 1 * 0.1 + 0.1 * 0.9 = 0.19 \tag{20}$$

- (b) What is the probability that a patient has disease A if we know that the patient has disease B
  - A and B are conditionally independent, so P(a|b) = P(a) = 0.19
- (c) What is the probability that a patient has disease A if we know that the patient has disease B AND symptom S

$$P(a|s,b) = \frac{P(a,s,b)}{P(s,b)}$$
(21)

$$= \frac{P(a)P(b)P(s|a,b)}{P(a)P(b)P(s|a,b) + P(\neg a)P(b)P(s|\neg a,b)}$$

$$= \frac{0.19 * 0.4 * 1}{0.19 * 0.4 * 1 + 0.81 * 0.4 * 0.8} = 0.2267$$
(23)

$$= \frac{0.19 * 0.4 * 1}{0.19 * 0.4 * 1 + 0.81 * 0.4 * 0.8} = 0.2267$$
 (23)