

Note: Page numbers followed by “f” indicate figures, and “t” indicate tables.

A

Absorption corrections

- attenuation coefficients, 231–232
- cylindrical geometry, 232–233
- equations, 230–231
- flat-plate reflection geometry
 - diffraction patterns, 235
 - incoming and outgoing beams, 233–234, 233f
- kapton foil, 235–236, 236f
- ratio, 238
- supported sample, 236–237, 237f
- flat-plate transmission geometries, 238–241
- and multiple-scattering
 - attenuation factors, 176–177
 - beam profile function, 177
 - isotropic approximation, 180
 - neutron absorption resonance, 177–178
 - symmetric flat-plate geometries, 177, 178f
 - X-ray double-scattering ratios, 179, 179f

Amorphous materials

- multicomponent glass structure, 456–457
- PDF analysis, 455–456
- structural changes
 - mechanical deformation, 459–464
 - structural relaxation, 457–459
- Anelastic deformation, metallic glass, 461–462
- Anisotropic PDF, 90–91, 108–109
- Anomalous X-ray scattering, 81–83
- Atomic displacement parameters (ADPs), nanoparticles, 412–414, 452
- Atomic form-factor
 - anomalous scattering factors, 187
 - coherent and Laue incoherent scattering, 187
 - definition, 186
 - Dirac–Slater atomic wavefunctions, 187
 - normalized data, 188, 188f
 - atomic pair distribution function. *See* Pair Distribution Function (PDF)

B

- Beevers–Lipson strips, 73–74, 74f, 103–104
- Big Box model

- description, 270–271
- DISCUS program, 280
- disordered systems, 279
- ergodicity, 276
- inverse problem, 278–279
- Metropolis simulated annealing algorithm, 276–277
- Monte-Carlo simulated annealing algorithm, 276–277
- reverse Monte Carlo (RMC) technique, 277–278, 279–280
- Biotic manganese oxides, 394–395

C

- Carbamazepine, local molecular material
 - packing
 - β -crystalline form, 445–446, 446f
 - crystalline polymorphs, 445–446
 - melt-quenching, 445–447, 445f, 446f, 448, 449f
 - total scattering reduced structure functions, 445–446, 445f
 - XRD fingerprints, 443–445, 443f
- Carbon nanostructures
 - disordered nanoporous carbon model, 388–391, 390f
 - fullerenes (C₆₀)
 - discovery, 386–387
 - graphitizing and nongraphitizing carbons, 388–391, 390f
 - point symmetry, 387
 - reduced structure function, 387, 388f
 - total scattering, 388
 - Warren line-shape, 388–391
- CdSe nanoparticles
 - archetypal quantum dot, 408–409
 - 1D diffraction patterns, 410f
 - 2D diffraction patterns comparison, 409f
 - PDFs, 422–423, 425f
- Cerium oxides
 - annealing effect, 383–384, 384f
 - automotive industry, 380

- Cerium oxides (*cont.*)
 - catalyst support, 380–386
 - fcc lattice, 383
 - Frenkel-type defects, 381–383, 382f
 - intermediate range order, 378–379
 - interstitial oxygen defects, 381–383, 382f
 - neutron PDF analysis, 381–383
 - oxygen storage capacity (OSC), 380–381
 - Rietveld analysis, 381–383
 - role of, 380
 - zirconia mixing, 385–386, 386f
- Chemical short-range order (CSRO), 397–399
- Chopper spectrometer
 - backscattering spectrometer IRIS, 301, 302f
 - description, 301, 301f
 - double-differential scattering cross-section functions, 302, 303f
- Clusters
 - CRN, 432–433
 - gold nanoparticle formation, 424–426
 - hydrothermal chemical reactions
 - PDF experiments, in situ setup, 426–427, 426f
 - SnO₂, time-dependent PDF, 426–427, 427f
 - SrFit, 427–428
 - independent neutron, X-ray and EXAFS datasets, 430, 431f
 - nanoporous solids
 - hydrogen in cyanide host, 431–432
 - mercury in silica, 434
 - MOFs, 432–433
 - sodium in silica, 434–437
 - ZIFs, 432–433
 - PDF, 423–424
 - Pt(II) ion complexes structures, 430
 - surfactant micelles structures
 - configurations, 429–430
 - EPSR modeling, 428–429
 - TAB, 428
- Colossal magnetoresistance (CMR)
 - phenomenon, 349, 350
- Complex materials
 - crystallographic analysis
 - (*see* Crystallographic analysis)
 - phase transitions
 - atomic bonds, hierarchy of, 354–355
 - disorder effect, 355, 356f
 - first-order transitions, 356–357
 - non-linearity, 356–357
 - structure
 - atomic and molecular level, 1–3
 - colossal magnetoresistive (CMR) oxides, 3
 - local crystallography methods, 7–10
 - nonlinear system, 4
 - PDF and total scattering methods, 11–22
 - periodic crystal, 4
 - real and reciprocal space, 10–11
 - shadow of Bragg's law, 6–7
 - translational symmetry, 5
- Complex modeling, PDF
 - automated trial-and-error methods, 280–281
 - components, 288
 - constrained reverse Monte Carlo (RMC) technique, 284–285
 - EPSR, 282–284
 - nanoparticle modeling, 285–288
 - powder diffraction, 281
 - schematic illustration, 281, 282f
 - Scherrer broadening, 281
 - SrFit, 288–289
- Complex oxides
 - disorder effect, 343–345
 - manganites
 - behaviors, 349
 - description, 349
 - Jahn–Teller distortions, 349–351
 - La_{1-x}A_xMnO₃, 349–350
 - locally ordered polaron model, 350–351, 351f
 - Pb(Zr_{1-x}Ti_xO₃) (PZT)
 - antiferroelectric ordering, 346–347
 - PDF, 345–346
 - phase transformations, 345–346
 - uses, 345
 - Pb(Mg_{1/3}Nb_{2/3})O₃ (PMN)
 - crystallographic study, 347
 - disorder effect, 347
 - local structure, 347–348
 - pulsed neutron PDF method, 347–348, 348f
 - vs. PZT, 347–348, 348f
- Compton-scattering correction, 184–186
- Continuous random network (CRN), 432–433
- Crystal defects
 - carbon nanostructures, 386–393
 - cerium oxides, 378–379, 380–386
 - characterization, 372
 - crystallographically challenged materials, 372–374, 373f
 - CSRO, 397–399
 - CuIr₂S₄, structural response, 396–397
 - earth science

- biotic manganese oxides, 394–395
 - ferrihydrite case, 393–394
 - gamma alumina, 375–376
 - lead tellurides, nanocluster-enhanced
 - thermoelectricity, 374–375, 375f
 - in operando* fuel cell device, 401–403
 - oxygen ion conductors, 377–378
 - PDF method
 - accuracy, 372
 - nanoporous carbon, 388–391, 392f
 - operating fuel cell, 401–403
 - r²-scaled, 392–393, 393f
 - pyrolyzed polyvinylidene chloride, structure
 - function, 388–391, 391f
 - silicon crystals, 372
 - in situ* RAPDF studies, 399–401
 - Crystallographic analysis
 - atomic displacements, 36–37
 - crystal symmetry, 7–8
 - diffuse scattering
 - Bragg scattering, 36–37
 - high-Q region, 35
 - Huang scattering, 38–39
 - phonon intensity, 36–37
 - point defects, 39
 - DW approximation, 34–35
 - limitations of methods
 - caveat emptor, 47–48
 - correlated parameters, 46–47
 - DW factor, 43–45
 - imperfect models, 45–46
 - pair correlation functions, 8
 - PDF, 8
 - powder diffraction, 9
 - pulsed spallation neutron and X-ray
 - synchrotron sources, 9–10
 - rietveld refinement method
 - powder diffraction, 39–40
 - programs, 41
 - R-factor, 39–40
 - scattering amplitude
 - Born approximation, 50–51
 - Bragg peak, 31–32
 - derivation, 49–50
 - kinematic approximation, 31
 - over-sampling, 31–32
 - scattering cross-section, 48–49
 - scattering intensity
 - geometry of powder diffraction and
 - definition of, 27–28, 28f
 - scattering cross-sections, 30
 - TEM, 28–29
 - single crystal Fourier analysis, 41–42
 - CSRO. *See* Chemical short-range order (CSRO)
 - CuIr₂S₄, structural response, 396–397
- ## D
- Data collection analysis
 - approximate method
 - absorption factor, 228–229
 - double-differential cross section, 229–230
 - measurements, 228
 - multiple scattering, 229
 - detector
 - flux normalization, 224–225
 - total measured intensity, 224
 - total single and multiple-scattering intensities, 223–224
 - double-differential cross section, 165
 - Bragg scattering, 217
 - definitions, 216–217
 - inverse problem, 226–227
 - and multiple scattering, 221–223
 - schematic illustration, 216–217, 217f
 - effective scattering volume, 164–165
 - hydrogen corrections in neutron
 - measurements, 211–213
 - measurements, 163–164
 - optimization, 245–248
 - PDFgetN
 - blended data, 252, 253f
 - colossal magnetoresistance effect, 248
 - datasets, 248, 248f
 - detector-bank information, 251, 251f
 - plot window, 251–252, 252f
 - reduced pair distribution function, 252–254, 254f
 - reduced structure function, 252, 253f
 - sample information window, 250–251, 250f
 - smoothing parameters, 249–250
 - special environment powder diffractometer (SEPD), 249–250
 - PDFs using ad hoc data corrections
 - Debye–Waller factor, 167–168
 - frequency Fourier component, 166
 - multiplicative correction term, 166
 - PDFgetX3, 165
 - X-ray PDF, 168, 168f
 - programs, 210–211
 - random error propagation
 - covariance, 243
 - detection process, 242–243

- Data collection analysis (*cont.*)
 - dimensionless correlation coefficient, 242
 - integrated peak intensities, 245
 - sine-waves, 245
 - small fluctuation, 244
 - statistical correlations, 243
 - variance, 241
 - single-scattering cross section, 164
 - attenuation factors, 219–220
 - detector profile, 220
 - sample surrounded by sample environment, 218–219, 219f
 - schematic illustration, 218, 218f
 - sound effect, 218–219
 - statistical uncertainties on PDF
 - Fourier transform, 214
 - powder pattern, 215–216
 - programs, 213
 - refined parameters, 215
 - stochastic process, 213
 - total scattering structure function
 - absorption and multiple-scattering corrections, 176–180
 - combining datasets, 204–206
 - data termination, 206–208
 - detector dark counts and dead-time corrections, 171–173
 - elastic, inelastic, coherent, and incoherent scattering, 169–170
 - electron-specific data corrections, 201–204
 - flux normalization, 175–176
 - Fourier transform, 208–209
 - instrument resolution function, 209–210
 - random error propagation, 173–175
 - real-time PDFs, 170–171
 - time-of-flight neutron-specific corrections, 189–193
 - X-ray-specific corrections, 180–193
 - Debye model
 - description, 265–267
 - nanoparticles (NP), 419–420, 420f
 - PDF peak widths
 - atomic separation, function of, 267–268, 269f
 - manganite system, 267
 - Morse potential, 265–267
 - phonon density of states, 265–267, 267f
 - phonon spectrum determination, 265–267
 - r*-dependence, 267–268, 268f
 - temperature dependence, 265–267, 266f
 - r*-dependence, 267–268, 268f
 - temperature dependence, 265–267, 266f
 - Debye–Scherrer geometry, 232–233
 - Debye–Waller factor, 305–306, 308–310
 - Differential PDFs (DDFs), 80–81
 - Diffuse scattering
 - Bragg scattering, 36–37
 - high-Q region, 35
 - Huang scattering, 38–39
 - phonon intensity, 36–37
 - point defects, 39
 - DISCUS modeling package, 421–422
 - Dynamic pair-density function (DPDF)
 - PMN relaxor ferroelectrics, 312–314
 - polycrystalline nickel, 311–312, 313f
 - temperature dependence, 314, 315f
 - Dynamic structure factor, inelastic scattering
 - derivation, 322–323
 - multi-phonon scattering, 306–308
 - single-phonon scattering, 303–306
- ## E
- Earth science, crystal defects
 - biotic manganese oxides, 394–395
 - ferrihydrite case, 393–394
 - Elastic deformation, metallic glass, 459–460
 - Electron PDF methods
 - beam damage, 151
 - camera length, 148
 - clusters in gas phase and amorphous films, 145–146
 - data collection modes, 148–149
 - detector, 149–150
 - dynamical scattering effects, 144–145
 - energy filtering, 150
 - gold nanoparticles, 146–147, 147f
 - multiple scattering, 150–151
 - powder average, 147–148
 - sample preparation, 147
 - sample support measurement, 150
 - Empirical potential structure refinement (EPSR), 282–284
 - Energy-dispersive X-ray diffraction measurements, 144
 - EPSR. *See* Empirical potential structure refinement (EPSR)
 - EPSR modeling, 428–429
 - Error analysis, PDF
 - absorption correction, 99
 - diagnostics, 95–96
 - effect of Q-resolution, 98–99
 - remedies, 100
 - statistical, 97–98
 - termination and normalization, 96–97

F

- Ferrihydrite, 393–394
- Fictive temperature, 458
- Fourier transform
 - data collection analysis
 - statistical uncertainties on PDF, 214
 - total scattering structure function, 208–209
 - discrete scattering, 62–63
 - dynamic structure factor, 57–58
 - graphical representation, 67*f*, 68
 - intensity distribution, 57
 - Laue monotonic scattering, 57–58
 - nanomaterials, 60
 - nickel, 60–61, 60*f*
 - NIMROD/NOMAD, 61
 - orientational averaging, 63–64
 - Patterson function, 61
 - RDF, 65–66
 - scattering power, 62
 - shapes, 68–69
 - small-angle scatterers, 66–67
 - X-ray form factors, 64
- Free-volume theory, 458
- Fuel cell device, 401–403
- Fullerenes (C₆₀)
 - discovery, 386–387
 - graphitizing and nongraphitizing carbons, 388–391, 390*f*
 - point symmetry, 387
 - reduced structure function, 387, 388*f*
 - total scattering, 388

G

- Gamma alumina, crystal defects, 375–376
- Generalized Patterson function, 41–42
- Giant magnetoresistance, 349

H

- Harris criterion, 355
- Heterogeneous nanoparticle modeling
 - DISCUS modeling package, 421–422
 - model-independent analysis, 420–421
 - PDF, gold nanoparticles, 420–421, 422*f*
- Homogeneous size-dependent strain, 416–417

I

- Inelastic scattering measurement
 - chopper spectrometer
 - backscattering spectrometer IRIS, 301, 302*f*

- description, 301, 301*f*
 - double-differential scattering cross-section functions, 302, 303*f*
- description, 297–298
- dynamic structure factor
 - derivation, 322–323
 - multi-phonon scattering, 306–308
 - single-phonon scattering, 303–306
- hybrid techniques, 320–322
- inelastic X-ray scattering (IXS), 299–300, 300*f*
- neutron triple-axis spectrometer, 298–299, 298*f*
- PDF
 - correlated dynamics, 308–310
 - neutron method, 319
 - phonon dispersions, 319–320
 - Placzek correction, 314–319
- Inhomogeneous strains, 418–419
- In operando* fuel cell device, 401–403

L

- Landau–Devonshire theory, 353
- Lead tellurides, nanocluster-enhanced thermoelectricity, 374–375, 375*f*
- Lead zirconate (PbZrO₃)
 - description, 333
 - high-temperature phase, 334–337
 - low-temperature phase, 333–334
 - Pb polarization, 337–338
 - symmetries, 333
- Lennard–Jones cluster, 290–291, 292*f*
- Lithium-nickelate (LiNiO₂)
 - description, 363–364
 - domain formation, 364–366
 - hidden phase transition, 364
 - local structure, 364
 - pulsed neutron PDF, 364, 365*f*
 - Rietveld analysis, 364
 - temperature dependence, 364, 365*f*
- Local molecular material packing
 - active pharmaceutical ingredient (API), 442, 443*f*
 - carbamazepine pharmaceutical
 - β-crystalline form, 445–446, 446*f*
 - crystalline polymorphs, 445–446
 - melt-quenching, 445–447, 445*f*, 446*f*, 448, 449*f*
 - total scattering reduced structure functions, 445–446, 445*f*
 - XRD fingerprints, 443–445, 443*f*
 - drug delivery, 442

Local molecular material packing (*cont.*)
 indomethacin samples, 443–445, 444f
 PDF
 atomic displacement factors, 452
 β -crystalline form, CBZ, 445–447, 446f
 drug development, 442–443
 indomethacin, 443–445, 444f
 Nyquist–Shannon sampling frequency, 449–450
 Pearson product-momentum correlation, 446–447
 role, 443–445
 SrFit modeling software, 451f, 452
 and total scattering PDF measurements, 445–446
 pigment industry, 441–442

M

Magnetic correlation functions
 neutron scattering, 86–87
 PDF, 87–89
 MCM-41, 434
 Metallic glass
 fast cooling, 458
 formation, 456
 isotropic and anisotropic structure function, 460, 461f
 reverse Monte Carlo (RMC) technique, 456–457
 structural changes
 anelastic deformation, 461–462
 elastic deformation, 459–460
 plastic deformation, 462–464
 structural relaxation
 description, 458
 inherent structure, 458
 PDF method, 458–459
 Metal organic frameworks (MOFs),
 nanoporous solids, 432–433
 Morse potential, 265–267
 Multicomponent glass structure,
 456–457
 Multicomponent systems
 Faber–Ziman partial structure functions, 78
 Morningstar–Warren approximation, 79
 partial pair distribution function, 77–78
 Warren chemical/compositional short-range
 order parameter, 80
 Multi-phonon scattering, 306–308
 Multiple-scattering and absorption corrections
 attenuation factors, 176–177
 beam profile function, 177

isotropic approximation, 180
 neutron absorption resonance,
 177–178
 symmetric flat-plate geometries, 177, 178f
 X-ray double-scattering ratios, 179, 179f

N

Nanocluster-enhanced thermoelectricity, lead
 tellurides, 374–375, 375f
 Nanoparticles (NP)
 ab initio solution, 416
 bond strain plot, 417–418, 418f
 Cd–Se bond length, 416–417, 417f
 CdSe nanoparticles
 archetypal quantum dot, 408–409
 1D diffraction patterns, 410f
 2D diffraction patterns comparison, 409f
 PDFs, 422–423, 425f
 clusters, gold nanoparticle formation,
 424–426
 Debye model, 419–420, 420f
 heterogeneous nanoparticles modeling
 DISCUS modeling package,
 421–422
 model-independent analysis, 420–421
 PDFs, gold nanoparticles, 420–421,
 422f
 homogeneous size-dependent strain,
 416–417
 inhomogeneous strains, 418–419
 size and shape
 characteristic functions, 411
 Q-space, 411
 small angle scattering (SAS), 412
 size-dependent bond stiffness, 419–420
 structure and defects
 atomic displacement parameters (ADPs),
 412–414
 hydrated ruthenia, X-ray PDF, 414–416,
 415f
 model fits comparison, 412–414, 413f
 nanoparticle gold, X-ray PDF, 414–416,
 415f
 RuO₂, 414–416
 wurtzit model, 412–414, 413f
 surface capping ligand dependence,
 422–423
 surface to volume ratio, 418–419
 temperature dependence, 419–420, 420f
 thermodynamic quantities, 417–418
 ZnS, 422–423, 424f
 Nanopolymorphism, 447–448

Nanoporous solids
 hydrogen in cynde host, 431–432
 mercury in silica, 434
 MOFs, 432–433
 sodium in silica, 434–437
 ZIFs, 432–433

Negative thermal expansion (NTE)
 $\text{Ag}_3[\text{Co}(\text{CN})_6]$, 340–341, 342f
 mechanism in covalent network, 338–340, 340f
 zinc cyanide, 340–341, 341f

Neutron scattering experiments
 beam monitor, 128–130
 diffractometers
 detector configurations, 120–123, 121f
 D4 layout, 125, 126f
 intensity vs. time-of-flight, 120, 120f
 NPDF tof diffractometer, 120–123, 122f
 PHAROS, 120–123
 Q-space resolution, 118–119
 SANDALS and NIMROD, 119–120
 schematic representation, 118, 119f
 source spectrum, 124, 124f
 SPICA, 120–123
 time-focusing process, 123–124
 types, 117–118
 well-shielded primary beam dump, 124–125
 measurement geometry, 130–131
 neutron detection, 127–128
 sources
 D4 hot moderator, 116
 ILL hot moderator, 116–117
 instruments suitable for PDF measurements, 117, 118t
 spallation, 116

Neutron triple-axis spectrometer, 298–299, 298f

Normalization errors, 96–97

Nyquist–Shannon sampling theory
 error correlations, 103
 local molecular material packing, 449–450
 modeling, 101
 noise and peak overlap, 101
 refined parameter quality factors, 102–103, 102f
 statement, 100–101

O

One-dimensional PDF and layer-layer correlations, 91–92

Oxygen ion conductors, crystal defects
 barium cerate, 378, 379f
 LAMOX, 377–378, 377f

P

Pair distribution function (PDF). *See*
also Dynamic pair-density function (DPDF)

ab initio nanostructure solution
 C_{60} problem, 290, 291f
 genetic algorithm, 290
 Lennard–Jones cluster, 290–291, 292f
 LIGA algorithm, 290–291, 292–293
 similarity constraints, 293
 simulated annealing algorithm, 289

alloy semiconductor, 11–12, 13f

anisotropic, 90–91, 108–109

anomalous X-ray scattering, 81–83

atom-pair probability distribution, 261–262

atom-pair separation, 261

Beevers–Lipson strips, 73–74, 74f, 103–104

bulk nanostructured thermoelectrics, 14–15

chemical short-range order, $\beta\text{-Na}_3\text{BiO}_4$, 20–22

CMR manganites, 15–17

coordination number, peak integrated intensity, 261

correlation functions, 73–74

crystalline single element material example, 260

in 3D, 89–90

data collection analysis
 Debye–Waller factor, 167–168
 frequency Fourier component, 166
 multiplicative correction term, 166
 PDFgetX3, 165
 X-ray PDF, 168, 168f

differential, 80–81

diffuse scattering, 442

dynamical disorder and symmetry lowering, silica, 17–19

dynamical scattering corrections, 77

electron PDF experiment (*see* Electron PDF methods)

error analysis
 absorption correction, 99
 diagnostics, 95–96
 effect of Q-resolution, 98–99
 remedies, 100
 statistical, 97–98
 termination and normalization, 96–97

femtosecond-pulsed laser, 76–77

- Pair distribution function (PDF) (*cont.*)
- as Fourier transform
 - discrete scattering, 62–63
 - dynamic structure factor, 57–58
 - graphical representation, 67f, 68
 - intensity distribution, 57
 - Laue monotonic scattering, 57–58
 - nanomaterials, 60
 - nickel, 60–61, 60f
 - NIMROD/NOMAD, 61
 - orientational averaging, 63–64
 - Patterson function, 61
 - RDF, 65–66
 - scattering power, 62
 - shapes, 68–69
 - small-angle scatterers, 66–67
 - X-ray form factors, 64
 - goodness-of-fit parameter, 269–270
 - graphene sheet production, 263–265, 264f
 - inelastic scattering measurement
 - correlated dynamics, 308–310
 - Einstein model, 308–310
 - neutron method, 319
 - phonon dispersions, 319–320
 - Placzek correction, 314–319
 - intermediate range structure, 442–443
 - isotopic substitution
 - total structure function, 83–84
 - vibrational frequency, 85–86
 - YBa₂Cu₃O_{6.93}, 84–85, 85f
 - local atomic density, 268–269, 270f
 - local bond length measurements, 263
 - local molecular material packing
 - atomic displacement factors, 452
 - β-crystalline form, CBZ, 445–447, 446f
 - drug development, 442–443
 - indomethacin, 443–445, 444f
 - Nyquist–Shannon sampling frequency, 449–450
 - Pearson product-momentum correlation, 446–447
 - role, 443–445
 - SrFit modeling software, 451f, 452
 - and total scattering PDF measurements, 445–446
 - and TSPDF measurements, 445–446
 - local structural research, 442–443
 - magnetic, 87–89
 - model calculation
 - Big Box, 276–280
 - complex, 280–289
 - Small Box, 271–276
 - molecular solids, 12–13
 - multicomponent systems
 - Faber–Ziman partial structure functions, 78
 - Morningstar–Warren approximation, 79
 - partial pair distribution function, 77–78
 - Warren chemical/compositional short-range order parameter, 80
 - neutron scattering, 86–87
 - Nobel prizes, 74–75
 - Nyquist–Shannon sampling theory
 - error correlations, 103
 - modeling, 101
 - noise and peak overlap, 101
 - refined parameter quality factors, 102–103, 102f
 - statement, 100–101
 - one-dimensional and layer-layer correlations, 91–92
 - and pair density function, $r(r)$, 69
 - peaks
 - integrated intensity, 263–265
 - position, 262–263
 - width of, 265–268
 - phase transitions, 352, 354
 - photoreceptor material, 75–76
 - programs, 23–24
 - quasicrystals, 343, 344f
 - radial distribution function $R(r)$, 260
 - radial distribution function, $R(r)$, 72–73
 - reduced pair distribution function, $G(r)$
 - average peak-peak amplitude, 70–72, 71f
 - correlation function and data modeling, 70
 - direct Fourier transform, 70
 - structural coherence, 70–72, 72f
 - weighted residuals function, 70
 - Rietveld analysis and atomic positions, 75
 - roles of, 442–445
 - termination ripples, 76
 - three-dimensional and powder, 95
 - total and differential PDF, 86
 - two-dimensional and intralayer correlation, 92–94
 - In_{1-x}Ga_xAs material, 262–263
 - X-ray absorption fine structure method, 106–107
- Patterson function, 41–42
- Pb(Mg_{1/3}Nb_{2/3})O₃ (PMN)
- DPDF, 312–314
 - elastic and instantaneous PDFs, 308–310, 310f
- PDF. *See* Pair distribution function (PDF)
- Perovskites

- BaTiO₃, 328–331, 329*f*, 330*f*
 - description, 326–327
 - LaMnO₃, 331–333
 - PbTiO₃, 331, 332*f*
 - space groups, 327
 - SrTiO₃, 327–328, 328*f*
 - Phase transitions
 - atomic polarizations, 354
 - complex materials
 - atomic bonds, hierarchy of, 354–355
 - disorder effect, 355, 356*f*
 - first-order transitions, 356–357
 - non-linearity, 356–357
 - ferroelectric, 351–352
 - PDF methods, 352, 354
 - and phonon soft modes, 352
 - second order, 352, 353
 - systems with competing interactions
 - lithium-nickelate, 363–366
 - relaxor ferroelectricity, 357–359
 - Phonon dispersions, 319–320
 - Placzek correction
 - description, 316–317
 - neutron scattering, 316–317
 - TOF measurement, 318
 - triple-axis spectrometer, 317
 - validity, 318–319
 - X-ray inelastic scattering measurements, 314–316
 - Plastic deformation, metallic glass, 462–464
 - Polarization correction, 183–184
 - Pt(II) ion complex structures, 430
- Q**
- Quasicrystals
 - atomic structure, 343
 - characterization, 342–343
 - description, 342–343
 - discovery, 342–343
 - 6D periodic crystallography, 342–343
 - PDF, 343, 344*f*
- R**
- Random error propagation
 - data collection analysis
 - covariance, 243
 - detection process, 242–243
 - dimensionless correlation coefficient, 242
 - integrated peak intensities, 245
 - sine-waves, 245
 - small fluctuation, 244
 - statistical correlations, 243
 - variance, 241
 - scattering process, 173–174
 - statistical errors, 174–175
 - variations method, 175
 - RAPDF method. *See* Rapid acquisition PDF (RAPDF) method
 - Rapid acquisition PDF (RAPDF) method
 - advantages, 138
 - capillary, 134–135
 - Compton and fluorescence, 137–138
 - data collection
 - 2D detectors, 190
 - modeling, 189, 190*f*
 - displex cold stage, 135–136
 - image plate detectors, 137–138
 - kapton tubing, 136
 - medium-angle scattering, 136–137
 - pictorial representation, 134*f*, 134
 - representative 2D diffraction pattern, 134, 135*f*
 - Relaxor ferroelectricity
 - description, 312–314
 - phase transitions, 357–359
 - AFE interaction, 359–360
 - Burns temperature, 362–363
 - phonons, 361
 - polar nanoregions, 361–362
 - random anisotropy, 360–361
 - random field, 360
 - spin glass origin, 357–359
 - PMN, DPDF, 312–314
 - Reverse Monte Carlo (RMC) technique, 456–457
 - Rietveld refinement method
 - powder diffraction, 39–40
 - programs, 41
 - R-factor, 39–40
- S**
- Single-phonon scattering, 303–306
 - Small angle scattering (SAS), 411, 412
 - Small Box model, 275–276
 - anharmonicity, 273–274
 - atomic potential, 274
 - crystallographic structure factor, 272
 - Debye–Waller factors, 272, 273–274
 - description, 270–271
 - graphical user interface driven program, 275–276
 - PDFgui code, 275–276
 - real-space Rietveld approach, 272–273

Small Box model (*cont.*)
 Rietveld full-profile fitting method, 271–272
 thermal factors, 273–274
 well ordered crystalline material, 273, 274
 Soft-mode theory, 352
 Spin glasses, 357–358
 Spin valves, 349
 SrFit
 complex modeling, 288–289
 molecular system, 451*f*, 452
 Statistical errors, 97–98
 Strain
 nanoparticles
 homogeneous size-dependent, 416–417
 inhomogeneous, 418–419
 Surfactant micellar structures
 configurations, 429–430
 EPSR modeling, 428–429
 TAB, 428

T

Termination errors, 96–97, 105–106
 Thermal diffuse scattering (TDS), 37–38
 Thompson scattering, 181–183
 Three-dimensional PDF and powder PDF, 95
 Total scattering methods
 accuracy of
 diffraction intensity, 115
 Q-values, 114–115
 source size, 115
 and atomic PDF analysis (*see* Pair distribution function (PDF))
 definition, 56
 electron PDF experiment
 beam damage, 151
 camera length, 148
 clusters in gas phase and amorphous films, 145–146
 data collection modes, 148–149
 detector, 149–150
 dynamical scattering effects, 144–145
 energy filtering, 150
 gold nanoparticles, 146–147, 147*f*
 multiple scattering, 150–151
 powder average, 147–148
 sample preparation, 147
 sample support measurement, 150
 instrument alignment, 153–156
 monochromatic vs. polychromatic diffraction, 113–114
 neutron scattering
 beam monitor, 128–130

 diffractometers, 117–126
 measurement geometry, 130–131
 neutron detection, 127–128
 sources, 116–117
 point-by-point data collection, 151–153
 single crystal and powder diffraction methods, 114
 X-ray scattering
 beam monitor, 142
 detection, 139–141
 diffractometer, 133–139
 energy-dispersive X-ray diffraction measurements, 144
 measurement geometry and sample issues, 142–143
 sources, 131–133
 Total scattering PDF (TSPDF), 445–446, 448, 450–452
 Total scattering structure function
 absorption and multiple-scattering corrections, 176–180
 combining datasets, 204–206
 data termination, 206–208
 detector dark counts and dead-time corrections, 171–173
 elastic, inelastic, coherent, and incoherent scattering, 169–170
 electron-specific data corrections, 201–204
 flux normalization, 175–176
 Fourier transform, 208–209
 instrument resolution function, 209–210
 random error propagation, 173–175
 real-time PDFs, 170–171
 time-of-flight neutron-specific corrections, 189–193
 X-ray-specific corrections, 180–193
 Two-dimensional PDF and intralayer correlation, 92–94

W

Well-ordered crystal structures
 lead zirconate (PbZrO₃)
 description, 333
 high-temperature phase, 334–337
 low-temperature phase, 333–334
 Pb polarization, 337–338
 symmetries, 333
 NTE materials, 338–341
 perovskites
 BaTiO₃, 328–331, 329*f*, 330*f*
 description, 326–327

LaMnO₃, 331–333
PbTiO₃, 331, 332*f*
space groups, 327
SrTiO₃, 327–328, 328*f*
Wurtzite model, 412–414, 413*f*

X

X-ray detection
 amorphous silicon detector, 139–140, 141*f*
 CCD, 139–140
 ESRF Frelon detector, 139–140
 gas and array detectors, 141
 MAR345, 139
 single-element/multi-element solid state
 detector (SSD), 140–141
X-ray PDF
 hydrated ruthenia, 414–416, 415*f*
 nanoparticle gold, 414–416, 415*f*
X-ray scattering experiments
 beam monitor, 142

detection, 139–141
diffractometer
 conventional point detection strategies,
 138–139
 RAPDF method, 133–138
energy-dispersive X-ray diffraction
 measurements, 144
measurement geometry and sample issues,
 142–143
sources
 Q-space, 131–132, 131*f*
 rotating anode instruments, 132
 synchrotron radiation, 132
 wiggler, 133
 X-ray generator, 131

Z

Zeolitic imidazolate frameworks (ZIFs),
 nanoporous solids, 432–433