

# $q$ -Hodge filtrations, Habiro cohomology & $ku$

---

Ferdinand Wagner

January 8, 2026

# $q$ -Hodge filtrations, **Habiro cohomology** & $ku$

---

Ferdinand Wagner

January 8, 2026



- 1 What's Habiro cohomology supposed to do?



- 1 What's Habiro cohomology supposed to do?
- 2  $q$ -Hodge filtrations & Habiro cohomology
- 3  $q$ -Hodge filtrations from homotopy theory



- 1 What's Habiro cohomology supposed to do?
- 2  $q$ -Hodge filtrations & Habiro cohomology
- 3  $q$ -Hodge filtrations from homotopy theory



**Disclaimer:** We'll focus on “algebraic” Habiro cohomology; Scholze’s “analytic” version will only show up in the end.



- 1 What's Habiro cohomology supposed to do?
- 2  $q$ -Hodge filtrations & Habiro cohomology
- 3  $q$ -Hodge filtrations from homotopy theory



**Disclaimer:** We'll focus on “algebraic” Habiro cohomology; Scholze’s “analytic” version will only show up in the end.



- 1 What's Habiro cohomology supposed to do?
- 2  $q$ -Hodge filtrations & Habiro cohomology
- 3  $q$ -Hodge filtrations from homotopy theory
- 4 Refined  $\mathrm{THH}/\mathrm{TC}^-$  & analytic Habiro cohomology



**Disclaimer:** We'll focus on “algebraic” Habiro cohomology; Scholze’s “analytic” version will only show up in the end.

# **1. What's Habiro cohomology supposed to do?**

---





Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ ,  $\dots$ )



Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , ...) by analysing the cohomology of the associated geometric objects.




Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , ...) by analysing the cohomology of the associated geometric objects.

Many cohomology theories:



Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , ...) by analysing the cohomology of the associated geometric objects.


Many cohomology theories:





Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , ...) by analysing the cohomology of the associated geometric objects.


Many cohomology theories:





Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , ...) by analysing the cohomology of the associated geometric objects.


Many cohomology theories:





Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , ...) by analysing the cohomology of the associated geometric objects.


Many cohomology theories:





Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , ...) by analysing the cohomology of the associated geometric objects.

Many cohomology theories:








Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , ...) by analysing the cohomology of the associated geometric objects.


Many cohomology theories:





Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , ...) by analysing the cohomology of the associated geometric objects.


Many cohomology theories:





Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , ...) by analysing the cohomology of the associated geometric objects.

Many cohomology theories:





**$q$ -de Rham complex** (Bhatt–Scholze). There is a functor

$$q\text{-}\Omega_{(-)}: \operatorname{Sm}_{\mathbb{Z}} \longrightarrow \mathcal{D}(\mathbb{Z}[[q-1]])$$



**$q$ -de Rham complex** (Bhatt–Scholze). There is a functor

$$q\text{-}\Omega(-): \underset{\substack{\text{smooth} \\ \mathbb{Z}\text{-algebras}}}{\mathrm{Sm}_{\mathbb{Z}}} \longrightarrow \underset{\substack{\text{derived category} \\ \text{of } \mathbb{Z}[[q-1]]}}{\mathcal{D}(\mathbb{Z}[[q-1]])}$$



**$q$ -de Rham complex** (Bhatt–Scholze). There is a functor

$$q\text{-}\Omega_{(-)}: \underset{\substack{\text{smooth} \\ \mathbb{Z}\text{-algebras}}}{\mathrm{Sm}_{\mathbb{Z}}} \longrightarrow \underset{\substack{\text{derived category} \\ \text{of } \mathbb{Z}[[q-1]]}}{\mathcal{D}(\mathbb{Z}[[q-1]])}$$

such that:

**1** At  $q = 1$ ,  $q\text{-}\Omega_R/(q-1) \simeq \Omega_R^*$ .



**$q$ -de Rham complex** (Bhatt–Scholze). There is a functor

$$q\text{-}\Omega(-): \underset{\substack{\text{smooth} \\ \mathbb{Z}\text{-algebras}}}{\mathrm{Sm}_{\mathbb{Z}}} \longrightarrow \underset{\substack{\text{derived category} \\ \text{of } \mathbb{Z}[[q-1]]}}{\mathcal{D}(\mathbb{Z}[[q-1]])}$$

such that:

- 1 At  $q = 1$ ,  $q\text{-}\Omega_R/(q-1) \simeq \Omega_R^*$ .
- 2 For all primes  $p$ ,  $(q\text{-}\Omega_R)_p^\wedge \simeq \widehat{\Delta}_{\widehat{R}_p[\zeta_p]}/(\mathbb{Z}_p[[q-1]], [p]_q)$ .



**$q$ -de Rham complex** (Bhatt–Scholze). There is a functor

$$q\text{-}\Omega_{(-)}: \underset{\substack{\text{smooth} \\ \mathbb{Z}\text{-algebras}}}{\mathrm{Sm}_{\mathbb{Z}}} \longrightarrow \underset{\substack{\text{derived category} \\ \text{of } \mathbb{Z}[[q-1]]}}{\mathcal{D}(\mathbb{Z}[[q-1]])}$$

such that:

- 1 At  $q = 1$ ,  $q\text{-}\Omega_R/(q-1) \simeq \Omega_R^*$ .
- 2 For all primes  $p$ ,  $(q\text{-}\Omega_R)_p^\wedge \simeq \Delta_{\widehat{R}_p[\zeta_p]}/(\mathbb{Z}_p[[q-1]], [p]_q)$ .
- 3 Rationally,  $(q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)}^\wedge \simeq (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q})[[q-1]]$





**$q$ -de Rham complex (Bhatt–Scholze).** There is a functor

$$q\text{-}\Omega(-): \underset{\substack{\text{smooth} \\ \mathbb{Z}\text{-algebras}}}{\text{Sm}_{\mathbb{Z}}} \longrightarrow \underset{\substack{\text{derived category} \\ \text{of } \mathbb{Z}[[q-1]]}}{\mathcal{D}(\mathbb{Z}[[q-1]])}$$

such that:

- 1 At  $q = 1$ ,  $q\text{-}\Omega_R/(q-1) \simeq \Omega_R^*$ .
- 2 For all primes  $p$ ,  $(q\text{-}\Omega_R)_p^\wedge \simeq \Delta_{\widehat{R}_p[\zeta_p]}/(\mathbb{Z}_p[[q-1]], [p]_q)$ .
- 3 Rationally,  $(q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)}^\wedge \simeq (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q})[[q-1]]$ ,  
 $(q\text{-}\Omega_R)_p^\wedge \left[\frac{1}{p}\right]_{(q-1)}^\wedge \simeq (\Omega_R^*)_p^\wedge \left[\frac{1}{p}\right] [[q-1]]$ .



**$q$ -de Rham complex (Bhatt–Scholze).** There is a functor

$$q\text{-}\Omega(-): \underset{\substack{\text{smooth} \\ \mathbb{Z}\text{-algebras}}}{\text{Sm}_{\mathbb{Z}}} \longrightarrow \underset{\substack{\text{derived category} \\ \text{of } \mathbb{Z}[[q-1]]}}{\mathcal{D}(\mathbb{Z}[[q-1]])}$$


such that:

- 1 At  $q = 1$ ,  $q\text{-}\Omega_R/(q-1) \simeq \Omega_R^*$ .
- 2 For all primes  $p$ ,  $(q\text{-}\Omega_R)_p^\wedge \simeq \Delta_{\widehat{R}_p[\zeta_p]}/(\mathbb{Z}_p[[q-1]], [p]_q)$ .
- 3 Rationally,  $(q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)}^\wedge \simeq (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q})[[q-1]]$ ,  
 $(q\text{-}\Omega_R)_p^\wedge \left[\frac{1}{p}\right]_{(q-1)}^\wedge \simeq (\Omega_R^*)_p^\wedge \left[\frac{1}{p}\right] [[q-1]]$ .



Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , ...) by analysing the cohomology of the associated geometric objects.

Many cohomology theories:





**Habiro cohomology** (Scholze's expectations). Should have coefficients not in  $\mathbb{Z}[[q-1]]$ , but in the *Habiro ring*



**Habiro cohomology** (Scholze's expectations). Should have coefficients not in  $\mathbb{Z}[[q-1]]$ , but in the *Habiro ring*

$$\mathcal{H} := \mathbb{Z}[q]$$



**Habiro cohomology** (Scholze's expectations). Should have coefficients not in  $\mathbb{Z}[\![q-1]\!]$ , but in the *Habiro ring*

$$\mathcal{H} := \mathbb{Z}[q]_{(q^m-1)}^\wedge$$



**Habiro cohomology** (Scholze's expectations). Should have coefficients not in  $\mathbb{Z}[[q-1]]$ , but in the *Habiro ring*

$$\mathcal{H} := \lim_{m \in \mathbb{N}} \mathbb{Z}[q]_{(q^m-1)}^\wedge.$$



**Habiro cohomology** (Scholze's expectations). Should have coefficients not in  $\mathbb{Z}[[q-1]]$ , but in the *Habiro ring*

$$\mathcal{H} := \lim_{m \in \mathbb{N}} \mathbb{Z}[q]_{(q^m-1)}^{\wedge}.$$

Moreover:

- 1 At  $q = 1$ , should get Hodge cohomology (*not* de Rham).





**Habiro cohomology** (Scholze's expectations). Should have coefficients not in  $\mathbb{Z}[[q-1]]$ , but in the *Habiro ring*

$$\mathcal{H} := \lim_{m \in \mathbb{N}} \mathbb{Z}[q]_{(q^m-1)}^\wedge.$$

Moreover:

- 1 At  $q = 1$ , should get Hodge cohomology (*not* de Rham).
- 2 Should recover prismatic cohomology for all primes  $p$ .



**Habiro cohomology** (Scholze's expectations). Should have coefficients not in  $\mathbb{Z}[[q-1]]$ , but in the *Habiro ring*

$$\mathcal{H} := \lim_{m \in \mathbb{N}} \mathbb{Z}[q]_{(q^m-1)}^{\wedge}.$$

Moreover:

- 1 At  $q = 1$ , should get Hodge cohomology (*not* de Rham).
- 2 Should recover prismatic cohomology for all primes  $p$ .
- 3 Relation to quantum invariants of 3-manifolds?



**Habiro cohomology** (Scholze's expectations). Should have coefficients not in  $\mathbb{Z}[[q-1]]$ , but in the *Habiro ring*

$$\mathcal{H} := \lim_{m \in \mathbb{N}} \mathbb{Z}[q]_{(q^m-1)}^{\wedge}.$$


Moreover:

- 1 At  $q = 1$ , should get Hodge cohomology (*not* de Rham).
- 2 Should recover prismatic cohomology for all primes  $p$ .
- 3 Relation to quantum invariants of 3-manifolds? E.g. *Habiro ring of a number field* (Garoufalidis–Scholze–Wheeler–Zagier).



Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , ...) by analysing the cohomology of the associated geometric objects.


Many cohomology theories:





Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , ...) by analysing the cohomology of the associated geometric objects.


Many cohomology theories:





Study equations over  $\mathbb{Z}$  or  $\mathbb{Q}$  (or  $\mathbb{F}_p$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , ...) by analysing the cohomology of the associated geometric objects.

Many cohomology theories:





**Habiro cohomology** (Scholze's expectations). Should have coefficients not in  $\mathbb{Z}[[q-1]]$ , but in the *Habiro ring*

$$\mathcal{H} := \lim_{m \in \mathbb{N}} \mathbb{Z}[q]_{(q^m-1)}^\wedge.$$

Moreover:

- 1 At  $q = 1$ , should get Hodge cohomology (*not* de Rham).
- 2 Should recover prismatic cohomology for all primes  $p$ .
- 3 Relation to quantum invariants of 3-manifolds? E.g. *Habiro ring of a number field* (Garoufalidis–Scholze–Wheeler–Zagier).



**Habiro cohomology** (Scholze's expectations). Should have coefficients not in  $\mathbb{Z}[[q-1]]$ , but in the *Habiro ring*

$$\mathcal{H} := \lim_{m \in \mathbb{N}} \mathbb{Z}[q]_{(q^m-1)}^\wedge.$$

Moreover:

- 1 At  $q = 1$ , should get Hodge cohomology (*not* de Rham).
- 2 Should recover prismatic cohomology for all primes  $p$ .
- 3 Relation to quantum invariants of 3-manifolds? E.g. *Habiro ring of a number field* (Garoufalidis–Scholze–Wheeler–Zagier).

⇒ **We should look for  $q$ -deformations of the Hodge filtration!**



## 2. $q$ -Hodge filtrations & Habiro cohomology

---



**$q$ -Hodge filtrations.** Let  $\mathrm{Sm}_{\mathbb{Z}}^{q\text{-Hdg}}$  be the category of pairs

$$\left( R, \quad \mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \right)$$



**$q$ -Hodge filtrations.** Let  $\mathrm{Sm}_{\mathbb{Z}}^{q\text{-Hdg}}$  be the category of pairs

$$\left( \begin{array}{c} R \\ \text{smooth over } \mathbb{Z} \end{array}, \quad \mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \right)$$



**$q$ -Hodge filtrations.** Let  $\mathrm{Sm}_{\mathbb{Z}}^{q\text{-Hdg}}$  be the category of pairs

$$\left( \begin{array}{c} R \\ \text{smooth over } \mathbb{Z} \end{array}, \begin{array}{c} \text{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \\ \text{filtered } (q-1)^*\mathbb{Z}[[q-1]]\text{-module} \end{array} \right)$$



**$q$ -Hodge filtrations.** Let  $\mathrm{Sm}_{\mathbb{Z}}^{q\text{-Hdg}}$  be the category of pairs

$$\left( \begin{array}{c} R \\ \text{smooth over } \mathbb{Z} \end{array}, \begin{array}{c} \mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \\ \text{filtered } (q-1)^*\mathbb{Z}[[q-1]]\text{-module} \end{array} \right)$$

such that:

$$\boxed{1} \quad \mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R / (q-1) \simeq \mathrm{fil}_{\mathrm{Hdg}}^* \Omega_R^*.$$



**$q$ -Hodge filtrations.** Let  $\mathrm{Sm}_{\mathbb{Z}}^{q\text{-Hdg}}$  be the category of pairs

$$\left( \begin{array}{c} R \\ \text{smooth over } \mathbb{Z} \end{array}, \begin{array}{c} \mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \\ \text{filtered } (q-1)^*\mathbb{Z}[[q-1]]\text{-module} \end{array} \right)$$

such that:

- 1  $\mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R / (q-1) \simeq \mathrm{fil}_{\mathrm{Hdg}}^* \Omega_R^*.$
- 2  $(\mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)}^{\wedge} \simeq \mathrm{fil}_{(\mathrm{Hdg}, q-1)}^* (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q})[[q-1]]$



**$q$ -Hodge filtrations.** Let  $\mathrm{Sm}_{\mathbb{Z}}^{q\text{-Hdg}}$  be the category of pairs

$$\left( \begin{array}{c} R \\ \text{smooth over } \mathbb{Z} \end{array}, \begin{array}{c} \mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \\ \text{filtered } (q-1)^*\mathbb{Z}[[q-1]]\text{-module} \end{array} \right)$$

such that:

- 1  $\mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R / (q-1) \simeq \mathrm{fil}_{\mathrm{Hdg}}^* \Omega_R^*.$
- 2  $(\mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)}^{\wedge} \simeq \mathrm{fil}_{(\mathrm{Hdg}, q-1)}^* (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q})[[q-1]]$

$$\left( \begin{array}{c} \text{Recall: } (q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)}^{\wedge} \simeq (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q})[[q-1]]. \end{array} \right)$$



**$q$ -Hodge filtrations.** Let  $\mathrm{Sm}_{\mathbb{Z}}^{q\text{-Hdg}}$  be the category of pairs

$$\left( \begin{array}{c} R \\ \text{smooth over } \mathbb{Z} \end{array}, \begin{array}{c} \mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \\ \text{filtered } (q-1)^*\mathbb{Z}[[q-1]]\text{-module} \end{array} \right)$$

such that:

- 1  $\mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R / (q-1) \simeq \mathrm{fil}_{\mathrm{Hdg}}^* \Omega_R^*.$
- 2  $(\mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)}^{\wedge} \simeq \mathrm{fil}_{(\mathrm{Hdg}, q-1)}^*(\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q})[[q-1]]$

$$\left( \begin{array}{c} \text{Recall: } (q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)}^{\wedge} \simeq (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q})[[q-1]]. \end{array} \right)$$





**$q$ -Hodge filtrations.** Let  $\mathrm{Sm}_{\mathbb{Z}}^{q\text{-Hdg}}$  be the category of pairs

$$\left( \begin{array}{c} R \\ \text{smooth over } \mathbb{Z} \end{array}, \begin{array}{c} \mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \\ \text{filtered } (q-1)^*\mathbb{Z}[[q-1]]\text{-module} \end{array} \right)$$

such that:

- 1  $\mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R / (q-1) \simeq \mathrm{fil}_{\mathrm{Hdg}}^* \Omega_R^*.$
- 2  $(\mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)}^{\wedge} \simeq \mathrm{fil}_{(\mathrm{Hdg}, q-1)}^* (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q})[[q-1]]$

$$\left( \begin{array}{c} \text{Recall: } (q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)}^{\wedge} \simeq (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q})[[q-1]]. \end{array} \right)$$



**$q$ -Hodge filtrations.** Let  $\mathrm{Sm}_{\mathbb{Z}}^{q\text{-Hdg}}$  be the category of pairs

$$\left( \begin{array}{c} R \\ \text{smooth over } \mathbb{Z} \end{array}, \begin{array}{c} \mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \\ \text{filtered } (q-1)^*\mathbb{Z}[[q-1]]\text{-module} \end{array} \right)$$

such that:

- 1  $\mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R / (q-1) \simeq \mathrm{fil}_{\mathrm{Hdg}}^* \Omega_R^*.$
- 2  $(\mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)}^{\wedge} \simeq \mathrm{fil}_{(\mathrm{Hdg}, q-1)}^* (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q})[[q-1]],$   
 $\mathrm{fil}_{q\text{-Hdg}}^* (q\text{-}\Omega_R)_p^{\wedge} \left[ \frac{1}{p} \right]_{(q-1)}^{\wedge} \simeq \mathrm{fil}_{(\mathrm{Hdg}, q-1)}^* (\Omega_R^*)_p^{\wedge} \left[ \frac{1}{p} \right] [[q-1]]$  for all  $p$ .



**$q$ -Hodge filtrations.** Let  $\mathrm{Sm}_{\mathbb{Z}}^{q\text{-Hdg}}$  be the category of pairs

$$\left( \begin{array}{c} R \\ \text{smooth over } \mathbb{Z} \end{array}, \begin{array}{c} \mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \\ \text{filtered } (q-1)^*\mathbb{Z}[[q-1]]\text{-module} \end{array} \right)$$

such that:

- 1  $\mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R / (q-1) \simeq \mathrm{fil}_{\mathrm{Hdg}}^* \Omega_R^*.$
- 2  $(\mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)}^{\wedge} \simeq \mathrm{fil}_{(\mathrm{Hdg}, q-1)}^* (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q})[[q-1]],$   
 $\mathrm{fil}_{q\text{-Hdg}}^* (q\text{-}\Omega_R)_p^{\wedge} \left[ \frac{1}{p} \right]_{(q-1)}^{\wedge} \simeq \mathrm{fil}_{(\mathrm{Hdg}, q-1)}^* (\Omega_R^*)_p^{\wedge} \left[ \frac{1}{p} \right] [[q-1]]$  for all  $p$ .

**Theorem (W. 2025).**

The forgetful functor  $\mathrm{Sm}_{\mathbb{Z}}^{q\text{-Hdg}} \rightarrow \mathrm{Sm}_{\mathbb{Z}}$  has *no* section



**$q$ -Hodge filtrations.** Let  $\mathrm{Sm}_{\mathbb{Z}}^{q\text{-Hdg}}$  be the category of pairs

$$\left( \begin{array}{c} R \\ \text{smooth over } \mathbb{Z} \end{array}, \begin{array}{c} \mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \\ \text{filtered } (q-1)^*\mathbb{Z}[[q-1]]\text{-module} \end{array} \right)$$

such that:

- 1  $\mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R / (q-1) \simeq \mathrm{fil}_{\mathrm{Hdg}}^* \Omega_R^*.$
- 2  $(\mathrm{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)}^{\wedge} \simeq \mathrm{fil}_{(\mathrm{Hdg}, q-1)}^* (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q})[[q-1]],$   
 $\mathrm{fil}_{q\text{-Hdg}}^* (q\text{-}\Omega_R)_p^{\wedge} \left[ \frac{1}{p} \right]_{(q-1)}^{\wedge} \simeq \mathrm{fil}_{(\mathrm{Hdg}, q-1)}^* (\Omega_R^*)_p^{\wedge} \left[ \frac{1}{p} \right] [[q-1]]$  for all  $p$ .

**Theorem (W. 2025).**

The forgetful functor  $\mathrm{Sm}_{\mathbb{Z}}^{q\text{-Hdg}} \rightarrow \mathrm{Sm}_{\mathbb{Z}}$  has *no* section, but over the full subcategory

$$\{R \mid p \in R^{\times} \text{ for all primes } p \leq \dim(R/\mathbb{Z})\} \subseteq \mathrm{Sm}_{\mathbb{Z}}$$

there exists a section.



**The  $q$ -Hodge complex.** Given  $(R, \text{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R)$ , define the  $q$ -Hodge complex

$$q\text{-Hdg}_R :=$$



**The  $q$ -Hodge complex.** Given  $(R, \operatorname{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R)$ , define the  $q$ -Hodge complex

$$q\text{-Hdg}_R := q\text{-}\Omega_R$$



**The  $q$ -Hodge complex.** Given  $(R, \text{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R)$ , define the  $q$ -Hodge complex

$$q\text{-Hdg}_R := q\text{-}\Omega_R \left[ \frac{\text{fil}_{q\text{-Hdg}}^n}{(q-1)^n} \mid n \geq 1 \right]_{(q-1)}^{\wedge}.$$



**The  $q$ -Hodge complex.** Given  $(R, \text{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R)$ , define the  $q$ -Hodge complex

$$q\text{-Hdg}_R := q\text{-}\Omega_R \left[ \frac{\text{fil}_{q\text{-Hdg}}^n}{(q-1)^n} \mid n \geq 1 \right]_{(q-1)}^{\wedge}.$$





**The  $q$ -Hodge complex.** Given  $(R, \text{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R)$ , define the  $q$ -Hodge complex

$$q\text{-Hdg}_R := q\text{-}\Omega_R \left[ \frac{\text{fil}_{q\text{-Hdg}}^n}{(q-1)^n} \mid n \geq 1 \right]_{(q-1)}^{\wedge}.$$

**Theorem** (W. 2025).

- 1 There exists a non-trivial factorisation

$$\text{Sm}_{\mathbb{Z}}^{q\text{-Hdg}} \xrightarrow{q\text{-Hdg}_{(-)}} \mathcal{D}(\mathbb{Z}[[q-1]])$$



**The  $q$ -Hodge complex.** Given  $(R, \text{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R)$ , define the  $q$ -Hodge complex

$$q\text{-Hdg}_R := q\text{-}\Omega_R \left[ \frac{\text{fil}_{q\text{-Hdg}}^n}{(q-1)^n} \mid n \geq 1 \right]_{(q-1)}^{\wedge}.$$

**Theorem** (W. 2025).

1 There exists a non-trivial factorisation

$$\begin{array}{ccc} & & \mathcal{D}(\mathcal{H}) \\ & & \downarrow (-)_{(q-1)}^{\wedge} \\ \text{Sm}_{\mathbb{Z}}^{q\text{-Hdg}} & \xrightarrow{q\text{-Hdg}_{(-)}} & \mathcal{D}(\mathbb{Z}[[q-1]]) \end{array}$$



**The  $q$ -Hodge complex.** Given  $(R, \text{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R)$ , define the  $q$ -Hodge complex

$$q\text{-Hdg}_R := q\text{-}\Omega_R \left[ \frac{\text{fil}_{q\text{-Hdg}}^n}{(q-1)^n} \mid n \geq 1 \right]_{(q-1)}^\wedge.$$

**Theorem (W. 2025).**

- 1 There exists a non-trivial **factorisation**

$$\begin{array}{ccc} & & \mathcal{D}(\mathcal{H}) \\ & \nearrow \text{Habiro–Hodge complex } q\text{-}\mathcal{H}\text{dg}_{(-)} & \downarrow (-)_{(q-1)}^\wedge \\ \text{Sm}_{\mathbb{Z}}^{q\text{-Hdg}} & \xrightarrow{q\text{-Hdg}_{(-)}} & \mathcal{D}(\mathbb{Z}[[q-1]]) \end{array}$$



**The  $q$ -Hodge complex.** Given  $(R, \text{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R)$ , define the  $q$ -Hodge complex

$$q\text{-Hdg}_R := q\text{-}\Omega_R \left[ \frac{\text{fil}_{q\text{-Hdg}}^n}{(q-1)^n} \mid n \geq 1 \right]_{(q-1)}^\wedge.$$

**Theorem** (W. 2025).

- 1 There exists a non-trivial factorisation

$$\begin{array}{ccc} & & \mathcal{D}(\mathcal{H}) \\ & \nearrow \text{Habiro–Hodge complex } q\text{-}\mathcal{H}\text{dg}_{(-)} & \downarrow (-)_{(q-1)}^\wedge \\ \text{Sm}_{\mathbb{Z}}^{q\text{-Hdg}} & \xrightarrow{q\text{-Hdg}_{(-)}} & \mathcal{D}(\mathbb{Z}[[q-1]]) \end{array}$$

- 2  $H^*(q\text{-}\mathcal{H}\text{dg}_R/(q^m-1)) \cong q\text{-}\mathbb{W}_m\Omega_R^*.$



**The  $q$ -Hodge complex.** Given  $(R, \text{fil}_{q\text{-Hdg}}^* q\text{-}\Omega_R)$ , define the  $q$ -Hodge complex

$$q\text{-Hdg}_R := q\text{-}\Omega_R \left[ \frac{\text{fil}_{q\text{-Hdg}}^n}{(q-1)^n} \mid n \geq 1 \right]_{(q-1)}^{\wedge}.$$

**Theorem (W. 2025).**

1 There exists a non-trivial factorisation

$$\begin{array}{ccc} & & \mathcal{D}(\mathcal{H}) \\ \text{Habiro–Hodge complex } q\text{-}\mathcal{H}\text{dg}_{(-)} & \nearrow & \downarrow (-)_{(q-1)}^{\wedge} \\ \text{Sm}_{\mathbb{Z}}^{q\text{-Hdg}} & \xrightarrow{q\text{-Hdg}_{(-)}} & \mathcal{D}(\mathbb{Z}[[q-1]]) \end{array}$$

2  $H^*(q\text{-}\mathcal{H}\text{dg}_R/(q^m-1)) \cong q\text{-}\mathbb{W}_m\Omega_R^*$ . In particular,

$$H^*(q\text{-}\mathcal{H}\text{dg}_R/(q-1)) \cong \Omega_R^*.$$



## Theorem (continued)

$$3 \quad q\text{-}\Omega_R \simeq L_{\eta(q-1)}(q\text{-Hdg}_R).$$



## Theorem (continued)

- 3  $q\text{-}\Omega_R \simeq \mathbf{L}_{\eta(q-1)}(q\text{-Hdg}_R).$
- 4 The *Habiro ring of a number field*  $F$  (Garoufalidis–Scholze–Wheeler–Zagier) can be recovered as

$$q\text{-}\mathcal{H}\text{dg}_{\mathcal{O}_F[1/\Delta]} \simeq \mathcal{H}_{\mathcal{O}_F[1/\Delta]}$$

( $\Delta$  = discriminant of  $F$ ).



## Theorem (continued)

- 3  $q\text{-}\Omega_R \simeq L_{\eta(q-1)}(q\text{-Hdg}_R)$ .
- 4 The *Habiro ring of a number field*  $F$  (Garoufalidis–Scholze–Wheeler–Zagier) can be recovered as

$$q\text{-Hdg}_{\mathcal{O}_F[1/\Delta]} \simeq \mathcal{H}_{\mathcal{O}_F[1/\Delta]}$$

( $\Delta$  = discriminant of  $F$ ).

If  $X$  is a smooth scheme over  $\mathbb{Z}$ , up to inverting all primes  $p \leq \dim(X/\mathbb{Z})$ , one can define *algebraic Habiro cohomology of  $X$*  as

$$R\Gamma_{\mathcal{H}}(X) := R\Gamma(X, q\text{-Hdg}_X).$$





## Theorem (continued)

- 3  $q\text{-}\Omega_R \simeq L_{\eta(q-1)}(q\text{-Hdg}_R)$ .
- 4 The *Habiro ring of a number field*  $F$  (Garoufalidis–Scholze–Wheeler–Zagier) can be recovered as

$$q\text{-Hdg}_{\mathcal{O}_F[1/\Delta]} \simeq \mathcal{H}_{\mathcal{O}_F[1/\Delta]}$$

( $\Delta$  = discriminant of  $F$ ).

If  $X$  is a smooth scheme over  $\mathbb{Z}$ , up to inverting all primes  $p \leq \dim(X/\mathbb{Z})$ , one can define *algebraic Habiro cohomology of  $X$*  as

$$R\Gamma_{\mathcal{H}}(X) := R\Gamma(X, q\text{-Hdg}_X).$$



Probably no six functor formalism/stacky approach.



## Theorem (continued)

- 3  $q\text{-}\Omega_R \simeq L_{\eta(q-1)}(q\text{-Hdg}_R)$ .
- 4 The *Habiro ring of a number field*  $F$  (Garoufalidis–Scholze–Wheeler–Zagier) can be recovered as

$$q\text{-Hdg}_{\mathcal{O}_F[1/\Delta]} \simeq \mathcal{H}_{\mathcal{O}_F[1/\Delta]}$$

( $\Delta$  = discriminant of  $F$ ).

If  $X$  is a smooth scheme over  $\mathbb{Z}$ , up to inverting all primes  $p \leq \dim(X/\mathbb{Z})$ , one can define *algebraic Habiro cohomology of  $X$*  as

$$R\Gamma_{\mathcal{H}}(X) := R\Gamma(X, q\text{-Hdg}_X).$$



Probably no six functor formalism/stacky approach. But there should be cohomology classes from quantum invariants (Garoufalidis–Wheeler).

### 3. $q$ -Hodge filtrations from homotopy theory

---



**Recall:** For an associative ring  $R$ , *Hochschild homology* of  $R$  is

$$HH(R) := \left| \quad \text{cyclic bar construction} \quad \right| .$$



**Recall:** For an associative ring  $R$ , *Hochschild homology* of  $R$  is

$$\mathrm{HH}(R) := \left| R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} R \otimes_{\mathbb{Z}}^L R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} R \otimes_{\mathbb{Z}}^L R \otimes_{\mathbb{Z}}^L R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} \cdots \right|.$$



**Recall:** For an associative ring  $R$ , *Hochschild homology* of  $R$  is

$$\mathrm{HH}(R) := \left| R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} R \otimes_{\mathbb{Z}}^L R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} R \otimes_{\mathbb{Z}}^L R \otimes_{\mathbb{Z}}^L R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} \cdots \right|.$$

There's an action  $S^1 \curvearrowright \mathrm{HH}(R)$ , put  $\mathrm{HC}^-(R) := \mathrm{HH}(R)^{hS^1}$  (Connes).



**Recall:** For an associative ring  $R$ , *Hochschild homology* of  $R$  is

$$\mathrm{HH}(R) := \left| R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} R \otimes_{\mathbb{Z}}^{\mathbb{L}} R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} R \otimes_{\mathbb{Z}}^{\mathbb{L}} R \otimes_{\mathbb{Z}}^{\mathbb{L}} R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} \cdots \right|.$$

There's an action  $S^1 \curvearrowright \mathrm{HH}(R)$ , put  $\mathrm{HC}^-(R) := \mathrm{HH}(R)^{hS^1}$  (Connes).

**Theorem** (... , Antieau).



**Recall:** For an associative ring  $R$ , *Hochschild homology* of  $R$  is

$$\mathrm{HH}(R) := \left| R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} R \otimes_{\mathbb{Z}}^L R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} R \otimes_{\mathbb{Z}}^L R \otimes_{\mathbb{Z}}^L R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} \cdots \right|.$$

There's an action  $S^1 \curvearrowright \mathrm{HH}(R)$ , put  $\mathrm{HC}^-(R) := \mathrm{HH}(R)^{hS^1}$  (Connes).

**Theorem** (... , Antieau).

If  $R$  is quasisyntomic, there is a complete exhaustive “motivic” filtration  $\mathrm{fil}_{\mathrm{mot}}^* \mathrm{HC}^-(R)$  with associated graded

$$\mathrm{gr}_{\mathrm{mot}}^i \mathrm{HC}^-(R) \simeq \mathrm{fil}_{\mathrm{Hdg}}^i \mathrm{dR}_R$$





**Recall:** For an associative ring  $R$ , *Hochschild homology* of  $R$  is

$$\mathrm{HH}(R) := \left| R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} R \otimes_{\mathbb{Z}}^{\mathbb{L}} R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} R \otimes_{\mathbb{Z}}^{\mathbb{L}} R \otimes_{\mathbb{Z}}^{\mathbb{L}} R \begin{smallmatrix} \leftarrow \\ \leftarrow \\ \leftarrow \end{smallmatrix} \cdots \right|.$$

There's an action  $S^1 \curvearrowright \mathrm{HH}(R)$ , put  $\mathrm{HC}^-(R) := \mathrm{HH}(R)^{hS^1}$  (Connes).

**Theorem** (... , Antieau).

If  $R$  is quasisyntomic, there is a complete exhaustive “motivic” filtration  $\mathrm{fil}_{\mathrm{mot}}^* \mathrm{HC}^-(R)$  with associated graded

$$\mathrm{gr}_{\mathrm{mot}}^i \mathrm{HC}^-(R) \simeq \mathrm{fil}_{\mathrm{Hdg}}^i \mathbf{dR}_R$$

(where  $\mathbf{dR}_{(-)}$  denotes the non-abelian derived functor of  $\Omega_{(-)}^*$ ).



**Recall:** For an associative ring  $R$ , *Hochschild homology* of  $R$  is

$$\mathrm{HH}(R) := \left| R \overset{\mathbb{L}}{\leftarrow} R \otimes_{\mathbb{Z}}^{\mathbb{L}} R \overset{\mathbb{L}}{\leftarrow} R \otimes_{\mathbb{Z}}^{\mathbb{L}} R \otimes_{\mathbb{Z}}^{\mathbb{L}} R \overset{\mathbb{L}}{\leftarrow} \cdots \right|.$$

There's an action  $S^1 \curvearrowright \mathrm{HH}(R)$ , put  $\mathrm{HC}^-(R) := \mathrm{HH}(R)^{hS^1}$  (Connes).

**Theorem** (... , Antieau).

If  $R$  is quasisyntomic, there is a complete exhaustive “motivic” filtration  $\mathrm{fil}_{\mathrm{mot}}^* \mathrm{HC}^-(R)$  with associated graded

$$\mathrm{gr}_{\mathrm{mot}}^i \mathrm{HC}^-(R) \simeq \mathrm{fil}_{\mathrm{Hdg}}^i \widehat{\mathrm{dR}}_R[2i]$$

(where  $\mathrm{dR}_{(-)}$  denotes the non-abelian derived functor of  $\Omega_{(-)}^*$ ).



**Even filtration** (Hahn–Raksit–Wilson). The motivic filtration agrees with the *even filtration* on  $HC^-(R)$ :



**Even filtration** (Hahn–Raksit–Wilson). The motivic filtration agrees with the *even filtration* on  $HC^-(R)$ :

$$\mathrm{fil}_{\mathrm{ev}}^* \mathrm{HH}(R) \quad := \quad \lim_{\substack{\mathrm{HH}(R) \rightarrow E \text{ even} \\ \mathbb{E}_\infty}} \tau_{\geq 2\star} ( E ).$$



**Even filtration** (Hahn–Raksit–Wilson). The motivic filtration agrees with the *even filtration* on  $HC^-(R)$ :

$$\mathrm{fil}_{\mathrm{ev}}^* \mathrm{HH}(R) \quad := \quad \lim_{\substack{\mathrm{HH}(R) \rightarrow E \text{ even} \\ \mathbb{E}_\infty}} \tau_{\geq 2\star} (E).$$



**Even filtration** (Hahn–Raksit–Wilson). The motivic filtration agrees with the *even filtration* on  $HC^-(R)$ :

$$\mathrm{fil}_{\mathrm{ev}, hS^1}^* HC^-(R) := \lim_{\substack{HH(R) \rightarrow E \text{ even} \\ S^1 \text{ equivariant } \mathbb{E}_\infty}} \tau_{\geq 2\star}(E^{hS^1}).$$



**Even filtration** (Hahn–Raksit–Wilson). The motivic filtration agrees with the *even filtration* on  $HC^-(R)$ :

$$\mathrm{fil}_{\mathrm{ev}, hS^1}^* HC^-(R) := \lim_{\substack{HH(R) \rightarrow E \text{ even} \\ S^1 \text{ equivariant } \mathbb{E}_\infty}} \tau_{\geq 2\star}(E^{hS^1}).$$

The even filtration works equally well for  $HC^-(-/ku)$ .



**Even filtration** (Hahn–Raksit–Wilson). The motivic filtration agrees with the *even filtration* on  $\mathrm{HC}^-(R)$ :

$$\mathrm{fil}_{\mathrm{ev}, hS^1}^* \mathrm{HC}^-(R) := \lim_{\substack{\mathrm{HH}(R) \rightarrow E \text{ even} \\ S^1 \text{ equivariant } \mathbb{E}_\infty}} \tau_{\geq 2\star}(E^{hS^1}).$$

The even filtration works equally well for  $\mathrm{HC}^-(-/\text{connective complex } K\text{-theory } \mathbf{ku})$ .





**Even filtration** (Hahn–Raksit–Wilson). The motivic filtration agrees with the *even filtration* on  $\mathrm{HC}^-(R)$ :

$$\mathrm{fil}_{\mathrm{ev}, hS^1}^* \mathrm{HC}^-(R) := \lim_{\substack{\mathrm{HH}(R) \rightarrow E \text{ even} \\ S^1 \text{ equivariant } \mathbb{E}_\infty}} \tau_{\geq 2\star}(E^{hS^1}).$$

The even filtration works equally well for  $\mathrm{TC}^-(\text{connective complex } K\text{-theory}/\mathrm{ku})$ .



**Even filtration** (Hahn–Raksit–Wilson). The motivic filtration agrees with the *even filtration* on  $HC^-(R)$ :

$$\mathrm{fil}_{\mathrm{ev}, hS^1}^* HC^-(R) := \lim_{\substack{HH(R) \rightarrow E \text{ even} \\ S^1 \text{ equivariant } \mathbb{E}_\infty}} \tau_{\geq 2\star}(E^{hS^1}).$$

The even filtration works equally well for TC<sup>-</sup>(-/ku)<sup>connective complex K-theory</sup>.

There's a version for  $\mathbb{E}_1$ -ring spectra too (Pstrągowski).



**Even filtration** (Hahn–Raksit–Wilson). The motivic filtration agrees with the *even filtration* on  $HC^-(R)$ :

$$\mathrm{fil}_{\mathrm{ev}, hS^1}^* HC^-(R) := \lim_{\substack{HH(R) \rightarrow E \text{ even} \\ S^1 \text{ equivariant } \mathbb{E}_\infty}} \tau_{\geq 2\star}(E^{hS^1}).$$

The even filtration works equally well for TC<sup>-</sup>(-/ku). connective complex K-theory

There's a version for  $\mathbb{E}_1$ -ring spectra too (Pstrągowski).

**Theorem** (W. 2025).

Let  $R$  be quasisyntomic,

(Raksit proved this first for  $\mathbb{S}[x]$ ; closely related work of Devalapurkar)



**Even filtration** (Hahn–Raksit–Wilson). The motivic filtration agrees with the *even filtration* on  $HC^-(R)$ :

$$\mathrm{fil}_{\mathrm{ev}, hS^1}^* HC^-(R) := \lim_{\substack{HH(R) \rightarrow E \text{ even} \\ S^1 \text{ equivariant } \mathbb{E}_\infty}} \tau_{\geq 2\star}(E^{hS^1}).$$

The even filtration works equally well for TC<sup>-</sup>(-/ku). connective complex K-theory

There's a version for  $\mathbb{E}_1$ -ring spectra too (Pstrągowski).

**Theorem** (W. 2025).

Let  $R$  be quasisyntomic,  $2 \in R^\times$ .

(Raksit proved this first for  $\mathbb{S}[x]$ ; closely related work of Devalapurkar)



**Even filtration** (Hahn–Raksit–Wilson). The motivic filtration agrees with the *even filtration* on  $\mathrm{HC}^-(R)$ :

$$\mathrm{fil}_{\mathrm{ev}, hS^1}^* \mathrm{HC}^-(R) := \lim_{\substack{\mathrm{HH}(R) \rightarrow E \text{ even} \\ S^1 \text{ equivariant } \mathbb{E}_\infty}} \tau_{\geq 2\star}(E^{hS^1}).$$

The even filtration works equally well for  $\mathrm{TC}^-(\text{connective complex } K\text{-theory}/\mathrm{ku})$ .

There's a version for  $\mathbb{E}_1$ -ring spectra too (Pstrągowski).

## Theorem (W. 2025).

Let  $R$  be quasisyntomic,  $2 \in R^\times$ . Suppose  $R$  lifts to an  $\mathbb{E}_2$ -ring spectrum  $\mathbb{S}_R$  such that  $\mathbb{S}_R \otimes \mathbb{Z} \simeq R$ .

(Raksit proved this first for  $\mathbb{S}[x]$ ; closely related work of Devalapurkar)



**Even filtration** (Hahn–Raksit–Wilson). The motivic filtration agrees with the *even filtration* on  $\mathrm{HC}^-(R)$ :

$$\mathrm{fil}_{\mathrm{ev}, hS^1}^* \mathrm{HC}^-(R) := \lim_{\substack{\mathrm{HH}(R) \rightarrow E \text{ even} \\ S^1 \text{ equivariant } \mathbb{E}_\infty}} \tau_{\geq 2*}(E^{hS^1}).$$

The even filtration works equally well for  $\mathrm{TC}^-(\text{connective complex } K\text{-theory}/\mathrm{ku})$ .

There's a version for  $\mathbb{E}_1$ -ring spectra too (Pstrągowski).

## Theorem (W. 2025).

Let  $R$  be quasisyntomic,  $2 \in R^\times$ . Suppose  $R$  lifts to an  $\mathbb{E}_2$ -ring spectrum  $\mathbb{S}_R$  such that  $\mathbb{S}_R \otimes \mathbb{Z} \simeq R$ . Then there exists a  $q$ -Hodge filtration  $\mathrm{fil}_{q\text{-Hdg}}^* q\text{-d}R_R$  such that

$$\mathrm{gr}_{\mathrm{ev}, hS^1}^i \mathrm{TC}^-(\mathrm{ku} \otimes \mathbb{S}_R/\mathrm{ku}) \simeq \mathrm{fil}_{q\text{-Hdg}}^i q\text{-d}R_R \quad .$$

(Raksit proved this first for  $\mathbb{S}[x]$ ; closely related work of Devalapurkar)



**Even filtration** (Hahn–Raksit–Wilson). The motivic filtration agrees with the *even filtration* on  $\mathrm{HC}^-(R)$ :

$$\mathrm{fil}_{\mathrm{ev}, hS^1}^* \mathrm{HC}^-(R) := \lim_{\substack{\mathrm{HH}(R) \rightarrow E \text{ even} \\ S^1 \text{ equivariant } \mathbb{E}_\infty}} \tau_{\geq 2*}(E^{hS^1}).$$

The even filtration works equally well for  $\mathrm{TC}^-(\text{connective complex } K\text{-theory}/\mathrm{ku})$ .

There's a version for  $\mathbb{E}_1$ -ring spectra too (Pstrągowski).

## Theorem (W. 2025).

Let  $R$  be quasisyntomic,  $2 \in R^\times$ . Suppose  $R$  lifts to an  $\mathbb{E}_2$ -ring spectrum  $\mathbb{S}_R$  such that  $\mathbb{S}_R \otimes \mathbb{Z} \simeq R$ . Then there exists a  $q$ -Hodge filtration  $\mathrm{fil}_{q\text{-Hdg}}^* q\text{-d}R_R$  such that

$$\mathrm{gr}_{\mathrm{ev}, hS^1}^i \mathrm{TC}^-(\mathrm{ku} \otimes \mathbb{S}_R/\mathrm{ku}) \simeq \mathrm{fil}_{q\text{-Hdg}}^i q\text{-}\widehat{\mathrm{d}R}_R[2i].$$

(Raksit proved this first for  $\mathbb{S}[x]$ ; closely related work of Devalapurkar)



**Theorem** (Antieau, Hahn–Raksit–Wilson).

Let  $R$  be quasisyntomic. Then

$$gr_{ev, hS^1}^i HC^-(R) \simeq fil_{Hdg}^i \widehat{dR}_R[2i].$$

$$\begin{array}{ccc} \text{deformation } ku \rightarrow \mathbb{Z} & \uparrow & \text{deformation } q\text{-}dR_R \rightarrow dR_R \end{array}$$

**Theorem** (W. 2025).

Let  $R$  be quasisyntomic,  $2 \in R^\times$ . Suppose  $R$  lifts to an  $\mathbb{E}_2$ -ring spectrum  $\mathbb{S}_R$  such that  $\mathbb{S}_R \otimes \mathbb{Z} \simeq R$ . Then there exists a  $q$ -Hodge filtration  $fil_{q\text{-}Hdg}^* q\text{-}dR_R$  such that

$$gr_{ev, hS^1}^i TC^-(ku \otimes \mathbb{S}_R/ku) \simeq fil_{q\text{-}Hdg}^i q\text{-}\widehat{dR}_R[2i].$$

(Raksit proved this first for  $\mathbb{S}[x]$ ; closely related work of Devalapurkar)





**Theorem** (W. 2025).

Let  $KU := ku[\beta^{-1}]$  be the periodic complex  $K$ -theory spectrum.



**Theorem** (W. 2025).

Let  $KU := ku[\beta^{-1}]$  be the periodic complex  $K$ -theory spectrum. Then

$$\mathrm{gr}_{\mathrm{ev}, hS^1}^0 \mathrm{TC}^-(KU \otimes \mathbb{S}_R / KU) \simeq q\text{-Hdg}_R .$$



## Theorem (W. 2025).

Let  $KU := ku[\beta^{-1}]$  be the periodic complex  $K$ -theory spectrum. Then

$$\mathrm{gr}_{\mathrm{ev}, hS^1}^0 \mathrm{TC}^-(KU \otimes \mathbb{S}_R / KU) \simeq q\text{-Hdg}_R .$$

The Habiro–Hodge complex is given by

$$\mathrm{THH}(KU \otimes \mathbb{S}_R / KU)$$



**Theorem** (W. 2025).

Let  $KU := ku[\beta^{-1}]$  be the periodic complex  $K$ -theory spectrum. Then

$$\mathrm{gr}_{\mathrm{ev}, hS^1}^0 \mathrm{TC}^-(KU \otimes \mathbb{S}_R / KU) \simeq q\text{-Hdg}_R .$$

The Habiro–Hodge complex is given by

$$\mathrm{THH}(KU \otimes \mathbb{S}_R / KU)^{C_m} ,$$

where  $(-)^{C_m}$  denotes genuine fixed points for  $C_m \subseteq S^1$ .



**Theorem** (W. 2025).

Let  $KU := ku[\beta^{-1}]$  be the periodic complex  $K$ -theory spectrum. Then

$$\mathrm{gr}_{\mathrm{ev}, hS^1}^0 \mathrm{TC}^-(KU \otimes \mathbb{S}_R / KU) \simeq q\text{-Hdg}_R .$$

The Habiro–Hodge complex is given by

$$(\mathrm{THH}(KU \otimes \mathbb{S}_R / KU)^{C_m})^{h(S^1 / C_m)} ,$$

where  $(-)^{C_m}$  denotes genuine fixed points for  $C_m \subseteq S^1$ .



**Theorem** (W. 2025).

Let  $KU := ku[\beta^{-1}]$  be the periodic complex  $K$ -theory spectrum. Then

$$\mathrm{gr}_{\mathrm{ev}, hS^1}^0 \mathrm{TC}^-(KU \otimes \mathbb{S}_R / KU) \simeq q\text{-Hdg}_R .$$

The Habiro–Hodge complex is given by

$$\lim_{m \in \mathbb{N}} (\mathrm{THH}(KU \otimes \mathbb{S}_R / KU)^{C_m})^{h(S^1 / C_m)} ,$$

where  $(-)^{C_m}$  denotes **genuine fixed points** for  $C_m \subseteq S^1$ .



**Theorem** (W. 2025).

Let  $KU := ku[\beta^{-1}]$  be the periodic complex  $K$ -theory spectrum. Then

$$\mathrm{gr}_{\mathrm{ev}, hS^1}^0 \mathrm{TC}^-(KU \otimes \mathbb{S}_R / KU) \simeq q\text{-Hdg}_R .$$

The Habiro–Hodge complex is given by

$$\mathrm{gr}_{\mathrm{ev}, S^1}^0 \left( \lim_{m \in \mathbb{N}} (\mathrm{THH}(KU \otimes \mathbb{S}_R / KU)^{C_m})^{h(S^1/C_m)} \right) \simeq q\text{-}\mathcal{H}\mathrm{dg}_R ,$$

where  $(-)^{C_m}$  denotes genuine fixed points for  $C_m \subseteq S^1$ .



**Theorem** (W. 2025).

Let  $KU := ku[\beta^{-1}]$  be the periodic complex  $K$ -theory spectrum. Then

$$\mathrm{gr}_{\mathrm{ev}, hS^1}^0 \mathrm{TC}^-(KU \otimes \mathbb{S}_R / KU) \simeq q\text{-Hdg}_R .$$

The Habiro–Hodge complex is given by

$$\mathrm{gr}_{\mathrm{ev}, S^1}^0 \left( \underbrace{\lim_{m \in \mathbb{N}} (\mathrm{THH}(KU \otimes \mathbb{S}_R / KU)^{C_m})^{h(S^1/C_m)}}_{\text{topological cyclonic homology } \mathrm{TCn}(KU \otimes \mathbb{S}_R / KU)} \right) \simeq q\text{-}\mathcal{H}\mathrm{dg}_R ,$$

where  $(-)^{C_m}$  denotes genuine fixed points for  $C_m \subseteq S^1$ .



## 4. Refined THH/TC<sup>-</sup> & analytic Habiro cohomology

---



**Problems:**



## Problems:

- 1 To define algebraic Habiro cohomology, need to invert “small” primes or an  $\mathbb{E}_2$ -lift to the sphere spectrum.



## Problems:

- 1 To define algebraic Habiro cohomology, need to invert “small” primes or an  $\mathbb{E}_2$ -lift to the sphere spectrum.

*“Solution”: Work with varieties over  $\mathbb{Q}$ .*



## Problems:

- 1 To define algebraic Habiro cohomology, need to invert “small” primes or an  $\mathbb{E}_2$ -lift to the sphere spectrum.

*“Solution”: Work with varieties over  $\mathbb{Q}$ .*

- 2 Recall:

$$(q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)} \simeq (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q}) \llbracket q-1 \rrbracket.$$



## Problems:

- 1 To define algebraic Habiro cohomology, need to invert “small” primes or an  $\mathbb{E}_2$ -lift to the sphere spectrum.

*“Solution”: Work with varieties over  $\mathbb{Q}$ .*

- 2 Recall:

$$(q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)} \simeq (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q}) \llbracket q-1 \rrbracket.$$

So for varieties over  $\mathbb{Q}$ , we only get de Rham information, no comparison maps to étale/singular cohomology with torsion coefficients.



## Problems:

- 1 To define algebraic Habiro cohomology, need to invert “small” primes or an  $\mathbb{E}_2$ -lift to the sphere spectrum.

*“Solution”:* Work with varieties over  $\mathbb{Q}$ .

- 2 Recall:

$$(q\text{-}\Omega_R \otimes_{\mathbb{Z}} \mathbb{Q})_{(q-1)} \simeq (\Omega_R^* \otimes_{\mathbb{Z}} \mathbb{Q}) \llbracket q-1 \rrbracket.$$

So for varieties over  $\mathbb{Q}$ , we only get de Rham information, no comparison maps to étale/singular cohomology with torsion coefficients.

*Solution:* Refined THH/TC<sup>-</sup>.



**Construction** (Efimov–Scholze). Let  $k$  be an  $\mathbb{E}_\infty$ -ring spectrum.





**Construction** (Efimov–Scholze). Let  $k$  be an  $\mathbb{E}_\infty$ -ring spectrum.

$$\mathrm{Mot}_k^{\mathrm{loc}} \xrightarrow[\otimes]{\mathrm{THH}(-/k)} \mathrm{Mod}_k^{\mathrm{BS}^1}$$



**Construction** (Efimov–Scholze). Let  $k$  be an  $\mathbb{E}_\infty$ -ring spectrum.

$$\text{Mot}_k^{\text{rigid}} \xrightarrow[\otimes]{\text{THH}(-/k)} \text{Mod}_k^{\text{BS}^1}$$



**Construction** (Efimov–Scholze). Let  $k$  be an  $\mathbb{E}_\infty$ -ring spectrum.

$$\begin{array}{ccc} \text{rigid} & & \text{not rigid} \\ \text{Mot}_k^{\text{loc}} & \xrightarrow[\otimes]{\text{THH}(-/k)} & \text{Mod}_k^{\text{BS}^1} \end{array}$$



**Construction** (Efimov–Scholze). Let  $k$  be an  $\mathbb{E}_\infty$ -ring spectrum.

$$\begin{array}{ccc}
 \text{rigid} & & \text{not rigid} \\
 \text{Mot}_k^{\text{loc}} & \xrightarrow[\otimes]{\text{THH}(-/k)} & \text{Mod}_k^{\text{BS}^1} \\
 & \searrow \text{THH}^{\text{ref}}(-/k) & \uparrow \\
 & & (\text{Mod}_k^{\text{BS}^1})^{\text{rig}}
 \end{array}$$



**Construction** (Efimov–Scholze). Let  $k$  be an  $\mathbb{E}_\infty$ -ring spectrum. Suppose  $k$  is complex orientable, choose complex orientation  $t \in \pi_{-2}(k^{hS^1})$ .

$$\begin{array}{ccc}
 \text{rigid} & & \text{not rigid} \\
 \text{Mot}_k^{\text{loc}} & \xrightarrow[\otimes]{\text{THH}(-/k)} & \text{Mod}_k^{BS^1} \\
 & \searrow \text{THH}^{\text{ref}}(-/k) & \uparrow \\
 & & (\text{Mod}_k^{BS^1})^{\text{rig}}
 \end{array}$$



**Construction** (Efimov–Scholze). Let  $k$  be an  $\mathbb{E}_\infty$ -ring spectrum. Suppose  $k$  is complex orientable, choose complex orientation  $t \in \pi_{-2}(k^{hS^1})$ .

$$\begin{array}{ccccc}
 \text{rigid} & & \text{not rigid} & & \\
 \text{Mot}_k^{\text{loc}} & \xrightarrow[\otimes]{\text{THH}(-/k)} & \text{Mod}_k^{BS^1} & \xrightarrow[\simeq]{(-)^{hS^1}} & \text{Mod}_k^\wedge[[t]] \\
 & \searrow \text{THH}^{\text{ref}}(-/k) & \uparrow & & \\
 & & (\text{Mod}_k^{BS^1})^{\text{rig}} & & 
 \end{array}$$



**Construction** (Efimov–Scholze). Let  $k$  be an  $\mathbb{E}_\infty$ -ring spectrum. Suppose  $k$  is complex orientable, choose complex orientation  $t \in \pi_{-2}(k^{hS^1})$ .

$$\begin{array}{ccccc}
 \text{rigid} & & \text{not rigid} & & \\
 \text{Mot}_k^{\text{loc}} & \xrightarrow[\otimes]{\text{THH}(-/k)} & \text{Mod}_k^{\text{BS}^1} & \xrightarrow[\simeq]{(-)^{hS^1}} & \text{Mod}_k^{\wedge}[\![t]\!] \\
 & \searrow \text{THH}^{\text{ref}}(-/k) & \uparrow & & \uparrow \\
 & & (\text{Mod}_k^{\text{BS}^1})^{\text{rig}} & \xrightarrow{\simeq} & \text{Nuc}(k[\![t]\!])
 \end{array}$$



**Construction** (Efimov–Scholze). Let  $k$  be an  $\mathbb{E}_\infty$ -ring spectrum. Suppose  $k$  is complex orientable, choose complex orientation  $t \in \pi_{-2}(k^{hS^1})$ .

$$\begin{array}{ccccc}
 \text{rigid} & & \text{not rigid} & & \\
 \text{Mot}_k^{\text{loc}} & \xrightarrow[\otimes]{\text{THH}(-/k)} & \text{Mod}_k^{\text{BS}^1} & \xrightarrow[\simeq]{(-)^{hS^1}} & \text{Mod}_k^{\wedge}[\![t]\!] \\
 & \searrow \text{THH}^{\text{ref}}(-/k) & \uparrow & & \uparrow \\
 & & (\text{Mod}_k^{\text{BS}^1})^{\text{rig}} & \xrightarrow[\simeq]{} & \text{Nuc}(k[\![t]\!])
 \end{array}$$

$\text{TC}^{-, \text{ref}}(-/k)$





$\mathrm{THH}^{\mathrm{ref}}$  can be nontrivial modulo  $p$ , even for rational inputs.



$THH^{\text{ref}}$  can be nontrivial modulo  $p$ , even for rational inputs. We can describe  $THH^{\text{ref}}(\mathbb{Q}) \otimes ku$



$THH^{\text{ref}}$  can be nontrivial modulo  $p$ , even for rational inputs. We can describe  $THH^{\text{ref}}(\mathbb{Q}) \otimes ku$ , or equivalently  $TC^{-,\text{ref}}(ku \otimes \mathbb{Q}/ku)$ :

$\mathrm{THH}^{\mathrm{ref}}$  can be nontrivial modulo  $p$ , even for rational inputs. We can describe  $\mathrm{THH}^{\mathrm{ref}}(\mathbb{Q}) \otimes ku$ , or equivalently  $\mathrm{TC}^{-,\mathrm{ref}}(ku \otimes \mathbb{Q}/ku)$ :

**Theorem** (Meyer–W. 2024).

$$\pi_* \mathrm{TC}^{-,\mathrm{ref}}(ku \otimes \mathbb{Q}/ku)$$

$\mathrm{THH}^{\mathrm{ref}}$  can be nontrivial modulo  $p$ , even for rational inputs. We can describe  $\mathrm{THH}^{\mathrm{ref}}(\mathbb{Q}) \otimes ku$ , or equivalently  $\mathrm{TC}^{-,\mathrm{ref}}(ku \otimes \mathbb{Q}/ku)$ :

**Theorem** (Meyer–W. 2024).

$$\pi_0 \mathrm{TC}^{-,\mathrm{ref}}(KU \otimes \mathbb{Q}/KU)$$

$\mathrm{THH}^{\mathrm{ref}}$  can be nontrivial modulo  $p$ , even for rational inputs. We can describe  $\mathrm{THH}^{\mathrm{ref}}(\mathbb{Q}) \otimes ku$ , or equivalently  $\mathrm{TC}^{-,\mathrm{ref}}(ku \otimes \mathbb{Q}/ku)$ :

**Theorem** (Meyer–W. 2024).

$$\pi_0 \mathrm{TC}^{-,\mathrm{ref}}(KU_p^\wedge \otimes \mathbb{Q}/KU_p^\wedge)$$

$\mathrm{THH}^{\mathrm{ref}}$  can be nontrivial modulo  $p$ , even for rational inputs. We can describe  $\mathrm{THH}^{\mathrm{ref}}(\mathbb{Q}) \otimes \mathrm{ku}$ , or equivalently  $\mathrm{TC}^{-,\mathrm{ref}}(\mathrm{ku} \otimes \mathbb{Q}/\mathrm{ku})$ :

**Theorem** (Meyer–W. 2024).

$$\pi_0 \mathrm{TC}^{-,\mathrm{ref}}(\mathrm{KU}_p^\wedge \otimes \mathbb{Q}/\mathrm{KU}_p^\wedge) \cong \mathcal{O}(Z^\dagger),$$

$\mathrm{THH}^{\mathrm{ref}}$  can be nontrivial modulo  $p$ , even for rational inputs. We can describe  $\mathrm{THH}^{\mathrm{ref}}(\mathbb{Q}) \otimes ku$ , or equivalently  $\mathrm{TC}^{-,\mathrm{ref}}(ku \otimes \mathbb{Q}/ku)$ :

**Theorem** (Meyer–W. 2024).

$$\pi_0 \mathrm{TC}^{-,\mathrm{ref}}(KU_p^\wedge \otimes \mathbb{Q}/KU_p^\wedge) \cong \mathcal{O}(Z^\dagger),$$

where

- $\mathcal{O}(-^\dagger)$  denotes a ring of overconvergent functions,



THH<sup>ref</sup> can be nontrivial modulo  $p$ , even for rational inputs. We can describe THH<sup>ref</sup>( $\mathbb{Q}$ )  $\otimes$  ku, or equivalently TC<sup>-,ref</sup>(ku  $\otimes$   $\mathbb{Q}$ /ku):


**Theorem** (Meyer–W. 2024).


$$\pi_0 \text{TC}^{-,\text{ref}}(\text{KU}_p^\wedge \otimes \mathbb{Q}/\text{KU}_p^\wedge) \cong \mathcal{O}(\textcolor{brown}{Z}^\dagger),$$

where

- $\mathcal{O}(-^\dagger)$  denotes a ring of overconvergent functions,
- $\textcolor{brown}{Z} \subseteq \text{Spa } \mathbb{Z}_p[[q-1]]$  denotes the subset

$$\text{Spa}(\mathbb{F}_p((q-1)), \mathbb{F}_p[[q-1]]) \cup \bigcup_{\alpha \geq 0} \text{Spa}(\mathbb{Q}_p(\zeta_{p^\alpha}), \mathbb{Z}_p[\zeta_{p^\alpha}]) .$$







spectrum of  $\pi_0 \mathrm{TC}^{-, \mathrm{ref}}(\mathrm{KU}_p^\wedge \otimes \mathbb{Q}/\mathrm{KU}_p^\wedge) \cong \mathcal{O}(Z^\dagger)$


 $q = 1$ 

spectrum of  $\pi_0 \mathrm{TC}^-(\mathrm{KU}_p^\wedge \otimes \mathbb{Q}/\mathrm{KU}_p^\wedge) \cong \mathbb{Q}_p[[q - 1]]$



spectrum of  $\pi_0 \mathrm{TC}^{-, \mathrm{ref}}(\mathrm{KU}_p^\wedge \otimes \mathbb{Q}/\mathrm{KU}_p^\wedge) \cong \mathcal{O}(Z^\dagger)$



spectrum of  $\pi_0 \mathrm{TC}^{-,\mathrm{ref}}(\mathrm{KU}_p^\wedge \otimes \mathbb{Q}/\mathrm{KU}_p^\wedge) \cong \mathcal{O}(Z^\dagger)$



**Analytic Habiro cohomology** (Scholze). Coefficients in the *analytic Habiro ring*  $\mathcal{H}^{\text{an}}$ .



**Analytic Habiro cohomology** (Scholze). Coefficients in the *analytic Habiro ring*  $\mathcal{H}^{\text{an}}$ . This is a sheaf on the stack of norms.





**Analytic Habiro cohomology** (Scholze). Coefficients in the *analytic Habiro ring*  $\mathcal{H}^{\text{an}}$ . This is a  $\mathbb{Z}((u))_{\blacksquare}$ -algebra,  $|u| = \frac{1}{2}$ .



**Analytic Habiro cohomology** (Scholze). Coefficients in the *analytic Habiro ring*  $\mathcal{H}^{\text{an}}$ . This is a  $\mathbb{Z}((u))_{\blacksquare}$ -algebra,  $|u| = \frac{1}{2}$ .

... via  $\text{THH}^{\text{ref}}$  (work in progress). Crucial observations:



**Analytic Habiro cohomology** (Scholze). Coefficients in the *analytic Habiro ring*  $\mathcal{H}^{\text{an}}$ . This is a  $\mathbb{Z}((u))_{\blacksquare}$ -algebra,  $|u| = \frac{1}{2}$ .

... **via  $\text{THH}^{\text{ref}}$**  (work in progress). Crucial observations:

- 1 (Scholze) The analytic de Rham stack lifts canonically to  $\mathbb{S}((u))_{\blacksquare}$ .



**Analytic Habiro cohomology** (Scholze). Coefficients in the *analytic Habiro ring*  $\mathcal{H}^{\text{an}}$ . This is a  $\mathbb{Z}((u))_{\blacksquare}$ -algebra,  $|u| = \frac{1}{2}$ .

... **via  $\text{THH}^{\text{ref}}$**  (work in progress). Crucial observations:

- 1 (Scholze) The analytic de Rham stack lifts canonically to  $\mathbb{S}((u))_{\blacksquare}$ .
- 2 (W.) Analytic Habiro ring can be recovered as

$$\mathcal{H}^{\text{an}} \simeq \pi_0 \text{TCn}^{\text{ref}}(\text{KU}((u))/\text{KU}[\![u]\!]) \left[ \frac{1}{u} \right].$$



**Analytic Habiro cohomology** (Scholze). Coefficients in the *analytic Habiro ring*  $\mathcal{H}^{\text{an}}$ . This is a  $\mathbb{Z}((u))_{\blacksquare}$ -algebra,  $|u| = \frac{1}{2}$ .

... via  $\text{THH}^{\text{ref}}$  (work in progress). Crucial observations:

- 1 (Scholze) The analytic de Rham stack lifts canonically to  $\mathbb{S}((u))_{\blacksquare}$ .
- 2 (W.) Analytic Habiro ring can be recovered as

$$\mathcal{H}^{\text{an}} \simeq \pi_0 \text{TCn}^{\text{ref}}(\text{KU}((u))/\text{KU}[[u]])\left[\frac{1}{u}\right].$$

$$\left( \text{Recall: } \text{TCn}(-) := \lim_{m \in \mathbb{N}} (\text{THH}(-)^{C_m})^{h(S^1/C_m)}. \right)$$



**Analytic Habiro cohomology** (Scholze). Coefficients in the *analytic Habiro ring*  $\mathcal{H}^{\text{an}}$ . This is a  $\mathbb{Z}((u))_{\blacksquare}$ -algebra,  $|u| = \frac{1}{2}$ .

... via  $\text{THH}^{\text{ref}}$  (work in progress). Crucial observations:

- 1 (Scholze) The analytic de Rham stack lifts canonically to  $\mathbb{S}((u))_{\blacksquare}$ .
- 2 (W.) Analytic Habiro ring can be recovered as

$$\mathcal{H}^{\text{an}} \simeq \pi_0 \text{TCn}^{\text{ref}}(\text{KU}((u))/\text{KU}[\![u]\!])\left[\frac{1}{u}\right].$$

## Question.

Can we compute this for bases other than KU (e.g. elliptic cohomology, MU,  $\mathbb{S}$ )?

**Thank you!**