

Embracing Lag Uncertainty: Bayesian Model Averaging in Set-Identified Structural VARs

Akın Aydın* Ferdinand Loser†

June 23, 2026

Seminar Evaluating Econometric Methods using Monte Carlo Simulations
(Prof. Derya Uysal)

Abstract

Structural Vector Autoregressions (VARs) identified via set restrictions are a cornerstone of modern empirical macroeconomics, yet the structural impulse response functions (IRFs) they produce remain highly sensitive to reduced-form lag order specification. Updating the foundational Monte Carlo analysis of Ivanov and Kilian (2005), this paper evaluates the efficacy of traditional Information Criteria (AIC, BIC, HQC) against modern Bayesian Model Averaging (BMA) techniques in recovering the true median target IRF. Utilizing a realistic data-generating process calibrated to a structural model of the global oil market, we analyze the relative Mean Squared Error (MSE) of competing specification strategies across varying finite sample sizes. Our simulations confirm that for single-model selection, the Akaike Information Criterion (AIC) consistently dominates stricter criteria by avoiding the severe underfitting bias that heavily distorts IRF inference. More crucially, we demonstrate that hedging lag specification risk through a joint-grid BMA approach—averaging across both lag orders and Bayesian VAR shrinkage hyperparameters—yields the most accurate IRF recovery in the majority of finite-sample scenarios. By dynamically applying tighter shrinkage to overparameterized specifications, this methodology mitigates lag uncertainty without compromising the precision of critical lower-order dynamics, offering a highly robust alternative to traditional single-model selection.

Keywords: Structural Vector Autoregression (SVAR), Set Identification, Bayesian Model Averaging (BMA), Model Uncertainty, Lag Order Selection, Monte Carlo

*Ludwig-Maximilian University of Munich, akin.aydin@campus.lmu.de

†Ludwig-Maximilian University of Munich, ferdinand.loser@campus.lmu.de

Simulation

JEL Codes: C11, C15, C32, C52

1 Introduction

Structural Vector Autoregressions (VARs) remain the cornerstone of a large part of empirical macroeconomics, especially for monetary policy, shock transmission, and financial markets. While forecasting may be the main goal for some researchers, many economists also use it to estimate the relationships among factors that determine macroeconomic movements. For this purpose, Impulse Response Functions (IRFs) are derived from the VAR parameters, establishing how a shock in one variable affects the development of another period, both contemporaneously and over horizons of several months or even years. Using this transformation, structural VARs are continually adapted to answer complex questions, such as how speculative trading and real-time news shocks drive global oil market dynamics (Kilian and Murphy, 2014; Gazzani et al., 2024; Castelnovo et al., 2024), how high-frequency monetary policy surprises transmit through corporate credit spreads (Gertler and Karadi, 2015; Ferreira et al., 2023), and how these policy shifts impact global commodity prices (Miranda-Pinto, 2023). The specification of the VAR model, as well as the method to decompose it into its structural form, are the two crucial specifications that a researcher has to decide on. Traditionally, VAR models were specified with a lag order determined by economic theory or an information criterion (IC) and then estimated by Ordinary Least Squares (OLS) or Maximum Likelihood (ML). The past 15 years have seen a surge in the popularity of Bayesian methods, in which not a single lag order is specified, but lag orders are weighted and/or shrunk based on a prior and a posterior probability distribution (Blake and Mumtaz, 2015). These methods enable the researcher not only to rely on a single specification but also to incorporate a larger model or set of models while still keeping variance low. The methodology for estimating IRFs has also advanced significantly since Sims (1980) showed that structural IRFs can be retrieved via Cholesky decomposition, i.e., a predetermined causal ordering of the relationships between variables. Newer methods typically require weaker assumptions and adopt a more agnostic approach to inferring economic relationships. The dominant approach nowadays is set identification using sign restrictions informed by economic theory (Uhlig, 2005; Rubio-Ramírez et al., 2010). Unlike in Cholesky decomposition, all variables can simultaneously react to each other’s shocks, and sign restrictions can reduce the set of possible IRFs to those that fit the researcher’s assumptions. This yields a set of admissible IRFs. Typically, the median target—the specific IRF closest to the point-wise median of the admissible set—is selected as the primary result (Fry and Pagan, 2011).

This study draws inspiration from Ivanov and Kilian (2005), the first paper to examine which VAR specification best retrieves the true IRF using Monte Carlo simulations. In 2005, this meant testing criteria for lag selection to assess how well they fit VAR models and retrieve structural IRFs via Cholesky decomposition. We aim to update this analysis to the more recent world of VAR methodology, where IRFs are identified via

sign restrictions and estimation relies heavily on Bayesian methods rather than solely on information criteria.. We construct a realistic DGP based on Kilian and Murphy (2014)’s study of the oil market, using the set identification method. Then we compare different methods across repeated simulations by how well they can retrieve the true IRF, using the median target of their identified set.

We then compare the information criteria AIC, BIC, and HQC with estimation using Bayesian Model Averaging (BMA) which doesn’t fix one lag order but weights a set of probable lag orders and can be combined with Bayesian Vector Autoregression (BVAR) to shrink down parameters of higher lag orders. Our results demonstrate that hedging lag selection risk through Bayesian Model Averaging (BMA) significantly improves the accurate recovery of median target Impulse Response Functions (IRFs). Particularly the joint-grid BMA approach dominates traditional AIC-selected models in most scenarios and is shown to diversify estimation weights across the most broad spectrum of lag orders. By dynamically applying tighter hyperparameter shrinkage (λ_1) to overparameterized specifications ($p > p_0$), this approach successfully mitigates the risk of lag order uncertainty while preserving precise parameter estimation for the crucial lower lags. Furthermore, our results confirm the results presented by Ivanov and Kilian (2005) when relying on single-model selection. In this case, the Akaike Information Criterion (AIC) consistently outperforms stricter criteria like BIC and HQC, as its milder penalization avoids the severe underfitting bias that heavily distorts IRF inference. As T grows, AIC also improves in comparison to BMA for a sufficiently low p_0 as this enables parameters to be estimated with less variance also in simple OLS.

The rest of the paper is structured as follows: Section 2 introduces the reader to the theoretical foundations needed to understand the methodology of this simulation study. It first introduces the VAR framework, Section 2.1, then explains the sign restriction approach for IRF decomposition, Section 2.2, and explains information criteria as well as the BVAR method, Section 2.3 and ???. We aim to both convey the intuition behind these methods as well as derive key properties. Section ??? explains the simulation design and Section 4 presents our results. Finally, the conclusion, Section 5, summarizes and discusses our key findings.

2 Theoretical Background

2.1 VAR Framework and IRFs

The reduced form of a vector autoregression, $\text{VAR}(p_0)$, establishes a relationship between current observations of a set of variables with the past observations of these variables up until a pre-determined lag p_0 :

$$Y_t = v + A_1 Y_{t-1} + A_2 Y_{t-2} + \dots + A_{p_0} Y_{t-p_0} + e_t \quad (1)$$

Here, Y_t is a $(K \times 1)$ vector of endogenous variables, v is a vector of intercepts, $A_1, \dots, A_{p_0}$ are $(K \times K)$ coefficient matrices, e_t is a white-noise innovation vector with $\mathbb{E}[e_t] = 0$ and covariance matrix $\mathbb{E}[e_t e_t'] = \Sigma$, and p_0 denotes the lag order. This is the standard reduced-form representation of a VAR process (Lütkepohl, 2005; Sims, 1980). However, reduced-form innovations are composed of structural shocks and are therefore uninformative about the effects of any individual shock. To identify the structural shocks of interest, restrictive assumptions must be imposed. The relationship between the two is given by:

$$e_t = B \epsilon_t \quad (2)$$

where ϵ_t denotes the vector of mutually uncorrelated structural shocks (that are normalized to unit variance, i.e., $\mathbb{E}[\epsilon_t \epsilon_t'] = I$) and B is a matrix satisfying $\Sigma = B B'$. Since there are infinitely many candidate matrices B (for any valid factor $\tilde{B}$ and any orthogonal matrix D , the product $\tilde{B}D$ also satisfies $\Sigma = (\tilde{B}D)(\tilde{B}D)'$), restrictive assumptions are required to single out a unique B . The literature offers several strategies for this purpose, including short-run (recursive) restrictions (Sims, 1980; Christiano et al., 1999), long-run restrictions (Blanchard and Quah, 1989), and sign restrictions (Faust, 1998; Uhlig, 2005; Rubio-Ramírez et al., 2010); see Kilian and Lütkepohl (2017) for a comprehensive treatment. Following Ivanov and Kilian (2005), we do not commit to a single scheme mentioned above but instead adopt, for each data-generating process, the identifying assumptions used in the original study on which it is based.

In general, interpreting the dynamics implied by VAR coefficients is difficult. Therefore, practitioners use Impulse Response Functions (IRF) to summarize the main dynamics that they want to explain. To obtain the coefficient set that constructs the impulse response function, we express the VAR model in terms of an infinite VMA process. Using (1):

$$Y_t = [I - A_1 L - A_2 L^2 - \dots - A_{p_0} L^{p_0}]^{-1} v + [I - A_1 L - A_2 L^2 - \dots - A_{p_0} L^{p_0}]^{-1} e_t \quad (3)$$

Which is equivalent to:

$$Y_t = \mu + \sum_{i=0}^{\infty} \Phi_i e_{t-i} \quad \text{and} \quad \Phi_0 = I_K \quad (4)$$

where $\mu = [I - A_1 - \dots - A_{p_0}]^{-1} v$ is the unconditional mean vector and the moving average coefficient matrices are obtained recursively as $\Phi_i = \sum_{j=1}^i A_j \Phi_{i-j}$ where $A_j =$

0 for $j > p_0$ (see appendix for detailed derivation). Finally, in order to obtain the structural impulse response functions, we plug the expression given in (2) into (4):

$$Y_t = \mu + \sum_{i=0}^{\infty} \Phi_i B \epsilon_{t-i} = \mu + \sum_{i=0}^{\infty} \Theta_i \epsilon_{t-i} \quad (5)$$

Here, the matrices Θ_i are the structural impulse response matrices, and they contain the coefficients required to construct the impulse response function. To illustrate, the equation for the first endogenous variable can be written as:

$$Y_{1t} = \mu_1 + \sum_{k=1}^K \theta_{1k,0} \epsilon_{kt} + \sum_{k=1}^K \theta_{1k,1} \epsilon_{kt-1} + \sum_{k=1}^K \theta_{1k,2} \epsilon_{kt-2} + \dots \quad (6)$$

where $\theta_{jk,i}$ denotes the response of variable j to structural shock k at horizon i , i.e., the (j,k) element of Θ_i .

The structural impulse responses in (6) depend on the model through two distinct channels. First, the moving average coefficients Φ_i are determined by the reduced-form autoregressive coefficients $A_1, \dots, A_{p_0}$, so the impulse responses depend on the lag order p_0 ; in practice, p_0 is unknown and must be estimated, which is the subject of the lag-order selection criteria discussed in Section 3. Second, the mapping from reduced-form to structural form impulse responses requires the identification of B introduced in (2). Since there are infinitely many candidate matrices B (for any valid factor $\tilde{B}$ and any orthogonal matrix D , the product $\tilde{B}D$ also satisfies $\Sigma = (\tilde{B}D)(\tilde{B}D)'$), restrictive assumptions are required to single out a unique B . The literature offers several strategies for this purpose, including short-run (recursive) restrictions, often computed by Cholesky decomposition (Sims, 1980; Christiano et al., 1999), long-run restrictions (Blanchard and Quah, 1989), and sign restrictions (Faust, 1998; Canova and Nicoló, 2002; Uhlig, 2005; Rubio-Ramírez et al., 2010); see Kilian and Lütkepohl (2017) for a comprehensive treatment. We calibrate our data-generating process to the model of Kilian and Murphy (2014) and accordingly employ a sign-restriction algorithm to identify the impact matrix (though we omit the elasticity bounds for computational feasibility). A key feature of this approach is that sign restrictions deliver only partial identification: rather than a single impact matrix, they identify a *set* of candidate matrices consistent with the a priori restrictions implied by economic theory. This raises several questions about the identification strategy, which are the focus of the next subsection.

Structural Vector Autoregressions (VARs) remain the cornerstone of a large part of empirical macroeconomics, especially for monetary policy, shock transmission, and financial markets. While forecasting may be the main goal for some actors, especially central banks or national economic institutes, many macroeconomists also use it to estimate the relationships among factors that determine macroeconomic movements. For this purpose, Impulse Response Functions (IRFs) are derived from the VAR parameters,

establishing how a shock in one variable affects the development of another period, both contemporaneously and over horizons of several months or even years. For example, it may be applied to study how shocks in speculative oil demand affect oil demand and supply flows (Kilian and Murphy, 2014) or how interest rate shocks impact credit costs (Gertler and Karadi, 2015) (change to add newer papers citations). The specification of the VAR model, as well as the method to decompose it into its structural form, are the two crucial specifications that a researcher has to decide on. Traditionally, VAR models were specified with a lag order determined by economic theory or an information criterion (IC) and then estimated by Ordinary Least Squares (OLS) or Maximum Likelihood (ML). The past 15 years have seen a surge in the popularity of Bayesian methods, in which not a single lag order is specified, but lag orders are weighted and/or shrunk based on a prior and a posterior probability distribution (CITATION). These methods enable the researcher not only to rely on a single specification but also to incorporate a larger model or set of models while still keeping variance low. The methodology for estimating IRFs has also advanced significantly since Sims (1980) showed that structural IRFs can be retrieved via Cholesky decomposition, i.e., a predetermined causal ordering of the relationships between variables. Newer methods typically require weaker assumptions and adopt a more agnostic approach to inferring economic relationships. The dominant approach nowadays is set identification using sign restrictions informed by economic theory (Uhlig, 2005; Rubio-Ramírez et al., 2010). Unlike in Cholesky decomposition, all variables can simultaneously react to each other's shocks, and sign restrictions can reduce the set of possible IRFs to those that fit the researcher's assumptions. In the end, there is a set of admissible IRFs of which the median target, the IRF overall closest to the median of all IRFs, is usually selected as the main result (Fry and Pagan, 2011).

This study draws inspiration from Ivanov and Kilian (2005), the first paper to examine which VAR specification best retrieves the true IRF using Monte Carlo simulations. In 2005, this meant testing criteria for lag selection to assess how well they fit VAR models and retrieve structural IRFs via Cholesky decomposition. We aim to update this analysis to the more recent world of VAR methodology, where IRFs are determined by set identification and estimation relies not only on information criteria but also on Bayesian methods. We construct a realistic DGP based on Kilian and Murphy (2014)'s study of the oil market, using the set identification method. Then we compare different methods across repeated simulations by how well they can retrieve the true IRF, using the median target of their identified set. This study compares the information criteria AIC, BIC, and HQC with BVAR estimates using different overall shrinkage hyperparameters λ_1 . We find that BVARs are a well-performing alternative to OLS-estimated VAR models with IC-selected lag orders, beating the best criterion, AIC, in most scenarios. We could now deduce a clear pattern for which λ_1 -values shall be optimally employed depending on sample size T and true lag order p_0 . We attribute this to the fact that both higher and lower

shrinkage can have the same overall effect: one reduces variance, the other reduces bias. Additionally, we suspect that our simulation still contains too much sampling variance, as computational restrictions did not allow for sufficient iterations. Optimization of λ_1 using marginal density distribution (MDD), which optimizes on out-of-sample performance, is found not to generally favor lower values of λ_1 , which is found to be suboptimal for the purpose of IRF decomposition. We therefore show that the goals of forecasting and IRF analysis do not necessarily work together in VAR estimation.

The rest of the paper is structured as follows: Section 2 introduces the reader to the theoretical foundations needed to understand the methodology of this simulation study. It first introduces the VAR framework, Section 2.1, then explains the sign restriction approach for IRF decomposition, Section 2.2, and explains information criteria as well as the BVAR method, Section 2.3 and ???. We aim to both convey the intuition behind these methods as well as derive key properties. Section ??? explains the simulation design and Section 4 presents our results. Finally, the conclusion, Section 5, summarizes and discusses our key findings.

2.2 Sign Restrictions

Point identification strategies, such as the Cholesky decomposition, impose an assumption restrictive enough to single out a unique impact matrix; in the Cholesky case, this is a recursive ordering that determines the contemporaneous causal structure among the endogenous variables. Uhlig (2005), however, criticizes such strategies on the grounds that these restrictive assumptions may not reflect the underlying economic reality, but instead may impose the very conclusions the researcher expects to find.

Uhlig (2005) proposes an alternative he terms an "agnostic" identification procedure. The agnostic approach imposes only minimal restrictions—and, crucially, leaves the response of the variable of central interest entirely unrestricted—so that the answer to the research question is determined by the data rather than by the identifying assumptions. The restrictions that are imposed are deliberately weak: they consist only of sign restrictions on a few responses that can be reasonably agreed upon across economists. This minimalism comes at a cost, however. Imposing only sign restrictions does not single out a unique impact matrix; instead, it identifies a set of candidate impact matrices consistent with the assigned signs. This set-identification (or partial identification) strategy is less restrictive, but it comes at the cost of ambiguity, since the object of interest is no longer a single value but a range. Moreover, as emphasized by Uhlig (2005), sign restrictions introduce a further choice that point-identifying schemes do not: the horizon over which the restrictions are imposed. In addition to specifying the sign of a response on impact, the practitioner must decide how many subsequent periods that sign is required to hold—the longer this horizon, the more candidate models are discarded and the narrower the

identified set becomes.

This trade-off is described by Manski (2003) as the Law of Decreasing Credibility: the credibility of inference decreases as the strength of the maintained assumptions increases. Point identification strategies impose stronger assumptions and are therefore more informative—they reduce the identified object to a single point—but they are less credible, since their validity rests entirely on assumptions that may not hold. Partial identification strategies, by contrast, are less informative—they deliver a trimmed range rather than a single element—but more credible, precisely because they rely on weaker and more defensible assumptions.

Here, we follow Kilian and Murphy (2014) with slight differences due to computational feasibility.¹ First, we fit VAR(p_0) model to the data of Kilian and Murphy (2014) and obtain the reduced-form autoregressive coefficients ($\hat{A}$) and the variance-covariance matrix of the reduced-form errors ($\hat{\Sigma}$). We then factor this covariance matrix using Cholesky decomposition ($\hat{\Sigma} = PP'$).² Next, following Rubio-Ramírez et al. (2010), we generate random orthogonal matrices: we draw a matrix of independent standard normal entries, compute its QR decomposition, and normalize the sign of each column so that the diagonal of (R) is positive. This procedure yields orthogonal matrices (Q) (with $QQ' = I_K$) that are uniformly distributed with respect to the Haar measure, ensuring that the space of admissible rotations is sampled without bias. For each draw, we form a candidate impact matrix ($\tilde{B} = PQ$) and check whether it satisfies the imposed sign restrictions; candidates that violate the restrictions are discarded. Finally, we retain a select number of admissible impulse response structures and select the median target as our true IRF (population response). For the sake of clarity, we find it appropriate to outline the step-by-step blueprint of this algorithm:

1. Draw a ($K \times K$) matrix W , where each element is drawn independently from a standard normal distribution: $W_{ij} \sim N(0, 1)$.
2. Compute the QR decomposition $W = QR$, where Q is an orthogonal matrix (the rotation matrix) and R is upper triangular. Normalize the sign of each column of Q so that the diagonal of R is positive, which renders the decomposition unique and ensures that Q is distributed according to the Haar measure.

¹Kilian and Murphy (2014) argue that agnostic sign restrictions alone are not sufficient to narrow the set of admissible models adequately, and they therefore impose additional quantitative restrictions; bounds on the implied oil supply and demand elasticities. Additionally, they set dynamic restrictions for a horizon of 12 months. We omit the elasticity restrictions and reduce the dynamic restrictions to 3 months for reasons of computational feasibility: in our finite samples, the acceptance rate under the elasticity bounds is prohibitively low, so that obtaining a sufficient number of admissible models in each of the many Monte Carlo iterations would require a computationally unachievable number of draws.

²Kilian and Murphy (2014) uses eigendecomposition here. However, it doesn't make any difference. Any two factors satisfying $BB' = \tilde{B}\tilde{B}' = \hat{\Sigma}$ differ only by an orthogonal rotation ($B = \tilde{B}D$). Since our procedure applies the full set of Haar-distributed rotations to the initial factor, the resulting set of admissible models is invariant to the initial decomposition. The only element we need is an initial factor to run the algorithm.

3. Construct the candidate impact matrix $\tilde{B} = PQ$, and then check whether $\tilde{B}$ satisfies the imposed restrictions.

4. Repeat this process by drawing new W matrices (i.e., obtaining new rotation matrices Q) until the desired number of admissible models is reached.

5. Among the retained models, select the median target m^* as the representative (true) model.

The selection of a representative model is a much-debated issue in the literature, particularly the comparison between the pointwise median and the median target. Since we must obtain a single population response rather than a set—otherwise the mean squared error (MSE) could not be computed against a unique true IRF—we adopt the approach most appropriate for our design. Fry and Pagan (2011) criticize the pointwise median for a compelling reason: the pointwise median does not correspond to any single admissible model, as it combines responses drawn from different models with different impact matrices. Consequently, this synthetic profile may appear plausible but need not correspond to any B satisfying $\hat{\Sigma} = BB'$. This implies that the shocks may not be properly isolated, so the pointwise median cannot be used for variance decomposition. To address this, Fry and Pagan propose the median target approach:

$$m^* = \arg \min_m \sum_{j,k,i} \left(\frac{\theta_{jk,i}^{(m)} - \text{med}_{jk,i}}{\sigma_{jk,i}} \right)^2, \quad (7)$$

$$\text{where } \text{med}_{jk,i} = \text{median}_m \{\theta_{jk,i}^{(m)}\}, \quad \sigma_{jk,i} = \text{std}_m \{\theta_{jk,i}^{(m)}\}.$$

Solving this minimization yields the admissible model whose impulse responses are closest to the pointwise median ($m^* \in M$). Since m^* is an actual admissible model rather than a synthetic construct, it satisfies $\hat{\Sigma} = BB'$ (which makes sure that shocks are orthogonalized) and can be used for variance decomposition. We adopt this approach due to the coherence concerns described above and take the median target as our representative true IRF whereby its specific impact matrix $\tilde{B}$ determines the DGP.³

Identification is an essential part of recovering the underlying impulse response function. As noted earlier, however, it is not the only factor that matters: the specification of the reduced-form VAR also plays a role. Recall that $\Phi_i = \sum_{j=1}^i A_j \Phi_{i-j}$ where $A_j = 0$ for $j > p_0$, so the moving-average coefficients, and hence the impulse responses, depend on the lag order (p_0). The practitioner must therefore choose the lag order carefully. We address this in the next section.

2.3 Information Criteria

The true lag order p_0 is a priori unknown when estimating a reduced-form VAR; it is a matter of specification. Adding too many lags may lead to overfitting, which inflates

³For a visualization of the IRFs selected for the DGP, see Figure ?? in the Appendix.

the variance of the impulse response estimates by requiring the estimation of additional parameters. Underfitting is also problematic, as it introduces bias. The bias-variance trade-off is therefore central to lag-order selection. Traditionally, information criteria are used to address this issue, and their working principle is essentially the same (Lütkepohl, 2005, chap. 4):

$$\text{IC}(p) = \ln |\hat{\Sigma}(p)| + \underbrace{c_T, \frac{K^2 p}{N}}_{\text{penalty term}} \quad (8)$$

Information criteria are formulas that embody the bias-variance trade-off described above. The $\ln |\hat{\Sigma}(p)|$ term is the log-determinant of the reduced-form error covariance matrix, estimated by quasi-maximum likelihood; this term penalizes underfitting, since a model with too few lags leaves $\hat{\Sigma}(p)$ large and thus raises $\text{IC}(p)$. The Penalty weight c_T depends on the specific criterion, and the penalty term increases with the lag order; the second component therefore penalizes overfitting. The lag order is estimated by:

$$\hat{p} = \arg \min p \text{IC}(p), \quad 1 \leq p \leq \bar{p} \quad (9)$$

Following Ivanov and Kilian (2005), we set the maximum lag order to $\bar{p} = 12$ for our monthly data, corresponding to one year. This choice exceeds the largest true lag order considered $p_0 = 10$, ensuring that all criteria can both identify and overfit relative to the true order, and it makes our results directly comparable to those of Ivanov and Kilian (2005).⁴

Although there are different types of $\text{IC}(p)$, three are mainly used:

$$\text{AIC}(p) = \ln |\hat{\Sigma}(p)| + \frac{2}{N}, K^2 p \quad (10)$$

The Akaike Information Criterion (AIC) (Akaike, 1974) is the least overfitting-averse of these three criteria. It uses $c_T = 2$ and tends to overfit in practice.⁵

$$\text{BIC}(p) = \ln |\hat{\Sigma}(p)| + \frac{\ln N}{N}, K^2 p \quad (11)$$

The Bayesian Information Criterion (BIC), also called the Schwarz Information Criterion (Schwarz, 1978), penalizes overfitting the most. It uses $c_T = \ln N$ and is a consistent

⁴We only include AIC, BIC and HQC as information criteria and omit the Likelihood Ratio tests and the Lagrange Multiplier test for lag selection as these have become practically irrelevant in the last 20 years for macroeconomic time series analysis and were also shown to perform poorly in Ivanov and Kilian (2005)

⁵In the literature, the AIC is generally regarded as prone to overfitting. In the VAR setting, however, this tendency is attenuated: adding one more lag requires estimating an entire $(K \times K)$ coefficient matrix, so the penalty grows by $\frac{2K^2}{N}$ at each step. Because each additional lag is this costly, even the AIC—the most lenient of the three criteria—can underfit in practice.

estimator of the true lag order $\left(\text{plim}_{N \rightarrow \infty} \hat{p}^{\text{BIC}} = p_0\right)$.

$$\text{HQC}(p) = \ln |\hat{\Sigma}(p)| + \frac{2 \ln \ln N}{N}, K^2 p \quad (12)$$

The Hannan-Quinn Information Criterion (Hannan and Quinn, 1979) is neither as harsh against overfitting as AIC nor as tolerant as BIC. It uses $c_T = 2 \ln \ln N$ as its penalty weight and, like BIC, is a consistent estimator of the true lag order.

The three criteria share the same fit term and differ only in the penalty weight c_T , which equals 2 for the AIC, $2 \ln \ln N$ for the HQC, and $\ln N$ for the BIC. A heavier penalty makes each additional lag more costly and therefore yields a more parsimonious model. Comparing the weights gives:

$$2; \leq; 2 \ln \ln N; \leq; \ln N \quad (13)$$

where the first inequality holds for $N \geq 16$ and the second for all N .⁶ Consequently, the implied lag-order estimates satisfy:

$$\hat{p}^{\text{BIC}}; \leq; \hat{p}^{\text{HQC}}; \leq; \hat{p}^{\text{AIC}} \quad (14)$$

so that the BIC is the most parsimonious criterion and the AIC the least, with the HQC in between (Ivanov and Kilian, 2005). Since $N \geq 16$ holds across all sample sizes in our study, this ordering applies throughout. As seen in Table 1, our Monte Carlo Simulation confirms this ordering. The penalization of BIC is evidenced to be so strong that the lag order selected, $\hat{p}^{\text{BIC}}$, always stays close to 2, regardless of the sample size T and the true lag order p_0 . $\hat{p}^{\text{HQC}}$ lies above $\hat{p}^{\text{BIC}}$ and increases more apparently with T ; it also does not come close to converging to the true lag order. On the other hand, $\hat{p}^{\text{AIC}}$ is consistently much closer to the true lag order and has already nearly reached p_0 at $T = 600$. Therefore, although AIC is not asymptotically consistent, it is the only IC that can come close to the true lag order for realistic sample sizes.

Lütkepohl (1985) compares information criteria based on their ability to recover the true lag order. Ivanov and Kilian (2005), by contrast, object to this approach on the grounds of the bias-variance trade-off discussed above. To see why, it is helpful to consider the decomposition of the mean squared error (MSE) of the estimated median target IRF $\hat{\theta}$ compared to the true DGP IRF θ :

$$\text{MSE}(\hat{\theta}) = E[(\hat{\theta} - \theta)^2] = E[(\hat{\theta} - E[\hat{\theta}] + E[\hat{\theta}] - \theta)^2] \quad (15)$$

⁶The weights coincide at $N = e^e \approx 15.2$ (for $2 = 2 \ln \ln N$) and never cross for $\ln N$ versus $2 \ln \ln N$, since $x - 2 \ln x > 0$ for all $x > 0$. The condition $\ln N \geq 2$ (BIC versus AIC) holds for $N \geq 8$. Therefore, $N \geq 16$ is the requirement for (14) to hold.

Table 1: Average Evaluated Lag Order ($\hat{p}$) selected by Information Criteria.

	Estimator	AIC	BIC	HQC
p_0	T			
4	240	2.99	1.95	2.06
	360	3.39	2.00	2.16
	480	3.61	2.01	2.40
	600	3.80	2.02	2.66
10	240	6.54	1.96	2.07
	360	8.53	2.00	2.23
	480	8.96	2.00	2.66
	600	9.16	2.02	4.03

which can be reformulated to:

$$MSE(\hat{\theta}) = E \left[\left(\hat{\theta} - E[\hat{\theta}] \right)^2 + \underbrace{2 \left(\hat{\theta} - E[\hat{\theta}] \right) \left(E[\hat{\theta}] - \theta \right)}_{=0} + \left(E[\hat{\theta}] - \theta \right)^2 \right] \quad (16)$$

and finally, after we distribute the expected value operator:

$$MSE(\hat{\theta}) = Var(\hat{\theta}) + Bias(\hat{\theta})^2 \quad (17)$$

we have derived an equation for the bias-variance tradeoff. In small samples, if underfitting produces a reduction in variance large enough to offset the resulting increase in bias, the MSE falls, and if we take the MSE as a proxy for accuracy, this matters more than a criterion's ability to recover the true lag order. The lag order selected by a criterion that underfits may therefore outperform the true lag order; one of the central remarks of Ivanov and Kilian (2005).

2.4 Bayesian Model Averaging (BMA)

Bayesian Model Averaging is an averaging mechanism designed to address the problem of model uncertainty (Hoeting et al., 1999; Steel, 2020). For any inference question, there may be more than one model that fits the data well, and committing to a single one ignores the uncertainty inherent in that choice. In our setting, the candidate models are reduced-form VARs of different lag orders. Bayesian Model Averaging retains all of them and forms a weighted average, with each model weighted by its posterior model probability. The averaged result is preferable for two reasons. First, it accounts for model uncertainty in addition to parameter uncertainty, rather than conditioning on a single model as if it were known to be correct. Second, averaging over all models in proportion to their posterior probabilities yields better average performance, as measured

by a logarithmic scoring rule, than any single model (Madigan and Raftery, 1994; Hoeting et al., 1999). In this section, we list four different Bayesian Model Averaging methods that we use as a metric (namely, Uniform prior-OLS, Geometric prior-OLS, Joint Grid, and Geometric Grid) and discuss their properties. To clarify, here is the blueprint of the process that we follow:

1. We estimate every reduced form lag order specification up to $\bar{p}$ (maximum lag order) using OLS, and different $(p; \lambda_1)$ combinations using the BVAR developed in Paper A.
2. Then we calculate the weight parameters of each model (we use the BIC/AIC for OLS estimations and the marginal data density (MDD) for the BVAR estimations).
3. We draw different amounts of IRFs based on the weights (Higher weight implies more draws from that specific model) and pool them.
4. We choose the median target IRF as discussed before.

This process raises several questions. How are the model weights calculated? Is averaging genuinely better than selecting a single model? What do the six BMA variants share, and how do they differ? We address these in turn.

General Bayesian Model Averaging Approach

We find it important to discuss the common properties of the four different BMA approaches listed above. We follow the canonical treatment of Hoeting et al. (1999) for this part. First, we start with the problem of choosing the right weight parameters (so that we can average the models based on the weights). Let Δ be our parameter of interest (Impulse response functions) and let $M_1, \dots, M_K$ be our candidate models (different reduced form VAR models). The posterior distribution of Δ given the data y (step 3 defined in the blueprint), averaged over all models, is:

$$p(\Delta | y) = \sum_{k=1}^K p(\Delta | M_k, y) p(M_k | y) \quad (18)$$

so this is the weighted average of the posterior distributions conditioned on the model $p(\Delta | M_k, y)$, weighted by $p(M_k | y)$ (posterior probability of model conditioned on data). By Bayes' theorem applied at the level of models, this weight is:

$$p(M_k | y) = \frac{p(y | M_k) p(M_k)}{\sum_{l=1}^K p(y | M_l) p(M_l)} \quad (19)$$

where $p(M_k)$ is the prior probability assigned to model M_k and $p(y | M_k)$ is its marginal data density—the likelihood integrated over the model’s parameters under their prior:

$$p(y | M_k) = \int p(y | \theta_k, M_k) p(\theta_k | M_k) d\theta_k. \quad (20)$$

The marginal data density (MDD or marginal likelihood) in (20) is central to our procedure, as it is the primary input to the model weights in (19).

These 3 equations defined above are the backbones of the Bayesian Model Averaging methodology. In practice, these three equations map directly onto the steps of our procedure. Equation (20)—the marginal data density—is evaluated first: for each candidate model, we obtain a measure of how well it explains the data once its parameters are integrated out. For the BVAR models, indexed by a lag order and tightness pair (p, λ_1) , this quantity is the marginal data density returned in closed form by the conjugate Normal-Wishart estimator; for the OLS models, indexed by lag order alone (p) , it is approximated by the Bayesian Information Criterion and Akaike Information Criterion⁷ Equation (19) then converts these marginal data densities into model weights: each is multiplied by its model prior $(p(M_k))$ —uniform or geometric—and normalized so that the weights across all candidate models sum to one (denominator). These weights are the posterior model probabilities. Finally, (18) is realized not by analytical combination but by Monte Carlo: from each model we draw admissible impulse responses in a number proportional to its weight, so that models the data favor contribute more draws, and we collect all draws into a single pool. This pool is the Monte Carlo representation of the model-averaged distribution in (18). From it, we select the median target as our single, coherent estimate of the impulse response, completing the averaging.

So far, we have described the common structure they share. However, we use four different BMA methods, and each has its own priors or estimation methods. We discuss the differences below.

Weights and Priors

We use two weight sources and two model-space priors (uniform prior and geometric prior). Each combination yields a different metric:

We find it fruitful to discuss model priors that we use $(p(M_k))$ first. The uniform prior is straightforward. It represents the belief that every candidate model has the same probability of being the true model (reflecting the absence of any prior preference among

⁷Steel (2020, chapt. 3.7) mentions that marginal data density is approximated by the Bayesian Information Criterion when the closed form of equation (20) does not exist, as BIC asymptotically converges to marginal data density. Also, Burnham and Anderson (2002) justifies the usage of AIC as an approximation of the marginal data density. Therefore, we use both criteria.

	Uniform prior	Geometric prior ($\theta = 0.5$)
OLS (BIC-weighted, p)	Uniform-OLS	Geometric-OLS
OLS (AIC-weighted, p)	Uniform-OLS	Geometric-OLS
BVAR (MDD-weighted, p, λ_1)	Joint Grid	Geometric Grid

lag orders):

$$p(M_1) = p(M_2) = \cdots = p(M_K) = \frac{1}{\bar{p}} \quad (21)$$

On the other hand, geometric prior constructs the prior as:

$$p(M_k) \propto \theta^p \quad \text{where} \quad \theta = 0.5 \quad (22)$$

We set $\theta = 0.5$, under which each additional lag halves the prior weight—equivalently, the prior demands that the marginal data density favor a longer model by at least a factor of two before the additional lag is preferred. This value is a deliberate, interpretable benchmark rather than an optimized choice: it imposes a moderate preference for parsimony, neither so weak as to coincide with the uniform prior nor so strong as to dominate the evidence from the marginal data density.

We use BIC and AIC for the models estimated via OLS. The reason is that it is not possible to calculate the marginal data density since OLS places no prior on the parameters: under the frequentist approach, each estimated parameter consumes one full degree of freedom, with no shrinkage. We therefore approximate it by the Schwarz criterion, which Schwarz (1978) and Raftery (1995) derive as an asymptotic approximation to the log marginal data density and which, in large samples, does not depend on a prior. For the BVAR models, by contrast, the Normal-Wishart prior—a normal prior on α and an Inverse-Wishart prior on Σ —supplies the joint density that (20) integrates over, and the conjugate structure makes the integral available in closed form, so the marginal data density is computed directly.

Weight Sources

By weight sources, we mean the way the marginal data density in (20) is obtained for each model class. As we discussed before, for models estimated by OLS, (20) was undefined because there were no priors $p(\theta_k | M_k)$. Therefore, for the OLS models, the marginal data density is approximated by $\exp(-\frac{1}{2}\text{IC}(p))$, since the log marginal data density is approximately $-\frac{1}{2}\text{IC}(p)$, where IC is either BIC or AIC. On the other hand, the BVAR method uses an inverse-Wishart joint prior (normally distributed prior for α and Inverse-Wishart distribution for prior Σ), which is described in Paper A and the appendix. Since the joint prior has a closed form, we can calculate the marginal data density ($m(y | p, \lambda_1)$) by plugging in the joint prior in (20) as $p(\theta_k | M_k)$.

Calculated Weights

By using (19) we calculate the weights for the models estimated via OLS:

$$w_{\text{OLS}}(p) = \frac{\exp\left(-\frac{1}{2}\text{IC}(p)\right) \pi(p)}{\sum_{p'=1}^{\bar{p}} \exp\left(-\frac{1}{2}\text{IC}(p')\right) \pi(p')} \quad (23)$$

where $\pi(p) = \frac{1}{\bar{p}}$ under the uniform prior and $\pi(p) \propto \theta^p(\theta = 0.5)$ under the geometric prior. For the BVAR models, indexed by the pair (p, λ_1) , the weight is:

$$w_{\text{BVAR}}(p, \lambda_1) = \frac{m(y | p, \lambda_1) \pi(p)}{\sum_{p'=1}^{\bar{p}} \sum_{\lambda'_1} m(y | p', \lambda'_1) \pi(p')} \quad (24)$$

where $m(y | p, \lambda_1)$ is the calculated marginal data density (20) with joint prior and $\pi(p)$ is the same priors that are described above.

The BVAR models are indexed by pairs (p, λ_1) , with the lag order ranging over $(p = 1, \dots, \bar{p})$ and the tightness over the discrete grid $\lambda_1 \in \{0.20, 0.40, 0.60, 0.80\}$. For every pair in this grid, we estimate the Normal-Wishart BVAR and compute its marginal data density. The pairs are then weighted by their posterior model probability (24); pairs whose normalized weight falls below a threshold of 0.01 are discarded, and the weights of the retained pairs are renormalized to sum to one. From each retained pair, we draw impulse responses in proportion to its weight, pool the draws, and select the median target.

Advantages of the Bayesian Model Averaging

First of all, Bayesian Model Averaging explicitly takes model uncertainty into account. Following Hoeting et al. (1999):

$$\text{Var}[\Delta | y] = \sum_k \left(\text{Var}[\Delta | y, M_k] + \hat{\Delta}_k^2 \right) p(M_k | y) - E[\Delta | y]^2 \quad (25)$$

Here, $\hat{\Delta}_k = E[\Delta | y, M_k]$. After decomposing $\text{Var}[\Delta | y]$, it is possible to see that BMA framework accounts for the model uncertainty:

$$\text{Var}[\Delta | y] = \underbrace{\sum_k \text{Var}[\Delta | y, M_k] p(M_k | y)}_{\text{within-model}} + \underbrace{\sum_k \left(\hat{\Delta}_k - E[\Delta | y] \right)^2 p(M_k | y)}_{\text{across-model}} \quad (26)$$

From (26) we see that if a single model dominated the posterior—so that $\hat{\Delta}_k = E[\Delta | y]$ for that model—the across-model term would vanish and $\text{Var}[\Delta | y]$ would reduce to the within-model variance alone. That is clearly an advantage of Bayesian Model Averaging

since it allows the practitioner to estimate the posterior variance in a more accurate way.

Second, model averaging has a formal optimality property: Madigan and Raftery (1994) show that, under a logarithmic scoring rule, the model-averaged predictive distribution outperforms any single model. While our study concerns estimation rather than forecasting, this result underpins the broader case for averaging. Hoeting et al. (1999) provide a concrete illustration in which even the highest-probability model carries only 17% of the posterior model probability, underscoring that model uncertainty is often substantial. Therefore, it makes sense to use Bayesian Model Averaging in order to account for model uncertainty. Our central motivation for using this approach is to assess whether accounting for model uncertainty leads to better MSE results or not.

Having established the theoretical framework and estimation methodology, we now turn to the simulation design to evaluate the empirical performance of the competing approaches.

3 Simulation Design

This Monte Carlo simulation study follows Ivanov and Kilian (2005) design with several alterations in order to make our extensions feasible and to fit the needs of our analysis. First of all, we only use one monthly DGP based on the empirical VAR study of Kilian and Murphy (2014). Therefore, we do not get estimations for quarterly VAR series or VEC series, and additionally, we do not have additional robustness by including several different series with different macroeconomic indicators and movements. The reason for this limitation is our tight time constraint. We use the dataset from Kilian and Murphy (2014) containing the following 4 variables: percent change in global oil production, real activity index from Kilian (2009), the log real price of oil, and changes in OECD crude oil inventories. Just as in the original study, we first estimate a VAR with a set lag order of these 4 variables and then use the sign restrictions approach explained earlier to identify structural IRFs and select the median target.

The DGP is formed by structural VAR model using the structural impact matrix $\tilde{B}$ that corresponds to the median target IRF as the parameter for the structural shocks e_t , which are drawn from an i.i.d. standard normal distribution. The formula for the DGP is represented by a structural VAR model using the estimated reduced form parameters A (for the lag coefficients y_{t-p}) and V (for the monthly dummy parameters d_t) and the structural impact matrix $\tilde{B}$ as follows:

$$y_t = Vd_t + Ay_{t-p} + \tilde{B}e_t \quad (27)$$

We estimate these parameters using the original data for two different lag orders, $p_0 \in \{4, 10\}$ and create DGPs with different sample sizes, $T \in \{240, 360, 480, 600\}$, where

$T = N + \bar{p}$. $\bar{p}$ is the set of lag orders that are considered for both the IC and the BVAR estimations; it is uniformly set to $\bar{p} = 12$ to ensure that the true lag order p_0 is part of the set. We restrict the lag order selection p_0 to only include one high and one low lag order and drop $T = 300$ in comparison to the setup of Ivanov and Kilian (2005) in order to save computational power for the sign restriction decomposition. For sampling, we use random draws of length p_0 from the original dataset as initial values, then apply the parametric bootstrap using the SVAR described above and generate sequences of lengths T .

On each of these samples, OLS VARs fitted by lag orders selected by both the three ICs as well as the BMA determined OLS VARs and BVARs, for both the geometric and BIC-based approach, are estimated, a fixed number of admissible IRFs found, and the median target IRF determined. We only include AIC, BIC and HQC as information criteria and omit the Likelihood Ratio tests and the Lagrange Multiplier test for lag selection as these have become practically irrelevant in the last 20 years for macroeconomic time series analysis and were also shown to perform poorly in Ivanov and Kilian (2005). The four different BMA estimations follow the methodology described in the previous section. The space of optimal shrinkage parameters for the implementation with BVAR is limited to $\lambda_1 \in \{0.05, 0.2, 0.35, 0.5, 0.65, 0.8, 0.95\}$ in order to make computation in the simulation feasible. It still captures a good range of typically used λ_1 -values, both tighter and looser specifications.

Finally, we estimate the MSE by calculating the geometric mean over the ratio of the squared errors of the median target IRFs $\hat{\theta}_j$ computed by method j in relation to the squared errors of the median target IRF by a model fit with the true lag order $\hat{\theta}_{p_0 i}$ over all Monte Carlo iterations from 1, ..., R :

$$MSE_j^{relative} = \exp \left(\frac{1}{n} \sum_{i=1}^R \ln \left(\frac{SE_{ji}}{SE_{p_0 i}} \right) \right) \quad (28)$$

The geometric mean is preferred over the arithmetic mean as the ratios are bounded on the left side at 0, but unbounded on the right side. Outliers on the right side are hence overweighed in the arithmetic mean. Normalization by the model fit with p_0 ensures that performance in each iteration is weighed by how well a correctly identified model performs.

For the computation of the DGP parameters, we set the number of selected admissible IRFs to 1 million in order to get a highly accurate representation of the real world movements. However, during the Monte Carlo simulation we have to reduce that number to 1,000, which is seen as the minimum requirement for accurate set identification, albeit still with a larger variance. We set the number of Monte Carlo iterations to $R = 500$. We understand the concern that this may be too little for eliminating all Monte Carlo biases. Yet we have to rely on such a small number as it is the absolute maximum that

our hardware allows for, even though we have optimized the code by employing parallel processing and delegation to C-based processes using the `numba` and `joblib` packages in `python`. All in all, this simulation still took 18 hours to complete with 6.5 million admissible IRFs drawn.

Finally, the squared error matrix of the selected median target IRF to the true DGP IRF is calculated, collapsed into a scalar total squared error by summation. Then we estimate the ratio of this squared error of a specific estimation approach to the model fitted by the true lag order p_0 . Therefore, the deviations of the true IRF are directly put into relation to how well a correctly specified model compares for the given sample. Finally, we estimate the MSE by calculating the geometric mean of these individual ratios. The geometric mean is preferred over the arithmetic mean as the ratios are bounded on the left side at 0, but unbounded on the right side. Outliers on the right side are hence overweighed in the arithmetic mean. The relative MSE for one method for one true lag order p_0 and one sample size T is given by the following formula:

$$MSE_{relative} = \exp \left(\frac{1}{n} \sum_{i=1}^n \ln(r_i) \right) \text{ where } r_i = \frac{TSE_i}{TSE_i^{p_0}} \quad (29)$$

While, for a single estimation it is possible to estimate millions of admissible IRFs, the Monte Carlo simulation approach doesn't allow for such a large number of draws. In establishing the median target IRF specific to the DGP, we set the number of discovered admissible IRFs to 1 million so as to have a very accurate median target IRF for the specific dataset. In the Monte Carlo Simulation, we reduce this number drastically to 100 as it has proven to be the absolute limit that still allows us to have an acceptable number of Monte Carlo iterations, 1000, for our set of different methods. Even with computational optimization by parallel processing, more draws are not applicable under our hardware constraints. For each individual iteration, the median target IRF is therefore less accurately determined. However, because we use relative MSEs as our performance measure and repeat the process 1000 times, this should not impact the interpretability of our results.

4 Results

First, we will look at the main comparison of performance between the different ICs and BVAR specifications in terms of recovering an accurate median target IRF. Table 2 displays the full set of relative MSEs. It reveals two main findings:

1. We confirm the results from Ivanov and Kilian (2005) about the performance of the different information criteria. AIC consistently outperforms both BIC and HQC, revealing that the strict penalization of the latter two towards low lag orders is biasing the IRF too much. On the other hand, AIC's milder lag penalization ensures a closer fit

to the true lag order p_0 for finite samples, thereby reducing bias. The reduction of the variance by selecting a p much lower than the true lag order selected by BIC and HQC, see Table 1, is not evidenced to reduce the coefficient variance enough to compensate for the additional bias.

2. Hedging the risk of lag selection by employing BMA is the best option in nearly all scenarios. For smaller sample sizes, $T = 240$ and $T = 360$, joint grid BMA is dominating all other methods by a solid margin. The OLS fitted BMA using AIC as the criterion is the best criterion in two scenarios, once without and once with geometric decay. However, for $p_0 = 10$ and $T = 600$, joint grid BMA's relative MSE is only slightly higher. Only for the highest sample size $T = 600$ and the lower true lag order $p_0 = 4$ does the conventional VAR fitted by an AIC-selected lag order perform the best. This may be explained by the finding that $\hat{p}^{AIC}$ has basically converged to p_0 at this sample size (See Table 1) while the sample size is already large enough in order to accurately determine the coefficients for four lags.

Table 3 makes the comparison to AIC for all different scenarios clearer by setting the MSE_{AIC} as the benchmark. Both OLS BMA with AIC as well as joint grid BMA outperform regular AIC in the majority of scenarios. All other versions of BMA are actually not consistently better than simple AIC. The sensitivity of the BMA hedging advantage to the exact specification requires further examination.

To understand this, we have to consider Figures 2 and 3 that present the weighting distribution of the different BMA methods for $p_0 = 4$. The reason for the BIC-weighted OLS BMA to fail is very apparent. The strong penalty of BIC forces it to rely very heavily on the lag order $p = 2$ in finite samples, thereby basically eliminating the diversification of BMA. As BIC generally doesn't produce models with good IRF inference, this also applies to these BMAs.

Once AIC is introduced to OLS BMA, the goal of spreading out the estimation on a selection of lag orders that are determined to be probable is achieved. AIC's weaker penalization of higher lag orders ensures that four to five different p s are assigned positive weights. The weighting becomes more concentrated on the true lag order $p_0 = 4$ as T increases. The introduction of the geometric decay parameter $\theta = 0.5$ distorts the weighting towards lower lag orders. This moves the estimation further away from the correctly fitted model, resulting in a larger underfitting bias. Table 3 shows that the reduction in parameter variance does not compensate for this bias.

Finally, the joint grid approach provides the most widely distributed weighting scheme. Figure 3 aggregates the weights from the grid at the lag order p -level, i.e., weights assigned to the combination of an individual p with different λ_1 -values are added up. For lower sample sizes, nearly all lag orders are considered in the BVAR estimated BMAs. Even for the largest $T = 600$, the highest weight is just above 50%. Hence, a wider diversification, also into the territory of overfitting lag orders $p > p_0$, can improve the fit of the determined

median target IRF if combined with BVAR estimation. The inclusion of higher lag orders at first seems counterproductive. Yet, due to the additional shrinkage of parameters with the BVAR estimation, it enables you to spread the risk of lag order uncertainty wider while maintaining accurate parameter estimation for the crucial lower lags. The combined optimization of weighting on lag order p as well as the hyperparameter λ_1 ensures that this shrinkage is amplified for those models that use overparameterized specifications. 4 illustrates how, as an example for the scenario of $T = 600$ and $p_0 = 4$, tighter average λ_1 -values are selected on average for the BVAR specifications of higher lag orders, i.e., the weights among different combinations of p and λ_1 are skewed more concentrated at the lower end of λ_1 as p increases above p_0 . Although MDD optimizes on Out-of-Sample performance and therefore may select lower than optimal tau-values for IRF accuracy (CITATION), it appears to be a good enough approximation to ensure good estimations for BMA. The shift towards lower lag orders introduced by the geometric decay parameter θ , on the other hand, cannot be evidenced to be of advantage for a BMA model estimated by the grid BVAR approach.


A caveat for the interpretation of these results, however, remains: the low number of Monte Carlo iterations and admissible IRFs drawn per model. We cannot exclude that some part of the variation in relative MSEs may be driven by pure sampling variation. For example, the general increase in relative MSEs across all methods at $T = 360$ and $p_0 = 4$ can barely be explained by the theoretical properties.

Table 2: Relative Mean Squared Error: Hard Selection vs. Bayesian Model Averaging (Including Informative Priors).


p_0	T	AIC	BIC	HQC	OLS BMA (BIC)	OLS BMA (Geom. BIC)	OLS BMA (AIC)	OLS BMA (Geom. AIC)	Joint Grid BMA	Geom. Grid BMA
4	240	1.083	1.108	1.129	1.100	1.163	1.082	1.175	0.900	1.127
	360	1.117	1.274	1.296	1.142	1.235	1.035	1.138	1.026	1.046
	480	0.986	1.133	1.124	1.148	1.150	1.031	0.996	0.898	1.130
	600	0.989	1.145	1.118	1.218	1.129	1.032	1.124	1.113	1.109
10	240	1.033	1.018	1.021	1.010	0.842	0.979	0.982	0.800	0.849
	360	0.811	0.904	0.878	1.163	1.057	0.893	0.884	0.787	0.993
	480	0.989	1.190	1.185	1.052	1.283	0.883	1.210	1.019	0.898
	600	1.036	1.260	1.259	1.293	1.100	0.883	0.875	0.878	0.964

Table 3: Relative Mean Squared Error of Bayesian Model Averaging Specifications vs. AIC Baseline.

p_0	T	OLS BMA (BIC)	OLS BMA (Geom. BIC)	OLS BMA (AIC)	OLS BMA (Geom. AIC)	Joint Grid BMA	Geom. Grid B
4	240	1.015	1.074	0.999	1.085	0.831	1.040
	360	1.023	1.106	0.927	1.019	0.919	0.937
	480	1.164	1.166	1.046	1.010	0.911	1.147
	600	1.231	1.141	1.043	1.136	1.125	1.122
10	240	0.978	0.815	0.948	0.950	0.775	0.821
	360	1.434	1.304	1.102	1.090	0.970	1.224
	480	1.064	1.297	0.893	1.224	1.031	0.908
	600	1.248	1.062	0.852	0.845	0.847	0.930




(a) $p_0 = 4$




(b) $p_0 = 10$


Figure 1: Performance of BMA vs Hard Selection at different sample sizes T .




(a) OLS BMA (BIC-weighted)



(b) OLS BMA (Geom. BIC-weighted, $\theta = 0.5$)




(c) OLS BMA (AIC-weighted)



(d) OLS BMA (Geom. AIC-weighted, $\theta = 0.5$)

Figure 2: Posterior Weight Evolution across the four OLS BMA architectures ($p_0 = 4$).



(a) Joint Grid BMA

(b) Geometric Grid BMA

Figure 3: Posterior Weight Evolution for the two BVAR-based BMAs ($p_0 = 4$).




Figure 4: Optimal λ_1 is adapted to the lag order p .

5 Conclusion

This paper revisits the fundamental challenge of lag order specification in Structural Vector Autoregressions, updating the foundational Monte Carlo analysis of Ivanov and Kilian (2005) to reflect modern empirical practices. Specifically, we evaluate the performance of traditional Information Criteria (IC) against modern Bayesian Model Averaging (BMA) techniques in recovering the true median target Impulse Response Function (IRF) within a set-identified framework using sign restrictions. Utilizing a realistic data-generating process based on the global oil market model of Kilian and Murphy (2014), we analyze the bias-variance trade-offs inherent in single-model selection versus model averaging across various finite sample sizes.

Our findings underscore two major practical implications for applied macroeconomists. First, when relying on single-model selection, the Akaike Information Criterion (AIC) remains the superior choice for IRF inference. Strict criteria such as the BIC and HQC penalize parameterization too heavily in finite samples, leading to a severe underfitting bias that heavily distorts structural dynamics. AIC’s milder penalization yields models that converge closer to the true lag order, confirming that avoiding underfitting bias is paramount for accurate IRF recovery, even if it comes at the cost of slightly higher parameter variance.

Second, and more importantly, our results demonstrate that hedging lag selection risk through Bayesian Model Averaging consistently outperforms single-model selection. In particular, the joint-grid BMA approach, which averages across both lag orders (p) and BVAR shrinkage hyperparameters (λ_1), emerges as the most robust methodology across the majority of sample sizes and true lag structures. This approach effectively mitigates model uncertainty by distributing weights across a broad spectrum of plausible lag orders. Crucially, it manages the risk of overfitting by dynamically assigning tighter shrinkage to overparameterized specifications ($p > p_0$), thereby preserving the precision of critical lower-lag estimates while still capturing higher-order dynamics. Conversely, introducing a geometric prior to force parsimony proved counterproductive, as it artificially shifted weight toward lower lags and reintroduced the very underfitting bias that BMA seeks to avoid. Only in scenarios with very large sample sizes ($T = 600$) and a low true lag order ($p_0 = 4$) does a simple AIC-selected OLS model rival the joint-grid BMA, as parameter variance naturally decreases.

We acknowledge that the severe computational intensity of set identification via sign restrictions imposes natural limitations on our simulation design, restricting the number of Monte Carlo iterations to $R = 500$ and admissible draws to 1,000 per iteration. While this introduces some sampling variation, the persistent outperformance of the joint-grid approach provides a compelling baseline. Future research could expand upon this framework by utilizing more efficient sampling algorithms or greater computational resources

to confirm these findings across a wider array of data-generating processes and variable dimensions.

Ultimately, this study advocates for a shift in standard empirical practice. Rather than staking IRF inference on a single, potentially misspecified lag order determined by an information criterion, researchers are well-advised to embrace Bayesian Model Averaging. By jointly averaging over lag structures and dynamically scaling parameter shrinkage, practitioners can significantly improve the credibility, robustness, and accuracy of their structural impulse response estimates.

References

References

- Akaike, H. (Dec. 1974). “A New Look at the Statistical Model Identification”. In: *IEEE Transactions on Automatic Control* 19.6, pp. 716–723. ISSN: 1558-2523. DOI: [10.1109/TAC.1974.1100705](https://doi.org/10.1109/TAC.1974.1100705). URL: <https://ieeexplore.ieee.org/document/1100705> (visited on 06/22/2026).
- Blake, Andrew and Haroon Mumtaz (Apr. 2015). “Applied Bayesian Econometrics for Central Bankers”. In: *Handbooks* 36.
- Blanchard, Olivier Jean and Danny Quah (1989). “The Dynamic Effects of Aggregate Demand and Supply Disturbances”. In: *American Economic Review* 79.4, pp. 655–673. URL: <https://ideas.repec.org/a/aea/aecrev/v79y1989i4p655-73.html> (visited on 06/21/2026).
- Burnham, Kenneth P. and David R. Anderson, eds. (2002). *Model Selection and Multimodel Inference*. New York, NY: Springer. ISBN: 978-0-387-95364-9. DOI: [10.1007/b97636](https://doi.org/10.1007/b97636). URL: <http://link.springer.com/10.1007/b97636> (visited on 06/21/2026).
- Canova, Fabio and Gianni De Nicoló (Sept. 1, 2002). “Monetary Disturbances Matter for Business Fluctuations in the G-7”. In: *Journal of Monetary Economics* 49.6, pp. 1131–1159. ISSN: 0304-3932. DOI: [10.1016/S0304-3932\(02\)00145-9](https://doi.org/10.1016/S0304-3932(02)00145-9). URL: <https://www.sciencedirect.com/science/article/pii/S0304393202001459> (visited on 06/22/2026).
- Castelnuovo, Efrem, Lorenzo Mori, and Gert Peersman (2024). *Commodity Price Shocks and Global Cycles: Monetary Policy Matters*. DOI: [10.2139/ssrn.4868719](https://doi.org/10.2139/ssrn.4868719). Social Science Research Network: [4868719](https://papers.ssrn.com/abstract=4868719). URL: <https://papers.ssrn.com/abstract=4868719> (visited on 06/23/2026). Pre-published.
- Christiano, Lawrence, Martin Eichenbaum, and Charles Evans (1999). “Monetary Policy Shocks: What Have We Learned and to What End?” In: *Handbook of Macroeconomics*. Vol. 1, Part A. Elsevier, pp. 65–148. URL: <https://EconPapers.repec.org/RePEc:eee:macchp:1-02> (visited on 06/21/2026).
- Faust, Jon (Dec. 1, 1998). “The Robustness of Identified VAR Conclusions about Money”. In: *Carnegie-Rochester Conference Series on Public Policy* 49, pp. 207–244. ISSN: 0167-2231. DOI: [10.1016/S0167-2231\(99\)00009-3](https://doi.org/10.1016/S0167-2231(99)00009-3). URL: <https://www.sciencedirect.com/science/article/pii/S0167223199000093> (visited on 06/21/2026).
- Ferreira, Thiago R. T., Daniel A. Ostry, and John Rogers (June 22, 2023). “Firm Financial Conditions and the Transmission of Monetary Policy”. In: URL: <https://www.federalreserve.gov/econres/feds/firm-financial-conditions-and-the-transmission-of-monetary-policy.htm> (visited on 06/23/2026).

- Fry, Renée and Adrian Pagan (2011). “Sign Restrictions in Structural Vector Autoregressions: A Critical Review”. In: *Journal of Economic Literature* 49.4, pp. 938–960. ISSN: 0022-0515. JSTOR: [23071662](https://www.jstor.org/stable/23071662). URL: <https://www.jstor.org/stable/23071662> (visited on 05/20/2026).
- Gazzani, Andrea, Fabrizio Venditti, and Giovanni Veronese (May 1, 2024). “Oil Price Shocks in Real Time”. In: *Journal of Monetary Economics* 144, p. 103547. ISSN: 0304-3932. DOI: [10.1016/j.jmoneco.2023.12.005](https://doi.org/10.1016/j.jmoneco.2023.12.005). URL: <https://www.sciencedirect.com/science/article/pii/S0304393223001630> (visited on 06/23/2026).
- Gertler, Mark and Peter Karadi (Jan. 1, 2015). “Monetary Policy Surprises, Credit Costs, and Economic Activity”. In: *American Economic Journal: Macroeconomics* 7.1, pp. 44–76. ISSN: 1945-7707, 1945-7715. DOI: [10.1257/mac.20130329](https://doi.org/10.1257/mac.20130329). URL: <https://pubs.aeaweb.org/doi/10.1257/mac.20130329> (visited on 04/25/2026).
- Hannan, E. J. and B. G. Quinn (1979). “The Determination of the Order of an Autoregression”. In: *Journal of the Royal Statistical Society. Series B (Methodological)* 41.2, pp. 190–195. ISSN: 0035-9246. JSTOR: [2985032](https://www.jstor.org/stable/2985032). URL: <https://www.jstor.org/stable/2985032> (visited on 06/22/2026).
- Hoeting, Jennifer A. et al. (1999). “Bayesian Model Averaging: A Tutorial”. In: *Statistical Science* 14.4, pp. 382–401. ISSN: 0883-4237. JSTOR: [2676803](https://www.jstor.org/stable/2676803). URL: <https://www.jstor.org/stable/2676803> (visited on 06/20/2026).
- Ivanov, Ventzislav and Lutz Kilian (Jan. 14, 2005). “A Practitioner’s Guide to Lag Order Selection For VAR Impulse Response Analysis”. In: *Studies in Nonlinear Dynamics & Econometrics* 9.1. ISSN: 1558-3708. DOI: [10.2202/1558-3708.1219](https://doi.org/10.2202/1558-3708.1219). URL: <https://www.degruyter.com/document/doi/10.2202/1558-3708.1219/html> (visited on 04/22/2026).
- Kilian, Lutz (June 2009). “Not All Oil Price Shocks Are Alike: Disentangling Demand and Supply Shocks in the Crude Oil Market”. In: *American Economic Review* 99.3, pp. 1053–1069. ISSN: 0002-8282. DOI: [10.1257/aer.99.3.1053](https://doi.org/10.1257/aer.99.3.1053). URL: <https://www.aeaweb.org/articles?id=10.1257/aer.99.3.1053> (visited on 06/19/2026).
- Kilian, Lutz and Helmut Lütkepohl (2017). *Structural Vector Autoregressive Analysis. Themes in Modern Econometrics*. Cambridge: Cambridge University Press. ISBN: 978-1-107-19657-5. DOI: [10.1017/9781108164818](https://doi.org/10.1017/9781108164818).
- Kilian, Lutz and Daniel P. Murphy (Apr. 1, 2014). “The Role of Inventories and Speculative Trading in the Global Market for Crude Oil”. In: *Journal of Applied Econometrics* 29.3, pp. 454–478. ISSN: 1099-1255. DOI: [10.1002/jae.2322](https://doi.org/10.1002/jae.2322). URL: <https://onlinelibrary.wiley.com/doi/10.1002/jae.2322> (visited on 05/12/2026).
- Lütkepohl, Helmut (1985). “Comparison of Criteria for Estimating the Order of a Vector Autoregressive Process”. In: *Journal of Time Series Analysis* 6.1, pp. 35–52. ISSN: 1467-9892. DOI: [10.1111/j.1467-9892.1985.tb00396.x](https://doi.org/10.1111/j.1467-9892.1985.tb00396.x). URL: <https://doi.org/10.1111/j.1467-9892.1985.tb00396.x>

- [//onlinelibrary.wiley.com/doi/abs/10.1111/j.1467-9892.1985.tb00396.x](https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1467-9892.1985.tb00396.x) (visited on 04/22/2026).
- Lütkepohl, Helmut (2005). *New Introduction to Multiple Time Series Analysis*. Berlin, Heidelberg: Springer. ISBN: 978-3-540-40172-8 978-3-540-27752-1. DOI: [10.1007/978-3-540-27752-1](https://doi.org/10.1007/978-3-540-27752-1). URL: <http://link.springer.com/10.1007/978-3-540-27752-1> (visited on 06/21/2026).
- Madigan, David and Adrian E. Raftery (Dec. 1, 1994). “Model Selection and Accounting for Model Uncertainty in Graphical Models Using Occam’s Window”. In: *Journal of the American Statistical Association* 89.428, pp. 1535–1546. ISSN: 0162-1459. DOI: [10.1080/01621459.1994.10476894](https://doi.org/10.1080/01621459.1994.10476894). URL: <https://doi.org/10.1080/01621459.1994.10476894> (visited on 06/23/2026).
- Manski, Charles F (2003). *Partial Identification of Probability Distributions*. Springer Series in Statistics. New York: Springer-Verlag. ISBN: 978-0-387-00454-9. DOI: [10.1007/b97478](https://doi.org/10.1007/b97478). URL: <http://link.springer.com/10.1007/b97478> (visited on 06/21/2026).
- Miranda-Pinto, Jorge (Oct. 2023). “Monetary Policy Transmission through Commodity Prices”. In: *IMF Working Papers* 2023.215, p. 1. ISSN: 1018-5941. DOI: [10.5089/9798400256691.001](https://doi.org/10.5089/9798400256691.001). URL: <https://elibrary.imf.org/openurl?genre=journal&issn=1018-5941&volume=2023&issue=215&cid=540373-com-dsp-crossref> (visited on 06/23/2026).
- Raftery, Adrian E. (1995). “Bayesian Model Selection in Social Research”. In: *Sociological Methodology* 25, pp. 111–163. ISSN: 0081-1750. DOI: [10.2307/271063](https://doi.org/10.2307/271063). JSTOR: [271063](https://www.jstor.org/stable/271063). URL: <https://www.jstor.org/stable/271063> (visited on 06/20/2026).
- Rubio-Ramírez, Juan F., Daniel F. Waggoner, and Tao Zha (Apr. 2010). “Structural Vector Autoregressions: Theory of Identification and Algorithms for Inference”. In: *Review of Economic Studies* 77.2, pp. 665–696. ISSN: 00346527, 1467937X. DOI: [10.1111/j.1467-937X.2009.00578.x](https://doi.org/10.1111/j.1467-937X.2009.00578.x). URL: <https://academic.oup.com/restud/article-lookup/doi/10.1111/j.1467-937X.2009.00578.x> (visited on 06/03/2026).
- Schwarz, Gideon (1978). “Estimating the Dimension of a Model”. In: *The Annals of Statistics* 6.2, pp. 461–464. ISSN: 0090-5364. JSTOR: [2958889](https://www.jstor.org/stable/2958889). URL: <https://www.jstor.org/stable/2958889> (visited on 06/21/2026).
- Sims, Christopher A. (1980). “Macroeconomics and Reality”. In: *Econometrica* 48.1, pp. 1–48. ISSN: 0012-9682. DOI: [10.2307/1912017](https://doi.org/10.2307/1912017). JSTOR: [1912017](https://www.jstor.org/stable/1912017). URL: <https://www.jstor.org/stable/1912017> (visited on 06/20/2026).
- Steel, Mark F. J. (Sept. 1, 2020). “Model Averaging and Its Use in Economics”. In: *Journal of Economic Literature* 58.3, pp. 644–719. ISSN: 0022-0515. DOI: [10.1257/jel.20191385](https://doi.org/10.1257/jel.20191385). URL: <https://pubs.aeaweb.org/doi/10.1257/jel.20191385> (visited on 06/20/2026).

Uhlig, Harald (Mar. 2005). “What Are the Effects of Monetary Policy on Output? Results from an Agnostic Identification Procedure”. In: *Journal of Monetary Economics* 52.2, pp. 381–419. ISSN: 03043932. DOI: [10.1016/j.jmoneco.2004.05.007](https://doi.org/10.1016/j.jmoneco.2004.05.007). URL: <https://linkinghub.elsevier.com/retrieve/pii/S0304393205000073> (visited on 06/11/2026).

Declaration of AI Usage

During the preparation of this seminar paper, I, [NAME], used Claude, Gemini, and ChatGPT for help with coding, research, and adapting writing to academic standards. I have reviewed and edited the content as needed and take full responsibility for the content.

[NAME]