

Self-Referential Dissipative Systems: A Materials Test Across Ten Substrates

Ferdinand M. Schessl, M.Sc.*

June 2026

In memory of my grandfather Franz Scheßl (1929–2024)

Abstract

The same sigmoid functional form recurs across ten physically heterogeneous substrates, read off from a single cumulative measurement quantity along an ordered load history. The finding is grounded in the tensile test: on fiber-parallel loaded spruce specimens (*Picea abies*, DIN 52188:1979-05, tension per DIN EN 408:2012-10), the nonlinearity residual $Q(x) = |\sigma - E\varepsilon| / \max$ of the stress-strain curve normalized to $[0, 1]$, taken over $x = \sigma / \sigma_{\max}$, fits a logistic sigmoid whose inflection location $\hat{a}$ reads the load-bearing capacity. Across $n = 230$ specimens this location correlates with fracture stress at Pearson $r = -0.83$ ($p < 10^{-30}$); this coupling is largely an arithmetic consequence of the onset normalization (instrument validation on known sorting physics, not an independent mechanism, §4.1). The mean inflection location $\tilde{a} = 0.567$ carries the fiber anisotropy, an early inflection signals lower fracture stress. The frame is materials testing applied isomorphically to ordered load histories: the specimen bears a load, accumulates damage, passes a yield point, and becomes irreversible. Every conversation, market trajectory, or autocratization episode is such a specimen, not a bag of features. The burden of proof is thus reversed: a falsification would have to exhibit a finding that violates the mechanics (§8).

The class of self-referential dissipative systems (SRDS) fixes five axioms A0–A4 (self-reference, dissipation, accumulation, transition, irreversibility); only A0 and A1 are primitive, A2–A4 follow as theorems under four regularity conditions L1–L4 (load-bearing L2, the single-peaked state-dependent damage rate, and L4, the return-and-elasticity protocol). Three core results carry it: T1 signs the conditional phase-space volume rate $\text{Tr } A$ through the sign of the windowed trend of mutual information; T2 secures, under contraction, a unique non-equilibrium steady state; T3 (Goldilocks principle) places optimal function near criticality; the inflection is the order parameter. A computer-assisted no-go theorem excludes a universal numeric inflection constant ($\hat{Q}(1/2) \in [1.8412, 1.8963]$, Arb certificate + Lean-4 machine-checked assembly over 12 declared project axioms, 2 numerical + 10 operator, public): what is universal is the functional form, not the numeric value; the inflection location is substrate-specifically measured, not a universal constant. Under substrate-specific normalization the class family carries across timber, multiaxial fatigue ($n = 914$, 136 materials; $\hat{a}$ reads the load-path non-proportionality), financial crashes (coherence beyond window anchoring; Bear2022 0/500 surrogates), lignin (boundary case without an inner $\hat{a}$; $\rho_S = -0.78$), V-Dem autocratization episodes ($n = 117$; Gompertz class gain $3.4\times$ against matched nulls), earthquakes ($n = 284$; instrument validation, $\rho_S = +0.673$ Bonferroni), battery degradation ($n = 26$, KS $D = 0.733$), colorectal carcinomas ($n = 579$; $\Delta\hat{a}$ = stage occupancy, not curve form), and music ($n = 2,840$; sequence-carried, median $R^2 = 0.975$). For conversation the load history (lag-1 $\bar{\rho}_1 = 0.928$) is signal, not a nuisance term, and five axes coupled via $\sigma = E\varepsilon$ are read anti-sparse: across $n = 202$ specimens $\hat{a}$ correlates univariately with manipulation ($\rho = -0.359$, cluster-robust), the chat-level ridge reaches $\rho = 0.700 \pm 0.076$, replicated across twelve corpora ($\approx 35,300$ dialogues from 14 sources). Synthetic null specimens separate sharply (KS $p < 10^{-3}$ on all five substrates tested), a pre-registered null control (ProsocialDialog) and two falsified positive predictions (Sotopia, Google

*M.Sc. Civil Engineering, Technical University of Munich (Alumnus). Correspondence: ferdinand.schessl@web.de. Companion repository: <https://github.com/FerdinandSchessl/srds-enk-companion>

COVID Mobility) map the scope conditions, six of the ten substrates remain Bonferroni-compliant under the safety factor γ_M . Lean-4 proofs, metric code, and the gold corpus are open and reproducible; the operational calibration thresholds remain withheld (responsible disclosure): no finding that carries the cross-substrate thesis depends on them. The structural load-bearing capacity is threshold-independent.


Keywords: large language models, conversation analysis, AI safety, cross-domain validation, self-referential dissipative systems, sigmoid inflection, falsifiability.

Contents

1	Question and Anchor	4
1.1	The Physical Experiment	4
1.2	The Paradigm: Materials Testing, Applied Isomorphically	4
1.3	The General Form	5
1.4	What This Paper Delivers	8
2	SRDS Framework and Mathematical Load-Bearing Capacity	8
2.1	Two Core Equations	8
2.2	Five Axioms A0–A4 + Four Regularity Conditions L1–L4	9
2.3	Two-Scale Mapping + Renormalization Operator	9
2.4	Aggregation Modes and Mode Mixture	10
2.5	Mathematical Load-Bearing Capacity: Birkhoff-Hopf, No-Go, T1–T3	11
3	Measurement Apparatus ENK (Material Level: Conversation as Specimen)	14
3.1	Embedding Geometry as Measurement System	14
3.2	Five Effective Dimensions: the Anti-Sparse Rule	14
3.3	Gold Reference Corpus + Cluster-Robust Inference	16
3.4	Crystallization Signature: Stiffening as Damage	16
3.5	Structural Axes: What the Five Main Axes Bear	17
3.6	Comparison with Message-Level Detectors	18
4	Cross-Domain: Ten Substrates, One Functional Form	18
4.1	Timber under Tensile Load: DIN EN 408 (Anchor)	18
4.2	Multi-Material Multiaxial Fatigue: $\hat{a}$ Reads the Non-Proportionality	19
4.3	Finance SPY: Crash S-Curve against Calm Random-Walk	19
4.4	Lignin: Concave Digestion Kinetics without an Interior $\hat{a}$ (Cross-Sectional)	19
4.5	V-Dem: Autocratization Trajectories (Two-Span Beam)	20
4.6	Political Speeches: Support Topology, Not an Effect Claim	20
4.7	Earthquakes USGS: Instrument Validation on the Sequence Statistics	21
4.8	Protein Melting Curves (Meltome Atlas): Instrument Validation across Organism Classes	21
4.9	Battery Degradation, CALB (real): Knee + Hysteresis Accumulation	21
4.10	Colorectal Carcinomas: TCGA (Stage-Occupancy Finding, Reduced)	22
4.11	LLM Conversations (ENK Gold)	22
4.12	Music: MetaMIDI (ℓ_2), Pre-Registered + Cross-Method-Validated	22
4.13	Materials-Class Hierarchy	23
4.14	Cross-Substrate Status and Mutual Support	24

5	Structural Scale, Boundary Conditions, and Addressee-Support	26
5.1	Five Scales, Five Sub-Systems: the Luhmann Cut	26
5.2	Five Support Types: Structural Mechanics of Conversation	27
5.3	Addressee-Support: the Measurement Configuration as an API Axis	28
5.4	Propaganda Mechanics under Missing Restoring Force	29
5.5	Empirical Truss Evidence: Institutional Support as Counterforce	30
5.6	Budget State Machine: Architecture without Disclosed Thresholds	30
5.7	Application Sketch: ℓ_3 -Trajectory Monitoring	31
6	Conversation Cross-Corpus and Cross-Method Validation	31
6.1	Twelve Conversation Corpora (plus Two Cross-Domain Extensions)	31
6.2	LegalCon as Showcase: Highest Singular Cross-Corpus Effect	32
6.3	Sign Paradox: Ductile vs. Brittle Fracture	33
6.4	Cross-Method Triangulation: Three Methodologically Disjoint Apparatuses	33
6.5	One Predicted Null and Two Scope Conditions: Lemma Violations as Construct Support	34
7	Cognition Bridge and External Frame Convergence	35
7.1	Cognition Bridge: G_f/G_c and the Sigmoid Mélange	35
7.1.1	Chen et al. 2026: Long-CoT as Molecular Structure	35
7.2	External Frame Convergence: The Gap That SRDS Fills	35
8	Falsification Anchors, State of Measurement, and Limits	37
8.1	Prospective Acceptance Rule: What Is Fixed Before Measurement	37
8.2	The Falsification Lattice	38
8.3	Three Falsification Layers	38
8.4	Fit $\leftrightarrow$ Counterpart: the Nulls Are the Proof	40
8.5	ENK against Message-Level Detectors	40
8.6	Limits	41
8.7	What Remains: The Structural Limit	41
9	What Remains When Something Is Missing	41
9.1	Summary of Findings	41
9.2	Open Replication Tasks	42
9.3	Data and Code Availability	42
9.4	Genealogy	44
9.5	Disclosure	44
9.6	Why Physics Does Not Stop at Self-Referential Systems	44
9.7	Closing Remark	45

The physical anchor: tensile test per DIN EN 408



Whoever yields early breaks at lower load. The inflection point $\hat{a}$ determines fracture stress.

Figure 1: Bone-shaped timber specimen (spruce, DIN 52188:1979-05, fiber-parallel) under tension per DIN EN 408:2012-10. Right: nonlinearity residual $Q(x)$ with sigmoid fit across $n = 230$ specimens; $\hat{a}$ correlates with fracture stress at $r = -0.83$, $p < 10^{-30}$, mean inflection $\tilde{a} = 0.567$.

1 Question and Anchor

1.1 The Physical Experiment

A spruce timber specimen, bone-shaped per DIN 52188:1979-05, with a cross-section of 50×10 mm in the test region, was tested fiber-parallel under tension per DIN EN 408:2012-10 (Figure 1). The stress-strain curve follows the familiar sequence of linear-elastic regime, yield point, plastic deformation, necking, and fracture; a material holds because it responds to stress: by yielding and redistributing load. Across $n = 230$ specimens we fit the nonlinearity residual $Q = |\sigma - E\varepsilon|/\max$ with a logistic sigmoid $Q(x)$ over the normalized stress $x = \sigma/\sigma_{\text{max}}$ and extract its inflection point $\hat{a}$ ($\tilde{a} = 0.567$); $\hat{a}$ correlates with fracture stress at Pearson $r = -0.83$, $p < 10^{-30}$. An early inflection corresponds to a lower fracture stress. This relationship is established in timber engineering. The question of this work is why the same functional form recurs in nine further substrates: metals under cyclic load, markets, lignin pulping, democratic indices, earthquake sequences, battery trajectories, tumor mutation loads, LLM conversations, musical note-adjacency networks.

1.2 The Paradigm: Materials Testing, Applied Isomorphically

The wood tensile test is not an illustration but the construct anchor. SRDS applies the rules of continuum mechanics and statics isomorphically to sequences under load: to a conversation, a market trajectory, an autocratization episode as much as to a test specimen. The claim is not that language *behaves like* a material, but that the same structural laws (stress-strain coupling, damage accumulation, support reaction, fracture as Mode-I separation) hold once a self-referential system takes up and redistributes load. Prevailing statistical evaluation practice treats such sequences as collections of independent features; by construction it cannot resolve the coupled material response (strain read along several axes of one $\sigma = E\varepsilon$ law) that SRDS measures.

A reversed burden of proof follows. We do not defend the mechanics case by case; it is the frame




Figure 2: Nonlinearity residual $Q(x)$ of timber (spruce, $n = 230$, $\tilde{a} = 0.567$) and cumulative semantic drift of ENK-Gold ($n = 202$, $\hat{a} = 0.232$).

within which we measure. A falsification would have to exhibit a finding that *violates* the mechanics: a yield-point location that says nothing about strength; a truss that bends under axial load; a load accumulation that spontaneously reverses. These falsification tests are stated explicitly in §8.

The frame also delimits *what holds when*. Four physical axes of demarcation: (i) **forced vs. free sequence**: under machine-imposed ordering (tensile test) a path surrogate is physically meaningless, the existence test being model comparison; under free ordering (conversation, market, earthquake) the surrogate is mandatory; (ii) **uniaxial vs. multiaxial**: inflection and amplitude are two descriptors of *one* stress–strain response, coupled through $\sigma = E\varepsilon$, hence not separable quantities but the same material response read along two axes; (iii) **support topology**: it decides, before any measurement, which axis bears load and which falls into the null member (§5); (iv) **no universal constant** (no-go theorem, §2.5): what is shared is the *functional form*, not the numeric value. The axes are not auxiliary assumptions but consequences of the applied mechanics; they fix the scope of each test before the measurement.

1.3 The General Form

The question is not “How does one detect manipulation in a chat?”, it is more general: what happens structurally when a self-referential system stands under load? Table 1 lists ten substrates across eight research fields (Figure 3; timber and ENK conversation compared directly in Figure 2): materials science (timber, metals, battery), geophysics (earthquakes), biochemistry (lignin), finance (SPY), democracy (V-Dem), oncology (colorectal carcinomas), language (LLM conversations), music (MetaMIDI). The cumulative response follows a sigmoid saturation function with well-defined inflection in each; the substrate-specific form within this class family (Logistic, Weibull, Gompertz, Hill) is resolved in §2.4. The carriers differ: the class family of the functional form is shared, the specific form within it is substrate-specific.

Table 1: Ten substrates, common sigmoid class family with substrate-specific normalization Π and form

Substrate	Π	n	Headline
Timber (<i>Picea abies</i> , DIN-EN-408)	$Q(x)$	230	$r(\hat{a}, \sigma_{\text{break}}) = -0.83$, $\tilde{a} = 0.567$ (continued on next page)

(continued)

Substrate	Π	n	Headline
Multi-Material Multiaxial Fatigue (136 materials)	$\sigma(\varepsilon)$	914	$\hat{a}$ reads non-proportionality; accumulation in the ENK band (44–62% Chelton)
Finance SPY (1999–2026)	cumsum	10+	crash coherence beyond window anchoring (class separation $p = 0.019$; Bear2022 0/500); crash $R^2 = 0.973$ vs. calm 0.483
Lignin hydrothermal (autohydrolysis) (SP-LCC)	$\sigma(\varepsilon)$	72 cond.	$\rho_S = -0.78$ (Weibull, mode ii)
V-Dem ERT-v14 autocratization	cumsum	117 (89 fitted)	class comparison: Gompertz $3.4\times$ against matched nulls; partition, material memory, hysteresis
Earthquakes USGS, 6 regions	cumsum	284 seq.	instrument validation on the sequence statistics: $\rho(\hat{a}, \text{mainshock}) =$ $+0.673$ Bonferroni KS $D = 0.733$ ($p < 10^{-13}$); holds against
Battery CALB (arXiv:2502.18807)	cumsum	26 cells	curvature-matched null ($D = 0.692$); knee localization + G7 hysteresis
Colorectal TCGA	cumsum	579	MSI–MSS $\Delta\hat{a} = -0.153 = \text{stage}$ occupancy, not curve form (reduced, \$4.9)
LLM conversations (ENK gold)	cumsum	202	$\rho = -0.359$ cluster-robust, ridge $\rho = 0.700$
Music (MetaMIDI + music21 + Tagtraum)	cumsum	2,840	6-genre $\tau = +1.000$ (sequence-carried, \$4.11); median $R^2 = 0.975$

What must redistribute, deforms; what can no longer redistribute, breaks. This path-dependence of load-bearing capacity corresponds to the institutional path-dependence reading in historical institutional economics (North, 1990; Pierson, 2000; Mahoney, 2000). The structural-mathematical statement that recurs across ten substrates is not confined to conversation analysis; it was merely measured there first. Across twelve conversation corpora ($\approx 35,300$ dialogues from 14 sources) the same mechanism replicates; ProsocialDialog serves as the pre-registered null control; Sotopia and Google COVID Mobility, pre-registered with directional positive hypotheses, were falsified and thereby map the scope conditions.

Sigmoid inflection $\hat{a}$ across four substrates




Figure 3: Sigmoid fit of cumulative response across four substrates: timber ($\hat{a} = 0.567$), financial crashes ($\hat{a} = 0.62$ DotCom / 0.97 GFC), V-Dem autocratization (89 episodes), LLM conversations ($\hat{a} = 0.232$). Full list in Table 1.

1.4 What This Paper Delivers

This paper delivers three contributions. (i) SRDS (Self-Referential Dissipative Systems) as a formal framework with five axioms A0–A4, three core theorems T1–T3, four aggregation modes (parallel-instantaneous, serial-cascaded, multiaxial, continuously-accumulating; substrates as mode mixtures, §2.4), and a no-go theorem (§2.5) against universal constants outside mirror symmetry, connected to formal mathematics (de Bruijn, 1968; Harrison et al., 2014; Mathlib Community, 2020). Full development and proofs are in the formal companion framework (Schessler, 2026b), the Lean-4 formalization in the companion `srds-enk-companion/lean4/PositivityProofs/`. (ii) ENK (Emergent Narrative Control, here operationalized as an embedding-geometry apparatus) as a special case for the conversation substrate: gold-annotated corpus $n = 202$, twelve external replication corpora, five measurement axes, cluster-robust inference at turn-pair level (cumulative family $\bar{\rho}_1 = 0.928$; Schessler, 2026a). (iii) Falsifiability via synthetic null specimens (1,000 per substrate, KS-test $p < 10^{-3}$ on all five substrates tested against AR(1)/RW/WN), a model comparison against the saturation class family (Gompertz/Weibull/Hill), three axiom-violation controls with *predicted* null signals, and Bonferroni-conservative reporting.

Under the conservative safety factor γ_M Bonferroni ($\gamma_M = \alpha/9 \approx 0.0056$; nine per-substrate tests, with multiaxial fatigue as the tenth substrate carrying separately on the accumulation axis; family and per-substrate test §8), six of the ten substrates are conformant (Timber, ENK Conversation, Earthquakes, Battery under caveat, Music (sequence-carried, median $R^2 = 0.975$), V-Dem via the class comparison with partition, material memory, and hysteresis). Multi-Material Multiaxial Fatigue carries on the accumulation axis (autocorrelation structure in the ENK band, the Chelton inflation band of the ENK conversation substrate at 44–62%; $\hat{a}$ reads the load-path non-proportionality) with pooled-damped cross-material outcome coupling; three carry nominally with named caveats (Lignin, Colorectal, Finance with anchoring-calibrated coherence); in addition, there are a pre-registered null control (ProsocialDialog) and two falsified positive predictions that map the scope conditions (Sotopia, Google COVID Mobility). The pattern is the finding: the signal appears *where the mechanics predicts it* and is absent *where the mechanics predicts its absence*. A predicted null is thus not a gap but a passed falsification test (§8).

Substrate-specific deviations of the inflection location are reported not as confirmation of a universal value but as the measured asymmetry of each carrier. Axiomatics, Lean-4 proofs, gold corpus, metric code, and cross-substrate raw data lie open (companion directory `srds-enk-companion/`); the operational calibration thresholds of the state machine remain withheld. For a deployed stability detector, publishing the exact trigger values would enable targeted evasion (responsible disclosure). The structural architecture, axioms, theorems, and Lean-4 proofs are fully published; no reported finding depends on the thresholds, since the structural load-bearing capacity is threshold-independent. Simon et al. (2026) has concurrently proposed a programmatically related discipline, *learning mechanics* (arXiv:2604.21691); it is the convergence anchor to which this work independently converges.

The following sections develop the formal framework (§2), the measurement apparatus (§3), the cross-domain findings (§4), the structural scale (§5), the cross-corpus methodology (§6), the external convergence (§7), and the falsification anchors (§8).

2 SRDS Framework and Mathematical Load-Bearing Capacity

2.1 Two Core Equations

The structural grammar of SRDS consists of two coupled recursions. The first describes the evolution of state under external excitation:

$$z_{t+1} = F_t(z_t, u_t).$$

The second describes the evolution of the operator that propagates the state:

$$F_{t+1} = \Phi(F_t, z_t, u_t).$$

The first equation is classical feedback structure (Wiener, 1948; Ashby, 1956). It alone does not suffice: a specimen remembers its loading history (Bauschinger effect), a conversation carries path-dependence. The second equation makes this memory explicit: the operator F_t is itself the state of a superordinate system that changes its own response function over the course of loading. A structural fixed point in the sense of autopoiesis (Maturana and Varela, 1980) lies where $\Phi(F^*, z^*, u^*) = F^*$.

2.2 Five Axioms A0–A4 + Four Regularity Conditions L1–L4

Table 2: Axioms of the SRDS class

#	Axiom	Form	Substrate analogy
A0	Self-reference	$F_{t+1} = \Phi(F_t, z_t, u_t)$	Material memory (Bauschinger)
A1	Dissipation	Lyapunov functional V : $V(F_{t+1}) \leq V(F_t)$ in expectation (Prigogine, 1980)	Heat export, local order through exported entropy
A2	Accumulation	Damage Σ monotone: $\Sigma(F_{t+1}) \geq \Sigma(F_t)$ (Kachanov, 1958; Lemaitre, 1985)	Damage parameter, crack-area fraction, budget accumulator
A3	Transition	$\sigma(\Sigma)$ possesses a unique inflection point $\hat{a}$ (sigmoid form)	Yield point, tipping point (Scheffer et al., 2009)
A4	Irreversibility	Trajectory with $\Sigma > \hat{a}$ returns with probability < 1 (Seifert, 2012)	Hysteresis, plastic deformation

Explicit reduction. A2, A3, A4 are, under regularity conditions L1–L4, theorems derived from A0+A1: L1 (Lyapunov semi-order), L2 (saturation bound), L3 (asymmetry measure in the sense of stochastic thermodynamics), L4 (return protocol + elasticity below yield). Concretely: A2 follows from A0+A1+L1+L2; A3 from A2+L2; A4 from A0+A1+L3 (unconditional) or +L4 (conditional) (reduction theorem, proved in full in the formal companion framework (Schessl, 2026b)). The reduction is made explicit so that the load-bearing capacity of each axiom can be tested per substrate, rather than assumed uniformly across all substrates. For the predicted null (ProsocialDialog) it is precisely this explicit reduction that explains the null finding (L1 violation, $\beta \rightarrow 0$, recovery balances damage); for Sotopia (A0, symmetry instead of asymmetric self-reference) and Google COVID Mobility (L3 / $\sigma \perp \varepsilon$, both time-driven) it explains *post-hoc* why pre-registered positive predictions were falsified and maps the scope conditions (§8).

2.3 Two-Scale Mapping + Renormalization Operator

SRDS separates two scales: material level (single specimen, load-bearing capacity on the sample) and structural level (system into which the specimen is built). The renormalization operator M links them:

$$M_{\ell \rightarrow \ell'} = N_{\ell'} \circ R_{\ell \rightarrow \ell'} \circ C_{\ell \rightarrow \ell'}.$$

C **coarse-graining:** micro-trajectories to blocks (windows, clusters, representative volumes). R **rescale:** effective parameters on the coarser scale follow parameter flow $\theta_\ell \mapsto \theta_{\ell'}$. N **normalization:** extracts dimensionless invariants so that the renormalized description sits on the same structural tier as the source description.

Five operative tiers carry the empirical work: token (ℓ_0 , word embeddings) $\rightarrow$ turn-pair (ℓ_1 , semantic velocity, frame drift) $\rightarrow$ chat (ℓ_2 , $\hat{a}$, budget accumulator) $\rightarrow$ corpus (ℓ_3 , $\hat{a}$ distribution) $\rightarrow$ society (ℓ_4 , democracy index trajectories). Under the five axioms the dynamics on each scale admit a common normal form $\dot{z} = F(z, \Sigma) - \nabla_z \Pi(z, \Sigma) + \Theta \cdot \nabla_z I(z)$, form-invariant under M (Kadanoff-Wilson signature). Micro-details drop out under coarse-graining; the structural grammar remains.

2.4 Aggregation Modes and Mode Mixture

Under load the sigmoid form of a substrate follows one of four aggregation modes of the micro-imperfection cascade. Loading meets micro-imperfections, which activate in one of four ways: in parallel (CLT approximation), serially in a cascade (weakest-link statistics), multiaxially (hybrid), or continuously (diffusive averaging without a discrete failure event). The classical materials-science ductile/brittle typology (Considère, 1885; Tresca, 1864; Mohr, 1900; Bridgman, 1944) is a binary projection of this finer partition; SRDS distinguishes the modes through the sigmoid class family on $\Pi_S(X, Y)$, the image of the raw trajectory (X, Y) under the substrate-specific normalization Π from Table 1 into $[0, 1]^2$ (X : load or ordering variable, Y : response variable): Logistic, Weibull, Gompertz, and Hill are data-borne form classes on a single substrate, not the result of separate tests.

Table 3: Aggregation modes and form class

Mode	Activation	Form class	Typical $\tilde{a}$	Empirical anchor
(i) parallel-instantaneous	all micro-elements simultaneously under load, abrupt failure event	Logistic (CLT)	≈ 0.55	Logistic dominant in 6/8 material substrates of the model comparison (NR60 100%, FKM 80%, battery 100%; median $R^2 \geq 0.94$); ENK 100% across 12 conversation corpora
(ii) serial-cascaded	micro-failures cascade hierarchically	Weibull (weakest-link via order statistics)	≈ 0.30	CCP60 (cold-castable polyether, soft PE; public Dryad dataset): Weibull AIC-win (0% Logistic, median $R^2 = 0.96$, $n = 9$), $\rho(\hat{a}, \text{outcome}) = +0.867$: weakest-link failure of the polymer chain network
(iii) multiaxial	tensile and compressive components simultaneously	hybrid (Hill/Gompertz)	—	Multi-Material Multiaxial Fatigue ($n = 914$, 136 materials, Materials Cloud public): non-proportional multiaxial load; form partition as refinement
(iv) continuously-accumulating	gradual micro-defect accumulation without discrete failure event	Gaussian/Logistic (diffusive averaging)	≈ 0.55	Battery single-cell Logistic (CALB, arXiv:2502.18807, $n = 26$)

The binary creep/brittle-fracture reading remains useful as operating-range heuristic: creep $\approx$ modes (i) and (iv) with late inflection and slowly growing accumulation; brittle fracture $\approx$ mode (ii) or abruptly-activating mode-(i) substrates with late asymmetry, where the accumulation unloads abruptly instead of growing. The concrete substrate evidence distributes across both modes (financial crashes §4, conversational corpora §6). The sign-inversion between covert and overt substrates is itself a measurement value, not an artifact. Hysteresis appears in substrates with a long time axis: V-Dem ($n = 117$) shows 67% recovery within a decade but never a full return, and the fatigue shift $\rho \approx -0.30$ ($p = 0.029$) documents that $\hat{a}$ migrates path-dependently.

Mode mixture as substrate property. A substrate need not occupy a single mode but can be a linear combination of several; the substrate-specific characteristic is then the mixture vector $\pi = (\pi_1, \pi_2, \pi_3, \pi_4)$ on the mode simplex Δ_4 . A focused load-case study measures this mixture for the conversation substrate under pushback/sycophancy loading (Schessler, 2026c). Mode identities (i)–(iv) remain universal: basis families in the Π -sigmoid class vector space; the renormalization operator R_η and the no-go theorem (§2.5) are not affected by mode mixture, the substrate-specific asymmetry now resides in the mixture vector as well as in the inflection location.

Where a substrate exhibits none of the four modes, the question is which axiom is violated, not whether the framework holds universally. Null findings (§8) follow from the same axioms as positive findings: both are direct consequences of the same reduction.

2.5 Mathematical Load-Bearing Capacity: Birkhoff-Hopf, No-Go, T1–T3

The mathematical load-bearing capacity of the framework rests on three building blocks. The first is the Birkhoff-Hopf contraction (projective contraction theory on monotone kernels; Birkhoff, 1957; Hopf, 1963; Seneta, 2006): For a Gaussian smoothing operator T_ε with kernel $\exp(-(x-y)^2/\varepsilon)$ on monotone profiles over $[0, 1]$ (Hilbert diameter $\Delta_\varepsilon = 2/\varepsilon$), in the limit $\varepsilon \rightarrow 0$:

$$1 - \kappa(\varepsilon) \sim 2e^{-1/\varepsilon}, \quad \varepsilon_n \asymp \frac{1}{\log n}.$$

The global contraction budget closes exponentially in $1/\varepsilon$, the local diffusive gap only linearly: local smoothing scales favorably, while global order grows disproportionately with span. This exponential degeneration form is the classical Hilbert-metric contraction bound of projective geometry; recent work on Sinkhorn iteration in continuous settings confirms this exponential form and sharpens the associated rates (Chizat et al., 2024; Eckstein, 2023, the latter for kernels of bounded growth not bounded away from zero). The $\varepsilon \rightarrow 0$ degeneration is thus a singular-limit phenomenon of the worst-case bound, not the operative convergence rate. Analytic proof (fixed-point existence, sharp decay, no-go) in the formal companion framework (Schessler, 2026b); the Lean-4 formalization in `srds-enk-companion/lean4/PositivityProofs/` checks the algebraic components (symmetry, two-point gauge, TP_2 kernel, spectral bound) and the certificate-carried assembly, while the operator’s fixed-point existence and smoothness enter there as a declared axiom (the operator construction is not yet available in Mathlib).

No-Go Theorem. Let R_η be the damped renormalization operator on anchored monotone profiles, with TP_2 -positive integral kernel and damping η toward a reference. If the entire damped operator (including the reference) commutes with the reflection $SQ(x) = 1 - Q(1 - x)$, the fixed point Q^* is unique, and its density has a unique non-degenerate maximum, then the inflection location is $\Sigma_c = 1/2$; mirror symmetry is thereby sufficient, not proved necessary. Conversely, in a reparametrization-closed class without such reparametrization-breaking structure no universal inflection constant exists; in one certified asymmetric direction the variable inflection location $\tau(\alpha)$ shifts continuously off the symmetric point (Σ_c denotes only the symmetry value $1/2$). The certified spectral bound $\rho(DR_\eta[Q^*]) < 0.45$ at the certified parameter point secures local geometric convergence; global geometric convergence follows from the separate sup-norm Banach bound. **Computer-assisted proof (Flyspeck class):** an Arb interval certificate ($\dot{Q}(1/2) \in [1.8412, 1.8963]$, 256-bit, Krawczyk + Anselone + box enclosure, reproducible

via `make verify`) plus the chain-rule assembly machine-checked in Lean 4 — both public. Sorry-free is not axiom-free here: per `#print axioms` the certified theorem rests on 12 declared project axioms (2 numerical Arb-certificate axioms + 10 IFT-bridge operator axioms, which encapsulate the Banach fixed-point / IFT / Bochner analysis not yet constructed in Lean; Zeidler-referenced), not on an existence proof carried there (audit inventory 18, raw declarations 27; reconciled in the companion `AxiomAudit.lean`). The general statement for every admissible asymmetry and all (ε, η) remains open; the no-go theorem as a non-existence statement does not require it: a single certified asymmetric direction where the location shifts suffices. Substrate-specific asymmetry is thereby a measurement task, not a defect of the framework.

Substrate-specific scatter of the inflection. Substrate-specific asymmetry shifts $\hat{a}$ in structured fashion away from the symmetry value $1/2$, depending on how early the substrate yields under load; the deviations are not artifacts but the measurement value of the actual asymmetry of the respective substrate. Empirically: timber $\tilde{a} = 0.567$ (fiber asymmetry), battery $\tilde{a} = 0.646$ (CALB dataset), DotCom $\tilde{a} = 0.62$, GFC $\tilde{a} = 0.97$ (brittle-fracture asymmetry of a liquidity shock), earthquakes $\tilde{a} = 0.452$, lignin (boundary case: inflection at the origin, outside the interior $(0, 1)$, since the broad β -O-4 strength distribution has no endurance limit). Even sunflower phyllotaxis, the Fibonacci pattern par excellence, yields, across eight heads (floret-anthesis accumulation; Dryad 10.25338/B8963G), a mean inflection $\tilde{a} \approx 0.36$, not $1/\varphi$: $\hat{a}$ measures a yield transition, not a phyllotaxis angle. That several substrates fall near $1/\varphi \approx 0.618$ is a measurement reading, not a law; the structural reading of this proximity follows below.

Structure determines the value, family-internally and algebraically. The non-universality of the no-go has a constructive obverse: *within* a structured family the characteristic scale value is not fitted but algebraically determined, namely the Perron-Frobenius eigenvalue of the symmetry-specific linear substitution matrix (length factor λ ; the prototile-count matrix carries λ^2). In aperiodic order (quasicrystals) this inflation eigenvalue is, in the standard irreducible cyclotomic cases, a Pisot number of degree $\phi_{\text{Eul}}(n)/2$ (the symmetry fixes the degree, the substitution rule the value): five-fold the golden ratio φ (minimal polynomial $x^2 - x - 1$), eight-fold the silver ratio $1 + \sqrt{2}$ ($x^2 - 2x - 1$), twelve-fold $1 + \sqrt{3}$ ($x^2 - 2x - 2$), seven-fold the irreducibly cubic Perron-Frobenius root $\lambda_7 \approx 2.247$ of $x^3 - 2x^2 - x + 1$ that breaks the quadratic case (Pautze, 2017; Nakakura et al., 2019; Hare et al., 2018). The value $1/\varphi \approx 0.618$ is the inverse scale of the five-fold golden eigenvalue φ (whose algebraic conjugate is $-1/\varphi$), *one* of this spectrum, not *the* universal constant. Across *genuinely disjoint* substrates that share no common structural parameter, no common structure-to-value law exists; only the form is universal. Aperiodic order is the test bench here, not a substrate of this study: the one field where “structure determines the value” can be checked against an exactly known algebraic ground truth, pre-registered and tested (Figure 4). This sharpens the no-go twofold: no universal *value* and no universal *law* across disjoint substrates, alongside hard family-internal determination.

The same “form universal, value local” structure carries classical fracture mechanics at the crack-tip singularity: the near-field is a normalized angular structure with amplitude set by the boundary-value problem (load, geometry, boundary conditions) — in the linear-elastic case the square-root-singular K -field intensity (Williams, 1957), in the power-law hardening case the J -controlled HRR field, whose angular structure is universal within the constitutive class and is parametrized by the hardening exponent n (Rice, 1968; Hutchinson, 1968; Rice and Rosengren, 1968). Universal is the form within the class, local the amplitude value — structurally the same separation the no-go states for the inflection location; a structural analogy, not a derivation.

Three core theorems T1–T3.

Aperiodic order: the inflation eigenvalue is algebraically determined




Figure 4: Aperiodic order as the constructive counterpart to the no-go: the inflation eigenvalue (Perron-Frobenius of the symmetry-specific linear substitution matrix) is an algebraically determined Pisot number of degree $\phi_{\text{Eul}}(n)/2$. Five-fold golden φ , eight-fold silver $1 + \sqrt{2}$, twelve-fold $1 + \sqrt{3}$ (quadratic); seven-fold irreducibly cubic ($x^3 - 2x^2 - x + 1$, breaks the quadratic case). The value $1/\varphi \approx 0.618$ is the inverse golden scale (algebraic conjugate of φ : $-1/\varphi$), one of a family-specific spectrum. Pure aperiodic order, no SRDS substrates.

#	Theorem	Statement
T1	Information-stability correspondence	$\text{sign}(\text{slope}_W I) = -\text{sign}(\text{Tr } A)$: over a finite window the mutual-information trend signs the conditional phase-space volume rate $\text{Tr } A = \sum_i \lambda_i$ (Lyapunov-exponent sum); the $\lambda_{\max}$ reading is the dominant-mode case
T2	Autopoietic stationarity	$\mathcal{F}(\mathcal{F}) = \mathcal{F}$ yields non-equilibrium steady state with $\dot{D} = 0$ (entropy production = dissipative export)
T3	Goldilocks principle	Optimal function in a neighborhood of the edges of criticality, not at the measure-zero point of exact criticality

T1 connects to the stochastic thermodynamics of differentiable systems. There, in these differentiable settings, dissipated entropy is coupled to phase-space contraction (the sum of Lyapunov exponents), and the entropy-flow rate is spectrally bounded by the extremal eigenvalues of the symmetric Jacobian part (Das and Green, 2025): the same sign-coupling between the phase-space volume rate (the Lyapunov-exponent sum) and the information/entropy flow that T1 posits (the $\lambda_{\max}$ reading being the dominant-mode case). An exact identity linking entropy production and mutual information for overdamped Langevin dynamics (Cho and Jeong, 2025, preprint) further supports the direction. These external results are convergence anchors, not load-bearing premises. The full proof of T1 stands on its own in the formal companion framework (Schessl, 2026b).

Three further works realize the same structure in mathematically disjoint formal languages: AdS-GNN (Zhdanov et al., 2025, conformal-equivariant operator geometry), Topological Deep Learning (Schmidt et al., 2026, topological vs. combinatorial generalization), mesh-agnostic operator compo-

sition (Yang et al., 2026). The class is consistently realized across three mathematical languages: functor-invariant, not number-faithful. Functor invariance here is a functorial realization in the category-theoretic framing (Mac Lane, 1971; Spivak, 2014).

A functor $\Psi : \text{SRDS} \rightarrow \text{Aggregate-Class}$ realizes the sigmoid form with substrate-specific inflection (functorial realization; full development and proofs in the formal companion framework (Schessl, 2026b)).

3 Measurement Apparatus ENK (Material Level: Conversation as Specimen)

3.1 Embedding Geometry as Measurement System

Conversations are measured with transformer embeddings: high-dimensional vectors in which related utterances lie close to each other. Sentence-transformers (Reimers and Gurevych, 2019; Mikolov et al., 2013 as token precursor) measure semantic similarity at sentence level. The primary encoder is all-MiniLM-L6-v2 (384 dimensions); all-mpnet-base-v2 (768 dimensions) provides the triangulation. Each turn-pair (u_t, a_t) is a measurement point; the ordered sequence along a conversation is the deformation curve of this specimen. Decreasing embedding variance *is* stiffening of the semantic space (a reduced degree of freedom); irreversibly growing cumulative displacement *is* plastic deformation of the specimen. The apparatus measures not a speaker’s intent but the specimen’s trajectory under load.

Two robustness anchors. First: format ablation (Markdown / code blocks / lists removed before recomputation of E_a) shows that 95% of the signal stems from semantic content, 5% from formatting ($n = 202$). Second: partial Spearman correlation against conversation length holds the signal strength stable ($\rho_{\text{partial}} = 0.485$ after controlling for $\log(\text{turn count})$). Length is, in the SRDS sense, loading duration, not a nuisance term: longer specimens accumulate more damage, as expected in any fatigue process.

3.2 Five Effective Dimensions: the Anti-Sparse Rule

The five measurement axes each capture a distinct component of the deformation response.

Table 4: Five effective dimensions of conversational deformation

Axis	Interpretation	Material analogue	Strongest metric	ρ (vs manipulation)
D1 Stiffness	Variance decrease as reduced degree of freedom	Strain hardening	$E_a = 1/\text{Var}(\text{sem velocity})$	+0.543*** (space-robust: MiniLM +0.543, MPNet +0.507)
D2 Asymmetry	User-AI imbalance in embedding space	One-sided load introduction (cantilever)	MAS (Meta-Asymmetry)	+0.443*** , ICC=0.954
D3 Accumulation	Irreversible cumulative deformation	Creep / fatigue	B_w (windowed budget)	+0.249** cluster-robust
D4 Brittle fracture	Abrupt break-out from directional flow	Brittle fracture	dir_consistency	-0.377*** cluster-robust
D5 Coupling	Dynamics-structure covariation	Cross-field coupling	xcorr(sv, wd)	+0.210**

Plus core construct $\hat{a}$ (sigmoid inflection on cumulative damage over cumulative strain): $\rho = -0.359***$ strain axis, Cross-Corpus 4/5 (Awry marginal). (Reported here is the correlation $\rho(\hat{a}, \text{mal-frac})$; the inflection location $\hat{a}$ itself is constructed substrate-specifically, §2.5.)




Figure 5: Five-axis radar of the ENK deformation profile (D1 stiffness / D2 asymmetry / D3 accumulation / D4 brittle fracture / D5 coupling) with cluster-robust confidence intervals, $n = 202$.

Anti-sparse rule. The five axes are coupled, not redundant. Hooke’s law $\sigma = E\varepsilon$ couples stress, stiffness, and strain analytically: the same coupling reappears in the ENK axes. D1 (stiffness) and D3 (accumulation) are two projections of the same response. Lasso or other L1 procedures would discard three of the five axes under multicollinearity penalties. But the axes carry the same material response from orthogonal projections, so discarding them loses information, not noise. This coupling is an analytic constraint of the linear-response (Hooke) regime of the dissipative dynamics (A0+A1), not statistical multicollinearity one could regularize away. That is the wrong paradigm: materials testing, not regression statistics over independent features. Autocorrelation is signal, not a nuisance term: measurement points on a stress-strain curve *are* correlated (lag-1 $\bar{\rho}_1 = 0.928$ cumulative family, see §3.3).

3.3 Gold Reference Corpus + Cluster-Robust Inference

$n = 202$ conversations were annotated, 11,639 turn-pairs, across four LLM families (ChatGPT / Claude / DeepSeek / Gemini) and five contributors. A structured reference-measurement protocol (Claude Opus 4.6) assigns 17 annotation fields to each turn-pair (chunk-overlap consistency $\alpha = 0.87$; parallel run with independent LLM: 93.2% binary label agreement). Manipulation incidence: 42% manipulative turn-pairs. Four-phase classification P1 baseline $\rightarrow$ P2 buildup $\rightarrow$ P3 TypX (manipulation inflection) $\rightarrow$ P4 consolidation. Construct reliability rests not on a single annotator cohort but on cross-corpus replication (§6, §8): the same signal across twelve independently collected corpora demonstrates reliability more dependably than a single inter-rater study.

Cluster-robust inference. Lag-1 autocorrelation of the cumulative accumulator family: mean $\bar{\rho}_1 = 0.928$, individual metrics up to $\rho = 0.944$. Block bootstrap (2,000 iterations, full conversations as blocks) restores the specimen as analysis unit. Cluster-robust block bootstrap corresponds to the cluster-correction method in econometrics (Cameron and Miller, 2015; Wooldridge, 2003): sequential measurements along a deformation curve are not independent, and naive pooled inference overstates the confidence. Under cluster-robust inference, 47 of 113 pooled-significant metric-target associations remain conformant at the safety factor γ_M ; the remaining 66 are not identifiable under the correct inference architecture. The pooled p -values overstate confidence by 42% corpus-wide, and by about 55% for the cumulative family (Schessl, 2026a, Eq. 10). The safety factor itself is defined over the nine per-substrate tests ($\gamma_M = \alpha/9$, §1); the association count here uses γ_M as a conservative per-association threshold and does not form a separate Bonferroni family over the 113 associations.

Multivariate chat-level ridge (cluster-robust block bootstrap). Ridge regression on 15 chat-wide summary quantities reaches $\rho = 0.700 \pm 0.076$ against the continuous manipulation degree. Lasso reaches $\rho = 0.686 \pm 0.073$, comparable, consistent with the anti-sparse rule.

LOPO batch diagnostics. Leave-one-person-out: macro- $\rho = 0.339$ vs length reference 0.174 ($\Delta = +0.165$); pooled $\rho = 0.552$ vs 0.410 ($\Delta = +0.142$, with autocorrelation contribution). LOPO is not a model-generalization metric but a batch-invariance test: does the measurement system carry across the identity of contributors? $\Delta > 0$ indicates substrate-independent signal. An underpowered batch is therefore an expandable measurement (§8.5), not a falsification: the signal exists, its sharpness grows with batch diversity.

3.4 Crystallization Signature: Stiffening as Damage

Three-state classification via three material analogues: h_{bond} (semantic bond decay rate between turns, analogous to H-bond decay in polymer mechanics), hydrophobic packing density (frequency of dense topic clusters in the embedding space), and mean semantic velocity (drift rate of the embedding trajectory). Classification by median split: *crystalline* (h_{bond} and packing both above median), *glassy* (h_{bond} below median AND semantic velocity below 25th percentile), *amorphous* (remainder).

Table 5: Manipulation rate per state ($n = 196$)

State	n	Manipulation rate	Binomial p (against base rate 42%)
crystalline	52	71.2%	0.0020
glassy	42	26.2%	0.0012
amorphous	102	44.1%	n.s.

$\chi^2 = 18.813$, $p = 0.0001$, Cramér’s $V = 0.310$, cluster-robust 95% CI [0.198, 0.377]. Pairwise Mann-Whitney crystalline vs glassy: $U = 1650$, $p < 0.0001$, $r_{rb} = -0.511$ (large effect).

Cross-corpus confirmation. h_{bond} decay rate inverts sign between covert and overt manipulation: ENK-Gold $\rho = +0.398^{***}$ (crystalline stiffening), GasConv $\rho = -0.250$ (glassy disruption). Fisher meta over eight external corpora: $\chi^2 = 68.96$, $p < 10^{-6}$. Manipulation stiffens the conversational space; it does not randomize it. This is compatible with neural collapse (Pappan et al., 2020), but at conversation level rather than representation level.

3.5 Structural Axes: What the Five Main Axes Bear

The five main axes are best-of-class of a broader bundle of validated measurement concepts. The full inventory carries 38 cluster-robust concepts out of 77 (49%; full inventory in companion work in preparation).

The selection of load-bearing sub-axes follows the structural-engineering measurement paradigms: load combinations, fracture mechanics, stability, multiaxial load response, bond hierarchy, and microstructure.

Load combinations. Eurocode load combinations (corresponds to the EN 1990 load combination convention $\eta = E_d/R_d$ in structural design; Eurocode 0). Permanent load $G_{\text{baseline}_{sv}} = -0.478^*$ (favorable initial configuration carries through); variable load $Q_{\text{mean}_{sv}} = +0.531^*$ (strongest Q metric); utilization ratio $\eta = E_d/R_d$ via $\sigma_{\text{user}}/(B + 1)$: $\rho = -0.403^{***}$.

Fracture mechanics. stress_conc (K_I): $\rho = -0.312$ hold-out replicates; K_{IC} v2: $\rho = -0.247^*$; Griffith energy balance turn-level: discriminability 0.878** (critical load index 6.78, separation value from the cross-corpus spread, not an operative trigger).

Stability: creep, not buckling. The Euler slenderness ratio $\lambda = n_{\text{turns}} \cdot \beta/\text{WD}$ (Euler, 1744) is, on this substrate, almost entirely determined by sequence length ($\rho(\lambda, n_{\text{turns}}) = 0.94\text{--}0.99$) and sign-labile after length control; elastic buckling (stability loss, second-order theory) is not the governing failure criterion here; as a second-order limit state it is examined separately (companion). The governing mode is creep: uniform dynamics correlates robustly with load after length control ($\rho_{\text{partial}}(\text{CV}_{sv}) = -0.36$, cluster-robust). Hall-Petch segment granularity (corresponds to the grain-size strength relation $\sigma_y = \sigma_0 + k/\sqrt{d}$; Hall, 1951; Petch, 1953): $\rho = +0.381^*$ inverted. Avrami kinetics of stiffening (corresponds to the phase-transformation kinetics in metallurgy; Avrami, 1939; Kolmogorov, 1937): double-logarithm plot shows two regimes with kink at mid sigmoid phase, replicated across seven corpora. A convergent result: Aguilera et al. (2025) report explosive phase transitions in neural networks on curved statistical manifolds with a hysteresis signature. This independently supports, from a statistical-physics substrate, the sigmoid-inflection form measured here.

Multiaxial load response. Mohr deviatoric $|\text{asymm}| = -0.246$, partial -0.259^{***} after B_w control (suppressor effect). Shear stress $G_{\text{shear}} = \text{DIV}/|\Delta\text{WD}|$: $\rho = +0.292$ cluster-robust. Second principal stress σ_2 via MAS, orthogonal to $|\text{asymm}|$ ($\rho = 0.042$).

Bond hierarchy (protein analogy). H-bond decay $\rho = +0.398^{***}$ (LOPO 6/6); hydrophobic packing $\rho = +0.230$ cluster-robust; disulfide (long-range references) $\rho = +0.215$ in 5/6 LOPO. Weak bonds react $1.4\times$ more strongly than strong ones, the same weak-vs-strong-bond dichotomy as §5.4 lignin and §3.4 crystallization.

Microstructure. Grain-boundary NC2 (Newman conductance on 2-cluster partition, cf. Newman and Girvan, 2004; not to be confused with Neural Collapse Pappan et al., 2020): $r = 0.997$ across

encoder spaces (space-robust). PCA isotropy: $\rho = -0.256^*$. Manipulation is more isotropic than nominal conversation: the semantic space loses its preferred directions.

Cross-scale bridge. $h_{\text{bond_decay}}$ (token scale, S2) $\rightarrow$ stress_conc (chat scale, S4) partial $\rho = -0.573^*$ across 5/5 persons (controlled for mal_frac and length). Strongest cross-scale evidence of the framework: what the token scale registers as bond decay carries through, structurally consistent, to chat-scale stress intensity.

Minimum sampling. $n_{\text{turns}} \geq 15$ as minimum loading duration for stable E_a determination, analogous to the Nyquist minimum-sampling theorem (sample-size lower bound for reconstructable inflection, not a frequency argument).

3.6 Comparison with Message-Level Detectors

Existing AI-safety systems operate message-wise (LlamaGuard (Inan et al., 2023), Perspective API, Detoxify (Hanu and Unitary team, 2020)). Head-to-head subtle manipulations ($n = 8$): SRDS 5/8, LlamaGuard 2/8, one tie. On ProsocialDialog ($n = 5,000$ raw conversations, of which 308 with $n_{\text{turns}} \geq 15$ per §3.5 minimum sampling) SRDS monitoring delivers $\rho = -0.008$ (zero false alarms; budget NORMAL in 307/308 specimens); LlamaGuard shows on the same data $\rho = +0.327^{***}$: systematic false alarms at therapeutic intensity.

A fire alarm measures acute, localized danger; a structural monitor measures cumulative load-bearing capacity. Both are needed, neither replaces the other. The complementary profile (SRDS dominant on covert cumulative deformation, message-wise detectors dominant on acute single violations) is the operational signature the framework predicts. The operational calibration of the state machine is described in §5 as architecture.

4 Cross-Domain: Ten Substrates, One Functional Form

Timber is the methodological anchor, not a singular case. The transfer to nine further substrates tests whether the *method* carries over: whether it adapts to a new carrier, not whether a particular number recurs. What is universal is the functional form of inflection; what is substrate-specific is the asymmetry of the carrier on which the load is redistributed (no-go theorem, §2.5). The normalization Π differs per substrate (nonlinearity residual for timber, $\sigma(\varepsilon)$ for metals, cumsum for sequences; §2.4): what recurs is the form class, not the numeric value. Each sub-section states the finding in one sentence and, where the mechanics predicts it, the *counterpart* at which the signal must be absent. The substrates support one another via shared mechanical axes (§4.13). Raw data in the companion `srds-enk-companion/`.

4.1 Timber under Tensile Load: DIN EN 408 (Anchor)

Spruce specimens in bone-shaped geometry per DIN 52188:1979-05, fiber-parallel under tension per DIN EN 408:2012-10 (Schessl, 2025). Across $n = 230$ specimens (sorting classes S7+S10+S13, balanced from $n = 319$): $r(\hat{a}, \sigma_{\text{break}}) = -0.83$, $p < 10^{-30}$ (Spearman $\rho = -0.99$). The construction itself carries this coupling: the absolute stress at the inflection, $\sigma_{\text{onset}} = \hat{a} \sigma_{\text{max}}$, follows the onset-strength line $\sigma_{\text{onset}} = 5.27 + 0.433 \sigma_{\text{max}}$ ($R^2 = 0.9925$), and since the inflection axis is normalized by σ_{max} , $\hat{a} \approx 0.433 + 5.27/\sigma_{\text{max}}$ is, up to residuals, a decreasing function of strength; the line alone forces Spearman -1.0 , so the near-perfect rank order follows arithmetically from this relation (decomposition in `analysis/timber_normalization_control.py`, companion). A permutation control (onset permuted independently of strength: $r \approx -0.74$) answers only the narrower question of whether the onset is independent of strength; it is not: it spreads (CV = 0.228), scales with strength, and the coupling holds within each grading class ($r = -0.83$ S7, -0.86 S10, -0.95 S13). The anchor’s role is therefore instrument validation against known sorting physics (§1.1: the relation is established in timber engineering); this substrate does not evidence a mechanism beyond that established relation. The class-specific inflection means decrease monotonically with strength: S7 $\hat{a} = 0.574 \pm 0.057$ ($n = 144$),

S10 0.557 ± 0.043 ($n = 71$), S13 0.539 ± 0.026 ($n = 15$): the best class (knot-poor) carries the lowest $\tilde{a}$, consistent with $r < 0$. $\tilde{a} = 0.567$ lies measurably away from the mirror-symmetry value $\Sigma_c = 0.5$ (no-go) and measures the fiber anisotropy.

The counterpart sits within the substrate itself (§8): the near-elastic cohort ($R^2 = 0.994$, no plastic regime) carries **no** yield signal: the plastic regime *gates* the signal.

4.2 Multi-Material Multiaxial Fatigue: $\hat{a}$ Reads the Non-Proportionality

$n = 914$ strain-controlled specimens from the Multiaxial Fatigue Deep Learning Dataset (Chen et al., 2024; 136 material entries across 40 base alloys; AZ31B-Mg, pureTi, 16MnR steel, 6061/7075-Al); 31 material entries with $n \geq 8$ for material-wise S-N sigmoid fits. The input per specimen is a closed cyclic load path, a **hysteresis loop** in stress space. $\hat{a}$ reads its **load-path non-proportionality**, the *established* damage driver in multiaxial fatigue (critical-plane / Fatemi-Socie): out-of-phase loading rotates the principal stress axes, activates more slip systems, raises the dissipated energy per cycle (Palmgren-Miner). The autocorrelation structure lies, at Chelton 49.7%, in the same band as the ENK conversation substrate (PASS); intra-material $\hat{a}$ carries up to $|r| = 0.82$ (PA38-T6-ASN1 $+0.82$ at $n = 7$, CuZn37-NPR2 -0.74 , CP-Ti -0.69 ; the intra-material couplings run over entries with $n \geq 5$, the material-wise S-N fits over $n \geq 8$); the sigmoid form is robust (74% $R^2 \geq 0.5$). $\hat{a}$ and the amplitude (VM amplitude $\rho = -0.717$) are coupled descriptors of *one* response ($\sigma = E\varepsilon$, §1.1b). Aggregated cross-material, $\hat{a} \leftrightarrow N_f$ is directionally correct ($\rho = -0.058$: earlier inflection $\rightarrow$ longer life) but **pooled-damped** across 40 materials $\times$ 48 load paths, a property of heterogeneous pooling, not a failure.

4.3 Finance SPY: Crash S-Curve against Calm Random-Walk

Daily returns of the S&P-500 ETF SPY over 27 years (1999–2026). Per episode (historically named, no shape-search algorithm), a sigmoid fits the cumulative drawdown, read as a deformation curve. Two controls calibrate the finding. (i) Calm phases are random-walk: mean $R^2 = 0.973$ during crash episodes versus 0.483 in length-matched calm phases (Mann-Whitney $p = 0.0001$); the contrast survives magnitude matching (overlap band $|\text{return}|$ 24–55%: 0.986 vs. 0.212). This is the core negative control against “everything saturates”: the sigmoid form is episode-selective, not an artifact. (ii) An anchoring null quantifies the share contributed by window choice: block permutation of the daily returns, deepest peak-trough window of equal duration, identical σ_t construction (500 draws). The crash episodes lead the exceedance ordering as a class (Bear2022: 0/500; GFC: 6.0%; COVID: 16.2%; exact class separation crash vs. correction $p = 0.019$), the corrections lie within the anchoring band (28–88%); DotCom (51%) sits in the band as the creep pole. The poles are marked by GFC 2008 ($\hat{a} = 0.97$, brittle fracture) and DotCom 2000 ($\hat{a} = 0.62$, creep); inflection position and exceedance measure different quantities: the position places the mode, the exceedance the coherence of the episode, and across the ten episodes the two are decoupled ($\rho_S = -0.18$, n.s.); no real inflection position leaves the anchoring distribution. After DotCom a permanent redistribution remains in the daily-return autocorrelation (hysteresis).

4.4 Lignin: Concave Digestion Kinetics without an Interior $\hat{a}$ (Cross-Sectional)

Hydrothermal treatment degrades lignin via hydrolysis. We plot β -O-4 concentration against the P-factor (time-temperature severity) over $n = 72$ process conditions (95 samples, SP-LCC dataset, Alopaeus et al., 2025). Weibull sigmoid (serial-cascaded, mode (ii), weakest-link failure): $\rho_S = -0.78$. The β -O-4 bonds form a *broad* strength distribution (phenolic linkages cleave well before the non-phenolic ones), so damage is concave from the origin ($m < 1$): **no endurance limit**, the weakest ether bonds yield at the slightest load (Wöhler fatigue side, Wöhler, 1870; Palmgren-Miner accumulation, §5). The tested severity window therefore opens beyond the inflection: the weakest β -O-4 bonds yield before the mildest measured condition, and the purely concave profile is the predicted signature of a measurement beyond the yield point (companion framework, Theorem 2.2(ii): observation window opening past the rate peak), not a framework boundary; the test path is milder severities or early-phase kinetics, which would

move the inflection into the window. An interior $\hat{a}$ is not defined on these data and is not reported as a number, independently confirmed via birch AqSO (Schlee et al., 2023: β -O-4 already strongly reduced at the mildest severity, $\approx 34/100$ Ar) and kraft softwood (Mattsson et al., 2017: reactive β -O-4 cleavage early in the cook). Weak bonds (β -O-4, 60 kcal/mol or 250 kJ/mol) yield before the strong ones (5-5 linkages): a weakest-link failure that repeats, on the chemical level, the bond hierarchy from §3.5 and the crystallization signature from §3.4. It is a cross-sectional dose-response signature ($\rho_S = -0.78$, $m < 1$): the broad defect distribution pushes the yield point to the origin, and an interior $\hat{a}$ is not defined. The class-ordering analogy to timber defect class and music genre is not a positive support here: both, like CRC, reduce to positive decompositions in §4.13 (timber circular, music sequence-carried).

The design is cross-sectional and the data provenance is open; the cross-sectional finding ($\rho_S = -0.78$, $m < 1$) is nominal under γ_M to that extent.

4.5 V-Dem: Autocratization Trajectories (Two-Span Beam)

ERT-v14 (Episodes of Regime Transformation, Maerz et al., 2024; V-Dem-based, Coppedge et al., 2024): $n = 117$ autocratization episodes, 89 fitted. A monotonicity null (length- and depth-matched monotone random declines, identical 4-parameter fit) reaches the same fit quality as the episodes (median R^2 0.989 vs. 0.983); the form information lies in the class comparison: against 445 matched null curves, the saturation class family wins 60.7% real vs. 42.5% ($z \approx 3.2$), the driver is Gompertz (29.2% vs. 8.5%, 3.4-fold, $z \approx 4.1$), and linear collapses to 11% real (null 31%); the family form (Gompertz) is substrate-specific (§2.4). In the increment-permutation null (real yearly increments, permuted order) the family preference stays at null level (60.7% vs. 54.0%, n.s.): the S-shape lives in the distribution of the yearly increments. In 60 of the 89 episodes the inflection $\hat{a}$ falls within the observation window, in the remaining 29 the fit extrapolates it beyond the window (16 past the episode end, where the transition is not yet reached; 13 before the episode start). $\hat{a}$ partitions into creep erosion ($\hat{a} < 0.5$; Hungary, Austria 1933) and sudden regime switch ($\hat{a} > 0.8$; India 1975, Czechoslovakia 1938). On the institutional scale, hysteresis appears: about two-thirds of states return within a decade to the pre-episode corridor, never fully (Wilcoxon $p < 10^{-4}$), the same loading-unloading asymmetry as battery and multiaxial fatigue (G7, §4.13). For states with multiple episodes, $\hat{a}$ shifts to earlier values with each subsequent one ($\rho \approx -0.30$, $p = 0.029$): the institution carries prior damage as material memory. The structural mechanics of these trajectories (support topology, truss buffering) follows in §5.

4.6 Political Speeches: Support Topology, Not an Effect Claim

Political speeches test the support topology. Bundestag speeches $n = 6,874$, National Socialist (NS) speeches $n = 11$, ParlaMint HU+PL+RS+SI (pilot) and a 17-country panel. The Bundestag null ($\hat{a}$, E_a without significant correlation) is the **predicted confirmation** of support topology. The speech corpus over the order of business is a truss, and a truss bears no bending: the load is *passed on*, not locally deformed (§5.2). This makes no claim about the effect of speeches, only about the chosen measurement scale. That the discourse load *reaches* society is shown by the $\ell_3 \rightarrow \ell_4$ force transfer: on the 17-country panel (238 country-years) the sv axis carries cluster-robustly with a phase lag ($\rho = -0.284$ at lag +2, $G = 17$; BCa-CI excludes zero); in the 4-country pilot the $\hat{a}$ -discourse axis is uniformly negative across all four clusters (HU -0.40 , PL -0.52 , RS -0.21 , SI -0.40 ; BCa-CI₉₅ $[-0.34, -0.15]$), directionally consistent but not threshold-capable at $G = 4$ (cluster permutation $p = 0.207$; no single cluster significant; tail-arm); the reliable inference is carried by the 17-country panel ($G = 17$). The institution itself is measured as a support (companion program $\ell_3 \rightarrow \ell_4$, 17-country panel + federal-state panel): its coupling to opinion salience is pinned — it transmits force (difference coupling in 19/22 countries, $p < 0.0001$) but no moment (lag peak at 0, no resolvable lead). Its characteristic is that of an emergency support: under full load all chambers respond (pandemic: 26/26), under partial load the load path follows the actuator architecture of the load type. Subordinate chambers track the load level sluggishly (level coupling 0.656 versus difference coupling 0.378, lag of about six months). The

institutional mechanics is being measured in a program of its own. NS speeches ($n = 11$) are illustrative (direction, not magnitude).

4.7 Earthquakes USGS: Instrument Validation on the Sequence Statistics

USGS-FDSN across six regions, $M \geq 2.5$ (2016–2021), 11,505 events $\rightarrow n = 284$ sequences (spatio-temporal clustering). Per sequence, a sigmoid fits the normalized cumulative energy release: median $R^2 = 0.965$, $\tilde{a} = 0.452$ ($\sigma = 0.199$). The coupling $\rho_S(\hat{a}, \text{mainshock position}) = +0.673$, $p < 0.0001$ (Bonferroni) validates the instrument semantics against known seismicity: the inflection of the cumulative energy curve follows the largest jump under any event ordering (ordering null with identical event statistics: mean $+0.739$; the real value lies 1.74 SD below it, consistent with the Omori aftershock structure). It is a property of the measurement construction, not an additional mechanical finding; early $\hat{a}$ mark foreshock-dominated, late $\hat{a}$ mainshock-dominated sequences (pattern of §4.1: anchor role). Against the generic AR(1)/RW/WN synthetics, the $\hat{a}$ distribution separates with KS- $D = 0.178$ ($p < 0.001$); against the stricter null with identical event statistics the distributional difference is weak ($D = 0.096$, $p = 0.013$). The magnitude *prediction* is weaker ($\rho_S(\hat{a}, \text{max magnitude}) = -0.176$, $p = 0.006$).

4.8 Protein Melting Curves (Meltome Atlas): Instrument Validation across Organism Classes

Meltome Atlas (Jarzab et al., 2020), 13 species with thousands of proteins each: per protein the cumulative melting curve (fraction no longer soluble over temperature) is fitted with the O6 sigmoid, $\hat{a}$ = inflection temperature on the absolute Celsius axis; species endpoint = median over the valid $\hat{a}$. The class variable is the organism's growth temperature (OGT), externally assigned by biology and **outside** the measurement-temperature axis (unlike timber, where the axis contains the class quantity). Pre-registered (median structure anchor before the fit; prior knowledge and window censoring declared). Across the 13 species-median $\hat{a}$, $\hat{a}$ rises with OGT: Spearman $+0.936$ (permutation $p < 10^{-4}$, one-sided), the thermophiles lie above the 37-degree species. The validation core: the fitted $\hat{a}$ matches the source's published melting point (Spearman $+0.951$, absolute agreement ~ 1 degree). **Declared limit:** the measurement windows co-shift with OGT per species (right-censoring of the thermophiles, many proteins do not fully melt within the fixed window); as a novel finding the line would thereby be circularly amplified. It therefore runs, isomorphic to §4.7, as instrument validation against known thermostability physics: the instrument reads real class physics on a non-circular axis without claiming a new discovery. Register/data: analysis/angriff_register/zeile10_meltome_validierung/.

4.9 Battery Degradation, CALB (real): Knee + Hysteresis Accumulation

$n = 26$ CALB cells from the BatteryLife corpus (Tan et al., 2025) (0/35/45 °C). $\hat{a}$ localizes the knee point: the empirical $\hat{a}$ distribution is strongly shifted against the synthetic null: KS $D = 0.733$, $p < 10^{-13}$ (mean 0.667 vs. null 0.541; $\tilde{a} = 0.646$). The separation also holds against a curvature-matched knee-free null (per-cell power-law fade fitted to the raw curve, AR(1) residuals, identical measurement chain): KS- $D = 0.692$ ($p < 10^{-11}$); split-half stability $\rho_S = 0.956$; the logistic-over-power gain lies above the null P95 in 17/26 cells (Mann-Whitney $p = 7.8 \times 10^{-5}$). Two interlocking findings: (i) the voltage hysteresis $V_C - V_D = 0.389$ V **grows with ageing**, $\rho(V_C - V_D, \text{cycle}) = +0.886$ over 92% of cells: irreversible, accumulating polarization, the same G7/Bauschinger mechanics as multiaxial fatigue and V-Dem (§4.13); (ii) the outcome coupling is mode-dependent and open ($\rho_S(\hat{a}, \text{EOL}) = +0.18$, n.s.; right-censored, only 7/26 cells reached 80% EOL). As with earthquakes, $\hat{a}$ localizes the failure knee reliably but does not predict lifetime strength.

4.10 Colorectal Carcinomas: TCGA (Stage-Occupancy Finding, Reduced)

TCGA Pan-Cancer Atlas *coadread_tcga_pan_can_atlas_2018* (Hoadley et al., 2018): $n = 579$, of which 76 with microsatellite instability (MSI; MANTIS > 0.4) and 503 stable (MSS); mean $R^2 \geq 0.96$. Stage-aggregated, MSI $\hat{a} = 0.286$ lies below MSS 0.439 ($\Delta\hat{a} = -0.153$). This headline is, however, **not curve-form-borne but stage-occupancy structure**: MSI concentrates in early stage (39 of 70 in Stage II), and a swap test ($\hat{a}$ from MSS per-stage curve means with MSI stage occupancy) reproduces the MSI value 0.286 exactly, while per-stage curve means shift only slightly counter. The per-patient bootstrap means lie at $\Delta = +0.033$, opposite to the stage-aggregated headline; the patient-bootstrap 95% CI of the difference is $[-0.199, +0.035]$, spanning 0, so the sign direction is aggregation-dependent. Sorting within a stage by damage itself (mutation count) flips the sign ($\Delta\hat{a} = +0.090$), a count-sorting artifact. As with the timber anchor (§4.1), this substrate thus establishes **no material-class $\rightarrow\hat{a}$ mechanism beyond the diagnosis epidemiology**; the positively shown finding is the decomposition itself (the subtype difference is stage occupancy, not yield form). The coupling of $\hat{a}$ to survival remains open given the small MSI cohort.

4.11 LLM Conversations (ENK Gold)

$n = 202$ gold-annotated conversations, 11,639 turn pairs, four LLM families, five contributors. Univariate $\rho(\hat{a}, \text{mal-frac}) = -0.359$ cluster-robust (canonical strain axis; on the cumulative companion axis -0.332 across the $n = 193$ validly fitted specimens); multivariate chat-level ridge $\rho = 0.700 \pm 0.076$ (§3.3). $\hat{a}$ lives on the cumulative axis and carries a size effect: $\rho_S(\hat{a}, \text{turn count}) = -0.928$; as specimen length sets the loading duration, this is the material law, not a nuisance quantity (the k_h class). The $\hat{a} \leftrightarrow \text{mal}$ coupling holds within all length classes (quartiles $-0.187 / -0.257 / -0.201 / -0.138$; partial -0.197) as well as in every contributor group and every model family (block permutation $p = 0.003$ and $p < 0.0005$). Cross-corpus: $\hat{a}$ carries across the manipulation-corpus specimen classes (GasConv, LegalCon, MentalManip, P4G; Awry marginal), strongest LegalCon ($\rho = -0.751$); the coupling strength depends on which strain axis is read (canonical cumulative strain §2.5 or the FID axis, named per corpus); 26/113 associations robust at γ_M under shuffle calibration (each test against its own label-shuffle null distribution, §8). The counterpart is path destruction: destroying the turn ordering (multivariate phase surrogate preserving spectra and marginals per metric, $n = 193$) lets $\hat{a} \leftrightarrow \text{mal}$ fall into the generic cloud. Empirically, the real path stands out with $z = -4.86$; a second scheme (AAFT with rank remap) converges ($z = -4.60$, §8). The amplitude axis itself cannot be partialled out: $\hat{a}$ and the embedding amplitude (std) are strongly $\sigma = E\varepsilon$ -coupled descriptions of one response ($\rho(\hat{a}, \text{std}) = +0.807$, G5). Stratifying by amplitude terciles (as timber is stratified by sorting classes, §4.1) accordingly yields no coherent residual coupling ($\rho(\hat{a}, \text{mal}) = -0.03 / +0.24 / +0.08$ per tercile, $n \approx 64$); this is the coupling predicted by G5, not a separable channel a partial could regularize away (anti-sparse, §3.2). The instrument that holds the amplitude fixed (marginals preserved) and destroys only the path is the phase surrogate ($z = -4.86$); the separable content of $\hat{a}$ lies there, in the stability within length classes, and in the cross-corpus replication across twelve corpora, not in an amplitude-adjusted partial correlation.

4.12 Music: MetaMIDI (ℓ_2), Pre-Registered + Cross-Method-Validated

Pre-registered on 16 May 2026 (hypotheses H1–H3, genre order per Di Marco fixed before analysis, with a built-in shuffle falsification; the pre-registration is public in the companion, *data/musik_companion/*), run 17 May 2026. $n = 2,840$ MIDI pieces: 1,974 from music21 (Cuthbert and Ariza, 2010) (Bach $n = 413$) plus 866 from the Lakh subset (Raffel, 2016) with Tagtraum genre match (Schreiber, 2015) (≥ 15 turns). $\hat{a}$ via sigmoid fit on cum-unique-edges against cum- $|\text{interval}|/12$. The genre hierarchy is perfectly monotone: Rap $\tilde{a} = 0.196 \rightarrow$ Electronic $\rightarrow$ Pop $\rightarrow$ Rock $\rightarrow$ Jazz $\rightarrow$ Bach 0.409, Mann-Kendall $\tau = +1.000$, $p = 0.003$ (14/15 pairwise Bonferroni-Holm significant); median $R^2 = 0.975$. Genre orders $\hat{a}$ monotonically; the pre-registered shuffle control, however, reduces this to a sequence-carried finding (§4.13, not an independent material-class $\rightarrow\hat{a}$ mechanism) and comes out




Figure 6: Genre hierarchy across six macro-genres: Rap ($\tilde{a} = 0.196$) $\rightarrow$ Electronic $\rightarrow$ Pop $\rightarrow$ Rock $\rightarrow$ Jazz $\rightarrow$ Bach (Classical, $\tilde{a} = 0.409$); Mann-Kendall $\tau = +1.000$, $p = 0.003$. $n = 2,840$ MIDI pieces.

mixed. The expectation was that under note-order shuffle the $\hat{a}$ distribution degenerates to a tight cluster near 0.5 and the genre separation vanishes; the fixed reduction clause was that genre separation surviving the shuffle reduces the finding to a vocabulary effect. What occurred: no 0.5 collapse (the $\hat{a}$ cluster at 0.27–0.34), the monotone genre ordering breaks (Mann-Kendall n.s.; Rap moves from the lowest value into the midfield), and the Jazz level stays separated under shuffle. Under the reduction clause this yields: the monotone ordering is sequence-carried, the Jazz level carries a vocabulary component (the static note multiset). The perfectly monotone six-genre ordering ($\tau = +1.000$) is an own finding of this work, not externally validated. At the two-group level there is a cross-method concordance: Di Marco et al. (2026) (Sci Rep), with an independent method (network efficiency rather than sigmoid inflection), ranks Classical and Jazz above the modern genres; a finer monotone ordering *within* the modern group is explicitly not reported by that source. The cross-method agreement thus carries the coarse two-group separation, not the fine six-genre ordering.

4.13 Materials-Class Hierarchy

The $\hat{a}$ values order the substrates along an axis of increasing fracture asymmetry, the same sorting logic as DIN 4074-1 (timber S7/S10/S13), DIN EN 10025 (steel yield strength), EN 206 (concrete strength class). The 1D $\hat{a}$ -value comparison is illustrative (no-go: $\hat{a} = f(\Pi, \text{form})$, not strictly comparable cross-substrate); the load-bearing statement is the within-substrate correlation, not the absolute value. The same normalization logic is documented outside these substrates: in ab-initio water dynamics, part of the differences between density functionals can be read as shifts along the phase diagram; comparisons relative to each functional’s own melting temperature reduce the spread of the dynamical observables (Montero de Hijos et al., 2026).

Table 6: Materials-class hierarchy with form annotation. The $\tilde{a}$ column reports the substrate-typical inflection location (median or mean per the convention used in each substrate’s §4.x); the 1D comparison is illustrative.

Substrate	$\tilde{a}$	Form class / mode	Characterization
Al-6061 (tensile tests, $n = 146$)	0.084	Gompertz (form exception)	ductile metal, operating-range ceiling (continued on next page)

(continued)

Substrate	$\tilde{a}$	Form class / mode	Characterization
ENK Chat AI	0.232 ($D = 0.944$)	Logistic / (iv) continuous	conversation (chat-AI turns)
Trecento (14th c., music21 (Cuthbert and Ariza, 2010), $n = 103$)	0.235	Logistic / (iv) continuous	early tonality development
ENK all turns (median)	0.268	Logistic / (iv) continuous	LLM substrate (median over all turns; headline channel Chat-AI 0.232)
Bach (Classical)	0.409	Logistic / (iv) continuous	high baroque
Earthquakes	0.452	Logistic / (i) parallel-instantaneous	foreshock-dominated
Timber (Picea abies, tension)	0.567	Logistic / (i) parallel-instantaneous	fiber anisotropy
Finance SPY DotCom	0.62	Logistic / (i) parallel-instantaneous	speculative asymmetry
Battery (CALB)	0.646	Logistic / (iv) continuous	knee localization (KS) + G7 hysteresis
Finance SPY GFC	0.97	Weibull / (ii) serial-cascaded	liquidity shock, late inflection
DP1180 (Numisheet)	0.975	Weibull / (ii) serial-cascaded	brittle fracture high-strength steel (KS✓; outcome underpowered, small n)

Form class and mode stem from the companion model comparison (AIC over Logistic/Gompertz/Weibull/Hill; `srds-enk-companion/data/model_comparison/`), in which Logistic dominates 6 of 8 substrates and Al-6061 (Gompertz) and CCP60 (Weibull) are the no-go-predicted exceptions. Each carrier yields along its own path; the load redistributes as soon as one axis reaches its material limit.

4.14 Cross-Substrate Status and Mutual Support

Table 7: Bonferroni status across the ten substrates; political speeches separately as the support bridge to §5 (not counted among the ten)

Status	Substrates
Bonferroni-conformant (γ_M)	Timber, ENK Conversation, Earthquakes, Battery (knee+hysteresis), Music, V-Dem (class comparison, Gompertz driver; partition/material memory/hysteresis)
Confirmation (accumulation/isomorphism axis)	Multi-Material Multiaxial Fatigue ($\hat{a}$ reads non-proportionality, Chelton in the ENK band; cross-material outcome pooled-damped)
Nominal, caveat named	Lignin (cross-sectional), Colorectal ($\Delta\hat{a}$ = stage occupancy, not curve form; §4.9), Finance (coherence anchoring-calibrated: class separation $p = 0.019$, Bear2022 $p < 0.002$)
Support prediction	Political speeches (speech-corpus truss null; bridge directionally confirmed, §5)
Instrument validation (not counted among the ten)	Protein melting curves, Meltome Atlas ($\hat{a}$ vs. OGT +0.936; $\hat{a}$ vs. source melting point +0.951; §4.7b)

Materials class hierarchy by $\hat{a}$




Figure 7: Materials-class hierarchy as a 1D $\hat{a}$ -axis: from ductile Al-6061 ($\hat{a} = 0.084$) to the brittle high-strength end ($\hat{a} \approx 0.97$). Shown are $\hat{a}$ -positions across material classes incl. reference specimens (Al-6061, DP1180, Trecento, financial crashes); V-Dem, CRC, multiaxial fatigue and lignin (no endurance limit, $m < 1$) have no comparable single $\hat{a}$ and are not plotted here. The dotted line marks the mirror-symmetric inflection value $\Sigma_c = 1/2$ (no-go), from which substrate-specific asymmetry is measured; the 1D comparison is illustrative.


In addition, there are a pre-registered null control (ProsocialDialog L1) and two falsified positive substrates that map the scope conditions (Sotopia A0, Google COVID Mobility $\sigma \perp \varepsilon$, §8). The substrates do not stand side by side but support one another via shared mechanical axes:

- **Hysteresis / irreversibility (G7):** battery voltage hysteresis (+0.886) · multiaxial-fatigue loop · V-Dem recovery (67%, never full): *one* Bauschinger/dissipation mechanics across three substrates.
- **Near-elastic gate:** timber near-elastic ($R^2 = 0.994$) · battery pre-knee: nonlinearity *gates* the yield signal; where there is no plastic regime, no signal.
- **Class ordering and instrument validation:** the apparent material-class $\rightarrow \hat{a}$ cases reduce consistently to positive decompositions: timber S7 $\rightarrow$ S13 follows arithmetically from the onset normalization (circular axis, §4.1), CRC MSI/MSS is stage occupancy (§4.9), music genre is sequence-carried (§4.11). Where the axis does *not* contain the class quantity, by contrast, the validated $\hat{a}$ role carries as instrument validation: earthquake mainshock (§4.7) and protein melting curves across OGT classes (§4.7b; Spearman +0.936 to OGT, +0.951 to the published melting point).
- **$\hat{a}$ localizes failure, not strength:** earthquake mainshock (+0.673) · battery knee (KS): valid localization, weaker strength prediction, same pattern.

The structural statement carries not via the recurrence of a number but via the recurrence of the functional form with substrate-specific asymmetry and the mutual support across these axes: the signal appears where the mechanics predicts it and is absent where it predicts its absence.

Luhmann Chain as Statics Diagram

Five Social Sub-Systems — Boundary Conditions Decide, Not Scale.



Scale grows from left to right (human dyad → society). Measurement load-bearing capacity follows the support type, not the scale — Speech-corpus truss carries only normal forces, hence null signal in $\hat{a}$ and E_a .

Figure 8: Luhmann chain as statics diagram: the five measured carriers of the societal sub-systems with their support type (clamped, cantilever, truss, two-span beam). The boundary condition decides, not the scale.

5 Structural Scale, Boundary Conditions, and Addressee-Support

The material level from §3 measures the deformation of the individual specimen. The structural level installs this specimen into a configuration with supports, load paths, and internal forces. The boundary-condition typology is the map of these configurations; it decides which of the five measurement axes remains load-bearing and which falls into a zero-force member under the given support. A null signal is then not measurement failure but a prediction of the framing.

Two levels, two operators. The material level (ℓ_2) operates on the stress response of the specimen itself; under multiaxial loading a principal-axis transformation (Mohr, §3) would be the canonical pre-processing, under uniaxial loading it degenerates to the identity. The structural level ($\ell_2 \rightarrow \ell_3$) operates on the topology of the support: whether a member is clamped, cantilever, truss, or two-span beam is not a material property but a structural configuration. The two levels are orthogonal: the material principal-axis choice decides which stress components enter the measurement signal; the support topology decides which measurement axes are activated at all. Confusing the two would confuse materials testing with structural engineering.

5.1 Five Scales, Five Sub-Systems: the Luhmann Cut

Table 8: Scale-transfer profile across five societal sub-systems

Scale	Corpus	n	Support type	E_a	Q_{sv}	$\hat{a}$	Sigmoid R^2
Dyadic Monologue (short)	ENK Gold	202	Clamped	+0.543***	+0.531***	-0.359***	—
	AgoraSpeech	169	Cantilever	+0.176*	-0.285***	-0.210*	—
Multi-party debate	IQ2	216	Cantilever	-0.042 n.s.	-0.081 n.s.	-0.064 n.s.	—
Speech corpus (truss system)	Bundestag*	4,251	Truss	+0.023 n.s.	-0.032***	+0.035 n.s.	—
Macro-institutional trajectory	V-Dem	117	Two-Span-Beam	+0.103 n.s.	+0.347***	—	0.983

All turn-pair statistics are computed cluster-robust; the pooled variants overstate the confidence of the cumulative accumulator family (mean lag-1 $\bar{\rho}_1 = 0.928$; corpus-wide by 42%, [Schessl, 2026a](#)). **Bundestag subset* $n = 4,251$ after minimum sampling $n_{\text{turns}} \geq 15$ (§3.5) from the corpus of $n = 6,874$ in §4.6.

The profile does not decay monotonically with scale: it orders by support and load-bearing capacity. The dyadic specimen carries all three quantities, the macro-institutional V-Dem delivers the strongest sigmoid R^2 of the work; the extended Bundestag corpus falls into the structural null range on E_a and $\hat{a}$ (truss, no moment). The five-row structure is motivated by the subsystem decomposition of systems theory ([Luhmann, 1984](#)): five carriers, not five arbitrary scales. The support gradient clamped $\rightarrow$ cantilever $\rightarrow$ truss $\rightarrow$ two-span beam is itself a mutual-support axis: the same mechanics (the boundary condition decides which axis bears) across four scales.

5.2 Five Support Types: Structural Mechanics of Conversation

In materials testing, the boundary condition decides which displacement quantities are measurable, which support reactions arise (V vertical, H horizontal, M moment), and which internal forces the system carries (N normal force, Q shear force, M bending moment). The five support types correspond to the standard statics typology ([Hibbeler, 2012](#); [Albert et al., 2018](#); [Krätzig et al., 2014](#)).

Table 9: Structural-mechanics signature of the five support types ($V/H/M$ = support reactions; $N/Q/M$ = internal forces)

Support type	Support reactions	Transferred internal forces	Static determinacy	Substrate
Clamped	$V + H + M$ at both ends	N, Q, M	statically indeterminate (3-fold)	ENK dyadic
Cantilever	$V + H + M$ at the clamped end, free end	N, Q, M (max. M at the support)	determinate	AgoraSpeech, IQ2
Truss	$V + H$ at hinged nodes, no M	only N (member normal forces)	determinate if $2j = m + r$	Bundestag (corpus as truss system)
Two-Span-Beam	V at three supports	N, Q, M with mid-support moment	indeterminate to 1st degree (residual stresses, hysteresis)	V-Dem trajectory
Pinned	$V + H$, no M	N, Q , no M	determinate with roller	Political speeches in general

Static determinacy as a diagnostic axis. Three states can be read off every substrate: *determinate* (support reactions uniquely determined from equilibrium, unique load path: cantilever speeches), *indeterminate* (redundant constraints produce residual stresses that carry hysteresis: doubly-clamped gold-baseline dyad, crystallized conversation, V-Dem recovery asymmetry), *underdetermined* (no unique load path under free movement: diffuse conversation, ProsocialDialog recovery).

Axis-to-internal-force. The split into internal forces $N/Q/M$ is the structural reading of a load response that is uniform at the material level: the micro-imperfection cascade (§2.4) carries every load down to the tensile separation of the smallest bond, and only the support topology decides which measurement axis breaks through. E_a (stiffness) and $\hat{a}$ (yield inflection) break through only where the support transmits a bending moment M , i.e. where bidirectional load redistribution through a counter-support is present; Q_{sv} carries the normal-force path N , which is retained even in a pure truss. Table 8 then reads in advance: where the counter-support is missing (cantilever) or the bending moment vanishes (truss), E_a and $\hat{a}$ fall into the zero-force member while Q_{sv} carries; only the two-sided clamping (clamped) bears all three.

Clamped: dyadic conversation. Both interlocutors deliver full support reactions; all five measurement axes are load-bearing ($E_a, Q_{sv}, \hat{a}$ conformant).

Cantilever: component, not structure. A single speech *taken on its own* is a cantilever: the speaker loads, but an internal counter-support is missing. The axes that measure bidirectional coupling fall into the zero-force member: a yielding without immediate counter-support *within the speech scale*. AgoraSpeech carries a weak residual signal from the paragraph structure; IQ2 falls cleanly into the zero-force member (all three axes n.s.).

Truss: speech corpus as truss system. The corpus over 4,251 speeches is plotted as a truss system: each speech a member, hinged via the agenda as a node. The support reactions carry only $V + H$, no moment; the truss transfers only normal forces. Axes that presuppose a bending moment or hysteresis become zero-force members on the truss. The predicted null on E_a and $\hat{a}$ is structural mechanics, not measurement failure: a truss bears no bending. The institution generating this corpus is measured as a support separately (§4.6): pinned coupling to opinion salience (force yes, moment no), emergency-support characteristic under full load, sluggish level tracking by subordinate chambers. The effect of the speeches runs via other truss systems (audience, voting behavior, V-Dem indices) and is measurable at the next scale.

Two-Span-Beam: V-Dem. The index trajectory over time has three load-bearing supports (initial, crisis, final constitution), indeterminate to 1st degree $\rightarrow$ residual stresses from path-dependent deformation. Observed: 67% of episodes show recovery within a decade, never identical to the initial state: the hysteresis is the residual-stress signature of the second support. The sigmoid fit reaches $R^2 = 0.983$.

Pinned: political speeches in general. Campaign speeches without a dynamic audience response carry $V + H$, no moment; a methodological intermediate stage between cantilever and truss.

5.3 Addressee-Support: the Measurement Configuration as an API Axis

The five support types are implicit in the SRDS frame. Formalizing them explicitly as a measurement axis makes the configuration visible for applications, analogous to construction-materials tables (S7/S10/S13 for timber per DIN 4074-1; XC1/XC2/XD3/XF4 as exposure classes per EN 206). The SRDS API specifies the addressee classes as an extension and implements them operationally.

Table 10: Five API axes for SRDS measurement configurations

API parameter	Values	Construction analogy	Measurement consequence
mode	dyad / multi-speaker / monologue	number of supports	which component topology
exposure	XC1 / XC2 / XD3 / XF4	exposure class (EN 206)	which load-case corridor
tier (preset)	default / therapy / negotiation / education	sorting class (S7/S10/S13)	which material expectation value
addressee	direct / external_modeled / collective_proxy	support type	which support type is realized
addressee_E	default / typical_individual / typical_collective	E-modulus of the support	which standard assumption for an invisible support

Consequence for the speech section §4.6. With the addressee-support model, a political speech is no longer “predicted null” on the speech-language scale but extended-measurable against a modeled societal support proxy: `mode: monologue, addressee: collective_proxy` (audience as collective support: voting behavior, media discourse, V-Dem indices), `addressee_E: typical_collective_<era>`. Under this configuration the Bundestag null becomes a cantilever-style measurement against the societal proxy. ParlaMint realizes exactly this configuration: speeches as an $\ell_3 \rightarrow \ell_4$ bridge against the V-Dem Polyarchy Index. The force transfer is **directionally confirmed**: on the 17-country panel (1,802,373 speeches, 238 country-years) the sv axis carries cluster-robustly ($\rho = -0.284$ at lag +2, $G = 17$; BCa-CI excludes zero); in the 4-country pilot the $\hat{a}$ -discourse axis is




Figure 9: Three spring-stiffness configurations with deformation trajectories: (a) clamped dyadic (elastic restoring), (b) cantilever + collective_proxy (damped hysteresis), (c) cantilever without support (pure creep).

negative across all four clusters (HU -0.40 , PL -0.52 , RS -0.21 , SI -0.40 ; BCa- CI_{95} $[-0.34, -0.15]$ excludes zero). Open is the full operationalization (triangulation), not the mechanism.

5.4 Propaganda Mechanics under Missing Restoring Force

The addressee-support configuration carries a mechanical explanation of the propaganda phenomenon. A one-sided loading without direct resistance corresponds to a structure without spring stiffness at the free end: the elastic bedding of structural statics (Winkler, 1867; Pasternak, 1954). The load is absorbed, but no restoring force returns the system; mechanically this implies creep instead of fracture. The same statement follows from the theory of communicative action (Habermas, 1981): without an answer-support the validity-checking counter-speech-act is missing, and validity claims are adopted adaptively instead of examined.

Table 11: Three configurations of spring stiffness at the support

Configuration	Spring stiffness	Response under sustained load	Example
Clamped (dyadic)	high (both sides)	elastic restoring, then plastic deformation with hysteresis	ENK gold-standard dialog
Cantilever with answer-support	medium (collective via addressee_E)	damped restoring, slower hysteresis	Bundestag debate with audience
Cantilever without support modeling	near zero	creep without restoring, permanent plastic deformation	NS propaganda to an unstructured audience

Empirical anchor: NS speeches show lower $\hat{a}$ (Cohen’s $d = -2.39$, $p < 10^{-4}$, $n = 11$ NS vs. Bundestag $n = 6,874$) and stronger frame drift, consistent with the creep model (early yielding without a moderating support). At $n = 11$ the effect size documents direction, not population magnitude; illustrative only. This mechanism is not rhetorical analysis but statics: a structure without spring stiffness at the load point deforms plastically under any one-sided excitation; a democracy with institutional supports carries a moment-load-bearing restoring force that an authoritarian configuration does not provide.

5.5 Empirical Truss Evidence: Institutional Support as Counterforce

The spring-stiffness reading makes a testable prediction: if the institutional truss (horizontal accountability: judiciary/legislature constraining the executive) acts *during* loading as a counterforce that mitigates the deforming load, then a stronger truss should make autocratization slower and costlier without altering the recovery. Across 115 V-Dem episodes in 87 countries (truss measure: judicial-checks index in the year before onset; convergent with the democratic prior history, $\rho = +0.57$), cluster-robust against country clusters:

- **More work to failure (toughness).** Stronger truss $\rightarrow$ higher toughness: $\rho = +0.55$ (judicial checks; cluster-CI $[+0.40, +0.67]$, $p < 10^{-4}$) and $+0.61$ (prior history; $[+0.47, +0.71]$).
- **Slower erosion (damping).** Stronger truss $\rightarrow$ lower steepness k : $\rho = -0.18$ (cluster-CI $[-0.35, -0.001]$, $p = 0.047$) and -0.23 ($[-0.40, -0.04]$, $p = 0.014$).

Both effects are cluster-robust (confidence intervals exclude zero). They are the mechanical signature of the counterforce, not intrinsic material grade: the truss mitigates the *net* load and distributes it across the strut system. Consistently, recovery is *not* truss-modulated. The 67% hysteresis (§5.2) occurs regime-independently (toughness $\leftrightarrow$ recovery $\rho = 0.14$ n.s.): the truss acts during loading, not during recovery. On the time-lag axis: across the 17-country panel (238 country-years) the discourse signals lead the polyarchy response with a phase lag of about two years, cluster-robust on the sv axis ($\rho = -0.284$ at lag $+2$, $G = 17$; BCa-CI excludes zero; a contiguous negative band over lags 0 to $+2$). This phase lag mirrors the phase shift of transient heat transmission through a storage mass.

5.6 Budget State Machine: Architecture without Disclosed Thresholds

At the structural scale, the damage accumulation $B(t)$ from §3.5 is transferred into a graduated state machine:

NORMAL $\rightarrow$ WATCH $\rightarrow$ WARNING $\rightarrow$ CRITICAL.

Two configurations operate in parallel: a **production configuration** (ultimate limit state, tuned to a zero false-alarm rate against prosocial reference specimens, strict γ_M) and a **detection configuration** (serviceability limit state, broader sensitivity, nominal γ_M). The structural architecture (four states,

two configurations, transition rules) is published. The operational calibration parameters (thresholds, contribution weights, state boundaries) remain withheld: for a deployed stability detector, publishing the exact trigger values would enable targeted evasion (responsible disclosure, §9). No reported finding depends on them: the structural load-bearing capacity is threshold-independent.

Two properties follow from the cumulative structure, not from a configuration: **hysteresis** (evidence accumulates faster than it decays: a loading-unloading asymmetry) and **fatigue memory** (specimens previously in elevated states re-escalate with less additional anomaly; load-bearing capacity does not fully return). Both correspond to established phenomena in materials testing (Palmgren-Miner damage accumulation, Bauschinger effect) and interlock with the G7 hysteresis axis across battery, multiaxial fatigue, and V-Dem (§4.13). The path-dependence is structural, not parametric.

5.7 Application Sketch: ℓ_3 -Trajectory Monitoring

Online hate speech often operates in cantilever mode (a single poster loads, with no direct counter-support in the thread) against a modeled collective-support proxy. Message-level detectors (Perspective API, Detoxify) measure the stress peak value, not the cumulative load-bearing capacity of the structure. ℓ_3 -trajectory monitoring (thread, account trajectory, subforum corpus) is the structural measurement complement: it is the structural monitor that observes a building over time, rather than a set of single smoke detectors.

Such an application targets trajectory stability, not intent: what a thread shows in cumulative deformation is a material statement about the structure, not about the inner stance of individual participants. Cohen’s κ is not the primary validation metric here: what is measured is continuous trajectory deformation, not categorical label agreement, and under the prevailing class imbalance the coefficient collapses paradoxically anyway (Feinstein and Cicchetti, 1990). The load-bearing capacity is established through cross-corpus replication (twelve corpora, §6), not through an inter-rater study. The structural monitor does not replace the smoke detector; it complements it by the second scale, just as a building simultaneously carries acute fire-hazard detectors and long-term settlement sensors (Structural Health Monitoring; Farrar and Worden, 2007; Sohn et al., 2003).

6 Conversation Cross-Corpus and Cross-Method Validation

The material level from §3 measures one conversation on the ENK gold corpus ($n = 202$). That the same deformation profile recurs across twelve external conversation corpora (per-corpus breakdown, including three pre-registered controls, in §6.5) forms the second support of the load-bearing claim. Cross-corpus replication corresponds to multi-cohort validation in clinical epidemiology (Steyerberg, 2009): a method survives when it yields the same measurement signature on disjoint samples under the same structural load.

6.1 Twelve Conversation Corpora (plus Two Cross-Domain Extensions)

Table 12: Cross-corpus validation across 12 corpora

Corpus	n	Mode	Headline statistic	Status
ENK-Gold (anchor)	202	mixed	$\hat{a}: \rho = -0.359^{***}$, ridge	PASS
BothBosu v3 (BothBosu, 2024)	1,245	covert	$\rho = 0.700 \pm 0.076$ $B_w \rho = +0.317$ $p < 10^{-6}$; permutation 19.2× null median	PASS

(continued on next page)

(continued)

Corpus	n	Mode	Headline statistic	Status
GasConv Gaslighting (Li et al., 2025)	3,998	overt	dir_consistency sign-inverted $p < 10^{-20}$	PASS
Awry (CGA Wikipedia) (Zhang et al., 2018)	3,421	mixed	cross-corpus replication	PASS
LegalCon (Sheshanarayana et al., 2025)	727	covert	NC2 $\rho = +0.719$, $p < 10^{-50}$ (highest singular effect); $\hat{a}$ $\rho = -0.751$	PASS
MentalManip (Wang et al., 2024)	1,294	covert	$\hat{a} \rho = -0.110^*$; bond hierarchy PASS	PASS
P4G Persuasion (Wang et al., 2019)	1,016	covert	$\hat{a} \rho = +0.071^*$; sign inversion (persuasion = later transition)	PARTIAL
SafeDialBench (Cao et al., 2025)	2,037	overt	NCD $d = +6.35$ (highest d value); $B_w d = +3.55$; $\det B_t d = +3.52$; 12/12 sig	PASS
CoSafe (Yu et al., 2024)	1,400	overt	BL-2 Cramér $V = 0.705$; dir_consistency $d = +3.17 / -3.80$; ANOVA NCD $H = 62.6^{***}$	PASS
ProsocialDialog	5,000	null result (therapeutic)	$ \rho < 0.10$ all metrics; budget NORMAL in 307/308 specimens	Null-PASS
Sotopia agent-agent	3,111	null result (symmetric)	38.7% sigmoid structure, A0 violation	Null-PASS
Google COVID Mobility	—	null result (externally imposed)	$R^2 > 0.9$ formal, yet no operator fixed point, no hysteresis	Null-PASS
DarkBench / WildJailbreak	—	skip	single-turn = hardness test, not tensile test	construct boundary

In addition, two extended cross-domain corpora from the §5 structure level: AnnoMI (Wu et al., 2022) ($n = 133$ Motivational-Interviewing transcripts, MAS $d = +1.02$ PASS cross-domain, $\hat{a}_{\text{AnnoMI}} = 0.33 \neq \hat{a}_{\text{Manipulation}} \sim 0.52$) and CMV Full (Tan et al., 2016) (Reddit Change-My-View, $n = 11,740$; 8/11 Bonferroni-sig; FID_max partial $\rho = +0.104/d = +0.232$; E_a partial -0.064^{***} survives length control).

Total: $\approx 35,300$ dialogues across 14 data sources (the 12 corpora of Table 12 plus two cross-domain extensions, AnnoMI and CMV Full): 8 validated, 3 null results, 1 construct boundary, 2 cross-domain extensions. The Sotopia count is the ENK subset with ≥ 8 turns from the cmu-lti/sotopia public release of 7,200 episodes ($n_{\text{Sotopia}} = 3,111$); AnnoMI contributes $n = 133$, and the remaining sources total 32,080.

6.2 LegalCon as Showcase: Highest Singular Cross-Corpus Effect

With NC2 $\rho = +0.719$, $p < 10^{-50}$, the LegalCon corpus ($n = 727$, plaintiff-defendant dialogues) carries the strongest singular cross-corpus effect of the entire validation. $\hat{a}$ as strain axis reaches

$\rho = -0.751$. NC2 (Newman conductance on 2-cluster partition) corresponds to grain-boundary density in microstructure analysis (Hall, 1951; Petch, 1953); within LegalCon it measures how the semantic space stiffens under adversarial cross-examination. Here the Hooke coupling runs inverted ($d = -0.881$, brittle-fracture mode), unlike covert manipulation, where it runs in the creep direction. The sign inversion is not a defect but a phase boundary: the same material shows different failure modes under different loading geometry.

6.3 Sign Paradox: Ductile vs. Brittle Fracture

Table 13: Sign inversion covert vs. overt, corresponds to the materials phenomenology of ductile vs. brittle failure modes (Considère, 1885; Bridgman, 1944)

Mechanism	Covert (BothBosu, P4G, MentalManip)	Overt (Awry, GasConv)	Material reading
B_{windowed}	+ (accumulates)	− (discharges)	creeping vs. brittle fracture
FID	+ (frame drifts slowly)	− (topic switches abruptly)	ductile vs. brittle
dir_consistency	− (zigzag, confusion)	+ (confrontation)	confusion vs. consistent confrontation
NC2	+ (expansion)	− (collapse)	stiffening vs. loss of cohesion

The paradox is a classification axis, not methodological ambiguity: the same functional form carries two failure modes (slow creep, abrupt brittle fracture), and the sign inversion tells them apart. This corresponds to the transition from ductile to brittle fracture characteristics in materials testing (Considère criterion for plastic instability; Mohr criterion for brittle fracture). That the same inversion appears between covert and overt manipulation is a measurement on the same substrate, not an analogy.

6.4 Cross-Method Triangulation: Three Methodologically Disjoint Apparatuses

On a D7 phase subset (P3 transition phase), three methodologically independent measurement apparatuses converge on the same phase structure. Method triangulation corresponds to the classical mixed-methods principle (Denzin, 1978; Flick, 1992): a finding holds when it remains replicable across different measurement axes with different formal preconditions.

Table 14: Three disjoint apparatuses on the same substrate

Apparatus	Formal basis	Finding across three phases P1→P3
TF-IDF/SVD norm statistics	linear algebra, singular-value spectrum	NC2-CoV: drop down to -32.5% at SVD-130, robust [80, 150]
Manifold learning TwoNN	Levina-Bickel MLE for intrinsic dimension (Levina and Bickel, 2004), GRIDE family (Denti et al., 2022); LLM-layer adaptation (Viswanathan et al., 2025)	intrinsic dimension monotonically falling
Topological data analysis	Vietoris-Rips filtration, persistence homology (Edelsbrunner and Harer, 2010); zigzag persistence on LLM trajectories (Gardinazzi et al., 2025)	persistence lifetime 1132 → 901 → 824 monotonically falling

The three apparatuses are methodologically disjoint: linear spectra, nonlinear manifold learning, topological filtration. They converge on the same structural statement: phase transitions in the D7 phase subset show a monotonically falling dimension / complexity signature. Cross-method convergence corresponds to the epistemic requirement of measurement robustness against method artifacts (Cronbach and Meehl, 1955, construct validity); no single apparatus proves the finding; the three together, each with its own falsification condition, supply the support.

Concrete drop signs and magnitudes are subset-specific (on a second subset NC2 shows the inverse direction), which supports the no-go statement from §2.5: the asymmetry is substrate-specific, not a universal constant. The triangulation runs on a DUA-restricted ENK subset (participant conversations); the three apparatuses rest on published methods (Table 14), scripts and aggregate results on request.

6.5 One Predicted Null and Two Scope Conditions: Lemma Violations as Construct Support

Three conversation corpora carry no load signal. ProsocialDialog was registered as a negative control *before* measurement (L1 violation, §2.2 Table 2): a genuine predicted null. Sotopia and Google COVID Mobility were pre-registered with directional positive hypotheses (29 March 2026) and falsified; the lemma violations (A0 and $L3/\sigma \perp \varepsilon$, respectively) account for the falsification *post hoc* and map the framework’s scope conditions.

Table 15: One pre-registered null control and two falsified positive substrates (scope conditions)

Negative corpus	Violated lemma	Consequence	Finding
ProsocialDialog ($n = 5,000$; Kim et al., 2022)	L1 (Lyapunov half-order) violated $\Rightarrow$ A2 not derivable	$\beta \rightarrow 0$, recovery balances damage	all SRDS metrics $ \rho < 0.10$; budget NORMAL in 307/308 specimens; LlamaGuard false alarm $\rho = +0.327^{***}$ on identical data
Sotopia agent-agent (Zhou et al., 2023)	A0 violated (symmetry instead of asymmetric self-reference)	no path asymmetry possible	38.7% sigmoid structure, fracture characteristic does not occur
Google COVID Mobility	L3 / $\sigma \perp \varepsilon$ violated	σ and ε both time-driven (circularity)	pre-registered-positive, falsified: raw $r = +0.592$ vanishes after calendar-time control ($R^2 > 0.9$ formal sigmoid, yet no operator fixed point)

Null and positive findings are situated under the same axioms: the same formal apparatus that measures load-bearing capacity on positive substrates *predicts* a null signal on a substrate pre-registered as a negative control (ProsocialDialog). This corresponds to Popper’s falsifiability criterion (Popper, 1959) in constructive form. Sotopia and Google COVID Mobility were pre-registered-positive, falsified, and map the scope conditions. Quantified in §8 as a balance: one predicted null plus two scope conditions.

In addition, the DarkBench/WildJailbreak skip: construct boundary, not measurement failure. Single-turn specimens correspond to a hardness test (Brinell, 1900; Smith and Sandland, 1922), not a tensile test (DIN EN 408). The ENK $\hat{a}$ methodology requires a loading history of at least 15 turns (minimum sampling from §3.5); single-turn lies below this minimum load duration. The skip is construct-consistent: a hardness test cannot be read as a tensile test.

7 Cognition Bridge and External Frame Convergence

7.1 Cognition Bridge: G_f/G_c and the Sigmoid Mélange

The sigmoid inflection $\hat{a}$ bridges to a cognitive carrier class; this is a theoretical extension, not yet empirically validated. The Cattell-Horn-Carroll model in intelligence psychology (Cattell, 1963; Horn and Cattell, 1967; McGrew, 2009) distinguishes fluid intelligence G_f (adaptive-elastic, new problems) from crystallized intelligence G_c (accumulated-brittle, experience-based). The life-span trajectory is empirically documented: G_f falls concavely from early adulthood, G_c rises convexly into late adulthood (Salthouse, 2004).

Material reading: G_f corresponds to elastic deformation (reversible, adaptive load redistribution), G_c to plastic deformation (accumulated, irreversible structure formation). The intersection of the two trajectories, the *mélange*, has sigmoid form: a transition from G_f -dominated to G_c -dominated with inflection point in middle adulthood. A sigmoid fits this *mélange* form ($R^2 = 0.853$). Unlike the empirical fits for V-Dem ($R^2 = 0.983$) and earthquakes ($R^2 = 0.965$), however, this fit measures agreement with the *theoretical* CHC *mélange*, not with longitudinal data. Validating this against longitudinal data (HRS/MIDUS) remains follow-up work. Here $\hat{a}$ marks the yield point at which accumulated experience (G_c , plastic) compensates for the slowing rate of adaptation (G_f , elastic): wisdom compensation in life-span psychology (Baltes and Staudinger, 2000; Baltes and Smith, 2008).

The bridge makes a cognitive carrier class visible as a *hypothesis*, alongside the empirically measured substrates, not as an eleventh validated substrate. Operationalizing the trajectory as a sigmoid is a hypothesis with a defined falsification condition (HRS/MIDUS longitudinal), not an analogy.

7.1.1 Chen et al. 2026: Long-CoT as Molecular Structure

A second bridge operationalizes the molecular structure of thought directly in the cs.CL substrate. Chen et al. (2026) (*The Molecular Structure of Thought*, ByteDance Seed / HIT / Peking University, arXiv:2601.06002) model long-chain-of-thought trajectories as molecule-like structures with three bond types:

Bond class (Chen 2026)	Materials-science reference	ENK analogue
Deep reasoning	covalent bond (~ 350 kJ/mol)	1 – sem velocity (long-range carrier structure)
Self-reflection	hydrogen bond (~ 12 kJ/mol)	h_{bond} decay ($\rho = +0.398^{***}$, LOPO 6/6)
Self-exploration	Van der Waals (~ 3 kJ/mol)	mean-cos far-field ($\rho = +0.133$, n.s., $p = 0.062$; theory-consistent weak for the weakest bond class)

Chen’s Effective Semantic Isomers carry the statement that *only bond configurations with fast entropy convergence* support stable long-CoT learning, while *structural competition* destabilizes training. This is structurally homomorphic to the damage signature measured here as B_w accumulation (§3.5). This corresponds to the Anfinsen hypothesis of protein folding (Anfinsen, 1973): just as the tertiary structure is encoded in the amino-acid sequence, the stable CoT trajectory is encoded in the bond configuration of the reasoning steps. The four-level hierarchy (sequence, secondary structure, tertiary folding, multimer; (Berg et al., 2019)) corresponds in the language substrate to the staggered levels from token sequence through local patterns and thematic clusters to multi-conversation behavior, paralleling the scale hierarchy ℓ_0 to ℓ_4 from §2.3.

7.2 External Frame Convergence: The Gap That SRDS Fills

Several independent groups arrive, under different disciplinary names, at structurally identical arguments. **These works do not confirm SRDS**: they show that the same class form arises independently

across disjoint fields. The list is not the load-bearing pillar of this work; it is evidence that the gap a cross-substrate frame closes is real. These convergence anchors satisfy the epistemic requirement that a theory be robust against observational artifacts (Hacking, 1983).

Table 16: Convergent independent work across disjoint fields

Anchor	Substrate	Statement	Reference to SRDS
Olmeda et al., 2026 (Nature Physics)	epigenome	self-similarity power-law $5/2$, renormalization-group schema	RG operator $M = N \circ R \circ C$ (§2.3) on a biological carrier
Simon et al., 2026 (<i>learning mechanics</i> , arXiv:2604.21691)	deep learning	discipline-naming in direct parallel to the frame presented here	seven desiderata, including <i>humble</i> , meets the construct-boundary concept (§8)
Zhdanov et al., 2025 (AdS-GNN)	conformal-equivariant geometry	operator composition on a mathematical carrier	functor invariance, not numerical fidelity (§2.5)
Schmidt et al., 2026 (Topological Deep Learning) Yang et al., 2026	topological vs. combinatorial generalization mesh-agnostic operator composition	topology as measurement substrate operator logic independent of the discretization mesh	functor $\Psi : \text{SRDS} \rightarrow$ aggregate class as functorial realization renormalization operator M carries under coarse-graining (§2.3)
Bakeer, 2026 (computational mechanics, arXiv:2605.28601)	spatial parameter identifiability	local information operators; two-span support coupling, beam kernels, “visible/invisible” directions	same statics vocabulary on a <i>spatial</i> rather than temporal carrier; published later and addressing a different problem: <i>existence</i> , not <i>confirmation</i>
Chen et al., 2026 (ByteDance Seed, arXiv:2601.06002)	cs.CL long-CoT	bond topology covalent / H-bond / vdW (§7.1)	mélange operator realized directly in the language substrate

The convergence rests not on count but on structural similarity across disjoint substrates. Olmeda (biological renormalization) and Simon (*learning-mechanics* naming) arrive via disjoint paths at the same form: renormalization operator as central measurement axis, construct-boundary concept as methodological stance. Three mathematical works (AdS-GNN, Topological Deep Learning, mesh-agnostic composition) realize the same functor on disjoint formal carriers; one cs.CL paper (Chen 2026) operationalizes the molecular structure of thought directly in the language substrate. The computational-mechanics anchor (Bakeer, 2026) applies local information operators to the *spatial* identifiability of distributed parameters: the same statics vocabulary (two-span support coupling, beam kernels, visible vs. locally invisible directions) on a disjoint, spatial carrier; later in time and on a different problem, hence convergence, not confirmation. The class SRDS is realized consistently in structure across three mathematical languages: functor-invariant, not number-faithful (§2.5).

Closest to this approach is Buehler’s materiomics line: hierarchical structural isomorphism across materials, proteins, music, and language, most recently a self-organized-critical graph-reasoning state with a dimensionless *Critical Discovery Parameter* (Buehler, 2025). The overlap is real and touches two substrates; so is the delta. That line reads *structural* similarity across a single substrate family and *seeks* a critical parameter, whereas SRDS fixes a *dynamical* damage class (load, accumulation, yield inflection $\hat{a}$, irreversibility) across ten physically disjoint substrates: with a synthetic null and pre-registered scope limits (§8), a computer-assisted no-go theorem with Lean-4 formalization (§2.5), and retained failures.

Where that line seeks such a critical parameter, the no-go theorem *excludes* a universal constant.

Beyond structural form convergence, the *methodology* of this cross-substrate move has its own name in philosophy of science: **patchwork concepts** (Wilson, 2006; Haueis, 2024). A concept that fragments into “patches” across scales and substrates (each with its own measurement technique, united by a *general reasoning strategy* rather than an identical core mechanism) is on this account not a mere analogy but a regular conceptual form. This covers exactly what SRDS does at the measurement level: the technique changes per substrate (tensile test, V-Dem fit, embedding geometry, capacity fade), the measured property (inflection $\leftrightarrow$ outcome) remains. Haueis’s third norm, *to check whether the property actually applies in each patch* (Haueis, 2024, pp. 749–750), coincides with the PARTIAL/FAIL discipline practiced here: the substrates without signal (ProsocialDialog, Sotopia, Mobility) *support* the conceptual form rather than weakening it. Independently, Rorot (2025) investigates the scale-free concept of *communication* across disciplines (corpus analysis over ~ 1.1 M papers) and defends the realist reading against the “mere metaphor” objection: there is “*no clear, arbitrary line*” between literal and metaphorical use, and the realist reading has “*firmer grounds than any anti-realist readings*” (pp. 265–266). This licenses the cross-substrate measurement step, not a shared mechanism: the *deeper* unification remains the RG folding onto atomic Mode-I tension (§2.3, §9.1).

8 Falsification Anchors, State of Measurement, and Limits

The burden of proof is reversed (§1.1b): the mechanics is not defended case by case; the skeptic would have to exhibit a finding that *violates it with adequate load and specimen count*. This section states the falsification tests (§8.1), the three falsification layers (§8.2), the fit $\leftrightarrow$ counterpart systematics (§8.3), the comparison against message-level detectors (§8.4), the limits (§8.5), and closes with the structural limit (§8.6).

The demarcation. A **falsification** is a finding that **violates a physical prediction with adequate load and specimen count** (sufficient statistical power to reach a yield signal): a severed yield-point/strength coupling, a bending truss, a self-reversing accumulation. **Not** a falsification is (a) an underpowered null (small sample, wide confidence interval), (b) a *moot* test (on forced sequences a path surrogate is physically meaningless, §1.1b), or (c) a near-elastic specimen without a plastic regime (no yield $\rightarrow$ no yield signal). An underpowered null is also **not** a “partial confirmation”. Each is named for what it is.

8.1 Prospective Acceptance Rule: What Is Fixed Before Measurement

The reversed burden of proof (§1.1b) would be immunizing were there no rule that fixes, *before* the analysis and from a substrate’s raw quantities, whether it is SRDS-capable, which axis bears load, and what counts as falsification. That rule is already fully in place; here it is gathered in one spot. It has three parts.

(a) Detectability conditions. A substrate supports a load-bearing measurement only when four things come together: a visible coupling (which quantity loads which), a markable material (a recoverable response, not just a topic), a load-free reference or control (a “before”), and sufficient resolution (Nyquist; too coarse a time axis denies the arrow of time). If one is missing, the substrate is describable but not demonstrable — the finding falls as *unpopulated*, not as a null.

(b) A-priori death conditions. Three conditions are fixed before measurement, and violating any of them drops the finding: (i) the **free-sequence surrogate** — every freely ordered substrate must really exceed the IAAFT surrogate (spectrum and marginals preserved, only the path destroyed), else it falls to generic form recurrence (passed: ENK $z = -4.86$; open per further substrate and named as open); (ii) the **pre-specified sign** — the sign is fixed a priori per load type (Mohr inversion), a PASS with the wrong sign is a fail, and Bonferroni runs against the *directed* family; (iii) the **pre-determined $\hat{a}$ band**, where the substrate structure predicts one (material class $\rightarrow\hat{a}$): a finding outside the band tears the prediction.

(c) **Front-end by stationarity class.** Which input feeds the state machine is fixed before measurement by the substrate’s stationarity class (same counter, different numerator): under recurrent rest with episodic load, the deviation from a fixed rest baseline; under sustained load without rest (unstable second moment), the residual against an adaptive or physics-based reference; under monotone drift, the shape feature of the cumulative curve (inflection point). The wrong choice produces a spurious null, not a real one.

These three parts fix SRDS-capability, mode, normalization, sign, and falsification criterion *before* the analysis. Their binding force shows in that the framework can fail against them, and does: Sotopia (A0 violation) and Google COVID Mobility ($\sigma \perp \varepsilon$ violation) were pre-registered as positive, tested, and falsified; the axiom reading of these two boundaries explains the falsification *post hoc* and is named as such (§8.2), not a pre-registered null. A framework that absorbed every outcome could not state this rule.

8.2 The Falsification Lattice

Each row names a physical rule, the finding that would *overturn* it, and the observation. None is overturned.

Table 17: Falsification lattice (rule $\rightarrow$ falsifier $\rightarrow$ observed)

#	Physical rule	What would falsify it	Observed
G1	forced $\rightarrow$ material law holds	timber $\hat{a}$ says nothing about strength	$r(\hat{a}, \sigma_{\max}) = -0.83$ (arithmetic consequence of onset normalization, §4.1; substantive: within-class $-0.83 / -0.86 / -0.95$, $CV_{\text{onset}} = 0.228$)
G2	free $\rightarrow$ the load <i>path</i> carries the signal	ENK $\hat{a} \leftrightarrow \text{mal}$ survives path destruction	$z = -4.86$ (stands out of the surrogate cloud)
G3	a truss bears no bending moment	strong $\hat{a}/E_a$ signal at the speech-corpus truss	$E_a + 0.023$ n.s., $\hat{a} + 0.035$ n.s. (null member)
G4	no universal constant (no-go)	<i>one</i> universal $\hat{a}$ value across all substrates	KS $\hat{a}_{\text{emp}} \neq 0.50$ in 10/10 distinct ($p < 0.01$; 7/10 at $p < 10^{-3}$)
G5	$\sigma = E\varepsilon$, axes coupled	$\hat{a}$ and amplitude perfectly orthogonal	$\rho(\hat{a}, \text{std}) = +0.807$, $z = +11.2$ (path-generated)
G6	“the sigmoid is trivial”	sigmoid wins also on random monotone curves	AICc: power law 9/10; sigmoid-4p nowhere a majority (Chat-AI 41.6% plurality); discrimination via KS of the $\hat{a}$ distribution, not the win rate
G7	damage accumulates irreversibly	budget decays faster than it accumulates	hysteresis: V-Dem 67% · battery $V_C - V_D + 0.886$ · multiaxial
G8	substrate-specific $\rightarrow$ $\hat{a} \leftrightarrow \text{outcome}$ not equal everywhere	permutation $r(\hat{a}, Y)$ identical everywhere	substrate-specific: timber, chat-AI, FKM, finance carry; near-elastic null (no yield) + DP1180 (underpowered) do not

8.3 Three Falsification Layers

Layer 1: synthetic null specimens. 1,000 specimens per substrate (AR(1)/RW/WN) analyzed with the same pipeline; on all five substrates tested against the synthetic null (Table 18) the Kolmogorov-

Smirnov test separates at $p < 10^{-3}$; the sectoral finance sub-structures bear the load more weakly the more loosely they couple to their index.

Table 18: KS-test against synthetic null, five substrates (all with $p < 10^{-3}$)

Substrate	KS- D	p	Component type
Timber DIN EN 408 ($n = 230$)	0.801	$< 10^{-3}$	timber tensile test
Al-6061 ($n = 146$)	1.000	$< 10^{-3}$	ductile metal ($\hat{a} = 0.084$)
DP1180 Numisheet	1.000	$< 10^{-3}$	high-strength steel
ENK Chat AI ($n = 202$)	0.944	$< 10^{-3}$	LLM conversation
Battery CALB ($n = 26$)	0.733	$< 10^{-13}$	Li-ion degradation

Layer 1b: model comparison against the saturation class family (K1). The sharper objection holds that a normalized monotone cumulative is almost always S-shaped, so the comparison against AR(1)/RW/WN is too weak; the *correct* null is some smooth saturation function. The model comparison across the closed class family (Logistic, Gompertz, Weibull CDF, Hill, AICc) settles this: on random monotone curves the 2-parameter power law wins in 9 of 10 domains; the 4-parameter sigmoid is not the best fit on those nine and only a plurality winner without a majority on the tenth (Chat-AI, 41.6% vs. 36.2% power law), hence not privileged against the correct null. The load-bearing discrimination is not the AICc win rate but the Kolmogorov-Smirnov test of the $\hat{a}$ distribution against the null (Chat-AI $D = 0.944$, $p < 10^{-3}$; §8.2, Table 18). On the empirical substrates Logistic is the best fit in 6 of 8, with substrate-specific form as a No-Go-predicted exception (Al-6061 $\rightarrow$ Gompertz, $\rho(\hat{a}, \sigma_{\max}) = -0.703$ on 146 validly fitted of 154 tensile specimens; CCP60 $\rightarrow$ Weibull, $\rho = +0.867$). The inflection–outcome relation is robust across all four model families (cross-model $\rho = 0.87$ –0.99). The corresponding pre-registration carried three primary form hypotheses, **none confirmed** (no universal Logistic dominance); the secondary hypotheses (inflection–outcome substance, cross-model robustness) are confirmed. That high fit quality alone does not discriminate between model classes is also documented in data-driven material modeling: six discovery frameworks reproduce the classic Treloar dataset equally well, with no single winner (arXiv preprint, [Holthusen et al., 2026](#)).

Layer 2: predicted null and empirically found scope conditions. ProsocialDialog was registered as a negative control *before* measurement (L1, $\beta \rightarrow 0$, §6.5) and carries the predicted null: no false alarm despite therapeutic intensity ($\rho = -0.008$). Sotopia and Google COVID Mobility, by contrast, were pre-registered with *directional positive hypotheses* (29 March 2026) and falsified; their failure empirically maps the framework’s scope conditions: Sotopia violates A0 (agent-agent symmetry instead of asymmetric self-reference); Google COVID Mobility violates the $\sigma \perp \varepsilon$ premise (mobility and cumulative stringency are both time-driven, a circular operationalization). Ductile metals additionally mark the operating-range ceiling. A *pre-registered* null is the sharpest falsification condition; a *falsified positive prediction* that reveals a scope limit is the second-sharpest; neither can be produced by a curve-fit artifact. The axiom reading of the two limits is a *post-hoc* explanation of the falsification, not a pre-registered null.

Layer 3: sealed test batch. Pre-registration (18 March 2026), six primary hypotheses under Holm-Bonferroni. On the strictly separated primary batch never used for calibration (batch A, $n = 36$) all six predictions lie in the predicted direction; under the conservative Holm correction **one** clears the single-metric threshold (H3, $r = +0.140$). **The decision rule $\geq 4/6$ was not reached** (an underpowered null finding under the §8.2 dividing line, not a partial confirmation). Alongside the six single metrics (primary hypotheses), the accumulator was pre-registered as a *meta-hypothesis*: the number of cluster-robust correlations lies on both batches above the pre-specified $2\times$ threshold against the nominal alpha expectation (batch A $4.5\times$, batch B $6.3\times$) and above the label-shuffle null (batch A: 25 observed against 14.4 ± 5.2 , $+2.6$ SD, $p_{\text{emp}} = 0.02$; batch B: 35 against 18.9 ± 6.1 , $+2.7$ SD, $p_{\text{emp}} < 0.02$).

Shuffle calibration is the standard procedure behind these counts: labels are permuted as whole conversation trajectories between conversations (50 draws, identical statistic per test); an association counts as calibrated-robust if its p lies below the 5-percent quantile of the test’s own null distribution — this yields the count of 26/113 at γ_M (§4.10; total count +2.6 SD above the null expectation of 33.3 ± 4.5). The inflection point $\hat{a}$ carries $\rho = -0.566$ on the second sealed batch ($n = 46$ valid of 50), a *non*-pre-registered additional finding, named as such (not “pre-reg confirmed”). That all six point in the predicted direction under limited test load and specimen count is the signature of a real effect at small sample size; a null effect would scatter the signs.

8.4 Fit↔Counterpart: the Nulls Are the Proof

Every positive fit has a *counterpart*: a predicted null where the signal must mechanically be absent. A frame that found signal everywhere could not produce these counterparts.

Table 19: Fit↔counterpart across substrates

Substrate	Positive fit	Counterpart (predicted null/boundary)	Demarcated
Finance	crash sigmoid $R^2 = 0.97$	calm = random walk $R^2 = 0.48$ ($p = 0.0001$)	“every system saturates” refuted
ENK	real path $z = -4.86$	surrogate cloud (path destroyed → generic)	path mechanism vs. generic accumulation
Timber (forced)	$\hat{a} \leftrightarrow \sigma_{\max} = -0.83$	near-elastic cohort, $R^2 = 0.994$ (no yield)	plastic regime <i>gates</i> the yield signal
Model form	sigmoid on real $\sigma(\varepsilon)$	power law 9/10 on random curves	sigmoid not a fit artifact (§8.2)
Structure	V-Dem two-span (class comparison, Gompertz driver)	speech-corpus truss $\hat{a}/E_a$ null (+0.035 n.s.)	support topology, not scale (§5)
Negative control	scam 19.2× (fires)	ProsocialDialog $\rho = -0.008$ (no false alarm)	dose by load, not everywhere
Pre-reg null / scope limit	(load corpora fire)	Prosocial L1 (pre-reg null, confirmed); Sotopia A0 / Mobility $\sigma \perp \varepsilon$ (positive pre-reg falsified)	one pre-reg null + two empirical limits

The pattern is invariant: the signal appears *where the mechanics predicts it* and is absent *where the mechanics predicts its absence*. The latter is the sharper statement: a curve-fit artifact produces no predicted null.

8.5 ENK against Message-Level Detectors

Head-to-head on subtle manipulations ($n = 8$): SRDS 5/8, LlamaGuard 2/8, one tie. On ProsocialDialog ($n = 5,000$, of which 308 with $n_{\text{turns}} \geq 15$) SRDS monitoring delivers $\rho = -0.008$ (zero false alarms; budget NORMAL in 307/308 specimens), while LlamaGuard shows on identical data $\rho = +0.327^{***}$: systematic false alarms at therapeutic intensity (Perspective API carries a comparable profile). This is not a detector competition but a pairing of complementary measurement classes: a fire alarm measures acute peaks, a structural monitor measures cumulative load-bearing capacity (Tresca peak value vs. fatigue analysis, [Wöhler, 1870](#)). Both are needed.

8.6 Limits

Eight limits, each named precisely and placed accordingly. The numbers stand unchanged; the placement only sorts each point as a sufficiently powered falsification, an underpowered measurement, or a construct property.

Table 20: Eight limits, placed

Limit	State	Placement
Bonferroni balance 6/10 conformant	six PASS, Fatigue accumulation-confirmation, three nominal, one predicted null + two scope limits	measurement balance (no re-count)
Annotator dominance 69%	LOPO macro $\rho = 0.339$ vs. length reference 0.174	underpowered measurement, extensible (more annotators)
Calibration thresholds withheld	architecture + axioms + Lean proofs open	disclosure choice (responsible disclosure, §9); structural capacity threshold-independent
Cross-temporal encoder coupling	addressable via <code>addressee_E</code> + epoch-specific encoders	open measurement task (ParlaMint shows multilingual scaling)
Encoder-specific signs	different encoders = different asymmetry measures	construct-internal prediction (no-go, §2.5), not a limit
Cross-group replication missing	four reproducibility layers documented (protocol/apparatus/corpus/method, §6.4)	underpowered measurement, cross-group = 5th layer (outreach)
$\hat{a}$ interpretation substrate-specific	no universal $\hat{a} \rightarrow \text{outcome}$ mapping	construct-internal prediction (universality classes, Wilson, 1979)
Sealed test batch: $\geq 4/6$ (batch A) not reached	accumulator meta holds (nominal basis $4.5 \times / 6.3 \times$; shuffle null $+2.6/+2.7$ SD); $\hat{a}$ on second batch $\rho = -0.566$	underpowered measurement (§8.2), not a falsification; extensible

In sum: predominantly state of measurement (extensible), plus two construct-internal predictions from the no-go theorem itself and one disclosure choice. No structural limit addressable by additional data, except the no-go (§8.6).

8.7 What Remains: The Structural Limit

After review of the limits, **one** structurally indispensable limit remains: the no-go theorem (§2.5). It rules out that Σ_c carries as a universal constant across all admissible kernels. This limit is not addressable by data or methods; it is *part of the theory itself* and is published in the companion. Structurally this corresponds to Gödel incompleteness ([Gödel, 1931](#)) in the sense in which A0 (§2.2) takes up self-reference: the frame formally specifies its own universality limit and makes it testable. The measurement grammar describes what the load does to the form, not which load a system *should* carry.

9 What Remains When Something Is Missing

9.1 Summary of Findings

This section draws the threads together. The framework comprises five axioms A0–A4, four regularity conditions L1–L4, three core theorems T1–T3, and one no-go theorem (computer-assisted proof:

$\dot{Q}(1/2) \in [1.8412, 1.8963]$). The proof is reproducible with `make verify`; the Lean-4 code is public and the certificate is load-bearing per `#print axioms`. The Lean axiom conservatively encodes $[1.75, 1.99] \supsetneq [1.8412, 1.8963]$, so zero is excluded. The measurement reduces to three steps: accumulate a cumulative strain coordinate, fit a sigmoid, and read off the inflection point $\hat{a}$. Synthetic reference specimens (1,000 per substrate, AR(1)/RW/WN) show that the functional form is not an artifact of fit freedom: on all five substrates tested against the synthetic null (Table 18) the KS test reaches $p < 10^{-3}$, and the model comparison against the saturation class family (Gompertz/Weibull/Hill) does not privilege the sigmoid on random curves (§8.2).

Six of ten substrates meet Bonferroni conformity under the conservative γ_M (timber, ENK conversation, earthquakes, battery under caveat, music, V-Dem sigmoid load-bearing capacity). Multi-Material Multiaxial Fatigue carries as a confirmation on the accumulation axis ($\hat{a}$ reads the load-path non-proportionality, cross-material outcome pooled-damped). Three show nominal signal with named caveats. The scope conditions are mapped by a pre-registered null control (ProsocialDialog L1, confirmed) and two falsified positive predictions (Sotopia A0, Google COVID Mobility $\sigma \perp \varepsilon$).

The main claim rests on empirical evidence. Timber opens with Pearson $r = -0.83$ over $n = 230$ specimens per DIN EN 408:2012-10; the same sigmoid class family recurs in ten substrates, with deviations in form class and inflection location that are themselves measurement values. The spread of inflection points is not deviation from an ideal but a measurement of the substrate’s actual asymmetry: timber $\tilde{a} = 0.567$ (fiber anisotropy), battery 0.646 (CALB, KS-validated), DotCom 0.62, GFC 0.97 (liquidity shock), lignin (inflection at the origin, no endurance limit), earthquakes 0.452, Bach 0.409. This scatter is consistent with the no-go theorem, which formally rules out a universal inflection constant: what is universally load-bearing is the functional form, not the numeric value. Mode-I universality anchors this mechanistically: bending produces compression above and tension below, yet the fracture at the crack tip is always atomic Mode-I tension; the mechanism does not rest on a single dataset, and each substrate carries its own falsification test. Experimentally, the concrete gap test quantifies the same Mode-I interpretation: crack-parallel compression there produces no leading Mode-I driving force but reorganizes the process zone, thereby modulating the effectively measured Mode-I fracture energy (by up to about a factor of two); reproduced mesoscopically in the Lattice Discrete Particle Model (LDPM) (Nguyen et al., 2020; Lyu et al., 2023). In polymer and glass networks the same tensile-terminal reading holds convergently: in a polymer a single covalent bond breaks under tension and releases the energy stored over the entire chain length, so the fatigue threshold rises the more the network deconcentrates the chain tension, whereas brittle silica concentrates the stress onto a few Si–O bonds (Steck et al., 2025).

9.2 Open Replication Tasks

Independent replication outside this research group remains open; two programs structure these tests.

Program A: Causal intervention. Separation of correlation and mechanism. The crystallization signature ($\chi^2 = 18.813$, $p = 0.0001$, Cramér’s $V = 0.310$) and the meta-Fisher across eight corpora ($\chi^2 = 68.96$, $p < 10^{-6}$) deliver structure; an intervention study decides whether they *carry* the load or only resonate: the RCT path of clinical medicine (Hill, 1965, Bradford-Hill criteria).

Program B: Multi-agent symmetry. Two discrete points are established: Sotopia (fully symmetric, 38.7% sigmoid structure, A0 null confirmed) and the ENK gold corpus (asymmetric human-machine dyads, $\rho = -0.359$, A0 active). What remains open is the continuous intermediate stage: slight asymmetries as a parametric gradient, A0 violation as a sweep rather than a single test.

9.3 Data and Code Availability

Table 21: Reproduction anchors (companion paths relative to `srds-enk-companion/`)

Path	Contents
<code>lean4/PositivityProofs/</code>	no-go theorem computer-assisted proof (Arb certificate [1.8412, 1.8963], certificate load-bearing per <code>#print axioms</code>)
<code>genealogy_of_frames.md</code>	dated precursor frame classes (priority anchor)
<code>data/earthquake, battery, crc, holz_master, lignin, musik_companion/</code>	cross-domain raw data and pipeline scripts
<code>data/model_comparison/, analysis/</code>	model comparison (K1, AIC over Logistic/Gompertz/Weibull/Hill) + KS null + permutation inference
<code>REPRODUCIBILITY_MANIFEST.md</code> (ENK subset, DUA, on request)	per table/figure: script path, raw data, external URLs cross-method triangulation NC2/TwoNN/Vietoris-Rips; methods published + aggregate finding in §6.4
Zenodo 10.5281/zenodo.18340365	timber DIN-52188 data (Schessl 2025)
Zenodo 10.5281/zenodo.5142664	MetaMIDI music source (Ens and Pasquier, 2021)

External substrates rely exclusively on publicly available raw data (USGS-FDSN, TCGA Pan-Cancer Atlas, Yahoo Finance, V-Dem ERT-v14, ParlaMint, Materials Cloud, BatteryLife). No data on identifiable persons outside the documented corpus; conversation data in the gold corpus were collected with consent and are pseudonymized. The central cross-substrate claim is independently reproducible via the twelve external conversation corpora ($\approx 35,300$ conversations from 14 sources, §6.1) without gold-corpus access.

9.4 Genealogy

The frame class is publicly documented with dates, a priority anchor, not an evidential support:

Document	Date	Repository
Master thesis (timber data anchor)	May 2025	Zenodo 10.5281/zenodo.18340365
Emergent Narrative Control	21 Aug 2025	Zenodo 10.5281/zenodo.17556213
Life as Organized Dissipation (bridge to A1)	26 Oct 2025	Zenodo 10.5281/zenodo.17445178
The Self-Reference Axiom (precursor to A0)	28 Oct 2025	Zenodo 10.5281/zenodo.17509367
ENK: Minimal-Disclosure Monitoring Interface	3 Feb 2026	Zenodo 10.5281/zenodo.18475921
Autocorrelation Blind-Spot	15 Apr 2026	arXiv:2604.14414

These precursor works are consolidated here into the class SRDS; an extended genealogy is documented in the companion as a public dating anchor (`genealogy_of_frames.md`).

9.5 Disclosure

Code availability. Lean-4 proofs and metric code are public (`github.com/FerdinandSchessler/srds-enk-companion`); the master-thesis timber raw data are archived on Zenodo (10.5281/zenodo.18340365).

Data availability. The ENK gold corpus is available on request under a data-use agreement (participant privacy compliance); cross-substrate raw data and reproduction scripts are documented in the companion (Table 21).

Responsible disclosure. The structural architecture of the state machine (four states, two configurations, transition rules), the axioms A0–A4, the theorems T1–T3, and the no-go theorem are fully published. Only the operational calibration thresholds remain withheld: for a deployed stability detector, publishing the exact trigger values would enable targeted evasion. No reported finding depends on them: the structural load-bearing capacity is threshold-independent; the withheld values are bound by a SHA-256 commitment in the companion (`COMMITMENT.md`), making their integrity and dating verifiable without disclosing them. The regime-bound design is not merely a disclosure choice but theoretically forced: a second no-go theorem in the formal companion framework (Schessler, 2026b) shows that under interventional monitoring (when the published verdict feeds back into the subsequent dynamics) no universal static classifier exists; the thresholds are calibration parameters of a regime, not universal constants, and only the state-machine architecture is part of the theory.

Use of generative AI. Generative AI (Claude Opus) was used both in the analysis (gold-annotation reference protocol, §3.3) and in drafting this manuscript; all content, numbers, references, and conclusions were manually checked by the author, verified against the raw data, and are fully the author’s responsibility. AI is not listed as an author.

Conflict of interest. The author declares no conflict of interest.

9.6 Why Physics Does Not Stop at Self-Referential Systems

A common reflex treats language as if mechanics made an exception for it: turns are sampled, permuted, and counted as independent observations. That is the category error. A conversation under load is *one* specimen with an ordered, continuous load history, not a bag of interchangeable tokens. Cutting a loaf into thirty slices does not yield thirty loaves but thirty cuts through *one* piece; the lag-1 autocorrelation ($\bar{\rho}_1 = 0.928$, §3.3) is the loaf as a continuum, not a nuisance term to be removed. That this treatment is physically correct is no special pleading of this work: geophysics has done the same for half a century.

Hasselmann (1976) reads climate as *integrated* weather: there the temporal autocorrelation (red noise) is the signature of accumulation, a measured value, not an artifact. The same structure carries the ENK accumulator (Palmgren-Miner damage, §5; Schessl, 2026a). What does *not* stop at self-referential systems is continuum mechanics: load, accumulation, yielding, fracture, RG-folded from the atomic Mode-I level up to society (§2.3, §9.1). What stops is only the *independence assumption* of inferential statistics, and for the loaf it never held. The substrate is language; the laws are the same.

9.7 Closing Remark

Simon et al. (2026) has concurrently proposed a programmatically related discipline under the name *learning mechanics*; its construct-boundary concept (*humble*: solid in what is described, explicit in what cannot be described) meets the one used here (§8.6). The frame is not ad hoc but structurally convergent; independent groups arrive at the same form on disjoint carriers: biological renormalization (Olmeda et al., 2026, Nature Physics), *learning mechanics* (Simon et al., 2026), three mathematical realizations of the same functor (AdS-GNN, Topological Deep Learning, mesh-agnostic composition), one cs.CL paper (Chen et al., 2026), and one computational-mechanics paper (Bakeer, 2026); §7.2. The work presented here operationalizes the class on language and tests it on ten substrates across eight research fields.

The no-go theorem (§8.6) formally bounds the universality limit and makes it testable. What must redistribute, deforms; what can no longer redistribute, breaks. What remains is the material.

References

- Miguel Aguilera, Pablo A. Morales, Fernando E. Rosas, and Hideaki Shimazaki. Explosive neural networks via higher-order interactions in curved statistical manifolds. *Nature Communications*, 16:6511, 2025. doi: 10.1038/s41467-025-61475-w. URL <https://www.nature.com/articles/s41467-025-61475-w>. arXiv preprint 2408.02326.
- Andrej Albert, Klaus-Jürgen Schneider, and Alfons Goris. *Bautabellen für Ingenieure*. Bundesanzeiger Verlag, 23rd edition, 2018. ISBN 978-3-8462-0880-9.
- Marie Alopaeus, Matthias Stosiek, Daryna Diment, Joakim Löfgren, MiJung Cho, Jarl Hemming, Teija Tirri, Andrey Pranovich, Patrik C. Eklund, Davide Rigo, Mikhail Balakshin, Chunlin Xu, and Patrick Rinke. SP-LCC – a dataset on the structure and properties of lignin-carbohydrate complexes from hardwood. *Scientific Data*, 12, 2025. doi: 10.1038/s41597-025-05327-8.
- Christian B. Anfinsen. Principles that govern the folding of protein chains. *Science*, 181(4096):223–230, 1973. doi: 10.1126/science.181.4096.223.
- W. Ross Ashby. *An Introduction to Cybernetics*. Chapman & Hall, 1956.
- Melvin Avrami. Kinetics of phase change. I. General theory. *The Journal of Chemical Physics*, 7(12): 1103–1112, 1939. doi: 10.1063/1.1750380. Vol. I (1939) and II/III follow 1940/41.
- Tammam Bakeer. Local information operators for spatial identifiability in distributed-parameter inverse problems in computational mechanics. *arXiv preprint arXiv:2605.28601*, 2026.
- Paul B. Baltes and Jacqui Smith. The fascination of wisdom: Its nature, ontogeny, and function. *Perspectives on Psychological Science*, 3(1):56–64, 2008. doi: 10.1111/j.1745-6916.2008.00062.x.
- Paul B. Baltes and Ursula M. Staudinger. Wisdom: A metaheuristic (pragmatic) to orchestrate mind and virtue toward excellence. *American Psychologist*, 55(1):122–136, 2000. doi: 10.1037/0003-066X.55.1.122.
- Jeremy M. Berg, John L. Tymoczko, Gregory J. Gatto, and Lubert Stryer. *Biochemistry*. W. H. Freeman, 9th edition, 2019.
- Garrett Birkhoff. Extensions of Jentzsch’s theorem. *Transactions of the American Mathematical Society*, 85:219–227, 1957. doi: 10.2307/1992971.
- BothBosu. Multi-agent scam conversation dataset. Hugging Face dataset, 2024. URL <https://huggingface.co/datasets/BothBosu/multi-agent-scam-conversation>.
- Percy W. Bridgman. The stress distribution at the neck of a tension specimen. *Transactions of the American Society for Metals*, 32:553–574, 1944.
- Johan August Brinell. Way of determining the hardness of bodies and some applications of the same. *Teknisk Tidskrift*, 30:69–87, 1900.
- Markus J. Buehler. Self-organizing graph reasoning evolves into a critical state for continuous discovery through structural-semantic dynamics, 2025. URL <https://arxiv.org/abs/2503.18852>. Chaos 35:113117.
- A Colin Cameron and Douglas L Miller. A practitioner’s guide to cluster-robust inference. *Journal of Human Resources*, 50(2):317–372, 2015. doi: 10.3368/jhr.50.2.317.

- Hongye Cao, Sijia Jing, Yanming Wang, Ziyue Peng, Zhixin Bai, Zhe Cao, Meng Fang, Fan Feng, Boyan Wang, Jiaheng Liu, Tianpei Yang, Jing Huo, Yang Gao, Fanyu Meng, Xi Yang, Chao Deng, and Junlan Feng. SafeDialBench: A fine-grained safety evaluation benchmark for large language models in multi-turn dialogues with diverse jailbreak attacks. *arXiv preprint arXiv:2502.11090*, 2025. URL <https://arxiv.org/abs/2502.11090>.
- Raymond B. Cattell. Theory of fluid and crystallized intelligence: A critical experiment. *Journal of Educational Psychology*, 54(1):1–22, 1963. doi: 10.1037/h0046743.
- Qiguang Chen, Yantao Du, Ziniu Li, Jinhao Liu, Songyao Duan, Jiarui Guo, Minghao Liu, Jiaheng Liu, Tong Yang, Ge Zhang, Libo Qin, Wanxiang Che, and Wenhao Huang. The molecular structure of thought: Mapping the topology of long chain-of-thought reasoning. *arXiv preprint arXiv:2601.06002*, 2026. URL <https://arxiv.org/abs/2601.06002>.
- Shuonan Chen, Yongtao Bai, Xuhong Zhou, et al. A deep learning dataset for metal multiaxial fatigue life prediction. *Scientific Data*, 11:1027, 2024. doi: 10.1038/s41597-024-03862-4.
- Lénaïc Chizat, Alex Delalande, and Tomas Vaškevičius. Sharper exponential convergence rates for Sinkhorn’s algorithm in continuous settings. *arXiv preprint arXiv:2407.01202*, 2024.
- Doohyeong Cho and Hawoong Jeong. Exact identity linking entropy production and mutual information. *arXiv preprint arXiv:2512.24877*, 2025.
- Armand Considère. Mémoire sur l’emploi du fer et de l’acier dans les constructions. *Annales des Ponts et Chaussées*, 9:574–775, 1885.
- Michael Coppedge, John Gerring, Carl Henrik Knutsen, Staffan I. Lindberg, Jan Teorell, et al. V-Dem codebook v14, 2024. URL <https://www.v-dem.net/data/the-v-dem-dataset/>.
- Lee J. Cronbach and Paul E. Meehl. Construct validity in psychological tests. *Psychological Bulletin*, 52(4):281–302, 1955. doi: 10.1037/h0040957.
- Michael Scott Cuthbert and Christopher Ariza. music21: A toolkit for computer-aided musicology and symbolic music data. In *Proceedings of the 11th International Society for Music Information Retrieval Conference (ISMIR)*, pages 637–642, 2010.
- Swetamber Das and Jason R. Green. Spectral bounds on the entropy flow rate and Lyapunov exponents in differentiable dynamical systems. *arXiv preprint arXiv:2501.04485*, 2025.
- Nicolaas Govert de Bruijn. AUTOMATH, a language for mathematics. T.H.-Report 68-WSK-05, Technische Hogeschool Eindhoven, 1968.
- Francesco Denti, Diego Doimo, Alessandro Laio, and Antonietta Mira. The generalized ratios intrinsic dimension estimator. *Scientific Reports*, 12:20005, 2022. doi: 10.1038/s41598-022-20991-1.
- Norman K. Denzin. *The Research Act: A Theoretical Introduction to Sociological Methods*. McGraw-Hill, 2nd edition, 1978.
- Niccolò Di Marco, Edoardo Loru, Alessandro Galeazzi, Matteo Cinelli, and Walter Quattrocioni. Decoding the evolution of melodic and harmonic structure of Western music through the lens of network science. *Scientific Reports*, 16:11121, 2026. doi: 10.1038/s41598-026-42872-7. arXiv:2501.07557.
- Stephan Eckstein. Hilbert’s projective metric for functions of bounded growth and exponential convergence of Sinkhorn’s algorithm. *arXiv preprint arXiv:2311.04041*, 2023.
- Herbert Edelsbrunner and John L. Harer. *Computational Topology: An Introduction*. American Mathematical Society, 2010.

- Jeff Ens and Philippe Pasquier. The MetaMIDI dataset, 2021.
- Leonhard Euler. *Methodus inveniendi lineas curvas maximi minimive proprietate gaudentes, sive solutio problematis isoperimetrici latissimo sensu accepti*. Marcum-Michaellem Bousquet & Socios, Lausanne and Geneva, 1744. Additamentum I: De curvis elasticis (column buckling).
- Charles R. Farrar and Keith Worden. An introduction to structural health monitoring. *Philosophical Transactions of the Royal Society A*, 365(1851):303–315, 2007. doi: 10.1098/rsta.2006.1928.
- Alvan R. Feinstein and Domenic V. Cicchetti. High agreement but low kappa: I. the problems of two paradoxes. *Journal of Clinical Epidemiology*, 43(6):543–549, 1990. doi: 10.1016/0895-4356(90)90158-L.
- Uwe Flick. Triangulation revisited: Strategy of validation or alternative? *Journal for the Theory of Social Behaviour*, 22(2):175–197, 1992. doi: 10.1111/j.1468-5914.1992.tb00215.x.
- Yuri Gardinazzi, Karthik Viswanathan, Giada Panerai, Alessio Ansuini, Alberto Cazzaniga, and Matteo Biagetti. Persistent topological features in large language models. In *Proceedings of the 42nd International Conference on Machine Learning (ICML)*, 2025. URL <https://arxiv.org/abs/2410.11042>. arXiv:2410.11042.
- Kurt Gödel. Über formal unentscheidbare sätze der Principia Mathematica und verwandter systeme I. *Monatshefte für Mathematik und Physik*, 38:173–198, 1931. doi: 10.1007/BF01700692.
- Jürgen Habermas. *Theorie des kommunikativen Handelns, Band 1: Handlungsrationalität und gesellschaftliche Rationalisierung*. Suhrkamp, 1981.
- Ian Hacking. *Representing and Intervening: Introductory Topics in the Philosophy of Natural Science*. Cambridge University Press, 1983. doi: 10.1017/CBO9780511814563.
- E. O. Hall. The deformation and ageing of mild steel: III. Discussion of results. *Proceedings of the Physical Society. Section B*, 64(9):747–753, 1951. doi: 10.1088/0370-1301/64/9/303.
- Laura Hanu and Unitary team. Detoxify. <https://github.com/unitaryai/detoxify>, 2020.
- Kevin G. Hare, Zuzana Masáková, and Tomáš Vávra. On the Spectra of Pisot-Cyclotomic Numbers. *Letters in Mathematical Physics*, 108(7):1729–1756, 2018. doi: 10.1007/s11005-018-1053-4. URL <https://arxiv.org/abs/1612.09285>.
- John Harrison, Josef Urban, and Freek Wiedijk. History of Interactive Theorem Proving. In Jörg Siekmann, editor, *Computational Logic, Vol. 9 of Handbook of the History of Logic*, pages 135–214. Elsevier, 2014. doi: 10.1016/B978-0-444-51624-4.50004-6.
- Klaus Hasselmann. Stochastic climate models part I. theory. *Tellus*, 28(6):473–485, 1976. doi: 10.1111/j.2153-3490.1976.tb00696.x.
- Philipp Haueis. A generalized patchwork approach to scientific concepts. *The British Journal for the Philosophy of Science*, 75(3):741–768, 2024. doi: 10.1086/716179.
- Russell C. Hibbeler. *Engineering Mechanics: Statics*. Pearson, 13th edition, 2012.
- Austin Bradford Hill. The environment and disease: Association or causation? *Proceedings of the Royal Society of Medicine*, 58(5):295–300, 1965. doi: 10.1177/003591576505800503.
- Katherine A. Hoadley, Christina Yau, Toshinori Hinoue, et al. Cell-of-origin patterns dominate the molecular classification of 10,000 tumors from 33 types of cancer. *Cell*, 173(2):291–304, 2018. doi: 10.1016/j.cell.2018.03.022.

- Hagen Holthusen, Moritz Flaschel, Denisa Martonová, and Ellen Kuhl. Benchmarking data-driven material models on the classic treloar dataset. 2026. doi: 10.48550/arXiv.2608.14063. arXiv-Preprint; Code und Daten: Zenodo, doi:10.5281/zenodo.21915635.
- Eberhard Hopf. An inequality for positive linear integral operators. *Journal of Mathematics and Mechanics*, 12(5):683–692, 1963.
- John L. Horn and Raymond B. Cattell. Age differences in fluid and crystallized intelligence. *Acta Psychologica*, 26(2):107–129, 1967. doi: 10.1016/0001-6918(67)90011-X.
- J. W. Hutchinson. Singular behaviour at the end of a tensile crack in a hardening material. *Journal of the Mechanics and Physics of Solids*, 16(1):13–31, 1968. doi: 10.1016/0022-5096(68)90014-8.
- Hakan Inan, Kartikeya Upasani, Jianfeng Chi, Rashi Rungta, Krithika Iyer, Yuning Mao, Michael Tontchev, Qing Hu, Brian Fuller, Davide Testuggine, Madian Khabza, et al. Llama guard: LLM-based input-output safeguard for human-AI conversations. *arXiv preprint arXiv:2312.06674*, 2023. URL <https://arxiv.org/abs/2312.06674>.
- Anna Jarzab, Nils Kurzawa, Thomas Hopf, et al. Meltome atlas—thermal proteome stability across the tree of life. *Nature Methods*, 17(5):495–503, 2020. doi: 10.1038/s41592-020-0801-4.
- L. M. Kachanov. Time of the rupture process under creep conditions. *Izvestiya Akademii Nauk SSSR, Otdelenie Tekhnicheskikh Nauk*, 8:26–31, 1958.
- Hyunwoo Kim, Youngjae Yu, Liwei Jiang, Ximing Lu, Daniel Khashabi, Gunhee Kim, Yejin Choi, Maarten Sap, et al. ProsocialDialog: A prosocial backbone for conversational agents. In *Proceedings of EMNLP 2022*, 2022. doi: 10.18653/v1/2022.emnlp-main.267.
- Andrei N. Kolmogorov. On the statistical theory of the crystallization of metals. *Bulletin of the Academy of Sciences of the USSR, Mathematics Series*, 1:355–359, 1937.
- Wilfried B. Krätzig, Reinhard Harte, Konstantin Meskouris, and Udo Wittek. *Tragwerke 1: Theorie und Berechnungsmethoden statisch bestimmter Stabtragwerke*. Springer, 5th edition, 2014. ISBN 978-3-662-43626-4.
- Jean Lemaitre. A continuous damage mechanics model for ductile fracture. *Journal of Engineering Materials and Technology*, 107(1):83–89, 1985. doi: 10.1115/1.3225775.
- Elizaveta Levina and Peter J. Bickel. Maximum likelihood estimation of intrinsic dimension. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 17, 2004.
- Wei Li, Luyao Zhu, Yang Song, Ruixi Lin, Rui Mao, and Yang You. Can a large language model be a gaslighter? In *The Thirteenth International Conference on Learning Representations (ICLR)*, 2025. URL <https://arxiv.org/abs/2410.09181>.
- Niklas Luhmann. *Soziale Systeme: Grundriß einer allgemeinen Theorie*. Suhrkamp, 1984.
- Yuhui Lyu, Madura Pathirage, Hoang T. Nguyen, Zdeněk P. Bažant, and Gianluca Cusatis. Dissipation mechanisms of crack-parallel stress effects on fracture process zone in concrete. *Journal of the Mechanics and Physics of Solids*, 181:105439, 2023. doi: 10.1016/j.jmps.2023.105439.
- Saunders Mac Lane. *Categories for the Working Mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer, 1971. doi: 10.1007/978-1-4612-9839-7.
- Seraphine F. Maerz, Amanda B. Edgell, Matthew C. Wilson, Sebastian Hellmeier, and Staffan I. Lindberg. Episodes of regime transformation. *Journal of Peace Research*, 61(6):967–984, 2024. doi: 10.1177/00223433231168192.

- James Mahoney. Path dependence in historical sociology. *Theory and Society*, 29(4):507–548, 2000. doi: 10.1023/A:1007113830879.
- The Mathlib Community. The Lean mathematical library (Mathlib). *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020)*, pages 367–381, 2020. doi: 10.1145/3372885.3373824.
- Cecilia Mattsson, Merima Hasani, Binh Dang, Maxim Mayzel, and Hans Theliander. About structural changes of lignin during kraft cooking and the kinetics of delignification. *Holzforschung*, 71(7–8): 545–553, 2017. doi: 10.1515/hf-2016-0190.
- Humberto R. Maturana and Francisco J. Varela. *Autopoiesis and Cognition: The Realization of the Living*, volume 42 of *Boston Studies in the Philosophy of Science*. D. Reidel, 1980. doi: 10.1007/978-94-009-8947-4.
- Kevin S. McGrew. CHC theory and the human cognitive abilities project: Standing on the shoulders of the giants of psychometric intelligence research. *Intelligence*, 37(1):1–10, 2009. doi: 10.1016/j.intell.2008.08.004.
- Tomas Mikolov, Kai Chen, Greg Corrado, and Jeffrey Dean. Efficient estimation of word representations in vector space. In *1st International Conference on Learning Representations (ICLR) Workshop*, 2013. URL <https://arxiv.org/abs/1301.3781>.
- Otto Mohr. Welche umstände bedingen die elastizitätsgrenze und den bruch eines materials? *Zeitschrift des Vereins Deutscher Ingenieure*, 44(45):1524–1530, 1900.
- P. Montero de Híjes, L. Neubeck, G. Kresse, and C. Dellago. Dynamical properties of ab initio water from machine-learning potentials. *The Journal of Chemical Physics*, 165(7):074502, 2026. doi: 10.1063/5.0348453.
- Joichiro Nakakura, Primož Ziherl, Junichi Matsuzawa, and Tomonari Dotera. Metallic-mean quasicrystals as aperiodic approximants of periodic crystals. *Nature Communications*, 10:4235, 2019. URL <https://doi.org/10.1038/s41467-019-12147-z>.
- M. E. J. Newman and M. Girvan. Finding and evaluating community structure in networks. *Physical Review E*, 69(2):026113, 2004. doi: 10.1103/PhysRevE.69.026113.
- Hoang T. Nguyen, Madura Pathirage, Masoud Rezaei, Mohammed Issa, Gianluca Cusatis, and Zdeněk P. Bažant. New perspective of fracture mechanics inspired by gap test with crack-parallel compression. *Proceedings of the National Academy of Sciences*, 117(25):14015–14020, 2020. doi: 10.1073/pnas.2005646117.
- Douglass C. North. *Institutions, Institutional Change and Economic Performance*. Cambridge University Press, 1990. doi: 10.1017/CBO9780511808678.
- Fabrizio Olmeda, Tim Lohoff, Ioannis Kafetzopoulos, Stephen J. Clark, Laura Benson, Fatima Santos, Felix Krueger, Simon Walker, Wolf Reik, and Steffen Rulands. Scaling and self-similarity in the formation of the embryonic epigenome. *Nature Physics*, 2026. doi: 10.1038/s41567-026-03263-x. URL <https://www.nature.com/articles/s41567-026-03263-x>.
- Vardan Papyan, X. Y. Han, and David L. Donoho. Prevalence of neural collapse during the terminal phase of deep learning training. *Proceedings of the National Academy of Sciences*, 117(40):24652–24663, 2020. doi: 10.1073/pnas.2015509117.
- Pinhas L. Pasternak. On a new method of analysis of an elastic foundation by means of two foundation constants. *Gosudarstvennoe Izdatel'stvo Literatury po Stroitel'stvu i Arkhitekture*, Moscow, 1954.

- Stefan Pautze. Cyclotomic Aperiodic Substitution Tilings. *Symmetry*, 9(2):19, 2017. URL <https://doi.org/10.3390/sym9020019>.
- N. J. Petch. The cleavage strength of polycrystals. *Journal of the Iron and Steel Institute*, 174:25–28, 1953.
- Paul Pierson. Increasing returns, path dependence, and the study of politics. *American Political Science Review*, 94(2):251–267, 2000. doi: 10.2307/2586011.
- Karl R. Popper. *The Logic of Scientific Discovery*. Hutchinson, 1959. Original: *Logik der Forschung*, Springer 1934.
- Ilya Prigogine. *From Being to Becoming: Time and Complexity in the Physical Sciences*. W. H. Freeman, 1980.
- Colin Raffel. *Learning-Based Methods for Comparing Sequences, with Applications to Audio-to-MIDI Alignment and Matching*. PhD thesis, Columbia University, 2016.
- Nils Reimers and Iryna Gurevych. Sentence-BERT: Sentence embeddings using Siamese BERT-networks. In *Proceedings of the 2019 Conference on Empirical Methods in Natural Language Processing*, pages 3982–3992, 2019. doi: 10.18653/v1/D19-1410.
- J. R. Rice. A path independent integral and the approximate analysis of strain concentration by notches and cracks. *Journal of Applied Mechanics*, 35(2):379–386, 1968. doi: 10.1115/1.3601206.
- J. R. Rice and G. F. Rosengren. Plane strain deformation near a crack tip in a power-law hardening material. *Journal of the Mechanics and Physics of Solids*, 16(1):1–12, 1968. doi: 10.1016/0022-5096(68)90013-6.
- Wiktor Rorot. *Scale-Free Communication? An Investigation of the Use of the Concept “Communication” in Biology and Cognitive Sciences*. PhD thesis, University of Warsaw, 2025.
- Timothy A. Salthouse. What and when of cognitive aging. *Current Directions in Psychological Science*, 13(4):140–144, 2004. doi: 10.1111/j.0963-7214.2004.00293.x.
- Marten Scheffer, Jordi Bascompte, William A. Brock, Victor Brovkin, Stephen R. Carpenter, Vasilis Dakos, Hermann Held, Egbert H. van Nes, Max Rietkerk, and George Sugihara. Early-warning signals for critical transitions. *Nature*, 461(7260):53–59, 2009. doi: 10.1038/nature08227.
- Ferdinand Maria Schessl. Empirische modellierung der verformung von holzproben unter zugbeanspruchung: Entwicklung einer sensorlosen abschätzungsmethode im Kontext thermomechanischer Untersuchungen. Master’s thesis, Technische Universität München, 2025. URL <https://zenodo.org/records/18340365>. Master’s thesis, project “Mechanical Properties of Wood for Fire Design”; Zenodo v2 22 January 2026.
- Ferdinand Maria Schessl. The autocorrelation blind spot: Why 42% of turn-level findings in LLM conversation analysis may be spurious. *arXiv preprint arXiv:2604.14414*, 2026a. URL <https://arxiv.org/abs/2604.14414>.
- Ferdinand Maria Schessl. The SRDS formal framework: A self-contained mathematical theory of critical transitions in self-referential dissipative systems, 2026b. URL <https://github.com/FerdinandSchessl/srds-enk-companion>. Companion document, available in the SRDS companion repository.
- Ferdinand Maria Schessl. Sycophancy as material failure under pushback loading: A multi-axis characterization across three loading cases and up to seventeen material charges. *arXiv preprint arXiv:2606.16617*, 2026c. URL <https://arxiv.org/abs/2606.16617>.

- Philipp Schlee, Dmitry Tarasov, Davide Rigo, and Mikhail Balakshin. Advanced NMR characterization of Aquasolv Omni (AqSO) biorefinery lignins/lignin–carbohydrate complexes. *ChemSusChem*, 16 (18):e202300549, 2023. doi: 10.1002/cssc.202300549.
- Johannes S. Schmidt, Martin Carrasco, Ernst Röell, Guy Wolf, Nello Blaser, and Bastian Rieck. No triangulation without representation: Generalization in topological deep learning. *arXiv preprint arXiv:2605.06467*, 2026. URL <https://arxiv.org/abs/2605.06467>.
- Hendrik Schreiber. Improving genre annotations for the million song dataset. In *Proceedings of the 16th International Society for Music Information Retrieval Conference (ISMIR)*, pages 241–247, 2015.
- Udo Seifert. Stochastic thermodynamics, fluctuation theorems and molecular machines. *Reports on Progress in Physics*, 75(12):126001, 2012. doi: 10.1088/0034-4885/75/12/126001.
- Eugene Seneta. *Non-negative Matrices and Markov Chains*. Springer Series in Statistics. Springer, 2nd, rev. edition, 2006. doi: 10.1007/0-387-32792-4. Cited as 2nd rev. ed. 2006; DOI resolves to Springer record dated 1981 for the original edition.
- Disha Sheshanarayana, Tanishka Magar, Ayushi Mittal, and Neelam Chaplot. CLAIM: An intent-driven multi-agent framework for analyzing manipulation in courtroom dialogues. In *Proceedings of the Third Workshop on Social Influence in Conversations (SICoN)*, pages 97–108, 2025. URL <https://aclanthology.org/2025.sicon-1.8/>.
- Jamie Simon, Daniel Kunin, Alexander Atanasov, Enric Boix-Adserà, Blake Bordelon, Jeremy Cohen, Nikhil Ghosh, et al. There will be a scientific theory of deep learning. *arXiv preprint arXiv:2604.21691*, 2026. URL <https://arxiv.org/abs/2604.21691>. Position paper proposing “learning mechanics” as a framework name.
- Robert L. Smith and George E. Sandland. An accurate method of determining the hardness of metals, with particular reference to those of a high degree of hardness. *Proceedings of the Institution of Mechanical Engineers*, 102(1):623–641, 1922.
- Hoon Sohn, Charles R. Farrar, François M. Hemez, Devin D. Shunk, Daniel W. Stinemates, Brett R. Nadler, Jerry J. Czarnecki, et al. A review of structural health monitoring literature: 1996–2001. Technical Report LA-13976-MS, Los Alamos National Laboratory, 2003.
- David I. Spivak. *Category Theory for the Sciences*. MIT Press, 2014. ISBN 978-0-262-02813-4.
- Jason Steck, Christine Heera Ahn, and Zhigang Suo. Polymers resist fatigue crack growth by deconcentrating stress. *Annual Review of Materials Research*, 55(1):333–358, 2025. doi: 10.1146/annurev-matsci-101922-122133.
- Ewout W. Steyerberg. *Clinical Prediction Models: A Practical Approach to Development, Validation, and Updating*. Springer, 2009. doi: 10.1007/978-0-387-77244-8.
- Chenhao Tan, Vlad Niculae, Cristian Danescu-Niculescu-Mizil, and Lillian Lee. Winning arguments: Interaction dynamics and persuasion strategies in good-faith online discussions. In *Proceedings of the 25th International Conference on World Wide Web (WWW)*, 2016. URL <https://doi.org/10.1145/2872427.2883081>.
- Ruifeng Tan, Weixiang Hong, Jiayue Tang, Xibin Lu, Ruijun Ma, Xiang Zheng, Jia Li, Jiaqiang Huang, and Tong-Yi Zhang. BatteryLife: A comprehensive dataset and benchmark for battery life prediction, 2025. URL <https://arxiv.org/abs/2502.18807>. KDD 2025.
- Henri Tresca. Mémoire sur l’écoulement des corps solides soumis à de fortes pressions. *Comptes Rendus Acad. Sci. Paris*, 59:754–758, 1864.

- Karthik Viswanathan, Yuri Gardinazzi, Giada Panerai, Alberto Cazzaniga, and Matteo Biagetti. The geometry of tokens in internal representations of large language models. *arXiv preprint*, 2025. URL <https://arxiv.org/abs/2501.10573>. arXiv:2501.10573.
- Xuewei Wang, Weiyan Shi, Richard Kim, Yoojung Oh, Sijia Yang, Jingwen Zhang, and Zhou Yu. Persuasion for good: Towards a personalized persuasive dialogue system for social good. In *Proceedings of the 57th Annual Meeting of the Association for Computational Linguistics (ACL)*, 2019. URL <https://aclanthology.org/P19-1566/>.
- Yuxin Wang, Ivory Yang, Saeed Hassanpour, and Soroush Vosoughi. MentalManip: A dataset for fine-grained analysis of mental manipulation in conversations. In *Proceedings of the 62nd Annual Meeting of the Association for Computational Linguistics (ACL)*, 2024.
- Norbert Wiener. *Cybernetics: Or Control and Communication in the Animal and the Machine*. MIT Press, 1948.
- M. L. Williams. On the stress distribution at the base of a stationary crack. *Journal of Applied Mechanics*, 24(1):109–114, 1957. doi: 10.1115/1.4011454.
- Kenneth G. Wilson. Problems in physics with many scales of length. *Scientific American*, 241(2): 158–179, 1979. doi: 10.1038/scientificamerican0879-158.
- Mark Wilson. *Wandering Significance: An Essay on Conceptual Behavior*. Oxford University Press, Oxford, 2006.
- Emil Winkler. *Die Lehre von der Elasticität und Festigkeit mit besonderer Rücksicht auf ihre Anwendung in der Technik*. H. Dominicus, Prag, 1867.
- August Wöhler. Über die Festigkeitsversuche mit Eisen und Stahl. *Zeitschrift für Bauwesen*, 20:73–106, 1870.
- Jeffrey M. Wooldridge. Cluster-sample methods in applied econometrics. *American Economic Review*, 93(2):133–138, 2003. doi: 10.1257/000282803321946930.
- Zixiu Wu, Simone Balloccu, Vivek Kumar, Rim Helaoui, Ehud Reiter, Diego Reforgiato Recupero, and Daniele Riboni. Anno-MI: A dataset of expert-annotated counselling dialogues. In *ICASSP 2022 – IEEE International Conference on Acoustics, Speech and Signal Processing*, 2022.
- Sunwoong Yang, Ricardo Vinuesa, and Namwoo Kang. Mesh-agnostic prediction of unsteady flow dynamics using graph U-Nets. *Expert Systems with Applications*, 2026. URL <https://www.sciencedirect.com/science/article/abs/pii/S0957417426007566>. Published March 2026; arXiv preprint 2406.03789 (extended preprint version: “Enhancing Graph U-Nets for Mesh-Agnostic Spatio-Temporal Flow Prediction”).
- Erxin Yu, Jing Li, Ming Liao, Siqi Wang, Zuchen Gao, Fei Mi, and Lanqing Hong. CoSafe: Evaluating large language model safety in multi-turn dialogue coreference. In *Proceedings of the 2024 Conference on Empirical Methods in Natural Language Processing (EMNLP)*, pages 17190–17205, 2024. URL <https://aclanthology.org/2024.emnlp-main.968/>.
- Justine Zhang, Jonathan P. Chang, Cristian Danescu-Niculescu-Mizil, Lucas Dixon, Yiqing Hua, Nithum Thain, and Dario Taraborelli. Conversations gone awry: Detecting early signs of conversational failure. In *Proceedings of the 56th Annual Meeting of the Association for Computational Linguistics (ACL)*, 2018. URL <https://aclanthology.org/P18-1125/>.
- Maksim Zhdanov, Nabil Iqbal, Erik Bekkers, and Patrick Forré. AdS-GNN: A conformally equivariant graph neural network. *arXiv preprint arXiv:2505.12880*, 2025. URL <https://arxiv.org/abs/2505.12880>. ICLR 2025 Workshop MLMP.

Xuhui Zhou, Hao Zhu, Leena Mathur, Ruohong Zhang, Haofei Yu, Zhengyang Qi, Louis-Philippe Morency, Yonatan Bisk, Daniel Fried, Graham Neubig, Maarten Sap, et al. Sotopia: Interactive evaluation for social intelligence in language agents. In *Proceedings of ICLR 2024*, 2023. URL <https://arxiv.org/abs/2310.11667>. arXiv:2310.11667.